

1 **Article**

2

3 **Determination of Some Variables Affecting Risk Factors of Coronary Heart Diseases in**  
4 **University Students**

5

6 Ali OZKAN<sup>1,2</sup>, Mutlu TURKMEN<sup>1,3</sup>, Taner BOZKUS<sup>1</sup>, Murat KUL<sup>1</sup>, Recep SOSLU<sup>4</sup>, Fatih  
7 YASARTURK<sup>1</sup>, Recep AYDIN<sup>1</sup>, Umit OZ<sup>1</sup>

8

9 <sup>1</sup>Bartın University, School of Physical Education and Sports, Bartın, Turkey; ali\_ozkan1@hotmail.com;  
10 mutluturkmen@bartin.edu.tr; tanerbozkus@yahoo.com; muratkul61@gmail.com; fatihyasarturk@gmail.com;  
11 g.recep.aydin@gmail.com; ozumut06@hotmail.com12 <sup>2</sup>Bartın University, Physical Education Research and Application Center, Bartın, Turkey,  
13 ali\_ozkan1@hotmail.com14 <sup>3</sup>Turkish Bocce Bowling and Darts Federation, Ankara, Turkey, mutluturkmen@bartin.edu.tr15 <sup>4</sup>Karamanoğlu Mehmetbey University, School of Physical Education and Sports, Karaman, Turkey,  
16 recepsosli@gmail.com

17

18

19 **Abstract:** The purpose of the present study was to determine the relationship between healthy  
20 lifestyle behaviors, physical fitness and coronary risk factors in university students. 320 male  
21 and female (n<sub>m</sub>:171; n<sub>f</sub>:149) students from a university participated in this study voluntarily.  
22 For the determination of body composition and Body Mass Indexes (BMI), subjects' height,  
23 body weight, and skinfold thickness were taken and body fat percentage (%Fat) was  
24 determined. Healthy lifestyle behaviors were determined using the healthy lifestyle behaviors  
25 questionnaire. Indicators of physical fitness included flexibility (sit-up) (F), muscle strength  
26 and endurance (isometric knee (KS), back strength (BS) and a total of shuttle (TS), sprint  
27 performance, BMI, and body fat percentage (%fat). Coronary heart disease risk factors  
28 included mean arterial blood pressure (systolic (SBP) and diastolic (DBP)), fasting blood  
29 levels of triglycerides (TG), total cholesterol (TC), hematocrit (HT), and hemoglobin (HM).  
30 Results indicated subjects have normal body mass index, body fat percentage, SBP, DBP, TG,  
31 TCF, BS, KS. The results of the Pearson Product Moment Correlation Analysis, indicated that  
32 SBP, DBP, TG, TCF, BS, KS for male and female was significantly correlated with flexibility  
33 (sit-up) (F), muscle strength and endurance (isometric knee (KS), back strength (BS) and total  
34 of shuttle (TS)), sprint performance. (p<0.01; p<0.05), In conclusion, the findings of the  
35 present study indicated that physical fitness and healthy lifestyle behaviors play a determinant  
36 role in coronary heart disease risk factors for male and female students from a university.

37 **Keywords:** coronary heart disease risk factors, healthy lifestyle behaviors, physical fitness

38

39

## 40 1. Introduction

41  
42 Coronary heart disease (CHD) continues to be a leading cause of morbidity and mortality  
43 among adults in Europe and North America [1]. For example, CHD is responsible for 29.500  
44 deaths in Australia annually. Ninety-three percent of women and 80% of men who die as a  
45 result of CHD are over the age of 65 [2]. Similar statistics are reported in Canada [3] and the  
46 United States, United Kingdom [4] and Turkey [5]. Risk factors have included blood pressure,  
47 cigarette smoking, cholesterol (TC), LDL-C, HDL-C, and diabetes. 2-4 Factors such as  
48 obesity, left ventricular hypertrophy, family history of premature CHD and physically inactive  
49 [3] have also been considered in defining CHD risk [1] and increased physical activity leads  
50 to greater cardiorespiratory fitness, decreased blood pressure and body weight and increased  
51 HDL-C, all of which lead to a more favorable CHD risk profile [6]. The study of adaptive  
52 functional changes of the human body induced by physical effort to optimise the physical  
53 wellness and to combat the metabolic, circulation and respiratory diseases are in the trends of  
54 interdisciplinary scientific research from medicine and physical activity[7,8,9].

55 A recent meta-analysis showed a significant protective effect of physical activity and physical  
56 health-related fitness on CHD [10,11]. Physical inactivity is associated with an increased risk  
57 of a wide variety of diseases like cardiovascular diseases, hypertension, type 2 diabetes,  
58 obesity, and depression. It has been stated the above-mentioned diseases which are associated  
59 with physical inactivity seldom manifest themselves before adulthood, however, promotion of  
60 physical activity may be important as physical inactivity may also predispose to a future  
61 sedentary lifestyle and hence have an increased risk for these diseases. Several reviews have  
62 indicated that the associations among physical activity fitness and CHD risk factors in youth  
63 have not been conclusively delineated [3, 11, 12, 13]. A number of more recent studies have  
64 examined these associations but results remain equivocal. For example, in the cardiovascular  
65 risk in young finns study (9 to 24 years age), active males had lower TG and higher HDL-C  
66 level than inactive males, while active females had lower TG levels than inactive females  
67 controlling for pubertal status [12]. For instance, Heggebo et al [14] reported that the results  
68 in terms of the anthropometrical measures, diastolic and systolic blood pressure were the  
69 major factors explaining the cardiorespiratory fitness. Hence, the purpose of the present study  
70 was to determine the relationship between healthy lifestyle behaviors, physical fitness and  
71 coronary risk factors in university students

72

## 73 2. Materials and Methods

### 74 2.1. Subjects

75 320 male and female ( $n_m:171$ ;  $n_f:149$ ) students attending to Bartın University in Turkey  
76 participated in this study voluntarily. Their mean age, height, body weight and body fat were  
77 20.89 (2.01) yrs, 168.04 (14.7) cm, 67.2 (16.25) kg, and 20.66 (7.01) respectively. All  
78 subjects gave their informed consent for inclusion before they participated in the study. The  
79 study was conducted in accordance with the Declaration of Helsinki and this study was  
80 approved by Bartın University Institutional Review Board, Ethics Committee and supported  
81 by Bartın University Scientific Research Projects Commission (Project No: 2016-SOS-A-  
82 006).

83

### 84 2.2. Anthropometric Measurements

85 Body height (cm), body mass (kg), and percentage of body fat (PBF) measurements were  
86 taken for each subject. The body height of the university students were measured by a  
87 stadiometer with an accuracy of  $\pm 1$  cm (SECA, Germany), and while electronic scales (Tanita  
88 BC 418, Japan) accurate to within 0.1 kg were used to measure body mass and percentage of  
89 body fat [12]. Skinfold thickness was measured with a Holtain skinfold caliper (Holtain, UK)  
90 which applied a pressure of 10 g/mm<sup>2</sup> with an accuracy of  $\pm 2$  mm. Gulick anthropometric  
91 tape (Holtain, UK) with an accuracy of  $\pm 1$  mm was used to measure the circumference of  
92 extremities. Diametric measurements were determined by Harpenden calipers (Holtain, UK)  
93 with an accuracy of  $\pm 1$  mm.

94

### 95 2.3. Health-promoting life-style profile scale

96 The participants were asked to provide information about the demographic factors, such as  
97 age, gender, and education. Health-promoting Life-style Profile Scale was used for collecting  
98 data on their health behaviors. The scale was developed by Walker et al. [16]. It is composed  
99 of 48 items and 6 subscales and consists of questions about health-promoting behaviors. The  
100 subscales were on self-actualization (SA), health responsibility (HR), exercise (E), nutrition  
101 (N), interpersonal support (IS), and stress management (SM). The total score reflects the  
102 healthy life-style behavior. Four more items were added to the scale, and now the scale is  
103 composed of 52 items [16]. Each respondent was asked to rate each item on Likerts' 1 to 4  
104 response scale where 1 corresponds to never, 2 sometimes, 3 often, 4 regularly. Alpha  
105 coefficient reliability of the scale was 0.92, and alpha coefficient reliability of the subscales

106 varied from 0.70 to 0.90. The reliability of the scale for Turkish population was tested by Esin  
107 [17] and Akça [18]. Alpha coefficient reliability of the scale was 0.91 in Esin's study and 0.90  
108 in Akça's study.

109

#### 110 **2.4. Flexibility measurement**

111 Flexibility was evaluated by the sit and reach test which is the most common flexibility test  
112 used in health-related fitness test batteries. The subjects sat with their feet approximately hip-  
113 width against the testing box. They kept their knees extended and placed the right hand over  
114 the left, and slowly reached forward as far as they could by sliding their hands along the  
115 measuring board. Reaches short of the toes were recorded as negative forward reach scores,  
116 and reaches beyond the toes were recorded as positive forward reach scores in centimeter to  
117 the nearest 0.5 cm using the scale on the box [19].

118

#### 119 **2.5. Strength measurement**

120 Isometric Dynamometer was used for the determination of knee, back, grip strength. Muscular  
121 strength was assessed using a Takei strength dynamometer (Takei Scientific Instruments,  
122 Tokyo, Japan).

123

#### 124 **2.6. Coronary Heart Disease Risk Factors**

##### 125 **2.6.1. Blood Pressure**

126 Systolic and diastolic blood pressures were measured with a sphygmomanometer (Erka  
127 perfect Aneroid, Germany) following the recommendations of the American Heart  
128 Association.

129

##### 130 **2.6.2. Lipids and Lipoproteins**

131 All venous and capillary blood lipid concentrations were determined in the morning after an  
132 overnight fasting period of at least eight hours. In each subject, venous and capillary blood  
133 samples were collected at the same time. Supplementary capillary sampling was performed on  
134 two consecutive days following the first collection. Capillary TC and TG concentrations were  
135 determined with the Accutrend® Plus using two drops of blood (15-40 µL) collected from  
136 different fingers, by using a lancing device (Accu-check® Softclix® Pro, Roche Diagnostics  
137 GmbH, Mannheim, Germany). The Accutrend® Plus test is a capillary serum test. It is based

138 on the retention of blood cells by filtration via a glass fibre fleece when a drop of blood is  
139 applied to the test strip. The enzymatic reaction that takes place in the underlying zone of the  
140 strip requires an adequate oxygen supply and results in the formation of a colored oxidation  
141 product. The reflectance of the strip (measured at 660 nm) is converted to concentration  
142 through a simple algorithm. The intra assay precision of Accutrend® Plus, as determined by  
143 the manufacturer, was 3.7% and 3.4% for TC and TG respectively. The inter-assay precision  
144 for Accutrend® Plus determined with control solution by the manufacturer was lower than  
145 5.0% for TC and 2.4% for TG [20].

146

### 147 **2.6.3. Hematocrit and Hemoglobin**

148 The Mission® Plus Hemoglobin (Hb) and Hematocrit Testing System (ACON Laboratories,  
149 Inc., US) are for the quantitative determination of hemoglobin and hematocrit in non-  
150 anticoagulated capillary whole blood or anticoagulated venous whole blood in EDTA (K2,  
151 K3, Na2) or sodium heparin. The testing system is designed for point-of-care use in primary  
152 care settings. Estimation of hematocrit is only for hemoglobin values from 12.3 to 17.5 g/dL  
153 (123 to 175 g/L). The Mission® Plus hemoglobin and hematocrit Control Solution is intended  
154 to validate hemoglobin and hematocrit testing using the Mission® Plus Hemoglobin (Hb)  
155 Testing System. Mission® Plus hemoglobin and hematocrit Testing System is for  
156 professional in vitro diagnostic use only. This device has not been evaluated for pediatric  
157 subjects.

158

### 159 **2.7. Statistical Analyses**

160 Means and standard deviations are given as descriptive statistics and the relationship among  
161 the relationship between healthy lifestyle behaviors, physical fitness, and coronary risk factors  
162 were evaluated by Pearson Product Moment Correlation analysis. All analyses were executed  
163 in SPSS for Windows version 21.0 and the statistical significance was set at  $p < .05$ .

164

165

### 166 **3. Results**

167 Body composition, physical fitness characteristics, the healthy lifestyle behaviors and  
168 coronary risk factors of university students are displayed in Table 1, 2 and 3 respectively.

169

170

**Table 1:** Body Composition Values of University Students

	Age (year)	Height (cm)	Body Weight (kg)	Fat (%)	BMI (kg/m <sup>2</sup> )
<b>Male (n:171)</b>	20.37	162.18	58.15	24.31	22.08
	± 1.53	± 5.56	± 8.99	± 6.15	± 3.18
<b>Female (n:149)</b>	21.53	175.16	78.29	16.35	24.86
	± 3.00	± 18.66	± 16.30	± 5.35	± 5.56
<b>Male and Female (n: 320)</b>	20.89	168.04	67.22	20.66	23.34
	± 2.37	± 14.76	± 16.25	± 7.01	± 4.61

171  
 172 Descriptive characteristics of the subjects across body composition are shown in Table 1.  
 173 According to this table, the highest rate was reached in % fat for male and the lowest rate in  
 174 female university students. Results indicated that the subjects have normal body mass index,  
 175 but high body fat percentage.

176  
 177

**Table 2:** Physical Fitness Characteristics Values of University Students

	BS (kg)	KS (kg)	RGS (kg)	LGS (kg)	F (cm)	TS (repeat)	10m (sec)	20m (sec)	30m (sec)
<b>Female (n:149)</b>	45.19	44.70	30.27	28.18	17.91	20.58	2.57	3.83	5.25
	± 6.52	± 28.33	± 5.96	± 4.29	± 8.49	± 7.29	± 0.28	± 0.32	± 0.43
<b>Male (n:171)</b>	110.84	103.03	44.81	42.15	21.67	32.26	2.14	3.31	4.55
	± 106.9	± 35.67	± 7.85	± 6.97	± 10.29	± 9.76	± 0.29	± 0.29	± 0.43
<b>Male and Female (n: 320)</b>	79.95	75.51	39.13	36.67	20.42	26.80	2.38	3.69	5.02
	± 87.13	± 43.92	± 10.21	± 9.27	± 9.81	± 10.53	± 0.35	± 0.41	± 0.55

178  
 179  
 180 Table 2 indicates that university students had good physical fitness performance. Sprint  
 181 performance and strength is fundamental activity for many sports and also sports performance  
 182 is the outcome of several variables, including physical fitness. Differentiated physical fitness  
 183 profiles might, therefore, be considered as a parameter of sports - specific demands, and as  
 184 such contribute to an enhanced knowledge of the level of performance [21].

185  
 186  
 187  
 188  
 189

**Table 3:** Coronary Risk Factors Values of University Students

	Heart Rate (rest) (beats per min)	SBP (mm Hg)	DBP (mm Hg)	TG (mg)	TC (mg/dL)	HGB (g/dL)	HT (%)
<b>Male (n:171)</b>	84.13 ± 10.72	112.87 ± 13.11	73.45 ± 11.81	101.3 ± 63.9	173.96 ± 32.29	12.99 ± 0.77	42.17 ± 5.68
<b>Female (n:149)</b>	91.30 ± 10.85	127.53 ± 11.70	78.66 ± 8.33	70.6 ± 38.2	173.31 ± 33.20	14.73 ± 0.97	48.83 ± 7.36
<b>Male and Female (n: 320)</b>	87.97 ± 11.43	120.61 ± 14.49	76.10 ± 10.44	85.9 ± 51.05	173.62 ± 32.72	13.89 ± 1.23	46.62 ± 7.53

190

191 According to Table 3, the mean values for HR, SBP, DBP, TG, TC, HGB, and HT were in  
 192 the normal range for both male and female students. In agreement with national and  
 193 international literature, the data from the present study shows a considerable prevalence of  
 194 cardiovascular risk factors among young adults. A family history of chronic diseases was  
 195 reported by many of the university students. Several studies have revealed a greater  
 196 prevalence of cardio vascular risk factors in relatives of individuals with  
 197 cardiovascular diseases and type 2 diabetes mellitus when compared with those without  
 198 family history of these diseases.

199

200

**Table 4:** Mean and standard deviation of scores of the healthy life-style behavior among the study participants

	Male and Female (n: 320)	Female (n:149)	Male (n:171)
<b>Self-actualization</b>	38.45 ± 7.10	36.02 ± 5.99	40.51 ± 7.33
<b>Health responsibility</b>	23.89 ± 7.17	21.10 ± 6.53	26.29 ± 6.85
<b>Exercise</b>	10.56 ± 4.05	10.25 ± 4.03	10.82 ± 4.06
<b>Nutrition</b>	17.66 ± 3.66	15.50 ± 2.86	19.52 ± 3.05
<b>Interpersonal support</b>	21.11 ± 3.65	20.07 ± 3.08	22.00 ± 3.58
<b>Stress management</b>	18.63 ± 4.42	17.22 ± 3.55	19.85 ± 4.75
<b>Total score of the healthy and life-style behavior</b>	130.36 ± 24.55	119.78 ± 20.54	139.12 ± 24.22

201

202 Table 4 shows the healthy life-style behaviors of the university students. According to this  
 203 table, the highest rate was reached in self-actualization sub-scale, and the lowest rate in  
 204 exercise. This finding depicts the contradictory attitude of university students towards  
 205 exercise.

206

207 Correlations between physical fitness and coronary risk factors of university students are  
 208 presented in Table 5.

209 **Table 5:** Correlations between physical fitness and coronary risk factors of university students

	BS	KS	F	TS	RPS	LPS	10m	20m	30m
HR	,244**	,430**	,409**	,178**	,202**	,195**	-,277**	-,408**	-,426**
SBP	,171**	,363**	,130*	,265**	,403**	,421**	-,296**	-,387**	-,340**
DBP	NS	,160**	NS	,151*	,246**	,254**	-,175*	NS	NS
TG	NS	,261**	NS	NS	,310**	,263**	-,203*	-,290**	-,290**
TC	NS	NS	NS	NS	NS	NS	NS	NS	NS
HT	,155*	,219**	NS	,212**	,374**	,366**	-,287**	-,275**	-,240*
HGB	,385**	,485**	NS	,367**	,482**	,565**	-,385**	-,316**	-,396**

210 \*\*p&lt;0.01, \* p&lt;0.05

211

212 As seen in Table 5, Results of Pearson's Product Moment Correlation Analyses indicated

213 significant positive correlations between physical fitness and coronary risk factors of

214 university students in the present study ( $p<0.05$ ). The findings of this study indicated that

215 physical fitness was an indicator of coronary risk factors of university students.

216

217 Correlations between physical fitness and coronary risk factors of male and female university

218 students are presented in Table 6 and 7.

219 **Table 6:** Correlations between physical fitness and coronary risk factors of male university students

	BS	KS	F	TS	RPS	LPS	10m	20m	30m
HR	,176*	,446**	NS	NS	,495**	NS	NS	-,404**	-,345*
SBP	NS	NS	NS	NS	NS	NS	NS	NS	NS
DBP	NS	NS	NS	NS	NS	NS	NS	NS	NS
TG	NS	NS	NS	NS	NS	NS	,326**	,316*	NS
TC	NS	NS	NS	NS	NS	NS	,326**	,316*	NS
HT	NS	NS	NS	NS	NS	NS	NS	NS	NS
HGB	,221*	NS	NS	NS	NS	NS	NS	NS	NS

221 \*\*p&lt;0.01, \* p&lt;0.05

222

223 As seen in Table 6, results of Pearson's Product Moment Correlation Analyses indicated

224 significant positive correlations between physical fitness and coronary risk factors of male

225 university students ( $p<0.05$ ). The findings of the present study indicated that physical fitness

226 was an indicator of coronary risk factors of male university students.

227

228

229

230

231

232

233

234

235

236

237

238



239

Table 7: Correlations between physical fitness and coronary risk factors of male university students

	<b>BS</b>	<b>KS</b>	<b>F</b>	<b>TS</b>	<b>RPS</b>	<b>LPS</b>	<b>10m</b>	<b>20m</b>	<b>30m</b>
<b>HR</b>	NS	NS	NS	NS	NS	NS	NS	NS	NS
<b>SBP</b>	<b>,364**</b>	<b>,265**</b>	NS	<b>,219**</b>	<b>,380**</b>	<b>,401**</b>	NS	NS	NS
<b>DBP</b>	NS	NS	NS	NS	<b>,184*</b>	<b>,253**</b>	NS	NS	NS
<b>TG</b>	NS	NS	NS	NS	<b>,319**</b>	<b>,337**</b>	NS	NS	NS
<b>TC</b>	NS	NS	<b>,171*</b>	NS	NS	NS	NS	NS	NS
<b>HT</b>	NS	NS	NS	<b>,193*</b>	<b>,394**</b>	<b>,408**</b>	NS	NS	NS
<b>HGB</b>	<b>,349**</b>	<b>,328**</b>	NS	<b>,307**</b>	<b>,458**</b>	<b>,580**</b>	NS	<b>,228*</b>	NS

\*\*p&lt;0.01, \* p&lt;0.05

240

241

242

243

As seen in Table 7, results of Pearson's Product Moment Correlation Analyses indicated significant positive correlations between physical fitness and coronary risk factors of female university students in the study subjects ( $p<0.05$ ). The findings of the present study indicated that physical fitness was an indicator of coronary risk factors of female university students in the Bartin University of Turkey.

248

Correlations between the healthy life-style behaviors and coronary risk factors of male and female university students are presented in Table 8 and 9.

250

251

Table 8: Correlations between healthy life-style behaviors and coronary risk factors of male university students

	<b>SA</b>	<b>HR</b>	<b>E</b>	<b>N</b>	<b>IS</b>	<b>SM</b>	<b>HLSB</b>
<b>HR</b>	<b>.208**</b>	<b>.245**</b>	NS	<b>.355**</b>	<b>.265*</b>	<b>.278**</b>	<b>.287**</b>
<b>SBP</b>	NS	NS	<b>.134*</b>	<b>.262**</b>	<b>.123*</b>	NS	NS
<b>DBP</b>	NS	NS	NS	NS	NS	NS	NS
<b>TG</b>	NS	<b>.247**</b>	<b>.125**</b>	<b>.225**</b>	<b>.187*</b>	<b>.162*</b>	<b>.214**</b>
<b>TC</b>	NS	<b>.126**</b>	<b>.205*</b>	NS	NS	NS	<b>.153*</b>
<b>HT</b>	<b>.128*</b>	<b>.248**</b>	<b>.297*</b>	<b>.293**</b>	<b>.189*</b>	<b>.176*</b>	<b>.234*</b>
<b>HGB</b>	<b>.248**</b>	<b>.282**</b>	<b>.239**</b>	<b>.369**</b>	<b>.215*</b>	<b>.232**</b>	<b>.258*</b>

\*\*p&lt;0.01, \* p&lt;0.05

252

253

254

As seen in Table 8, results indicated significantly positive correlations between healthy life-style behaviors and coronary risk factors of male university students in the study subjects ( $p<0.05$ ). The findings of the present study indicated that healthy life-style behaviors was an indicator of coronary risk factors of male university students in the Bartin University of Turkey.

259

260

261

262

263

264 Table 9: Correlations between healthy life-style behaviors and coronary risk factors of female university students

	SA	HR	E	N	IS	SM	HLSB
HR	.210**	.203**	NS	.334**	.214*	.267**	.275**
SBP	NS	NS	.145*	.245**	.153*	NS	NS
DBP	NS	NS	NS	NS	NS	NS	NS
TG	.102*	.252**	.152**	.227**	.179*	.141*	.206**
TC	NS	NS	.148*	NS	NS	NS	NS
HT	.149*	.256**	.298*	.289**	.176*	.149*	.211**
HGB	.217*	.276**	.248**	.387**	.146*	.228**	.295**

265 \*\*p&lt;0.01, \*p&lt;0.05

266

267 As seen in Table 9, results indicated significantly positive correlations between healthy life-  
 268 style behaviors and coronary risk factors of female university students in the study subjects  
 269 (p<0.05). The findings of the present study indicated that healthy life-style behaviors were an  
 270 indicator of coronary risk factors of female university students in the Bartın University of  
 271 Turkey.

272

273

274 **4. Discussion**

275 The findings of the present study indicated that physical fitness plays a determinant role in  
 276 coronary heart disease risk factors for male and female students from a university. In addition,  
 277 PAL was found to be an important factor in coronary heart disease risk factors of university  
 278 students. Coronary heart disease risk factors (CHDRF) included mean arterial blood pressure  
 279 (systolic (SBP) and diastolic (DBP)), fasting blood levels of triglycerides (TG), low-density  
 280 lipoprotein cholesterol (LDL-C), high-density lipoprotein cholesterol (HDL-C) and total  
 281 cholesterol (TC) in this present study [1]. Further, the pattern of loadings in boys and girls  
 282 remarkably similar in this sample (Table 3) suggesting that sex differences in the relationship  
 283 between PAL and CHDRF are small if they exist. And also the findings of the present study  
 284 are consistent with previous studies that have demonstrated relationships between PAL and  
 285 CHDRF, particularly for blood lipids [3, 12]. Although physical fitness (PF) was the best  
 286 predictor of SBP, DBP, TG, LDL-C, HDL-C, TC, while PAL were not shown to be  
 287 significant predictors. The result indicate that F, KS, BS, TS, VO<sub>2max</sub>, BMI and %fat are  
 288 important determinants of CHDRF in university students, with PF exhibiting a slightly  
 289 stronger relationship than PAL. However, this result must be tempered by the limitations of  
 290 the study. Although physical record used is a reliable measure of habitual activity levels, the  
 291 error associated with the indicators of physical activity is undoubtedly greater than error

292 associated with the measurements of PF [3]. Another explanation for the greater relationship  
293 between fitness and CHDRF may be genetics. Perhaps genes which are influencing physical  
294 fitness also influence CHDRF (genetic pleiotropy) [3]. Eisenmann et al [11] reported that a  
295 significant relationship between adolescent cardiorespiratory fitness and adult body fatness  
296 and a lack of an association between adolescent cardiorespiratory fitness and adult  
297 cholesterol, blood pressure, and glucose levels. Adolescent body fatness is moderately related  
298 to selected adult CHDRF and this could be influencing the pattern of loading for PF,  
299 independent of chronological age [3].

### 300 **5. Conclusion**

301 To sum up, an important prevalence of cardiovascular risk factors was observed in the  
302 university students included in the present study. Considering that some of the cardiovascular  
303 risk factors are modifiable by changes in lifestyle, educational programs aimed at motivating  
304 the adoption of healthy lifestyle choices would be helpful, especially in upcoming health care  
305 professionals, as it is them who will be taking care of the health of the population in the future  
306 [22].

307

308 **Acknowledgments:** This study was approved by Bartin University Institutional Review  
309 Board, Ethics Committee and supported by Bartin University Scientific Research Projects  
310 Commission (Project No: 2016-SOS-A-006).

311

### 312 **Author Contributions:**

313 Ali Ozkan was the primary one shaping the main text and Mutlu Turkmen formatted the last  
314 version of the text in English. All the other authors contributed to the article equally in  
315 gathering data, scanning the literature, formatting the research, and finally revising the text.

316

### 317 **Corresponding Author:**

318 Bartin University, School of Physical Education and Sports, Bartin, Agdaci Kampusu,  
319 74100, TURKEY, E-mail: mutlurkmen@bartin.edu.tr; Tel: +905325057565

320

321 **Conflicts of Interest:** The authors declare no conflict of interest.

322

323

324

325

326 **References**

327

328 1. Wilson, P.W.F.; Agostino, R.B.D.; Levy, D.; Belander, A.M.; Silbershatz, H.; Kannel,  
329 W.B. Prediction of coronary heart Disease Using Risk Factor Categories. *Circulation*.  
330 **1998**, 97:1837-1847; DOI: <https://doi.org/10.1161/01.CIR.97.18.1837>.

331 2. Knight, S.; Bermingham, M.A.; Mahajan, D. Regular non-vigorous physical activity  
332 and cholesterol levels in the elderly. *Gerontology*. **1999**, 45: 213-219; DOI:  
333 <https://doi.org/10.1159/000022090>.

334 3. Katzmarzyk, P. T.; Malina, R. M.; Bouchard, C. Physical activity, physical fitness, and  
335 Coronary heart disease risk factors in youth: The Quebec family study. *Preventive*  
336 *Medicine*. **1999**, 29; 555-562; DOI: <https://doi.org/10.1006/pmed.1999.0592>.

337 4. Flack, J.; Yunis, C. Therapeutic implications of the epidemiology and timing of  
338 myocardial infarction and other cardiovascular diseases. *J. Hum. Hypertens.* **1997**. 11;  
339 23-28; DOI:10.1038/sj.jhh.1000396.

340 5. Onat, A. Risk factors and cardiovascular disease in Turkey. *Atherosclerosis*. **2001**,  
341 156(1):1-10. DOI: [https://doi.org/10.1016/S0021-9150\(01\)00500-7](https://doi.org/10.1016/S0021-9150(01)00500-7).

342 6. Hokanson, J.E.; Kamboh, M.I.; Scarboro, S.; Eckel, R.H.; Hamman, R.F. (2003).  
343 Effects of the hepatic lipase gene and physical activity on coronary heart disease risk.  
344 *Am. Journal of Epidemiol.* **2003**, 158(9); 836-843; DOI:  
345 <https://doi.org/10.1093/aje/kwg230>.

346 7. Dusa, F.S.; Badau, A.; Badau, D.; Trambitas, C.; Brinzaniuc, K. Investigating the  
347 Deformation Parameters of PVC Fitness Balls in Relation to the Height and Body  
348 Mass Index of the Users. *Mater. Plast.* **2017**, 54, 606 -609.

349 8. Badau, A. Study of somatic, motor and functional effects of practicing initiation  
350 programs in water gymnastics and swimming by students of physical education and  
351 sports. *Physical Education of Students*, **2017**, 21; 158–164;  
352 DOI:10.15561/20755279.2017.0402.

353 9. Badau, D.; Prebeg, G.; Mitić, D.; Badau A. Fitness index and VO<sub>2</sub>max of physical  
354 education students, *Ovidius University Annals, Series Physical Education and Sport,*  
355 *Science, Movement and Health*, 2015, 15; 246-251.

356 10. Williams, P.T. (2001). Physical fitness and activity as separate heart disease risk  
357 factors: a meta-analysis. *Med Science Sports Exerc.* **2001**, 33:754-761; ISSN: 0195-  
358 9131.

- 359 11. Eisenmann, J.C.; Wickel, E.E.; Welk, G.J.; Blair, S.N. Relationship between  
360 adolescent fitness and fatness and cardiovascular disease risk factors in adulthood: the  
361 aerobics center longitudinal study (ACLS). *Am. Heart J.* **2005**, 149:46-53;  
362 DOI: <https://doi.org/10.1016/j.ahj.2004.07.016>.
- 363 12. Raitakari, O.T.; Taimela, S.; Porkka, K.V.K.; Telema, R.; Valimaki, I.; Akerblom,  
364 H.K.; Viikari, J.S.A. Associations between physical activity and risk factors for  
365 coronary heart disease: The Cardiovascular Risk in Young Finns Study. *Med. Sci.*  
366 *Sports Exerc.* **1997**, 29(8): 1055-1061; DOI: 10.1097/00005768-199708000-00011.
- 367 13. Sørensen, L.E.; Pekkonen, M.M.; Mannikkö, K.H.; Louhevaara, V.A.; Smolander, J.;  
368 Alen, M.J. Associations between work ability, health-related quality of life, physical  
369 activity and fitness among middle-aged men. *Applied Ergonomics.* 2008, 39: 786-791;  
370 DOI: <https://doi.org/10.1016/j.apergo.2007.11.001>.
- 371 14. Klasson-Heggebo, L.; Andersen, L.B.; Wennlöf, A.H.; Sardinha, L.B.; Harro, M.;  
372 Froberg, K.; Anderssen, S.A. Graded association between cardiorespiratory  
373 fitness, fatness and blood pressure in children and adolescents. *Br j Sports Med.* **2006**,  
374 40:25-29; DOI: <http://dx.doi.org/10.1136/bjism.2004.016113>.
- 375 15. Lohman, T.G.; Slaughter, M.H.; Boileau, R.A.; Bunt, J.; Lussier, L. Bone  
376 mineral measurements and their relation to body density in children,  
377 youth, and adults. *Human Biology.* **1984**, 56, 667-679; ISSN: 00187143.
- 378 16. Walker, S.N.; Sechrist, K.R.; Pender, N.J. The health-promoting lifestyle profile:  
379 development and psychometric characteristics. *Nursing research.* **1987**, 36(2), 76-81;  
380 DOI: <http://dx.doi.org/10.1097/00006199-198703000-00002>.
- 381 17. Esin, M.N. Evaluation and promotion of health behaviors of industry workers. Istanbul  
382 University Health Science Institute an Unpublished PhD Thesis, Istanbul, (in Turkish),  
383 1997.
- 384 18. Akca, S. Evaluation of health behaviors among lecturers and of affecting factors. Ege  
385 University Health Science Institute an Unpublished PhD Thesis, Izmir (in Turkish),  
386 1998.
- 387 19. Kayıhan, G.; Ersöz, G.; Özkan, A. The relationship between selected  
388 physical-physiological parameters and efficiency of pistol shooting. *Policing: An*  
389 *International Journal of Police Strategies & Management.* **2013** 36(4):  
390 819-832; DOI:<https://doi.org/10.1108/PIJPSM-03-2013-0034>.

- 391 20. Scafoglieri, A.; Tresignie, J.; Provyn, S.; Clarys, J.P.; Bautmans, I. Reproducibility,  
392 accuracy, and concordance of Accutrend® Plus for measuring circulating lipid  
393 concentration in adults. *Biochemia Medica: Biochemia Medica*. **2012**, *22*(1), 100-108.
- 394 21. Van de Vliet, P.; Rintala, P.; Fröjd, K.; Verellen, J.; Van Houtte, S.; Daly, D.J.;  
395 Vanlandewijck, Y.C. Physical fitness profile of elite athletes with intellectual  
396 disability. *Scandinavian Journal of Medicine & Science in Sports*. **2006**, *16*(6), 417-  
397 425; DOI: <https://doi.org/10.1111/j.1600-0838.2006.00539.x>
- 398 22. Costa Silva Zemdegs, J.; Barreto Corsi, L.; De Castro Coelho, L.; Duarte Pimentel, G.;  
399 Toyomi Hirai, A.; Sachs, A. Lipid profile and cardiovascular risk factors among first-  
400 year Brazilian university students in São Paulo. *Nutricion hospitalaria*. **2011**, *26*(3):  
401 553-559; DOI:10.3305/nh.2011.26.3.4660.
- 402
- 403