

NOX Inhibition Improves β -Adrenergic Stimulated Contractility and Intracellular Calcium Handling in the Aged Heart

Álvaro Valdés¹, Guillermo Barrios¹, Nikol Ponce¹, Raul A Dulce², and Daniel R González¹.

1 Departamento de Ciencias Básicas Biomédicas, Facultad Ciencias de la Salud, Universidad de Talca, Talca, Chile.

2 Interdisciplinary Stem Cell Institute, Miller School of Medicine, University of Miami, Miami, Florida, USA

Corresponding autor

Daniel R González, PhD.

Departamento de Ciencias Básicas Biomédicas,
Facultad de Ciencias de la Salud, Universidad de Talca,
Avenida Lircay s/n
Talca 3460000, Chile.

Phone +56-71-2-418856

Email:dagonzalez@utalca.cl

Talca, Chile

Abstract

Cardiac aging is characterized by alterations in contractility and intracellular calcium ($[Ca^{2+}]_i$) homeostasis. It has been suggested that oxidative stress may be involved in this process. We and others have reported that in cardiomyopathies the NADPH oxidase (NOX)-derived superoxide is increased, with a negative impact on $[Ca^{2+}]_i$ and contractility.

We tested the hypothesis that in the aged heart, $[Ca^{2+}]_i$ handling and contractility are disturbed by NOX-derived superoxide. Contractility was evaluated isolated hearts, challenged with isoproterenol. To assess $[Ca^{2+}]_i$, isolated cardiac myocytes were field-stimulated and $[Ca^{2+}]_i$ was monitored with fura-2. Cardiac concentration-response to isoproterenol was depressed in aged compared to adults hearts ($p < 0.005$), but was restored by NOX inhibitors apocynin and VAS2870. In isolated cardiomyocytes, apocynin increased the amplitude of $[Ca^{2+}]_i$ in aged myocytes ($p < 0.05$). Time-50 $[Ca^{2+}]_i$ decay was increased in aged myocytes ($p < 0.05$) and reduced towards normal by NOX inhibition. In addition, we found that myofilaments Ca^{2+} sensitivity was reduced in aged myocytes ($p < 0.05$), and further reduced by apocynin. Finally SERCA levels but not phospholamban were reduced in aged hearts ($p < 0.05$).

In conclusion, β -adrenergic-induced contractility was depressed in aged hearts, and NOX inhibition restored back to normal. Moreover, altered Ca^{2+} handling in aged myocytes was also improved by NOX inhibition. These results suggest a NOX-dependent effect in aged myocytes at the level of Ca^{2+} handling proteins and myofilaments.

Keywords: NADPH oxidases, apocynin, VAS2870, aged heart, cardiomyocytes, isoproterenol, TBARS.

1. Introduction

Aged population is increasing worldwide. Cardiovascular diseases are the leading cause of death worldwide and this becomes even more pronounced in the elderly population. In the heart, aging increases the risk of ischemic disease, atrial fibrillation and heart failure. Cardiac aging involves a progressive degeneration and decline in function in absence of major cardiovascular risks, which makes the heart more vulnerable to stress and contributes to increased cardiovascular mortality and morbidity. Cardiac aging includes hypertrophy, fibrosis and apoptosis of myocytes [1] and at the functional level, the loss of the cardiac reserve [2].

In the heart, the aging process is associated with an increase in the production of reactive oxygen species (ROS) [3]. Traditionally, the increase ROS observed in the aged heart has been attributed to mitochondria since ROS are a byproduct of the mitochondrial respiratory chain. It has been hypothesized that accumulation of these ROS are a cause of aging due to cumulative cellular damage they produce in time. Although this theory is attractive and there is experimental data in support of it, other sources of ROS cannot be discarded [4].

Recent reports suggest that the nicotinamide adenine dinucleotide phosphate (NADPH) oxidases may be relevant in cardiac aging [5, 6]. NADPH oxidases are a family of enzymes that generate reactive oxygen species ($O_2^{\cdot -}$ and H_2O_2) through electron transfer from NADPH to molecular oxygen. Seven family members of these oxidases are known, each based on a distinct catalytic subunit (Nox 1–5 and Dual oxidase 1–2). This oxidase is a multi-sub unit enzyme composed of a membrane-bound heterodimeric subunit (p22^{phox} and gp91^{phox}) and several cytoplasmic regulatory sub units including Rac1, p47^{phox} and p67^{phox}. NADPH oxidases are crucial regulators of redox signaling through tightly regulated production of low ROS levels in the vicinity of target proteins. More recently, NOX activity has been hypothesized to contribute to

cardiac aging [7]. NOX has been also implicated in the cardiovascular remodeling of the aging heart [8].

Several key protein involved in excitation–contraction coupling can undergo redox-sensitive alterations in activity, which contribute to myocardial contractile dysfunction. ROS generated by up regulated NADPH oxidase modulate Ca^{2+} transients [9], and myofilament function via post translational modifications in the aged heart [10]. Altered $[\text{Ca}^{2+}]_i$ regulation is directly responsible for the attenuated contractility and relaxation observed in aged myocardium, which correlated with impairment of both cell shortening and re-lengthening [11].

The aging heart is characterized by changes in the Ca^{2+} handling of cardiac myocytes [12]. Particularly, Ca^{2+} removal from the cytosol by re-uptake into the sarcoplasmic reticulum (SR) is decelerated in aged cardiac myocytes, favoring the development of diastolic dysfunction [13]. As a consequence, left ventricular (LV) diastolic dysfunction is a common condition in the elderly and manifested as exercises intolerance, dyspnea and fatigue [14].

The SR Ca^{2+} –ATPase (SERCA2a) and its inhibitory protein, phospholamban (PLB), are important regulators of SR Ca^{2+} re-uptake. SERCA2a, which couples the hydrolysis of one ATP molecule to transport Ca^{2+} into the SR, remove Ca^{2+} from the cytosol to facilitate relaxation. PLB is a negative regulatory protein of SERCA2a [15, 16]. Dephosphorylated PLB inhibits SERCA activity, whereas PLB phosphorylation, by Ca^{2+} /calmodulin or cAMP dependent protein kinases, relieves this inhibition. Decreased SERCA2a pumping rate, mRNA and protein levels and decreased phospholamban have been reported in senescent rat and human myocardium, while others have reported no change in mRNA and protein level of both SERCA2a and PLB with age [17].

Since we previously described a role for NOX in the dystrophic cardiomyopathy [18], the aim of the present study was to determinate the effect of NOX inhibition on the aged cardiomyocyte contractile function.

2. Materials and methods

2.1. Animals

According to the rat lifespan, we worked with adult and aged animals [19]. All procedures were performed in conform to the *NIH Guide for the Care and Use of Laboratory Animals*. The protocol was approved by the Animal Care Committee of Universidad de Talca.

To study changes in left ventricular function with age, studies were performed in Langendorff isolated perfused beating hearts. Briefly, male and female rats were injected intraperitoneally with thiopental (10 mg/Kg) and subsequently with heparin (100 U/Kg). With the animal was under deep anesthesia, (checked as the complete absence of sensitive reflexes by the investigator), the thorax was rapidly opened, the heart excised, and the aortic root was cannulated to initiate retrograde perfusion. A Krebs–Henseleit buffer was used as the perfusate (in mM: NaCl, 118; KCl, 4.7; CaCl₂, 1.75, MgSO₄, 1.2; KH₂PO₄ 1.2, NaHCO₃ 25; glucose, 11) equilibrated with 95% O₂ y 5% CO₂ (37°C). A polyvinyl balloon–tipped catheter made in our laboratory from polyethylene tubing (PE–50) was inserting into the left ventricle via the mitral valve through a small opening in the left atrium. The balloon was filled with saline solution to adjust the end diastolic pressure (EDP) at 5 mmHg. This balloon was used to monitor left ventricular pressure via a pressure transducer (Ohmeda Instruments P23 XL, Madison, WI, USA) positioned at the level of the heart. A second transducer, connected to a side arm of the aortic perfusion catheter, was used to monitor coronary perfusion pressure. Hearts were paced at a rate of 300 beats/min with an electrical stimulator Grass S6 (Quince, MA, USA). Left ventricular peak systolic pressure (LVPsP), maximum rate of contraction (+dP/dt) and maximal rate of relaxation (-dP/dt) were monitored using a Powerlab System (AD Instruments, Castle Hill, Australia), used as the index of global myocardial performance. All data were continuously acquired by Lab Chart software version 7 for Windows.

2.2 Isolated heart experimental protocols

All hearts were subjected to a stabilization period (15–20 min) before baseline measurement. After that, hearts were perfused with isoproterenol (Sigma–Aldrich, St. Louis, USA) at increasing concentrations (1 nM–1 μ M range). A group of aged hearts was treated with apocynin (Sigma) 100 μ M a not specific inhibitor of NADPH oxidase, before and during the isoproterenol challenge. Another group of aged hearts was treated identically but with the specific NADPH inhibitor VAS2870 [20], 20 μ M (Merck, Darmstadt, Germany).

2.3 Isolation of rat ventricular myocytes

Myocytes were isolated from the left ventricle of male rats as previously described [21]. Briefly, heart were rapidly removed from an anesthetized rat and mounted onto a Langendorff system. After being perfused with a modified Krebs–Henseleit buffer (mM: NaCl, 118; KCl, 4.7; MgSO₄, 1.2; KH₂PO₄ 1.2, NaHCO₃ 25; glucose, 11; butanedione monoxide 10; taurine 5) equilibrated with 95% O₂ y 5% CO₂ (37 °C) for 10 min, the heart was digested for ~75 min with collagenase type II 1 mg/ml (Worthington, NJ, USA) and protease IV 0.1 mg/ml (Sigma–Aldrich, St. Louis, USA) in the modified Krebs–Henseleit buffer. The digested heart was then removed from the cannula, and the left ventricle was cut into small pieces in the modified Krebs–Henseleit buffer. Tissue pieces were centrifuged, and the pellet of cells was resuspended in a modified Tyrode solution (Ca²⁺ free). Extracellular Ca²⁺ was added incrementally back to 1.3 mM over period of 60 min. Only rod–shaped cells with clear striation pattern without blebs or other morphological alterations were selected for mechanical and intracellular Ca²⁺ studies.

2.4 Measurement myocytes shortening/relengthening

Mechanical properties of cardiomyocytes were assessed using a video–based sarcomere length monitoring system (Ion Optix, Milton, MA, USA). In brief, myocytes were transferred to a Lucite chamber on the stage of an inverted microscope (Nikon TE 200, Tokyo, Japan) and superfused

at 37 °C with a Tyrode solution containing 1.3 mM Ca²⁺. Myocytes were field-stimulated and sarcomere length was monitored in real time using an IonOptix iCCD camera and specialized data acquisition software (IonWizard 6.3 SarcLen Acquisition System, IonOptix, Inc). The myocytes were field stimulated to contract at a frequency of 0.5, 2, 3 and 4 Hz using a pair of platinum wires placed on opposite sides of the chamber and connected to a MyoPacer Field Stimulator (Ion Optix, Milton, MA, USA). The myocyte being studied was displayed on the computer monitor using an IonOptix MyoCam CCD100M camera side-mounted onto the microscope. Changes in the average sarcomere length were determined by fast Fourier - transform of the Z-line density trace to the frequency domain using the acquisition software noted above. Changes in sarcomere length during shortening and re-lengthening was captured and converted to digital signal. Sarcomere shortening and re-lengthening were assessed using the following indices: peak shortening (PS), percentage of shortening (%S = $\Delta L / L_0 \times 100$), time to 50% peak shortening (TS₅₀) and time to 50% peak re-lengthening (TR₅₀) and Tau constant of relaxation. Twitch amplitude was computed as the difference between diastolic and peak systolic sarcomere lengths. Percentage of sarcomere shortening was expressed as the ratio of absolute twitch amplitude to diastolic sarcomere length.

2.5 Intracellular calcium

Cardiomyocytes were loaded with fura 2-AM (1 μM from Invitrogen, USA) in the dark for 15 min, and fluorescence measurements were recorder with a dual excitation fluorescence photomultiplier system (Ionoptix, Milton, MA, USA). Cardiomyocytes were placed in a Lucite chamber on the stage on a Nikon Eclipse E200 inverted microscope and imaged through a Fluor 40X oil objective. Intracellular Ca²⁺ concentration ([Ca²⁺]_i) was determined by exciting with light from a Xenon lamp passed alternately through 365 and 380 nm filters. The emission fluorescence was reflected through a barrier filter (510 ± 15 nm) to a photomultiplier tube. The fura-2 emission ratio at 360/380 nm represents an non-calibrated signal of [Ca²⁺]_i,

2.5.1. Analysis of Ca^{2+} transient data.

This was determined as the percentage of change between the levels of $[Ca^{2+}]_i$ in systole and diastole: $[Ca^{2+}]_i = 100 \times (F - F_0)/F_0$. The time course of Ca^{2+} fluorescence signal decay (the duration at which Ca^{2+} transient decays 50%) was calculated to determine $[Ca^{2+}]_i$ clearing rate. The time constant (τ) for myocyte re-lengthening and the Ca^{2+} transient decay is calculated by fitting the decay of the transients to the exponential curve: $Y(t) = a \cdot e^{-\lambda t} + b$. Where a =twitch amplitude or Ca^{2+} transient; b =baseline value; λ is calculated and $\tau = 1/\lambda$.

2.5.2 Assessment of calcium myofilament responsiveness

Changes in myofilament responsiveness to Ca^{2+} were assessed using the steady-state relation between cell length and $[Ca^{2+}]_i$ in intact single cardiac myocytes tetanized by high-frequency (40 Hz) stimulation after exposure to thapsigargin (0.2 μ mol/L for 15 minutes), as described previously [22]. Thapsigargin disables the SR and thus enables tetanization of otherwise intact myocytes. High-frequency electrical stimulation after thapsigargin treatment results in the effective summation of the repetitive transmembrane flux of Ca^{2+} current (because of the absence of sarcoplasmic reticulum Ca^{2+} sequestration and periodic release) to achieve a steady-state level of myoplasmic Ca^{2+} substantially elevated above resting levels, at the point that Ca^{2+} influx is balanced by the rate of Ca^{2+} extrusion (via Na/Ca exchanger). With this approach, the $[Ca^{2+}]_i$ can be reversibly clamped near peak systolic levels during the tetanic contracture for 10 to 20 seconds and then rapidly returned to resting levels on cessation of electrical stimulation via normal Na/Ca exchange mechanisms. Thus, changes in the degree of cell shortening between tetani clamped at the same $[Ca^{2+}]_i$ level can be attributed to changes in the relative myofilament responsiveness to Ca^{2+} .

Phase loop analysis. In addition, the myofilaments dynamics was studied by plotting the levels of $[Ca^{2+}]_i$ against its corresponding value for sarcomere shortening, to obtain information of the activating and the relaxation phases of the myofilaments, as has been previously reported [23, 24]. The relationship sarcomere shortening- $[Ca^{2+}]_i$ relationships were determined experimentally and fitted to the Hill equation to obtain Sarc Shortening mx, or maximal $[Ca^{2+}]_i$ activating shortening and ECa_{50} and the Hill slope as follow.

$$Sarcomere\ Shortening = Min + \frac{(Max - Min)}{1 + 10^{(LogEC50 - [Ca^{2+}]_i) \cdot Hill\ Slope}}$$

2.6. Oxidative stress analysis

Oxidative stress was evaluated using the levels of thiobarbituric acid reactive substances (TBARS). TBARS were quantified using heart homogenate supernatants, that were mixed with SDS (8% w/v), thiobarbituric acid (0.8% w/v), and acetic acid (20% v/v) and heated for 60 min at 90°C. Precipitated material was removed by centrifugation, and the absorbance of the supernatant was determined at 530 nm. Levels of TBARS were calculated using a calibration curve with malondialdehyde (MDA, Sigma Aldrich, Saint Louis, USA) in a Multiskan GO Microplate spectrophotometer (Thermo Scientific, Rockford, USA).

2.7. Western Blotting

Tissues samples from the heart ventricle were removed and homogenized in a lysis buffer containing 50 mM Tris (pH 7.4), 30 mM NaCl, 2 mM EDTA, 0.1 % SDS and 1% protease inhibitor cocktail (Sigma–Aldrich, St. Louis, USA). Samples were then centrifuged at 1500 rpm for 10 min at 4 oC. The protein concentration of the supernatant was determined by the bicinchoninic acid assay using bovine serum albumin as standard (Thermo Scientific, Rockford, USA). Same amount of the protein (100 µg protein/lane) and pre stained molecular weight marker (PageRuler™ Plus Thermo Scientific, Rockford, USA) were separated on 15% SDS–

polyacrylamide gel, then were transferred to nitrocellulose membranes (0.2 μm pore Bio-Rad Laboratories, Hercules, CA, USA). Membranes were incubated for 1 hour in a blocking solution containing 5% non-fat dry milk in TBT-T buffer, and were then incubated with anti-SERCA2a (Santa Cruz Biotechnologies, Santa Cruz, C, 1:1000), anti-PLB (Badrilla, Leeds, UK, 1:1000), anti p47phox (Cell Signaling Technology, Danvers, MA), 1:1000, and anti-GADPH (Santa Cruz) antibodies at 4°C overnight. After washing blots to remove excess primary antibody binding, blots were incubated for 1 hour with horseradish peroxidase (HRP)-conjugated secondary antibody (Santa Cruz, 1:2000). Antibody binding was detected using enhanced chemiluminescence reagent (SuperSignal® West Femto Maximum Sensitivity Substrate, Thermo Scientific, Rockford, USA) and film were digitalized and scanned and the intensity of immunoblot bands detected with Image J software version 1.49v (Wayne Rasband, National Institute of Health, USA).

2.8 Statistical Analysis

Sample size (animals) was estimated by performing power analysis based on previous data for contractility parameters, to achieve an 80% power (20% β -error) with α -error < 5%. Results are expressed as mean \pm standard deviation or mean \pm standard error where indicated. Comparison between two groups were made using the unpaired, two tails *t* student test. Analysis of concentration-response curves and shortening-frequency response experiments were performed using two-way analysis of variance (ANOVA) with Bonferroni as post hoc analysis and one way analysis with Neuman-Keuls post-hoc analysis, where appropriate. Isoproterenol concentration-response curves were fitted to the Hill equation as described above. For non-parametric data, Kruskal-Wallis test was applied, with Dunn's multiple comparisons test as post-hoc analysis. Statistical significance was set at a value of $p < 0.05$. These analyses were performed using the GraphPad Prism 5 software (San Diego, CA).

3. Results

3.1. General characteristic of adult and senescent rats

The baseline characteristics of the animals in the study and hemodynamic parameters are presented in Table 1. Body weight was increased in senescent compared to adult rats. Heart weight was also increased in aged rats, emphasizing that in the rat, aging is associated with cardiac hypertrophy. Nevertheless, aged hearts displayed diminished ventricular function evaluated as developed pressure, contractility (dp/dt_{max}) and lusitropy (dp/dt_{min}).

3.2. Isoproterenol response in the isolated heart

To assess cardiac ventricular function in the aged myocardium, we submitted isolated hearts from adult and aged rats to stimulation with increasing concentrations of isoproterenol, a β -adrenergic agonist. We observed that adult hearts displayed a positive inotropic and lusitropic response to increasing concentrations of isoproterenol, while aged hearts showed a significantly diminished response (Figure 1). Next, we treated aged hearts with NOX inhibitors. Apocynin (100 μ M) produced a complete recovery of developed pressure, dP/dt_{max} and dP/dt_{min} in aged hearts. On the other hand, apocynin had no effect on adult hearts (supplementary material). A more selective NOX inhibitor, VAS2870 (20 μ M) also produced a significant recovery of ventricular function in aged hearts, although not as complete as apocynin. These results suggest for a role of NOX in the diminished cardiac reserve in the aged rat heart.

3.3 Isolated cardiomyocytes

Next, to gain more insights into the cellular mechanisms involved in the effect of the NOX inhibitors and to rule out the influence of vascular effects, we designed experiments in isolated cardiomyocytes.

Baseline characteristic of isolated myocytes are indicated in Table 2. Regarding sarcomere shortening, aged myocytes presented prolonged time 50 to peak shortening (TS_{50}) and prolonged T_{50} to re-lengthening.

3.3.1. Sarcomere Shortening.

Isolated cardiomyocytes were submitted to a train of increasing frequency rates of stimulation, from 0.5 to 4 Hz (Figure 2). In adult myocytes, the amplitude of this response in sarcomere shortening is usually negative. This was the case for both adult and aged myocytes, although many aged myocytes could not complete the train of stimulation (not shown), not being able to reach 4 Hz. There were no differences in shortening amplitude between adult and aged myocytes and this amplitude was not further changed by apocynin treatment. The kinetic parameters T_{50} of peak shortening was increased in aged myocytes ($p < 0.05$), and was not improved by apocynin treatment. The same was obtained T_{50} of relaxation.

3.3.2. Intracellular Ca^{2+} Transients.

In parallel to myocyte shortening, we evaluated intracellular Ca^{2+} in isolated myocytes (Figure 3). $[Ca^{2+}]_i$ transients amplitude was slightly increased in aged cardiomyocytes ($p < 0.05$) and were further increased by apocynin treatment. T_{50} to peak Ca^{2+} was increased in aged myocytes ($p < 0.05$), and was not improved by the apocynin treatment. T_{50} to peak relaxation was increased in aged myocytes ($p < 0.05$) and reduced towards normal by apocynin treatment. These results suggest impairments in the release and reuptake of cytosolic Ca^{2+} in aged myocytes.

3.3.3. Myofilaments Ca^{2+} Sensitivity.

The diminished degree of contraction associated with sustained Ca^{2+} transients in aged myocytes suggests reduced Ca^{2+} sensitivity in the aged heart cells. We tested this hypothesis using a protocol treating the myocytes with thapsigargin, a pharmacological SERCA blocker.

After incubation with thapsigargin, myocytes were submitted to tetanic stimulation (40 Hz) to evaluate the myofilaments Ca^{2+} sensitivity (figure 4A). While the amplitude of the level of $[\text{Ca}^{2+}]_i$ evoked by stimulation was not different between adult and aged cardiomyocytes, the tetanic contraction was reduced in aged myocytes ($p < 0.05$), suggesting reduced Ca^{2+} sensitivity.

To further investigate the sensitivity to Ca^{2+} and myofilament interaction in these cardiomyocytes, we plotted sarcomere shortening versus $[\text{Ca}^{2+}]_i$ to obtain the hysteresis loops (Figure 4B). Slower relaxation is associated with sensitization of myofilaments to Ca^{2+} and increased affinity. Accordingly, we analyzed sarcomere shortening– $[\text{Ca}^{2+}]_i$ hysteresis loops of cardiomyocyte contractions induced by field-stimulation at either 1 or 4 Hz, in order to detect differences in responsiveness to Ca^{2+} . In general, the morphology of the loops was greatly affected in cardiomyocytes from aged rats. We observed a general frequency-dependent myofilament desensitization to Ca^{2+} (shift to the right of the loops) from 1 to 4 Hz, which was more pronounced in the aged groups (either in the absence or presence of apocynin) compared to the adult group (Figure 4C). From the analysis of the relaxation phases (segment transitioning between the maximum and minimum sarcomere length shortening), it was observed a differential behavior depending on the stimulation rate (Figure 4C). At 1 Hz, the aged myocytes exhibited increased sensitivity compared to adult controls, however this difference was inverted by increasing frequency (4 Hz) (Figure 4D). This substantial reduction in myofilament sensitivity to Ca^{2+} in the aged group by increasing pacing is consistent with our results in tetanized cardiomyocytes shown in Figure 4A. The treatment of aged myocytes with apocynin induced a robust desensitizing effect at the studied frequencies (Figure 4C and D).

The relationship Ca^{2+} –sarcomere shortening was further assessed through the analysis of the force activation phase of the sarcomere shortening– $[\text{Ca}^{2+}]_i$ loops (which is located between the maximum $[\text{Ca}^{2+}]_i$ and the maximum contraction as indicated in Figure 4C by lines), in order to evidence the effect of aging on the cross-bridge cycling. At 1 Hz, the slopes of the activation phase (actSlope) obtained by linear regression of the shortening– $[\text{Ca}^{2+}]_i$ pairs within the

mentioned segment of the loop (Figure 4C, bar graphs), were virtually similar in both groups, adults and aged. However, the increasing stimulation induced a decrease in the force-activation slope in myocytes from the aged rats, which might indicate deteriorated cross-bridge cooperativity. Consistently, apocynin promoted a reduction in the force activation slope at 1 Hz and did not correct the impaired myofilament interaction at 4 Hz in aged cardiomyocytes. However this treatment was able to preserve the contractile performance in these cells by increasing the systolic $[Ca^{2+}]_i$ and consequently the $[Ca^{2+}]_i$ amplitude.

3.4 Cardiac Oxidative Stress

NOX-derived oxidative stress was investigated by two methods (figure 5). First, we evaluated the expression of the p47^{phox} protein, one of the key regulators of NOX2. Levels of p47^{phox} were increased in aged hearts compared to the adult ones ($p < 0.05$). In addition we evaluated the concentration of thiobarbituric acid (TBARS), a marker of lipid peroxidation. TBARS concentration was increased in aged hearts, and was reduced towards in the aged hearts that were treated with apocynin ($p < 0.05$). This confirms that use of the NOX inhibitor reduced the cardiac oxidative stress.

3.5 Ca^{2+} -handling proteins levels.

Finally, since the kinetic parameters of Ca^{2+} release and re-uptake were altered in aged myocytes, we decided to look into the levels of Ca^{2+} handling proteins in cardiac homogenates (figure 6). SERCA2 and phospholamban (PLB) expression in adults and aged hearts were assessed by Western blotting. SERCA2 levels were reduced in aged hearts, while those of phospholamban were not different between groups. Consequently, the ratio of SERCA to PLB was decreased in aged hearts ($p < 0.05$). This suggests that in the aged rat myocardium, the capacity for Ca^{2+} reuptake is diminished.

4. Discussion

In the aged myocardium, increased ROS are associated with a decrease in function, as well as on ventricular remodeling. This ROS production has been ascribed to the mitochondria.

In the heart NOX2 and NOX4 are expressed. NOX2 is located in the sarcolemma while NOX4 is located in the mitochondrial matrix [25]. It is possible that in the aged rat heart, both isoforms may have contributed to the increased oxidative stress, since apocynin completely recovered ventricular function in the isolated heart, while VAS2870, a relatively NOX2 specific inhibitor restores partially this function.

Here we observed that the aged heart presents a diminished response to β -adrenergic stimulation compared to adult heart and for the first time show that acute NOX inhibition is able to restore this fundamental mechanism of cardiac inotropism. This impairment was reversed pharmacologically by the use of NOX inhibitors. At the subcellular level, we observed that aged myocytes presented a slightly increase in amplitude of Ca^{2+} transients and reduced myofilaments Ca^{2+} sensitivity. Apocynin, a NOX inhibitor increased the amplitude of these transients. Our results suggest that this agent improved the function of the RyR2 and SERCA2, improving the Ca^{2+} cycling, although the molecular mechanism remains to be determined.

There is general agreement that in the aged myocyte, the activating L-type Ca^{2+} is reduced [26], the activity of the RyR2 is reduced (at the level of Ca^{2+} sparks) [27] [28] and SR Ca^{2+} re-uptake is slowed [29], in part by reduced SERCA expression, making Ca^{2+} transients smaller in amplitude and slower [13]. Several of these observations have been confirmed in the present work. NOX inhibition may have influenced both SERCA and RyR2, in a redox-sensitive manner, since all the pharmacological maneuvers performed in this study were acute. Particularly, RyR2 is extremely redox sensitive [30], due to its large number of cysteines residues [31, 32].

Our results agree with those observed by Rueckschloss *et al* [33], who observed similar effects of NOX inhibition in aged compared to young mouse myocytes. Nevertheless, in our model we evaluated in the intact heart the β -adrenergic response. Furthermore, contrary to Rueckschloss *et al*, we observed significant changes in the Ca^{2+} handling proteins SERCA and in the ratio SERCA/phospholamban.

These results suggest that NOX-derived superoxide may have produced oxidative modifications on Ca^{2+} handling and myofilaments proteins. Cooper *et al* showed that in the aged rabbit heart, RyR2 is oxidized, although in that model the source of ROS was mitochondrial [34]. In the aged mouse, Andersson *et al* [35] found increased S-nitrosylation of RyR1. On the other hand, Qin *et al* found in the aged mouse heart that SERCA is oxidized by sulfonation at cysteine 674, decreasing its activity, although the source of ROS was not identified [36].

This increased activity in NOX activity in the aged heart has been proposed to depend on nitric oxide 1 (NOS1) [37] and on the renin-angiotensin system [8].

4.1 Study Limitations

The present study has several limitations. First, the inhibitors used are not selective for the different NOX isoforms, and hence it is not possible to distinguish whether the effect observed is due to NOX2 or NOX4, which are the main isoforms present in the cardiac myocytes. Second, although it is shown that NOX inhibition reduced the oxidative stress, the exact redox modifications on target proteins such as the excitation-contraction coupling machinery or the myofilaments responsible for the effects observed were not identified.

5. Conclusions

Aged rat hearts presented a diminished response to isoproterenol, which was restored upon pharmacological NOX inhibition. In aged myocytes, twitch contractions, presented slowed kinetics, and reduced myofilaments Ca^{2+} -sensitivity. NOX inhibition increased $[\text{Ca}^{2+}]_i$ transients amplitude and improved Ca^{2+} kinetics. Aged rat hearts presented lower levels of SERCA2.

Understanding how NOX activity influences the aged myocardium function at the cellular level may help to develop strategies for the prevention and treatment of cardiovascular disease in this population. Pharmacological targeting of NOX in aging may help to improve cardiac function.

Acknowledgements.

Funded by grant FONDECYT 1150662 (Daniel R Gonzalez), and Program of Excellence Interdisciplinary Research QuimBio and Interdisciplinary Excellence Research Program on Healthy Aging (PIEI-ES). Universidad de Talca

Declaration of conflict of interest: None.

Legends to Figures

Figure 1. *Heart function in the Langendorff preparation.*

Adult and aged isolated rat hearts were challenged with increasing concentrations of isoproterenol. Contractility was evaluated as developed pressure, dP/dt_{max} and lusitropy as dP/dt_{min} . Aged hearts were treated with NOX inhibitors apocynin (100 $\mu\text{mol/L}$) or VAS2870 (20 $\mu\text{mol/L}$). N=5 hearts each group. Values are presented as mean \pm standard error; $n = 5$. * $p < 0.05$, ** $p < 0.01$ and *** $p < 0.001$ versus corresponding control.

Figure 2. *Contractility in isolated myocytes.* Isolated cardiomyocytes stimulated at increasing frequencies. The left panels show representative traces of original trains of stimulation in adult ($n=19$), aged ($n=22$) and aged myocytes treated with apocynin 50 $\mu\text{mol/L}$ ($n=16$). The lower panel shows single twitch traces from each group (superimposed for comparison), of cardiomyocytes stimulated at 0.5, 1 and 2 Hz. The graphics to the right depict the analysis of sarcomere shortening, time to reach 50% of peak shortening (T_{50} peak contraction) and time to reach 50% (T_{50} relaxation). * $p < 0.05$ vs. adults.

Figure 3. *Intracellular Ca^{2+} transients.* $[\text{Ca}^{2+}]_i$ transients in response to increase in frequency of stimulation from 0.5 to 4 Hz. Upper left panel, representative traces of intracellular calcium in myocytes from adult ($n= 17$), aged ($n= 23$) and aged myocytes treated with apocynin 50 $\mu\text{mol/L}$ ($n=18$). The numbers on the traces indicate the frequency of stimulation. The lower panel shows representative traces with more temporal resolution depicting transients evoked at 0.5, 1 and 2 Hz. The graphs depict the amplitude of $[\text{Ca}^{2+}]_i$ transients, time 50 to reach peak $[\text{Ca}^{2+}]_i$ (T_{50} peak $[\text{Ca}^{2+}]_i$), and time 50 to reach relaxation (T_{50} baseline $[\text{Ca}^{2+}]_i$). * $p < 0.05$ vs. adult, *** $p < 0.001$ vs. adult.

Figure 4. Myofilaments Sensitivity. A, Using thapsigargin to block SERCA activity, myocytes were submitted to tetanic stimulation (40 Hz) to evaluate the myofilaments Ca^{2+} sensitivity.

Left panels: contraction (up) and the $[\text{Ca}^{2+}]_i$ evoked by tetanic stimulation. Right, pooled data for the amplitude of contraction (evaluated as degree of sarcomere shortening) and the amplitude of the cytosolic Ca^{2+} concentration evoked by tetanic stimulus in adult and aged cardiomyocytes. ***, $p=0.0002$. N= 20 cells from adult (n=3) hearts, 29 cells from aged hearts (n=3).

B. Phase loop analysis for the sarcomere shortening- $[\text{Ca}^{2+}]_i$ relationship. Example of hysteresis analysis of Ca^{2+} vs. sarcomere shortening for a normal twitch. The segment from a-b represents the Ca^{2+} activating phase and the b-c segment represents cross-bridge cycling, and b-c segment represent the relaxation phase. C, Hysteresis loops used to obtain the slope of the activation phase at 1 and 4 Hz. D, sarcomere shortening- $[\text{Ca}^{2+}]_i$ in the activating phase fitted to the Hill equation to obtain the ECa_{50} , the Ca^{2+} concentration that activates the 50% of the maximal shortening (normalized). The filled symbols represent determinations obtained at 1 Hz of stimulation; the open symbols represent determinations obtained at 4 Hz of stimulation.

N= 14 adults, 16 aged and 11 aged + apocynin cells for experiments at 1 Hz; 9 adult, 9 aged and 7 aged + apocynin cell for experiments at 4 Hz. * $p<0.001$ vs. adult. † $p<0.001$ vs. aged.

Figure 5. Oxidative Stress.

NOX-derived oxidative stress was investigated by two methods. Upper panel, Western blot for p47^{phox}, one of the activating subunits of NOX2, in adults (n=4) and aged hearts (4) and the quantification normalized to GAPDH levels (mean \pm standard deviation). The lower graph depicts the concentration (mean \pm standard deviation) of thiobarbituric acid (TBARS), a marker of lipid peroxidation, in cardiac homogenates from adult (n=4), aged (n=5) and aged hearts treated with apocynin 100 $\mu\text{mol/L}$ (n=5). * $p < 0.05$ vs. adult and aged plus apocynin.

Figure 6. Calcium handling proteins expression.

Left, representative images of Western blots for SERCA2 and phospholamban (PLB), from adult, aged and aged hearts treated with apocynin (100 $\mu\text{mol/L}$). The graphs to the right depict the densitometry analysis (mean \pm standard deviation), normalized to GAPDH levels. N= 5 hearts each group * p <0.05 vs. adults.

Table 1. Baseline characteristics of isolated hearts. Mean \pm standard deviation

	Adults	Aged	P value
N (hearts)	12	19	-
Age (months, range)	5 - 6	20-24	-
Heart weight (g)	1.45 \pm 0.08	2.19 \pm 0.13	0.00069551
Body weight (g)	344.4 \pm 50	702.5 \pm 82	<0.0001
Developed pressure (mmHg)	108.3 \pm 53.5	41.4 \pm 19.1	0.02956983
dP/dt max (mmHg/s)	2808 \pm 1269	1170 \pm 815	0.0281473
dP/dt min (-mmHg/s)	-2054 \pm 988	-743 \pm 650	0.02598013
Coronary perfusion pressure (mmHg)	60.5 \pm 34	37.9 \pm 9.9	0.17102549

Table 2. Baseline values for sarcomere shortening and intracellular calcium of isolated myocytes (mean \pm standard deviation)

Sarcomere Shortening	Adults	Aged	p value
N (hearts)	5	4	-
N (cells)	17	22	-
Sarcomere Length (μm)	1.74 \pm 0.066	1.73 \pm 0.077	0.289
Contraction Amplitude (μm)	0.189 \pm 0.056	0.17 \pm 0.07	0.185
% Peak to baseline	10.8 \pm 3.1	9.76 \pm 4.1	0.185
Time to peak 50%	0.072 \pm 0.013	0.107 \pm 0.04	<0.005
Time to 50% baseline	0.379 \pm 0.12	0.542 \pm 0.09	<0.005
Tau relaxation	0.193 \pm 0.407	0.186 \pm 0.08	0.407
<i>Intracellular Ca²⁺</i>			
Diastolic [Ca ²⁺] _i (fura ratio)	1.2 \pm 0.18	1.06 \pm 0.05	0.0051
Systolic [Ca ²⁺] _i (fura ratio)	0.20 \pm 0.12	0.27 \pm 0.2	0.0916
% Peak to baseline	17.1 \pm 11.1	26.7 \pm 20.7	0.044
Time to peak 50% [Ca ²⁺] _i	0.04 \pm 0.005	0.05 \pm 0.002	0.0067
Time to 50% baseline [Ca ²⁺] _i	0.25 \pm 0.08	0.35 \pm 0.01	<0.005
Tau [Ca ²⁺] _i decay	0.26 \pm 0.148	0.25 \pm 0.06	0.420

Figure 1




FIGURE 2





Figure 3




FIGURE 4




FIGURE 5





FIGURE 6



References

- [1] Boyle AJ, Shih H, Hwang J, Ye J, Lee B, Zhang Y et al. Cardiomyopathy of aging in the mammalian heart is characterized by myocardial hypertrophy, fibrosis and a predisposition towards cardiomyocyte apoptosis and autophagy. *Exp Gerontol* 2011;46:549-59.
- [2] Lakatta EG, Levy D. Arterial and cardiac aging: major shareholders in cardiovascular disease enterprises: Part II: the aging heart in health: links to heart disease. *Circulation* 2003;107:346-54.
- [3] Raju SV, Barouch LA, Hare JM. Nitric oxide and oxidative stress in cardiovascular aging. *Sci Aging Knowledge Environ* 2005;2005:re4.
- [4] Dai DF, Chen T, Johnson SC, Szeto H, Rabinovitch PS. Cardiac aging: from molecular mechanisms to significance in human health and disease. *Antioxid Redox Signal* 2012;16:1492-526.
- [5] Lee Y, Kwak HB, Hord J, Kim JH, Lawler JM. Exercise training attenuates age-dependent elevation of angiotensin II type 1 receptor and Nox2 signaling in the rat heart. *Exp Gerontol* 2015;70:163-73.
- [6] Oudot A, Martin C, Busseuil D, Vergely C, Demaison L, Rochette L. NADPH oxidases are in part responsible for increased cardiovascular superoxide production during aging. *Free Radic Biol Med* 2006;40:2214-22.
- [7] Krause KH. Aging: a revisited theory based on free radicals generated by NOX family NADPH oxidases. *Exp Gerontol* 2007;42:256-62.
- [8] Wang M, Zhang J, Walker SJ, Dworakowski R, Lakatta EG, Shah AM. Involvement of NADPH oxidase in age-associated cardiac remodeling. *J Mol Cell Cardiol* 2010;48:765-72.
- [9] Prosser BL, Ward CW, Lederer WJ. X-ROS signaling: rapid mechano-chemo transduction in heart. *Science* 2011;333:1440-5.
- [10] Li SY, Du M, Dolence EK, Fang CX, Mayer GE, Ceylan-Isik AF et al. Aging induces cardiac diastolic dysfunction, oxidative stress, accumulation of advanced glycation endproducts and protein modification. *Aging Cell* 2005;4:57-64.
- [11] Grandy SA, Howlett SE. Cardiac excitation-contraction coupling is altered in myocytes from aged male mice but not in cells from aged female mice. *Am J Physiol Heart Circ Physiol* 2006;291:H2362-H2370.
- [12] Fares E, Howlett SE. Effect of age on cardiac excitation-contraction coupling. *Clin Exp Pharmacol Physiol* 2010;37:1-7.
- [13] Janczewski AM, Lakatta EG. Modulation of sarcoplasmic reticulum Ca(2+) cycling in systolic and diastolic heart failure associated with aging. *Heart Fail Rev* 2010;15:431-45.

- [14] Dai X, Hummel SL, Salazar JB, Taffet GE, Zieman S, Schwartz JB. Cardiovascular physiology in the older adults. *J Geriatr Cardiol* 2015;12:196-201.
- [15] Kranias EG, Hajjar RJ. Modulation of cardiac contractility by the phospholamban/SERCA2a regulatome. *Circ Res* 2012;110:1646-60.
- [16] MacLennan DH, Kranias EG. Phospholamban: a crucial regulator of cardiac contractility. *Nat Rev Mol Cell Biol* 2003;4:566-77.
- [17] Feridooni HA, Dibb KM, Howlett SE. How cardiomyocyte excitation, calcium release and contraction become altered with age. *J Mol Cell Cardiol* 2015;83:62-72.
- [18] Gonzalez DR, Treuer AV, Lamirault G, Mayo V, Cao Y, Dulce RA et al. NADPH oxidase-2 inhibition restores contractility and intracellular calcium handling and reduces arrhythmogenicity in dystrophic cardiomyopathy. *Am J Physiol Heart Circ Physiol* 2014;307:H710-H721.
- [19] Andreollo NA, Santos EF, Araujo MR, Lopes LR. Rat's age versus human's age: what is the relationship? *Arq Bras Cir Dig* 2012;25:49-51.
- [20] ten FH, Huntgeburth M, Wingler K, Schnitker J, Baumer AT, Vantler M et al. Novel Nox inhibitor VAS2870 attenuates PDGF-dependent smooth muscle cell chemotaxis, but not proliferation. *Cardiovasc Res* 2006;71:331-41.
- [21] Gonzalez DR, Treuer AV, Castellanos J, Dulce RA, Hare JM. Impaired S-nitrosylation of the ryanodine receptor caused by xanthine oxidase activity contributes to calcium leak in heart failure. *J Biol Chem* 2010;285:28938-45.
- [22] Dulce RA, Yiginer O, Gonzalez DR, Goss G, Feng N, Zheng M et al. Hydralazine and organic nitrates restore impaired excitation-contraction coupling by reducing calcium leak associated with nitroso-redox imbalance. *J Biol Chem* 2013;288:6522-33.
- [23] Ramirez-Correa GA, Cortassa S, Stanley B, Gao WD, Murphy AM. Calcium sensitivity, force frequency relationship and cardiac troponin I: critical role of PKA and PKC phosphorylation sites. *J Mol Cell Cardiol* 2010;48:943-53.
- [24] Tong CW, Gaffin RD, Zawieja DC, Muthuchamy M. Roles of phosphorylation of myosin binding protein-C and troponin I in mouse cardiac muscle twitch dynamics. *J Physiol* 2004;558:927-41.
- [25] Ago T, Matsushima S, Kuroda J, Zablocki D, Kitazono T, Sadoshima J. The NADPH oxidase Nox4 and aging in the heart. *Aging (Albany NY)* 2010;2:1012-6.
- [26] Howlett SE. Age-associated changes in excitation-contraction coupling are more prominent in ventricular myocytes from male rats than in myocytes from female rats. *Am J Physiol Heart Circ Physiol* 2010;298:H659-H670.
- [27] Howlett SE, Grandy SA, Ferrier GR. Calcium spark properties in ventricular myocytes are altered in aged mice. *Am J Physiol Heart Circ Physiol* 2006;290:H1566-H1574.

- [28] Weisleder N, Ma J. Altered Ca²⁺ sparks in aging skeletal and cardiac muscle. *Ageing Res Rev* 2008;7:177-88.
- [29] Isenberg G, Borschke B, Rueckschloss U. Ca²⁺ transients of cardiomyocytes from senescent mice peak late and decay slowly. *Cell Calcium* 2003;34:271-80.
- [30] Donoso P, Finkelstein JP, Montecinos L, Said M, Sanchez G, Vittone L et al. Stimulation of NOX2 in isolated hearts reversibly sensitizes RyR2 channels to activation by cytoplasmic calcium. *J Mol Cell Cardiol* 2014;68:38-46.
- [31] Gonzalez DR, Treuer A, Sun QA, Stamler JS, Hare JM. S-Nitrosylation of cardiac ion channels. *J Cardiovasc Pharmacol* 2009;54:188-95.
- [32] Treuer AV, Gonzalez DR. Nitric oxide synthases, S-nitrosylation and cardiovascular health: from molecular mechanisms to therapeutic opportunities (review). *Mol Med Rep* 2015;11:1555-65.
- [33] Rueckschloss U, Villmow M, Klockner U. NADPH oxidase-derived superoxide impairs calcium transients and contraction in aged murine ventricular myocytes. *Exp Gerontol* 2010;45:788-96.
- [34] Cooper LL, Li W, Lu Y, Centracchio J, Terentyeva R, Koren G et al. Redox modification of ryanodine receptors by mitochondria-derived reactive oxygen species contributes to aberrant Ca²⁺ handling in ageing rabbit hearts. *J Physiol* 2013;591:5895-911.
- [35] Andersson DC, Betzenhauser MJ, Reiken S, Meli AC, Umanskaya A, Xie W et al. Ryanodine receptor oxidation causes intracellular calcium leak and muscle weakness in aging. *Cell Metab* 2011;14:196-207.
- [36] Qin F, Siwik DA, Lancel S, Zhang J, Kuster GM, Luptak I et al. Hydrogen peroxide-mediated SERCA cysteine 674 oxidation contributes to impaired cardiac myocyte relaxation in senescent mouse heart. *J Am Heart Assoc* 2013;2:e000184.
- [37] Villmow M, Klockner U, Heymes C, Gekle M, Rueckschloss U. NOS1 induces NADPH oxidases and impairs contraction kinetics in aged murine ventricular myocytes. *Basic Res Cardiol* 2015;110:506.