

Type of the Paper (Article)

# PPAR $\gamma$ Expression is Diminished in Macrophages of Recurrent Miscarriage Placentas

Theresa Maria Kolben<sup>1</sup>, Elisabeth Rogatsch<sup>1</sup>, Aurelia Vattai<sup>1</sup>, Anna Hester<sup>1</sup>, Christina Kuhn<sup>1</sup>, Elisa Schmoeckel<sup>2</sup>, Sven Mahner<sup>1</sup>, Udo Jeschke<sup>1\*</sup> and Thomas Kolben<sup>1</sup>

<sup>1</sup> Department of Obstetrics and Gynecology, University Hospital, LMU Munich, Marchioninstr. 15, 81377 Munich, Germany; Theresa.kolben@med.uni-muenchen.de, elisabeth.rogatsch@icloud.com, Aurelia.vattai@med.uni-muenchen.de, anna.hester@med.uni-muenchen.de, Christina.kuhn@med.uni-muenchen.de, sven.mahner@med.uni-muenchen.de, Thomas.kolben@med.uni-muenchen.de

<sup>2</sup> Department of Pathology, LMU Munich, Marchioninstr. 27, 81377 Munich, Germany; elisa.schmoeckel@med.uni-muenchen.de

\* Correspondence: udo.jeschke@med.uni-muenchen.de; Tel.: +49-89-4400-74775

**Abstract:** PPAR $\gamma$  belongs to the group of nuclear receptors which is expressed in the trophoblast and together with other factors is responsible for the maintenance of pregnancy. Apart from that PPAR $\gamma$  is also a main factor for macrophage polarization. The aim of this study was to investigate the combined expression pattern and frequency of PPAR $\gamma$  under physiological circumstances and in spontaneous and recurrent miscarriages in the trophoblast and in maternal macrophages of the decidua. Human placental tissues of the first trimester (15 physiologic pregnancies, 15 spontaneous abortion & 16 recurrent miscarriage placentas) were analyzed for expression of the nuclear receptor PPAR $\gamma$ . Expression changes were evaluated by immunohistochemistry and RT-PCR in trophoblast and in maternal macrophages of the decidua. Maternal macrophages were identified by double immunofluorescence using CD68 as marker for macrophages. The intermediate villous trophoblast revealed a significantly lower PPAR $\gamma$  expression in spontaneous and recurrent abortion. Maternal macrophages express PPAR $\gamma$ . Their number is significantly enhanced in the decidua of spontaneous miscarriages whereas in recurrent miscarriages maternal macrophages seem to express PPAR $\gamma$  only in very few cases. PPAR $\gamma$  is associated with an M2 polarization state that is common for decidual macrophages. The lack of PPAR $\gamma$  in recurrent miscarriage decidual macrophages seems to be associated with a specific inflammatory response against the fetus.

**Keywords:** PPAR $\gamma$ ; first trimester placenta; decidual macrophages; miscarriage

## 1. Introduction

Miscarriage, which is defined as either spontaneous or recurrent is a common disorder in pregnancy[1]. It is affecting 25-50% of all reproductive aged women. Immunologic, endocrine and metabolic mechanisms are involved in the success of human pregnancy and disturbances in any of these processes can lead to fetal loss [2]. Established risk factors are fetal chromosomal or endocrine disorders for spontaneous miscarriages and the antiphospholipid syndrome, thrombophilia or maternal anatomical malformations, especially for recurrent pregnancy losses[3]. In nearly 50% of affected patients, however, the cause of miscarriage remains unknown [4].

Peroxisome proliferator-activated receptor gamma (PPAR $\gamma$ ) belongs to the family of nuclear receptors[5] which are key players in maintaining pregnancy[6,7]. PPAR $\gamma$  together with its heterodimer binding partner retinoid X receptor  $\alpha$  (RXR $\alpha$ ) are involved in cell proliferation, cell differentiation, and organogenesis[8]. RXR $\alpha$  is upregulated in extravillous trophoblast in recurrent miscarriages in humans [9]. RXR $\alpha$  plays a pivotal role in the receptor family, due to its ability to form heterodimers with other nuclear receptors. Heterodimer partners are e.g. peroxisome proliferator-activated receptor (PPAR), thyroid hormone receptor (TR), and liver X receptor (LXR) [2,10-12]. Especially the expression of the isoform PPAR $\gamma$  is linked to trophoblast invasion[13] and downregulation of the isoform RXR $\alpha$  seems to protect from apoptosis in human trophoblasts[9].

Not only trophoblast cells express PPAR $\gamma$  but also macrophages [14]. Macrophages play a key role in immune response and they can respond to environmental stimuli by acquiring specific phenotypes [15]. In response to external cues they will undergo classical M1 activation with high levels of inflammation and microbicide as well as anti-tumor activity. Alternatively the M2 pathway contains mostly parasite containment, tissue remodeling and most importantly in this case immunomodulatory functions like pregnancy [16-18].

Our former studies showed that the number of decidual macrophages is increased at the fetomaternal interface of preeclampsia placentas [19] and also in spontaneous miscarriage cases. An additional finding was the FasL-positivity of these macrophages [20]. Therefore the aim of this study was a phenotype characterization of macrophage populations in abortive placental tissue, its PPAR $\gamma$  expression and the characterization of PPAR $\gamma$  expressing trophoblast sub-types.

## 2. Results

### 2.1. Immunohistochemistry

#### 2.1.1. PPAR $\gamma$ - expression in the trophoblast

The expression of PPAR $\gamma$  in the nucleus and cytoplasm of trophoblast cells was analyzed in tissue from healthy pregnancies, SM, and RM (Figure 1a-d). The intermediate villous trophoblast (IVT) revealed a significantly lower expression in the group with recurrent abortions (Figure 1b) than in the group with healthy placentas (Figure 1a, IRS 8 versus 12,  $p=0.01$ ). There was a significant downregulation of PPAR $\gamma$  in the IVT of spontaneous abortions (Figure 1c, IRS 9 versus 12,  $p=0.001$ ) compared to the control group. Briefly, the staining results are shown in the boxplot in Figure 1d.

#### 2.1.2. CD68 positive decidual macrophages in the decidua

CD68 positive macrophages were investigated in the placenta of healthy pregnancies, SM, and RM (Figure 2a-d). The number of CD68 positive macrophages was low in the decidua basalis of control specimens (Figure 2a). The macrophages were slightly increased in decidua basalis RM samples, but without statistical significance ( $p=0.181$ ) (Figure 2b; median number of macrophages = 21 versus 16). Decidual macrophages were significantly increased in the decidua basalis SM group ( $p=0.013$ ) (Figure 2c; median number of macrophages 32 versus 16). A summary of the staining results is shown in Figure 2d.

### 2.2. Double Immunofluorescence


#### 2.2.1. Identification of PPAR $\gamma$ -expressing cells in the decidua basalis

Decidua basalis tissue of regular first trimester pregnancies, spontaneous and recurrent miscarriages was double stained using antibodies against PPAR $\gamma$  (green staining), and CD68 (red staining). Nuclear staining appeared in blue. PPAR $\gamma$  + CD68 double immunofluorescence staining was performed to investigate the macrophage expression of PPAR $\gamma$  in RM, SM and control groups. PPAR $\gamma$  staining in the cytoplasm of normal decidual cells is shown in Figure 3a. Figure 3b presents the cytoplasmic staining of PPAR $\gamma$  positive cells from the same area. The depiction of CD68 and PPAR $\gamma$  is represented as a co-expression by triple filter excitation in Figure 3c. Both markers are ubiquitously expressed in the healthy placenta (Figure 3a-c). A large number of CD68 positive macrophages was observed in RM samples (Figure 3d), although with almost no PPAR $\gamma$  expression (Figure 3e). Triple filter excitation demonstrates (Figure 3f) a nearly absence of PPAR $\gamma$  in macrophages of recurrent miscarriages. A large population of CD68 positive macrophages (Figure 3g) and PPAR $\gamma$  expressing cells (Figure 3h) was detected in SM samples. Figure 3i shows a strong co-expression of both markers. We identified CD68 positive macrophages also expressing PPAR $\gamma$  in the healthy and in the spontaneous miscarriage placenta. In the group of recurrent miscarriages only very few PPAR $\gamma$  expressing macrophages were detected.


### 2.3. Evaluation of PPAR $\gamma$ -expression with Real-Time RT-PCR (TaqMan)

PPAR $\gamma$  mRNA (PPARG) expression was analyzed in placental tissue from spontaneous, recurrent abortions and healthy controls by quantitative RT-PCR. PPAR $\gamma$  was significantly downregulated in SM (1.8-fold;  $p=0.010$ ) and in RM (1.5-fold;  $p=0.004$ ) compared to the control group (Figure 4).


### 2.4. Figures, Tables and Schemes




**Figure 1.** Immunohistochemical staining of PPAR $\gamma$  in the villous trophoblast. PPAR $\gamma$  expression is found with high distribution and intensity in intermediate villous trophoblastic cells (IVT) of first-trimester placentas. The control group showed the strongest expression pattern (a). PPAR $\gamma$  was significantly downregulated ( $p = 0.01$ ) in trophoblastic tissue of recurrent miscarriage (b). PPAR $\gamma$  expression in the IVT of spontaneous miscarriage tissue (c) was significantly decreased compared to the control ( $p = 0.001$ ). The boxplot summarizes the statistical data of the immunohistochemical staining results (d). Scale is 200  $\mu$ m. The insert picture is 100  $\mu$ m scaled.



**Figure 2.** Immunohistochemical staining of decidual macrophages with CD68 as a marker for macrophage positivity. Decidual macrophages were increased in RM and SM samples compared to the control group (a). In recurrent miscarriage specimens, the decidual macrophages tended to be upregulated ( $p = 0.181$ ) (b). In spontaneous miscarriage samples, the population of macrophages was significantly higher compared to the control ( $p = 0.013$ ) (c). Summary of staining results of CD68 positive decidual macrophages (d). Scale is 200 µm. The insert picture is 100 µm scaled.



**Figure 3.** Double Immunofluorescence of CD68 & PPAR $\gamma$ . CD68 (red staining), PPAR $\gamma$  (green staining), nuclear staining (blue). CD68 (a) and PPAR $\gamma$  (b) are expressed in the decidua of healthy placenta, co-expression presented as triple filter yellow (c). High distribution of CD68 positive macrophages was found in recurrent miscarriages (d). PPAR $\gamma$  positive cells are shown in (E). Triple filter excitation showed an absence of PPAR $\gamma$  positive macrophages (f). Both markers (CD68, g & PPAR $\gamma$ , h) are co-expressed in the group of spontaneous abortions (i). All pictures are 40 x lens.



**Figure 4.** Results of PPAR $\gamma$  mRNA expression analysis with TaqMan RT-PCR from trophoblastic tissue. PPAR $\gamma$  mRNA expression was significantly downregulated in the miscarriage groups (spontaneous ( $p= 0.01$ ) and recurrent miscarriage ( $p= 0.004$ )) compared to the healthy controls. This bar graph shows the mean of relative PPARG expression, therefore the presentation of error bars is not appropriate.

**Table 1.** Amount of slides used for immunohistochemical staining for each gestational week.

Gestational age	Normal pregnancy	Spontaneous miscarriage	Recurrent miscarriage
4th week	0	0	1
7th week	2	1	3
8th week	4	5	5
9th week	2	2	4
10th week	3	3	0
11th week	0	3	2
12th week	3	1	1
13th week	1	0	0
	<b>n=15</b>	<b>n=15</b>	<b>n=16</b>

**Table 2.** Demographic and Clinical Characteristics of the Study population. Values are Mean  $\pm$  S.D

Characteristics*	Normal pregnancy n= 15	Spontaneous miscarriage n= 15	Recurrent miscarriage n= 16	P value (Kruskal Wallis Test)
Maternal age (years)	31.18 $\pm$ 8.06 (18.7-43.3)	37.8 $\pm$ 4.51 (29.2-43.2)	35.76 $\pm$ 4.8 (29.5- 46.9)	0.049
Gestational age (weeks)	9.53 $\pm$ 1.95 (7-13)	8.4 $\pm$ 1.89 (7-12)	9.3 $\pm$ 1.49 (4-12)	0.276
Gravidity	4 $\pm$ 1.8 (1-7)	1.6 $\pm$ 0.9 (1-4)	3.1 $\pm$ 1.1 (2-5)	0.001
Parity	2 $\pm$ 1.1 (0-4)	0.9 $\pm$ 0.8 (0-2)	0.3 $\pm$ 0.6 (0-2)	0.003

\*Mean, standard deviation, range

**Table 3.** Antibodies used for immunohistochemical characterization (a) and double immunofluorescence (b) of placental tissue samples

Antibody	Isotype	Clone	Dilution	Source
PPAR $\gamma$ <sup>a,b</sup>	rabbit IgG	polyclonal	1:500 in PBS <sup>a</sup> 1:500 in Dako <sup>b</sup>	Abcam Serotec, Cambridge; UK DAKO (S322); Carpenteira, CA; USA
CD 68 <sup>a,b</sup>	mouse IgG1	monoclonal	1:8000 in PBS <sup>a</sup> 1:8000 in Dako <sup>b</sup>	Sigma Aldrich (CL1346), St Louis; USA DAKO (S322); Carpenteira, CA; USA
Cy 3 <sup>b</sup>	goat IgG anti-mouse	polyclonal	1:500 <sup>b</sup>	Dianova, Hamburg; Germany
Cy 2 <sup>b</sup>	goat IgG anti-rabbit	polyclonal	1:100 <sup>b</sup>	Dianova, Hamburg; Germany

### 3. Discussion

Within this study we could show that PPAR $\gamma$  is downregulated in the intermediate villous trophoblast (IVT) in both, spontaneous and recurrent miscarriage placentas. The downregulation of PPAR $\gamma$  was confirmed by RT-PCR in both miscarriage pregnancy cases.

In addition, we showed that in recurrent miscarriages decidua basalis macrophages are nearly PPAR $\gamma$ -negative, whereas in normal controls and surprisingly also in spontaneous miscarriages decidua basalis macrophages are all PPAR $\gamma$ -positive.

PPAR $\gamma$  as nuclear receptor is already known to be essential for the maturation of alternatively activated M2 macrophages [21]. M2 macrophages and decidual macrophages have mainly immune regulatory and homeostatic properties [22]. These macrophages have little in common with pro-inflammatory M1 macrophages, which is in line with the role for decidual macrophages in establishing and sustaining fetal tolerance [23].

In addition, rosiglitazone as a selective peroxisome PPAR $\gamma$  agonist has been shown to induce an M2 macrophage polarization via activating the PPAR $\gamma$  pathway [24,25]. The activation of PPAR $\gamma$  suppresses gene transcription by interfering with signal transduction pathways, such as the NF- $\kappa$ B, STAT, and AP-1 pathways that are involved in pro-inflammatory immune responses [26-28]. It is a striking result of this study, that we could identify the loss of PPAR $\gamma$  in decidual macrophages of patients with recurrent miscarriages but not in patients with spontaneous miscarriages and of course in normal control placentas.

In spontaneous miscarriages, we identified a significant increase of decidual macrophages, although PPAR $\gamma$  positive. In former studies, we could show that these macrophages express FasL [20]. The expression of FasL on decidual macrophages had been already described before [29].

The role of the Fas/FasL system in the conditions of spontaneous abortion and pregnancy had been described for T cell apoptosis by decidual and trophoblast cells earlier [30,31]. Our group was able to describe an increased expression of FasL in decidual macrophages of spontaneous miscarriages [20]. Therefore we speculated that FasL expression by macrophages could be a part of an M2-like polarization. FasL-expressing macrophages could induce apoptosis to Fas-bearing activated T-cells reducing potentially harmful immune responses against the semi-allogenic embryo. We further assumed that macrophages might mediate the T-cell triggered trophoblast apoptosis highlighting an alternative way of inducing apoptosis in spontaneous miscarriage [15].

Placental growth is exponential in the first trimester of pregnancy and involves coordinated events in trophoblast and mesenchyme, one of these is the differentiation of progenitor cytotrophoblast cells into intermediate villous trophoblast cells (IVT). These IVT are programmed to either fuse with the syncytium, or are transferred to extravillous trophoblast cells [32]. We found a downregulation of PPAR $\gamma$  in the IVT compartment in both spontaneous and recurrent miscarriage on protein as well as on mRNA-level. Already decades ago, the natural binding partner of PPAR $\gamma$ ,

the RXR $\alpha$  was described in this trophoblast compartment: RAR and RXR, both types of receptors were present in the proliferative intermediate villous trophoblast [33]. Later, RXR $\alpha$  was found to play a crucial role in pregnancy and is a key regulator of apoptosis in trophoblasts of patients with recurrent miscarriages [9]. PPAR $\gamma$  on the other hand was also described to be dysregulated in different trophoblast compartments of the miscarriage placenta [34,35], although PPAR $\gamma$  expression in the IVT was never investigated before. Interestingly, Fournier et al. described that activation of PPAR $\gamma$  induces accumulation of lipids, villous trophoblast differentiation and inhibits trophoblast invasiveness[36]. In addition, the expression of PPAR $\gamma$  is downregulated by stimulation of trophoblast cells with either arachidonic acid or 15d-PGJ2 [37]. Because we identified a downregulation of PPAR $\gamma$  in the IVT of miscarriages and a missing PPAR $\gamma$  expression in macrophages of recurrent miscarriage cases, we might speculate that PPAR $\gamma$  ligands (e.g. prostaglandins) are released to a higher extent under these pathological circumstances.

## 4. Materials and Methods

### 4.1. Patient data

The Institutional Review Board of the Ludwig-Maximilian-University, Munich, (Number of approval: 337-06) approved this study. All women signed an informed consent allowing analysis of all clinical and laboratory data mentioned in this study. Placental tissue from spontaneous miscarriages (SM) (n=15) and recurrent miscarriages (RM) (n=16) at gestational weeks 4 to 13 was obtained at the Department of Obstetrics and Gynecology, LMU Munich. Placental tissue from legal terminations of healthy pregnancies (n=15) served as control group. The tissue was collected at a private practice clinic in Munich, Germany. The control group specimens were confirmed as healthy by a blinded independent pathologist. All placental material was acquired by dilatation and curettage, without any prior pharmaceutical induction. In cases of SM and RM, the operation was performed within 24 h after diagnosis. Instantly, after the uterine curettage, the obtained tissue was either frozen or formalin fixed for further analysis. All patients included had an inconspicuous family and medical history, which was obtained systematically. Patients with common disorders, autoimmune diseases, thrombophilia and microbiological infections (Bacteria and Chlamydia trachomatis) were excluded. Chromosomal abnormalities were ruled out by karyotype analysis in all samples, as described recently [11,38]. Table 1 summarizes the number of samples used for immunohistochemical staining for each gestational week. Table 2 shows the demographic and clinical characteristics of the study population.

### 4.2. Immunohistochemistry

Formalin fixed tissue slides were embedded in paraffin wax for immunohistochemistry. Samples were deparaffinized in xylol for 20 min and rinsed in 100% ethanol. Methanol/H<sub>2</sub>O<sub>2</sub> incubation for 20 min was performed to inhibit endogenous peroxidase reaction. Afterwards, the specimens were rehydrated in deescalating alcohol gradients, starting with 100% ethanol and ending with distilled water. The samples were cooked in a pressure pot, containing a sodium citrate buffer (pH 6.0), which consisted of 0.1 mM citric acid and 0.1 mM sodium citrate in distilled water. Subsequently, samples were washed in PBS twice and incubated with a blocking solution (reagent 1, ZytoChem Plus HRP Polymer System (Mouse/Rabbit), Zytomed Berlin; Germany) for 5 minutes. Incubation with the primary antibody was performed with each section for 16 hours at 4°C. All antibodies used are listed in Table 3. Following every subsequent step, samples were washed twice in PBS (pH 7.4). Blocking solutions, containing post block (reagent 2) for 20 min and HRP-Polymer (reagent 3) for 30 min, were applied. The chromogen-substrate staining was carried out using the Liquid DAB+ Substrate Chromogen System (Dako Scientific, Glostrup; Denmark), 1 min for CD68 and 2 min for PPAR $\gamma$ . The reaction was stopped by applying distilled water. Finally, tissue samples were counterstained with Hemalaun for 2 min and blued in tap water. Specimens were dehydrated in an ascending alcohol gradient and cover slipped with Eukitt® quick hardening mounting medium (Sigma Aldrich, St. Louis, Missouri; USA). Positive control (human colon tissue) as well as negative

control staining was carried out as described previously [10,37]. All slides were analyzed using the microscope Leitz Wetzlar (Germany; Type 307-148.001 514686). The immunoreactive score (IRS) was used for evaluation of the intensity and distribution pattern of antigen expression. This semi-quantitative score is calculated as follows: the optical staining intensity (grades: 0 = none, 1 = weak, 2 = moderate, 3 = strong staining) is multiplied by the total percentage of positively stained cells (0 = none, 1 ≤ 10%, 2 = 11% - 50%, 3 = 51% - 80% and 4 ≥ 81 % of the cells). This multiplication has a minimum of 0 and a maximum of 12. Analysis of all slides was performed independently by two experienced staff members. Total number of macrophages in a magnification field of 40x lens was calculated three times each in 3 different areas of the decidua basalis. The median number was calculated.

### 4.3. Immunofluorescence

#### 4.3.1. Evaluation of PPAR $\gamma$ expressing cells as macrophages

For the visualization of PPAR $\gamma$ -expressing cells in the trophoblast, tissue samples of spontaneous miscarriages, recurrent miscarriages, both first-trimester abortion placentas, and healthy controls (first trimester) were used. The antibodies used are shown in Table 3. Double immunofluorescence staining for PPAR $\gamma$  and CD68 as a specific macrophage marker was performed to identify expression patterns in the nucleus and the cytoplasm. Each specimen was incubated overnight at 4°C with polyclonal anti-PPAR $\gamma$  rabbit IgG and monoclonal anti-CD68 mouse IgG1. Polyclonal Cy-2 and polyclonal Cy-3 conjugated antibodies (Dianova, Hamburg; Germany) were used as secondary antibodies. Incubation was performed for 30 min at room temperature. Samples were fixed with Vectashield® mounting medium with DAPI (Vector Laboratories; Burlingame, CA; USA) and analyzed with the Axioskop fluorescent photomicroscope (Zeiss; Oberkochen; Germany). Images were taken with the AxioCam camera system (Zeiss CF20DXC).

### 4.4. Evaluation of PPAR $\gamma$ with Real Time RT-PCR (Taq Man)

#### 4.4.1. RNA Extraction from placental tissue

mRNA extraction was accomplished using the placental tissue of 5 women with spontaneous abortion, 6 women with recurrent miscarriage and 4 healthy controls from the 7th to the 12th week of gestation. RNA extraction using 10 mg tissue of each sample was accomplished with RNeasy® Lipid Tissue Mini Kit (Qiagen, Hilden; Germany) according to the manufacturer's protocol.

#### 4.4.2. Reverse Transcription

According to the protocol reverse transcription (RT) was carried out with the High Capacity cDNA Reverse Transcription Kit (Applied Biosystems™, Fisher Scientific Company; Canada) and placed in a mastercycler® gradient (Eppendorf, Hamburg; Germany). RT conditions were as follows: 10 min at 25°C, 2 h at 37°C, 5 min at 85°C and continued by a hold step at -20°C.

#### 4.4.3. Real-Time Reverse Transcription-PCR

After conversion of RNA to cDNA, PCR was performed on all samples individually. Real-Time Reverse Transcription PCRs were covered with optical caps in optical 96-well (Applied Biosystems™, Fisher Scientific Company; Canada) reaction microtiter plates. Each reaction was accomplished with a volume of 20 $\mu$ l, including 1 $\mu$ l cDNA, 8 $\mu$ l H<sub>2</sub>O (DEPC treated DI water; Sigma, Taufkirchen; Germany) and 10 $\mu$ l TaqMan® Fast Universal PCR Master Mix 2x (Applied Biosystems, Nr. 4367846; 50ml). The total contained 1 $\mu$ l TaqMan® Gene Expression Assay 20x (HS01115513\_m1 for PPAR $\gamma$ , Applied Biosystems). The temperature protocol was as follows: 20 s at 95°C, 40 cycles of amplification, denaturation for 3 s at 95°C and denaturation plus annealing process for 30 s at 60°C.

Processing the PCR assays was performed using the 7500 Fast Real-Time PCR System (Applied Biosystems), and quantification was accomplished by the  $2^{-\Delta\Delta CT}$  method using  $\beta$ -actin as housekeeping gene (Applied Biosystems, Hs\_9999903\_m1).

#### 4.5. Statistics

Analysis of the collection and statistical data was processed with the SPSS software version 24 (SPSS; Chicago; IL; USA) and Excel version 12.3.1 (Microsoft Windows 2016; Redmond, WA; USA). The Mann-Whitney U signed-rank test was used for the comparison of two independent groups. P-values <0.05 were considered to be statistically significant.

## 5. Conclusions

Mouse knockout models showed that PPAR $\gamma$  is essential for placentation. PPAR $\gamma$  depletion leads to fetal loss in early pregnancy due to the missing PPAR $\gamma$  expression and extended placental defects [39]. In addition, decidual M1-like macrophage polarization events are associated with PPAR $\gamma$  modulation strategies [14]. Therefore the PPAR $\gamma$  pathway is a new molecular target for future preventive strategies for the treatment of spontaneous and recurrent miscarriages.

**Author Contributions:** Conceptualization, T.M.K., T.K. and U.J.; Methodology, C.K.; Software, A.H.; Validation, A.V., C.K. and U.J.; Formal Analysis, C.K.; Investigation, E.R.; Resources, S.M.; Data Curation, E.S.; Writing-Original Draft Preparation, E.R.; Writing-Review & Editing, U.J.; Visualization, E.R.; Supervision, U.J.; Project Administration, T.K.; Funding Acquisition, S.M.

**Funding:** This research received no external funding.

**Acknowledgments:** We thank Simone Hofmann for technical support.

**Conflicts of Interest:** The authors declare no conflict of interest.

## Abbreviations

MDPI	Multidisciplinary Digital Publishing Institute
PPAR	Peroxisome-Proliferator-activated Receptor
SM	Spontaneous miscarriage
RM	Recurrent miscarriage

## References

1. Makrigiannakis, A.; Petsas, G.; Toth, B.; Relakis, K.; Jeschke, U. Recent advances in understanding immunology of reproductive failure. *Journal of reproductive immunology* **2011**, *90*, 96-104.
2. Knabl, J.; Vattai, A.; Huttenbrenner, R.; Hutter, S.; Karsten, M.; Jeschke, U. Rxralpha is upregulated in first trimester endometrial glands of spontaneous abortions unlike lxr and ppargamma. *European journal of histochemistry : EJH* **2016**, *60*, 2665.
3. Toth, B.; Jeschke, U.; Rogenhofer, N.; Scholz, C.; Wurfel, W.; Thaler, C.J.; Makrigiannakis, A. Recurrent miscarriage: Current concepts in diagnosis and treatment. *Journal of reproductive immunology* **2010**, *85*, 25-32.
4. Arck, P.C.; Rucke, M.; Rose, M.; Szekeres-Bartho, J.; Douglas, A.J.; Pritsch, M.; Blois, S.M.; Pincus, M.K.; Barenstrauch, N.; Dudenhausen, J.W., *et al.* Early risk factors for miscarriage: A prospective cohort study in pregnant women. *Reproductive biomedicine online* **2008**, *17*, 101-113.
5. Pestka, A.; Fitzgerald, J.S.; Toth, B.; Markert, U.R.; Jeschke, U. Nuclear hormone receptors and female reproduction. *Current molecular medicine* **2013**, *13*, 1066-1078.
6. Guibourdenche, J.; Alsat, E.; Soncin, F.; Rochette-Egly, C.; Evain-Brion, D. Retinoid receptors expression in human term placenta: Involvement of rxr alpha in retinoid induced-hcg secretion. *J Clin Endocrinol Metab* **1998**, *83*, 1384-1387.

7. Toth, B.; Hornung, D.; Scholz, C.; Djalali, S.; Friese, K.; Jeschke, U. Peroxisome proliferator-activated receptors: New players in the field of reproduction. *American journal of reproductive immunology* **2007**, *58*, 289-310.
8. Arck, P.; Toth, B.; Pestka, A.; Jeschke, U. Nuclear receptors of the peroxisome proliferator-activated receptor (ppar) family in gestational diabetes: From animal models to clinical trials. *Biology of reproduction* **2010**, *83*, 168-176.
9. Pestka, A.; Toth, B.; Kuhn, C.; Hofmann, S.; Wiest, I.; Wypior, G.; Friese, K.; Jeschke, U. Retinoid x receptor alpha and retinoids are key regulators in apoptosis of trophoblasts of patients with recurrent miscarriages. *Journal of molecular endocrinology* **2011**, *47*, 145-156.
10. Knabl, J.; Pestka, A.; Huttenbrenner, R.; Plosch, T.; Ensenaer, R.; Welbergen, L.; Hutter, S.; Gunthner-Biller, M.; Jeschke, U. The liver x receptor in correlation with other nuclear receptors in spontaneous and recurrent abortions. *PPAR research* **2013**, *2013*, 575604.
11. Vattai, A.; Ziegelmuller, B.; Kost, B.; Kuhn, C.; Hofmann, S.; Bayer, B.; Anslinger, K.; Jeschke, U.; Ditsch, N. The expression of thyroid hormone receptors (thr) is regulated by the progesterone receptor system in first trimester placental tissue and in bewo cells in vitro. *European journal of obstetrics, gynecology, and reproductive biology* **2015**, *195*, 31-39.
12. Chen, Y.; Moradin, A.; Schlessinger, D.; Nagaraja, R. Rxralpha and lxr activate two promoters in placenta- and tumor-specific expression of plac1. *Placenta* **2011**, *32*, 877-884.
13. Hutter, S.; Knabl, J.; Andergassen, U.; Jeschke, U. The role of ppars in placental immunology: A systematic review of the literature. *PPAR research* **2013**, *2013*, 970276.
14. Xu, Y.; Romero, R.; Miller, D.; Kadam, L.; Mial, T.N.; Plazyo, O.; Garcia-Flores, V.; Hassan, S.S.; Xu, Z.; Tarca, A.L., *et al.* An m1-like macrophage polarization in decidual tissue during spontaneous preterm labor that is attenuated by rosiglitazone treatment. *Journal of immunology* **2016**, *196*, 2476-2491.
15. Hutter, S.; Heublein, S.; Knabl, J.; Andergassen, U.; Vrekoussis, T.; Makrigiannakis, A.; Friese, K.; Mayr, D.; Jeschke, U. Macrophages: Are they involved in endometriosis, abortion and preeclampsia and how? *Journal of Nippon Medical School = Nippon Ika Daigaku zasshi* **2013**, *80*, 97-103.
16. Slukvin, II; Breburda, E.E.; Golos, T.G. Dynamic changes in primate endometrial leukocyte populations: Differential distribution of macrophages and natural killer cells at the rhesus monkey implantation site and in early pregnancy. *Placenta* **2004**, *25*, 297-307.
17. Heublein, S.; Vrekoussis, T.; Kuhn, C.; Friese, K.; Makrigiannakis, A.; Mayr, D.; Lenhard, M.; Jeschke, U. Inducers of g-protein coupled estrogen receptor (gper) in endometriosis: Potential implications for macrophages and follicle maturation. *J Reprod Immunol* **2013**, *97*, 95-103.
18. Stremmel, C.; Schuchert, R.; Wagner, F.; Thaler, R.; Weinberger, T.; Pick, R.; Mass, E.; Ishikawa-Ankerhold, H.C.; Margraf, A.; Hutter, S., *et al.* Yolk sac macrophage progenitors traffic to the embryo during defined stages of development. *Nature communications* **2018**, *9*, 75.
19. Petsas, G.; Jeschke, U.; Richter, D.U.; Minas, V.; Hammer, A.; Kalantaridou, S.; Toth, B.; Tsatsanis, C.; Friese, K.; Makrigiannakis, A. Aberrant expression of corticotropin-releasing hormone in pre-eclampsia induces expression of fasl in maternal macrophages and extravillous trophoblast apoptosis. *Molecular human reproduction* **2012**, *18*, 535-545.

20. Guenther, S.; Vrekoussis, T.; Heublein, S.; Bayer, B.; Anz, D.; Knabl, J.; Navrozoglou, I.; Dian, D.; Friese, K.; Makrigiannakis, A., *et al.* Decidual macrophages are significantly increased in spontaneous miscarriages and over-express fasl: A potential role for macrophages in trophoblast apoptosis. *International journal of molecular sciences* **2012**, *13*, 9069-9080.
21. Tontonoz, P.; Spiegelman, B.M. Fat and beyond: The diverse biology of ppargamma. *Annu Rev Biochem* **2008**, *77*, 289-312.
22. Gustafsson, C.; Mjosberg, J.; Matussek, A.; Geffers, R.; Matthiesen, L.; Berg, G.; Sharma, S.; Buer, J.; Ernerudh, J. Gene expression profiling of human decidual macrophages: Evidence for immunosuppressive phenotype. *PloS one* **2008**, *3*, e2078.
23. Svensson, J.; Jenmalm, M.C.; Matussek, A.; Geffers, R.; Berg, G.; Ernerudh, J. Macrophages at the fetal-maternal interface express markers of alternative activation and are induced by m-csf and il-10. *Journal of immunology* **2011**, *187*, 3671-3682.
24. Lehmann, J.M.; Moore, L.B.; Smith-Oliver, T.A.; Wilkison, W.O.; Willson, T.M.; Kliewer, S.A. An antidiabetic thiazolidinedione is a high affinity ligand for peroxisome proliferator-activated receptor gamma (ppar gamma). *The Journal of biological chemistry* **1995**, *270*, 12953-12956.
25. Bouhleb, M.A.; Derudas, B.; Rigamonti, E.; Dievart, R.; Brozek, J.; Haulon, S.; Zawadzki, C.; Jude, B.; Torpier, G.; Marx, N., *et al.* Ppargamma activation primes human monocytes into alternative m2 macrophages with anti-inflammatory properties. *Cell Metab* **2007**, *6*, 137-143.
26. Lavinsky, R.M.; Jepsen, K.; Heinzl, T.; Torchia, J.; Mullen, T.M.; Schiff, R.; Del-Rio, A.L.; Ricote, M.; Ngo, S.; Gemsch, J., *et al.* Diverse signaling pathways modulate nuclear receptor recruitment of n-cor and smrt complexes. *Proceedings of the National Academy of Sciences of the United States of America* **1998**, *95*, 2920-2925.
27. Ricote, M.; Huang, J.; Fajas, L.; Li, A.; Welch, J.; Najib, J.; Witztum, J.L.; Auwerx, J.; Palinski, W.; Glass, C.K. Expression of the peroxisome proliferator-activated receptor gamma (ppargamma) in human atherosclerosis and regulation in macrophages by colony stimulating factors and oxidized low density lipoprotein. *Proceedings of the National Academy of Sciences of the United States of America* **1998**, *95*, 7614-7619.
28. Ricote, M.; Li, A.C.; Willson, T.M.; Kelly, C.J.; Glass, C.K. The peroxisome proliferator-activated receptor-gamma is a negative regulator of macrophage activation. *Nature* **1998**, *391*, 79-82.
29. Wongweragiat, S.; Searle, R.F.; Bulmer, J.N. Expression of fas/fas ligand by decidual leukocytes in hydatidiform mole. *Biology of reproduction* **2001**, *64*, 784-789.
30. Hammer, A.; Blaschitz, A.; Daxbock, C.; Walcher, W.; Dohr, G. Fas and fas-ligand are expressed in the uteroplacental unit of first-trimester pregnancy. *American journal of reproductive immunology* **1999**, *41*, 41-51.
31. Hammer, A.; Dohr, G. Expression of fas-ligand in first trimester and term human placental villi. *Journal of reproductive immunology* **2000**, *46*, 83-90.
32. Aplin, J.D. Developmental cell biology of human villous trophoblast: Current research problems. *Int J Dev Biol* **2010**, *54*, 323-329.
33. Tarrade, A.; Rochette-Egly, C.; Guibourdenche, J.; Evain-Brion, D. The expression of nuclear retinoid receptors in human implantation. *Placenta* **2000**, *21*, 703-710.
34. Toth, B.; Bastug, M.; Mylonas, I.; Scholz, C.; Makovitzky, J.; Kunze, S.; Thaler, C.; Friese, K.; Jeschke, U. Peroxisome proliferator-activated receptor-gamma in normal human pregnancy and miscarriage. *Acta histochemica* **2009**, *111*, 372-378.
35. Toth, B.; Bastug, M.; Scholz, C.; Arck, P.; Schulze, S.; Kunze, S.; Friese, K.; Jeschke, U. Leptin and peroxisome proliferator-activated receptors: Impact on normal and

- disturbed first trimester human pregnancy. *Histology and histopathology* **2008**, *23*, 1465-1475.
36. Fournier, T.; Guibourdenche, J.; Handschuh, K.; Tsatsaris, V.; Rauwel, B.; Davrinche, C.; Evain-Brion, D. Ppargamma and human trophoblast differentiation. *Journal of reproductive immunology* **2011**, *90*, 41-49.
37. Knabl, J.; Huttenbrenner, R.; Hutter, S.; Gunthner-Biller, M.; Vrekoussis, T.; Karl, K.; Friese, K.; Kainer, F.; Jeschke, U. Peroxisome proliferator-activated receptor-gamma (ppargamma) is down regulated in trophoblast cells of gestational diabetes mellitus (gdm) and in trophoblast tumour cells bewo in vitro after stimulation with ppargamma agonists. *Journal of perinatal medicine* **2014**, *42*, 179-187.
38. Ziegelmueller, B.; Vattai, A.; Kost, B.; Kuhn, C.; Hofmann, S.; Bayer, B.; Toth, B.; Jeschke, U.; Ditsch, N. Expression of thyroid hormone receptors in villous trophoblasts and decidual tissue at protein and mrna levels is downregulated in spontaneous and recurrent miscarriages. *The journal of histochemistry and cytochemistry : official journal of the Histochemistry Society* **2015**, *63*, 511-523.
39. Barak, Y.; Nelson, M.C.; Ong, E.S.; Jones, Y.Z.; Ruiz-Lozano, P.; Chien, K.R.; Koder, A.; Evans, R.M. Ppar gamma is required for placental, cardiac, and adipose tissue development. *Mol Cell* **1999**, *4*, 585-595.