

Article

The Effects of Birth Year, Age and Sex on Hemagglutination Inhibition Antibody Responses to Influenza Vaccination

Ewan P Plant ^{1*}, Angelia A Eick-Cost ², Hussein Ezzeldin ³, Jose L Sanchez ⁴, Zhiping Ye ⁵ and Michael J Cooper ⁶

¹ Division of Viral Products, Center for Biologics Evaluation and Research, US Food and Drug Administration, Silver Spring, MD., USA; ewan.plant@fda.hhs.gov

² Armed Forces Health Surveillance Branch, Public Health Division, J3/CSA, Defense Health Agency; and Cherokee Nation Technology Solutions, Silver Spring, MD., USA; angelia.a.cost.ctr@mail.mil

³ Office of Biostatistics and Epidemiology, Center for Biologics Evaluation and Research, US Food and Drug Administration, Silver Spring, MD., USA; husein.ezzeldin@fda.hhs.gov

⁴ Armed Forces Health Surveillance Branch, Public Health Division, J3/CSA, Defense Health Agency, Silver Spring, MD., USA; jose.l.sanchez76.civ@mail.mil

⁵ Division of Viral Products, Center for Biologics Evaluation and Research, US Food and Drug Administration, Silver Spring, MD., USA; zhiping.ye@fda.hhs.gov

⁶ Armed Forces Health Surveillance Branch, Public Health Division, J3/CSA, Defense Health Agency, Silver Spring, MD., USA; current address: National Institutes of Health, Rockville, MD., USA; michael.cooper3@nih.gov

* Correspondence: ewan.plant@fda.hhs.gov; Tel.: +1-240-402-7319

Abstract: The first exposure to influenza is thought to impact subsequent immune responses later in life. The consequences of this can be seen during influenza epidemics and pandemics with differences in morbidity and mortality for different birth cohorts. With the development of new universal vaccines there is need for better understanding of how vaccine responses are affected by early exposures to influenza viruses. In this analysis of hemagglutination inhibition (HI) antibody responses in two cohorts of military personnel we noticed differences related to birth year. These data suggest that HI antibody production, in response to influenza vaccination, is affected by influenza activity in the years following birth. The magnitude of this antibody response is associated with, among other factors, the influenza strain that circulated following birth.

Keywords: influenza; serum; IgG; humoral antibody; original antigenic sin; hemagglutinin

1. Introduction

Inactivated influenza vaccines have been widely shown to reduce the morbidity and mortality associated with influenza. The level of hemagglutination inhibition (HI) antibodies correlated with protection has been established in several studies [1-4]. However, HA antibody levels to specific viruses don't afford broad protection and have been shown to wane after vaccination [5]. Regular seasonal vaccination is known to boost HI antibodies to the vaccine strains but the observed increase is reduced with additional vaccinations and age [6-9]. In general, the time taken for these antibodies to decrease to levels not associated with protection has not been well studied. Some studies have indicated the benefits of vaccination can extend into the following season [5, 10-14] while another study indicated that such antibody levels drop below accepted protective levels within a year [15]. Measurements of HI titers 180 days post-vaccination suggest that a significant portion of vaccinated subjects retain protective titers up to this point [16], but it is known that a reduction in vaccine effectiveness occurs later in the season, or in between seasons [17-20]. This has implications for those travelling between the northern and southern hemispheres, military personal posted to different

regions of the globe, and at risk people vaccinated early during a season when influenza virus infections emerge late in the season.

Some studies, with sex-stratified data have found that women develop higher HI antibody levels toward some influenza vaccine antigens following vaccination [21]. Higher titers have been reported for women in subjects that seroconverted after the emergence of the 2009 pandemic H1N1 strain [15]. However, after six months the antibody levels were similar between the male and female populations. In a previous analysis using Department of Defense Serum Repository (DoDSR) samples we observed that the antibody titers between two weeks and six months after vaccination were often lower for women [22]. It was not clear if this was due to unmeasured confounders related to the cohort or to actual sex differences in antibody waning.

It has been reported that antibody titers toward the first influenza strains encountered early in life are boosted by subsequent infections with antigenically drifted viruses (reviewed in [23]). In contrast to the anamnestic response to a previously encountered strain, serum from the secondary response to an antigenically drifted strain is more cross-reactive [24]. There is evidence that infection, or vaccination with a new strain, provides a significant boost to pre-existing antibodies levels [22, 25, 26]. It is thought that the antigenic distance between the strains also affects the generation of the antibody response [27]. The impact that circulating influenza viruses during early childhood have on subsequent risk of infection and antibody responses are well documented [23, 28, 29], but the impact of prior immunity to circulating influenza viruses on vaccine response is not known; and this is a critical question that needs to be investigated for the development of universal vaccines [30].

The initial focus of this study was to investigate antibody waning in men and women after influenza vaccination. Antibody titers, 2 to 6 weeks post-vaccination, towards the A(H1N1), A(H3N2), and B influenza virus strains included in the vaccine, were assessed by HI assay. Titers from a second blood draw taken during the following year were also measured. Analyses of factors that might affect the antibody response, such as prior vaccination or infection, year of birth, preparation for deployment, and length of time in military service, were assessed for impact upon HI titer. The results led us to hypothesize that influenza viruses encountered in the first years of life affect vaccine responses later in life in a strain specific manner.

2. Materials and Methods

The study population consisted of active component service members who received an inactivated trivalent influenza vaccine in 2003, 2010, or 2011. The vaccination was required to occur between September 1st and December 15th of the year of interest, and the individuals could only receive one influenza vaccination per year. To reduce potential confounders, service members were excluded if their length of time in the military was greater than 10 years before the post-vaccination sample, or if they had a medical encounter for an influenza-like illness (ILI) or pneumonia and influenza (PI) for the time period of the study.

Serum samples were obtained from the Department of Defense Serum Repository (DoDSR), maintained by the Armed Forces Health Surveillance Branch of the Defense Health Agency, in accordance with the relevant guidelines and regulations. The first serum collection window was two to six weeks after receipt of the influenza vaccination. The second serum sample was collected during the year following the first sample. Three different bins were used to define the window for the second serum collection; 20% 3-4 months post-vaccination, 30% 5-7 months post-vaccination, 40% 8-10 months post-vaccination, and 10% 11-12 months post-vaccination. The rationale for selecting more samples 5-10 months post-vaccination was based on publications indicating that protective HI titers were maintained up to 6 months before dropping to non-protective levels [15, 16].

The WHO recommendations for influenza vaccine formulation for the Northern hemisphere from 1990 forward were analyzed. Consecutive seasons with no strain changes were selected for additional inspection. This allowed us to expand the possible pool of available subjects and to reduce confounding effects associated with a specific season. Two groups, each comprised of 80 female and 80 male subjects, were selected. The first group was vaccinated in 2003; the second group

was vaccinated in 2010 or 2011 (see Table 1). The female and male groups were matched, where possible, with regard to the timing of the blood draws.

Hemagglutination Inhibition (HI) assays were performed using standard methods. Each volume of serum was treated with 3 volumes of receptor destroying enzyme (Accurate Chemical, NJ) at 37°C overnight, which was then inactivated by heating in a 56°C water bath for 30 minutes. Six volumes of PBS were added to make the final sera concentration of 1:10. After RDE treatment, 50 microliters of each diluted serum was added to the first well of a 96-well plate. 25 microliters of PBS was added to the next 11 wells and serial 1:2 dilutions of the sera were made using robotic diluters. 25 microliters of virus titrated to a final HA titer of 4 was added to the first 11 wells and the plates incubated at room temperature for 30 minutes. The last well was left as a no-sera control. Freshly diluted turkey red blood cells (0.5%) for H1N1 and B antigens, or guinea pig red blood cells (1.0%) for H3N2 antigens, were added to all wells and the plates incubated for an additional hour. After incubation, the last dilution of sera that completely inhibited agglutination was recorded.

Anti-human IgG Quantitation (AHQ) dip and read biosensors were used with a Pall ForteBio Octet QKe system to measure the relative quantity of IgG in the sera samples. A standard curve was made using dilutions of a sera sample that gave a high read and the samples tested were measured relative to that. Samples were measured in duplicate and averaged.

Geometric mean titers (GMT) were calculated from the log values of the HI titers. Univariate analyses were performed using GraphPad Prism 6. Unless noted otherwise, the data for age, time of sample collection, HI titers and time in service were rank ordered and the Pearson correlation coefficients were calculated in excel for correlation analyses. The R package nlme[31] was used to conduct stepwise multivariate regression analyses to explore the independent effects of sex, age, years in service, deployment status, prior vaccination, and race on the post-vaccination response. A linear-mixed model (LMM) was used to study these independent effects on the post-vaccination response to the H1N1, H3N2, and B antigens for the 2003 and 2011 cohorts. The LMM (fit using restricted maximum likelihood) included random effect for sex, years of service, deployment status and hierarchical effects for age within years of service.

3. Results

Study Population

The study subjects are from a highly vaccinated active component service member population with near complete capture of medical encounters. Serum samples from military personnel are routinely collected upon entry to military service and at other times such as before and after deployment. Subjects identified for the study had a blood draw 2 to 6 weeks after seasonal influenza vaccination (mean of 31.7 days with a standard deviation of +/- 6.2 days), and a second blood draw later that year. Because the ratio of men to women is higher in the military, samples from women were identified first, followed by samples from men matched for timing of the second blood draw and age. The age of the subjects ranged from 18 to 41 at the time of vaccination with an average of 26.7 years (standard deviation of +/- 5.5). The race was predominantly White (53%), followed by Black (25%), Hispanic (12%), Asian/Pacific Islander (3%), Other (3%), American Indian/Alaskan Native (2%), and Unknown (2%). Two cohorts of subjects vaccinated during different influenza seasons were identified to maximize the available sera and to reduce possible confounders associated with a particular season. The first cohort was vaccinated in 2003 and all but six members of the second cohort were vaccinated in 2011. These members included three men and three women; each vaccinated in 2010. The influenza vaccine strains did not change from 2010 to 2011. The years each cohort was vaccinated, and the strains included in the vaccines, are presented in Table 1.

Table 1. Experimental Cohorts, Virus Vaccine Strains and Years Vaccinated. See Materials and Methods section for more details.

Cohort	Vaccine Strains	Years Vaccinated
--------	-----------------	------------------

2002/03	A/New Caledonia/20/1999 (H1N1), A/Moscow/10/1999 (H3N2), B/Hong Kong/330/2001 (Victoria)	2003
2010/11	A/California/7/2009 (pdmH1N1), A/Perth/16/2009 (H3N2), B/Brisbane/60/2008 (Victoria)	2010 or 2011

Comparison of HI Titers for Males and Females

Geometric mean titers (GMTs) were compared between men and women for samples drawn 2 to 6 weeks post-vaccination. There was a trend of lower GMTs for vaccine antigens among women, with the exception of the H3N2 antigen in the 2011 cohort (Tables 2 and 3). The difference in post-vaccination titers did not reach statistical significance (Mann Whitney rank test).

Table 2. HI Data for the 2002/03 Cohort. The names of the antigens used in the HI assay are listed in the first column. The geometric mean titers immediately after vaccination (S1) and later in the year (S2) are shown for each antigen for each group (Men or Women). The number of samples for each group of 80 that had a decrease in titer is shown in the last column.

Antigen	Group	GMT (S1)	GMT (S2)	Number of Samples with Decline
A/New Caledonia/20/1999 (H1N1)	Men	66	66	17
	Women	63	53	30
A/Moscow/10/1999 (H3N2)	Men	41	52	22
	Women	38	51	20
B/Hong Kong/330/2001 (Victoria)	Men	33	33	21
	Women	32	25	24

Table 3. HI Data for the 2010/11 Cohort. The names of the antigens used in the HI assay are listed in the first column. The geometric mean titers immediately after vaccination (S1) and later in the year (S2) are shown for each antigen for each group (Men or Women). The number of samples for each group of 80 that had a decrease in titer is shown in the last column.

Antigen	Group	GMT (S1)	GMT (S2)	Number of Samples with Decline
A/California/07/2009 (pdmH1N1)	Men	130	79	38
	Women	111	55	50
A/Perth/16/2009 (H3N2)	Men	67	40	41
	Women	76	38	53
B/Brisbane/60/2008 (Victoria)	Men	41	31	32
	Women	40	33	29

Changes in titers between the first and second serum samples were also compared. There was a significant ($P < 0.0001$, Wilcoxon matched-pairs rank test) decrease in GMTs for the influenza A viruses among men and women in the 2011 cohort (Figure 1). The decrease was also significant for the influenza B virus for men ($P < 0.05$, Wilcoxon matched-pairs rank test) but not for women in the 2011 cohort. There were no significant decreases in titers for men in the 2003 cohort but there was a significant decrease in titers toward the B virus for women in the 2003 cohort ($P < 0.01$, Wilcoxon matched-pairs rank test) (Figure 1). There were increases in GMT for the H3N2 antigen for both men and women in the 2003 cohort, but these did not reach a level of statistical significance. Further inspection of the raw data revealed that many of the subjects had higher titers in the second blood

draw, suggesting an intervening infection or variability in the assay. The titer from the second blood draw was lower for more female than male subjects (Tables 2 and 3). Therefore the data were reanalyzed including only subjects with lower titers in the second blood draw. The trend for lower GMTs for women toward the vaccine antigens remained with the exception of the influenza B strains (data not shown).




Figure 1. The geometric mean titers for both sera samples from the (a) 2003 and (b) 2011 cohorts are shown. The S1 and S2 titers for each group and each antigen are linked by a line. The antigens tested with the 2003 cohort were A/New Caledonia/20/1999 (H1N1), A/Moscow/10/1999 (H3N2) and B/Hong Kong/330/2001 (Victoria lineage). The antigens tested with the 2011 cohort were A/California/07/2009 (pdmH1N1), A/Perth/16/2009 (H3N2) and B/Brisbane/60/2008 (Victoria lineage). The difference between male and female titers (S1 or S2) did not reach the level of significance (Mann Whitney test). The difference between the S1 and S2 titers for each group was evaluated using the Wilcoxon matched-pairs rank test and the p-values are indicated at the end of each line: n.s., not significant; *, $p \leq 0.05$; **, $p \leq 0.01$; and ****, $p \leq 0.0001$.

The antibody decline over time for each of the cohorts was also evaluated (Figure S1). The GMT values for the 3-4, 5-7, 8-10 and 11-12 month data points were determined for each cohort. The initial decline in HI titer for both of the influenza A viruses in the 2011 cohort subjects was greater than that for the influenza B antigen. This pattern was not observed for the 2003 cohort in which there was an increase in GMT values for the 3 to 4 month data points and less decline overall. This suggests differences in rate of decline depend on the antigen and the season. The pattern of decline was similar for men and women in the respective seasons with one exception. In the 2003 cohort, the titers 11 to 12 months after vaccination increased for men for all three antigens, but not for women. However, the number of subjects in these groups was small and the difference between men and women was not found to be statistically different (Mann Whitney rank test).

Age

It is known that HI titers post-vaccination negatively correlate with increasing age [6, 7], even within a young adult population [22]. There were negative correlations between age and HI titer for most antigens. The negative correlations were statistically significant (Spearman rank correlation coefficient in excel) for all antigens for men in the 2003 cohort (A/New Caledonia/20/1999, -0.541 , $p < 0.001$; A/Moscow/10/1999, -0.320 , $p < 0.01$; and B/Hong Kong/330/2001, -0.310 , $p < 0.01$) but only for the H1N1 antigen for women (A/New Caledonia/20/1999, -0.482 , $p < 0.001$). The negative correlations reached significance for only one antigen in the 2011 cohort: the A(H3N2) antigen for women (A/Perth/16/2009, -0.243 , $p < 0.05$). There were positive correlations between age and HI titer for the B antigen for both sexes in the 2011 cohort but it only reached significance for men (B/Brisbane/60/2008; 0.267 , $p < 0.01$). Analysis of each cohort using a linear-mixed model (LMM) revealed a negative correlation between age and HI titer for the influenza A antigens in the 2003 cohort with sex having minimal effect. In the 2011 cohort there was a positive correlation between age and HI titer for the B antigen with sex having a minimal effect.

Negative correlations between HI titer and age were also observed with the samples collected later in the year. In the 2003 cohort, significant negative correlations were observed for all three antigens for men (A/New Caledonia/20/1999, -0.423 , $p < 0.001$; A/Moscow/10/1999, -0.289 , $p < 0.016$; and B/Hong Kong/330/2001, -0.272 , $p < 0.05$) but only for the A(H1N1) antigen for women (A/New Caledonia/20/1999, -0.483 , $p < 0.001$). The negative correlation for the A(H1N1) antigen reached statistical significance for men in the 2011 cohort (A/California/07/2009, -0.376 , $p < 0.001$). The correlation between age and titer for the B antigen was positive for both men and women in the 2011 cohort (0.257 and 0.236 respectively, $p < 0.05$).

Prior Seasonal Vaccination

Medical records indicated that many of the subjects had received an influenza vaccination the year prior to the study. Approximately 40% of men and 35% of women in the 2003 cohort had received seasonal influenza vaccinations in 2002. In the 2011 cohort, 81% of men and 64% of women received seasonal vaccinations the prior season. Subjects in the 2011 cohort could have received different types of influenza vaccines. The CVX codes for prior season vaccination differentiated between Live Attenuated Influenza Vaccines (LAIV) and Inactivated Influenza Vaccines (IIV) allowing for a more refined analysis. Analyses on the data were performed to determine if there were any detectable effects of vaccination the prior season, by type, on HI titers.

Analysis of the 2003 cohort showed that the HI titers for the A(H1N1) antigen were slightly lower in subjects that had been vaccinated the prior year (Figure 2). This was observed for both men and women but was not found to be statistically significant (Mann Whitney test rank test).




Figure 2. The geometric mean titers (with 95% confidence interval) for previously unvaccinated subjects are compared to titers of those vaccinated the previous season. Results are shown for the (a) 2003 and (b) 2011 cohorts. The viruses used for testing HI titer are listed on the x axis: NC/20, A/New Caledonia/20/1999 (H1N1); MS/10, A/Moscow/10/1999 (H3N2); HK/330, B/Hong Kong/330/2001 (Victoria lineage); CA/7, A/California/07/2009 (pdmH1N1); PH/16, A/Perth/16/2009 (H3N2); and BR/60, B/Brisbane/60/2008 (Victoria lineage). The difference between the vaccinated and unvaccinated groups was evaluated using the Mann Whitney rank test. n.s., not significant; and *, $p < 0.05$.

Analysis of the 2011 cohort indicated that prior vaccination with IIV resulted in lower HI titers toward both the A(H1N1) and A(H3N2) components of the vaccine. This difference was not statistically significant (Mann Whitney rank test). In contrast, the titers for those vaccinated with LAIV the prior year were similar to, or higher than, the titers of those who were not vaccinated the prior year. The GMTs for the A(H1N1) strain was significantly higher than the titers for those vaccinated with an IIV ($P < 0.05$, Mann Whitney rank test). There was no difference in the influenza B titers regardless of prior seasonal vaccination status, reflecting the higher conservation of influenza B HA and, perhaps

a higher incidence of cross-reactive antibodies [32]. These results are in line with other studies indicating that individuals who have been vaccinated previously have lower titers than those only vaccinated in the current season, unless the prior vaccination was an LAIV [6, 7, 33-35].

Receipt of multiple annual influenza vaccinations is known to blunt the HI antibody response [6-8]. Our study protocol only included vaccination history for the prior year (see Methods section). Because the military has strict vaccination requirements we used the length of military service as a proxy for multiple vaccinations. Sera from men and women in the 2003 cohort who had been in military service for 8 to 9 years were compared to those who had been in military service for only 1 to 2 years (Figure 3). In this cohort, subjects with a shorter time in the service had higher titers toward all three antigens, the difference being significant for the A(H1N1) and A(H3N2) antigens ($P < 0.05$ and $P < 0.01$ respectively, Wilcoxon matched-pairs rank test).




Figure 3. The geometric mean titers (with 95% confidence interval) for sera from personnel with short and long service are shown. The data for (a), the 2003 cohort, are from 48 subjects with 1-2 years of service and 19 subjects with 8-9 years of service. The data for (b), the 2011 cohort, are from 63 subjects with 1-2 years of service and 23 subjects with 8-9 years of service. The viruses used for testing HI titer are listed on the x axis: NC/20, A/New Caledonia/20/1999 (H1N1); MS/10, A/Moscow/10/1999 (H3N2); HK/330, B/Hong Kong/330/2001 (Victoria lineage); CA/7, A/California/07/2009 (pdmH1N1); PH/16, A/Perth/16/2009 (H3N2); and BR/60, B/Brisbane/60/2008 (Victoria lineage). The difference between the short and long service groups was evaluated using the Mann Whitney rank test. n.s., not significant; *, $p < 0.05$; and **, $p < 0.01$.

Sera from men and women in the 2011 cohort who had been in military service for 8 to 9 years were compared to those who had been in military service for only 1 to 2 years (Figure 3). In this cohort, subjects with a shorter time in the service had higher titers toward all three antigens, the difference being significant for the A(H1N1) and A(H3N2) antigens ($P < 0.01$ and $P < 0.05$ respectively, Wilcoxon matched-pairs rank test). Analysis of each full cohort using LMM revealed a negative correlation between age and years in service for both cohorts. In the model the HI titers toward both influenza A antigens were explained by age for the 2003 cohort and years in service for the 2011 cohort. The impact of each variable was significant for the influenza B antigen in the 2011 cohort but neither variable had a significant effect on the 2003 cohort HI titers.

A subset of samples from the 2011 cohort was analyzed for HI titers toward the older antigens. Sera from 12 men and 12 women from the 2011 cohort that had been in military service 8 to 9 years were compared to sera from personnel, of a similar age, who had been in military service for only 1 to 3 years (Figure S2). Lower titers were observed for the subset with longer military service, but the differences did not reach statistical significance. In contrast, when the HI titers toward older

antigens were measured, the titers were higher for personnel that had served longer (Figure S2). This result is similar to other reports describing an increase in antibody titer toward previously encountered antigens after infection or vaccination [25, 26, 36]. Total IgG levels in the samples in the 2011 service subset were compared. The GMTs were similar for the shorter and longer military service groups. However, when the samples were stratified by sex, the IgG titers for women were found to be significantly lower than those of men (Figure S3). There was no correlation between IgG levels and age, or timing of the blood draw (Spearman test in prism).

Deployment

We analyzed deployment as a variable. Approximately 38% of the 320 subjects (60 men and 60 women) were deployed between the time of vaccination and the final blood draw (Figure 4). The GMT value for the first blood draw was significantly higher for the deployed personnel compared to the non-deployed group (Mann Whitney rank test). The A(H3N2) titers remained higher in deployed subjects when different sub-populations were analyzed. The difference was significant for the 2011 A(H3N2) antigen (A/Perth/16/2009), but not for the 2003 A(H3N2) antigen (A/Moscow/10/1999), when the 2003 and 2011 cohorts were analyzed separately. The difference remained significant for men, when the two sexes were analyzed separately, but not for women. There was no significant difference in the HI titers in samples collected later in the year between subjects who were or were not deployed. These results suggest that something associated with the preparation for deployment primes the immune system, and results in greater production of antibodies toward the A(H3N2) vaccine component in the 2 to 6 weeks following vaccination.




Figure 4. The geometric mean titers (with confidence interval) with deployment as a variable is shown. The viruses used for testing HI titer for 2003 cohort subjects were A/New Caledonia/20/1999 (H1N1), A/Moscow/10/1999 (H3N2), and B/Hong Kong/330/2001 (Victoria lineage). The viruses used for testing HI titer for 2011 cohort subjects were A/California/07/2009 (pdmH1N1), A/Perth/16/2009 (H3N2), and B/Brisbane/60/2008 (Victoria lineage). The difference between the two groups was evaluated using the Mann Whitney rank test. n.s., not significant; and **, $p < 0.01$.

Year of Birth

In general, titers decreased gradually with increasing military service (Figure S4). However, there were two notable exceptions. First, in the 2003 cohort, the GMT titers for the A(H1N1) antigen were noticeably higher for those with 1 to 4 years of military service compared to those with 5 to 8 years. This was not apparent for the A(H3N2) and B antigens. The birth years for the subjects in the 2003 cohort flanked the emergence of an A(H1N1) strain in 1977 after a 20-year absence. Second, the titer for the A(H1N1) antigen in the 2011 cohort were highest in those with two years of service. This suggests something prior to military service reduced the response to vaccination for those in the

first year of service, or something during the initial year of service, such as vaccination or exposure to the 2009 pdmH1N1 strain, increased responses for those already in military service. Interestingly, the titers for the A(H1N1) antigen increased from 3 to 5 years of service while the titers for the B antigen decreased over this time. This indicated that the trends were unique for each antigen.

It is hypothesized that the first exposure to an influenza virus influences the later response of the immune system. This effect is often referred to as antigenic sin [23, 28, 29]. The A(H1N1) viruses began circulating in 1977 after a 20-year absence. Approximately one half of the subjects in the 2003 cohort were born before 1977. The GMT toward the A(H1N1) vaccine component of those born before 1977 were compared to those born after 1977. The titers for those born after 1977 were significantly higher (Mann Whitney rank test, $p < 0.0001$) suggesting that exposure to an A(H1N1) virus early in life resulted in a more robust antibody production with A(H1N1) vaccination later in life (Figure S5).

Table 4. Influenza Vaccine Strain Recommendations from the WHO and Influenza Research Database websites (<http://www.who.int/influenza/vaccines/virus/recommendations/en/> and https://www.fludb.org/brc/vaccineRecommend.spg?decorator=influenza#1974-1975_Northern%20Hemisphere). Changes from the previous recommendation are shown in bold.

Year	H1N1	H3N2	B Virus
1974-75		A/Port Chalmers/1/73	B/Hong Kong/5/72
1975-76		A/Port Chalmers/1/73	B/Hong Kong/5/72
1976-77		A/Victoria/3/75	B/Hong Kong/5/72
1977-78		A/Victoria/3/75	B/Hong Kong/5/72
1978-79	A/USSR/90/77	A/Texas/1/77	B/Hong Kong/5/72
1979-80	A/Brazil/11/78	A/Texas/1/77	B/Hong Kong/5/72
1980-81	A/Brazil/11/78	A/Bangkok/1/79	B/Singapore/222/79
1981-82	A/Brazil/11/78	A/Bangkok/1/79	B/Singapore/222/79
1982-83	A/Brazil/11/78	A/Bangkok/1/79	B/Singapore/222/79
1983-84	A/Brazil/11/78	A/Philippines/2/82	B/Singapore/222/79
1984-85	A/Chile/1/83	A/Philippines/2/82	B/USSR/100/83
1985-86	A/Chile/1/83	A/Philippines/2/82	B/USSR/100/83
1986-87	A/Chile/1/83	A/Mississippi/1/85	B/Ann Arbor/1/86
1987-88	A/Taiwan/1/86	A/Leningrad/360/86	B/Ann Arbor/1/86
1988-89	A/Taiwan/1/86	A/Sichuan/2/87	B/Victoria/2/87
1989-90	A/Taiwan/1/86	A/Shanghai/11/87	B/Yamagata/16/88
1990-91	A/Taiwan/1/86	A/Shanghai/16/89	B/Yamagata/16/88
1991-92	A/Taiwan/1/86	A/Beijing/353/89	B/Panama/45/90
1992-93	A/Texas/36/91	A/Beijing/353/89	B/Panama/45/90

The effect of antigenicity of viruses circulating early in life on vaccine response was evaluated. Data was analyzed by the year the subjects were born and the viruses known to be circulating during the first years of life. The subjects were grouped by age, binning for each year of age from 19 to 29 separately for both the 2003 and 2011 cohorts. Each of these bins contained results for 6-13 subjects. The geometric mean titers were determined for each age group and graphed (Figure S6). We used information from the WHO vaccine strain recommendations, MMWR reports, and other publications [37-40] to identify the dominant circulating strains and severity of the influenza season for each year considered (Tables 4 and 5). The years in which the vaccine strains remained the same were used as an indicator that the circulating strains remained antigenically similar (vertical lines in Figure S6).

Table 5. The dominant circulating virus type for each northern hemisphere season ending in the years 1973 to 1993 is shown with references. An additional virus is listed if it contributed to more than 15% of the positive isolates for that season (data from Thompson et al., 2003).

Year	Dominant Virus Type	References
1973	H3N2	[37, 39, 40]
1974	H3N2	[37, 39, 40]
1975	H3N2	[37, 39, 40]
1976	H3N2	[37, 38, 40]
1977	B (+ 25% H3N2)	[37, 38, 40]
1978	H3N2 (+ 16% H1N1)	[38, 40]
1979	H1N1	[38, 40]
1980	B virus	[38, 40]
1981	H3N2 (+ 22% H1N1)	[40]
1982	B virus (+ 24% H1N1)	[40]; <u>MMWR 1982; 31(27): 375-9</u>
1983	H3N2	[40]; <u>MMWR 1983; 32(29): 373-7</u>
1984	H1N1 (+ 45%B virus)	[40]; <u>MMWR 1984; 33(29): 417-21</u>
1985	H3N2	[40]; <u>MMWR 1985; 34(28): 440-4</u>
1986	B Virus (+ 24% H2N3)	[40]; <u>MMWR 1986; 35(29): 475-9</u>
1987	H1N1	[40]; <u>MMWR 1988; 37(31): 466,468-70, 475</u>
1988	H3N2	[40]; <u>MMWR 1988; 37(32): 497-503</u>
1989	B Virus (+ 44% H1N1)	[40]; <u>MMWR 1989; 38(11): 183-5</u>
1990	H3N2	[40]; <u>MMWR 1990; 39(17): 293-6</u>
1991	B Virus	[40]; <u>MMWR 1991; 40(14): 231-3, 239-40</u>
1992	H3N2 (+ 18% H1N1)	[40]; <u>MMWR 1992; 41(18): 315-7, 323</u>
1993	B Virus	[40]; <u>MMWR 1992; 41(18): 315-7, 323</u>

The results from individuals born in seasons with antigenically similar viruses were grouped and GMTs calculated (Figure 5). Statistical analyses were performed to determine if the titers for each group were significantly different from the prior group (Mann Whitney rank tests). As expected, the GMT towards the A/New Caledonia/20/1999 (H1N1) antigen was lower for those born in 1977 or earlier than those born in 1978 in the 2003 cohort ($p < 0.05$, Figure 5). There was no significant difference in titer for those born when the A/USSR/90/1977 and A/Brazil/11/1978 viruses circulated. However, the GMT for those born when the A/Chile/1/1983 strain circulated was significantly higher than the GMT for those born when the A/Brazil/11/1978 virus circulated ($p < 0.001$). These data suggest that the antigenicity of influenza viruses circulating early in life affect vaccine response.




Figure 5. The geometric mean titers (with confidence interval) towards the vaccine antigens for the 2003 and 2011 cohorts based on antigens circulating after the year of birth. The difference between sequential groups, born when different viruses circulated, was evaluated using the Mann Whitney rank test. *, $p < 0.05$; **, $p < 0.01$; and ***, $p < 0.001$. The viruses used for HI titer are A/New Caledonia/20/1999 (H1N1), A/Moscow/10/1999 (H3N2), and B/Hong Kong/330/2001 (Victoria lineage) for the 2003 cohort. The viruses used for testing HI titer are A/California/07/2009 (pdmH1N1), A/Perth/16/2009 (H3N2), and B/Brisbane/60/2008 (Victoria lineage) for the 2011 cohort. The H1N1 strains from the birth year are USSR (A/USSR/90/1977), BZ/11/78 (A/Brazil/11/1978), CL/1/83 (A/Chile/1/1983), Tai/1/86 (A/Taiwan/1/1986) and TX/36/91 (A/Texas/36/1986). The H3N2 strains from the birth year are PC/1/73 (A/Port Charlmers/1/1973), VC/3/75 (A/Victoria/2/1975), TX/1/77 (A/Texas/1/1977), BK (A/Bangkok/1/1979), PL (A/Philippines/2/1982), MS (A/Mississippi/1/1985), LN (A/Leningrad/360/1986), SC (A/Sichuan/2/1987), 87 (A/Shanghai/11/1987), 89 (A/Shanghai/16/1989), and BJ (A/Beijing/353/1989). The B antigens from the birth year are HK/5/72 (B/Hong Kong/5/1972), SG/79 (B/Singapore/222/1979), 100/83 (B/USSR/100/1983), AA/86 (B/Ann Arbor/1/1986), VC/87 (B/Victoria/2/1987), YM/88 (B/Yamagata/16/1988) and PN/90 (B/Panama/45/1990).

There were other instances where the titer toward the influenza A vaccine antigens was significantly different after birth year was taken into consideration (that is, the antigenicity of the viruses circulating the year after birth changed) (Figure 5). The change in titer depended on the cohort being analyzed. Subjects from both the 2003 and 2011 cohorts were born during the period when the

A/Brazil/11/1978 or A/Chile1/1983 viruses circulated (but the subjects from the 2011 cohort were 8 years older when they were vaccinated than those in the 2003 cohort). While the titers toward the A/ New Caledonia/20/1999 (H1N1) vaccine antigen were significantly higher in the subjects born when the A/Chile/1/1983 strain circulated in the 2003 cohort, the titers toward the A/California/07/2009 (pdmH1N1) vaccine antigen were slightly lower, but not significantly different, in the 2011 cohort (Figure 5). A similar pattern was observed for the A(H3N2) antigens. In the 2003 cohort, the GMT toward the A/Moscow/10/1999 vaccine antigen was significantly higher for those born when the A/Philippines/2/1982 virus circulated compared to those born when the A/Bangkok/1/1979 virus circulated. Conversely, the GMT toward the A/Perth/16/2009 vaccine antigen was significantly higher for those born when the A/Bangkok/1/1979 virus circulated in the 2011 cohort. These results suggest a role for age and/or the vaccine antigen, in addition to the antigenicity of the virus circulating after birth.

HI data in the context of which type of virus circulated after the year of birth, and whether the season was mild or severe, was also evaluated. There were several seasons where a particular type of virus circulated (© symbol in Figure S6) and it was described as a severe influenza season (darker shading of bars in Figure S6). Many of these seasons correlated with higher titers after vaccination (for example: A(H1N1) viruses in 1978 and 1983, A(H3N2) viruses in 1980 and 1984, and B viruses in 1976 and 1979 for the 2003 cohort; A(H3N2) viruses in 1982, 1987, 1989 and 1991, and B viruses in 1985 and 1990 for the 2011 cohort). However, as mentioned above, this effect also depended on the vaccine antigen being tested: A(H3N2) titers were low in the 2003 cohort for subjects born in 1982 and high for those born in 1984, but this pattern was reversed in the 2011 cohort (Figure 5). This effect could also be due by the age of the subject at the time of vaccination (subjects born in 1984 were 19 years old in the 2003 cohort, or 27 years old in the 2011 cohort).

4. Discussion

This study contributes to existing knowledge about how age and sex affect the response to influenza vaccination. In addition, the analyses reveal correlations between the HI titer after vaccination and other variables. An increase in antibody response among personnel that were deployed after vaccination was observed. This suggests that something in the preparation for deployment better primed those personnel for vaccination. Subclinical infections prior to deployment are one possible explanation but, while MMWR reports indicate the H3N2 viruses circulated at that time [41, 42], there could be other factors, such as concomitantly administered vaccines, we have not considered. Analysis of the data presented here, in the context of other variables such as prior vaccination and year of birth, have led us to hypothesize that influenza infections early in life have an impact on vaccine response later in life.

Interest in the impact of sex on clinical trials has increased in recent years [43]. The effect sex has in the response towards influenza infection has been reviewed [44, 45], but fewer publications investigate the impact sex has on antibody response to influenza vaccination [21, 46-48]. There is also a dearth of information about the impact sex may have on the rate of HI antibody waning between men and women. In some instances, a stronger decline in antibody concentrations has been observed in women [49]. Here, as in our previous study [22], higher GMTs for men toward most of the vaccine antigens are reported. This trend did not reach the level of statistical significance and contrasts with some published research for influenza vaccinations [46, 50]. It may be due to some aspect of military service, as has been reported for other vaccines [51]. Alternatively, as demonstrated in the literature, the trends could be antigen or season specific. For example, Furman et al., did not observe a difference between the sexes by HI titer for the A(H1N1) antigen, but Hsu et al., observed a difference for the pdmH1N1 antigen [15, 52]. The blood draws in our previous study occurred up to six months after vaccination, prompting an assessment of the differences in antibody waning in this study. However, the data from this study, with blood draws between two and six weeks after vaccination, still indicated a trend for lower titers in young adult women. Additionally, there was a difference in the amount of total IgG between the sexes. A greater number of blood samples collected from women later in the year showed a decrease in HI titer than those collected

from men. These results leave questions about sex based differences in antibody levels, IgG concentrations and HI antibody waning after influenza vaccination unanswered. More research in this area is required to shed light on these issues.

A major limitation of this study design is the reliance on historic blood samples rather than designing a clinical trial with fixed sample collection times. This meant that antibody levels were not assessed at multiple time points from individuals throughout the year. Only two samples were collected from each subject and the timing of sample collection varied. Subjects were binned based on the second blood draw. This clustering may have hindered the detection of differences in the rate of antibody waning between men and women. This disadvantage was offset by the ability to include larger cohorts and compare different seasons. This study describes other effects, such as the effect of prior vaccination, without season-specific confounders. Another limitation of the study was the low number of subjects born in particular years which limited the statistical rigor of our analysis of year-of-birth effects. For the analysis birth cohorts born before and after the emergence of the A(H1N1) viruses in 1977, the p-value dropped from < 0.0001 to < 0.05 when the titer of single year birth cohorts were compared (Figure S5 and Figure 5). However, even with the limited number of subjects in each birth cohort, significant differences were observed supporting the hypothesis that birth year has an impact on vaccine response.

The effect birth year has on susceptibility to influenza disease, and for vaccine effectiveness, is of great interest [20, 23, 53]. In addition to factors related to an aging immune system, it appears that exposure to particular viral serotypes in the years immediately after birth also affects a person's response to vaccination [54]. In a recent study by Skowronski et al., they noted lower vaccine efficacy for the pdmH1N1 virus in the birth cohort for the years 1957 to 1967, that were likely primed with an A(H3N2) antigen rather than an A(H1N1) antigen [20]. Similarly, lower A(H1N1) antibody titers in subjects born prior to the reemergence of A(H1N1) viruses in the late 1970s were observed in this study. In another study, Skowronski et al., reported differences in the percentage of B/Victoria and B/Yamagata infections by year of birth [55]. When information about the viruses that circulated following the birth year of the subjects in this study was included, we observed differences in HI antibody titers that appear to be due to the antigenicity of the virus circulating in the first years of life, not just the presence or absence of a virus. These differences support the notion that the environment plays a significant role in priming and shaping the immune response [56] and the recommendation that antigenic sin be considered in the development of immunization strategies [23].

There is variability in the amplitude and direction of antibody response after repeat annual seasonal influenza vaccination, and the responses to the influenza B strains differ from the influenza A strains (reviewed in [57]). It has been reported that as pre-seasonal immunity toward influenza B viruses increases, vaccine effectiveness is less impacted by strain mismatches [58]. This is perhaps expected because of the generation of cross-reactive antibodies after influenza B infection [59]. In studies with repeatedly vaccinated individuals it is not clear if the lower subsequent HI titers correlate with vaccine efficacy because other components of the immune response are not measured [32]. This may depend on antigenic distance and the relative abundance of cross-reactive antibodies. In our current analyses, a decrease in HI titers for the pdmH1N1 component in subjects with prior vaccination was observed. Skowronski et al., reported lower vaccine efficacy for pdmH1N1 viruses for subjects that had received both current and prior season vaccination compared to those with just the current season vaccination [20]. This would suggest a correlation between HI titers, prior vaccination, and vaccine efficacy. Skowronski et al., did not observe a difference in efficacy against influenza B for those previously vaccinated and we did not observe a difference in titers for the influenza B component between those with or without prior vaccination. The differences between influenza A and B strains suggest that the HI antibody landscape develops differently for influenza A and B strains and that this difference may affect vaccine efficacy.

Responses to the same influenza vaccine antigen can vary from season to season. The responses to vaccination for a predominantly pediatric population were recorded for 10 years from 1982 until 1992 [60]. In that study, there was variation in vaccine response each year, with a significant increase

in influenza B antibodies over time, even when the vaccine antigen didn't change. Interestingly, the yearly variation in GMTs for the B component was greater than for the A(H1N1) component, which was greater than for the A(H3N2) component. More peaks and troughs in response to vaccination were observed for the B, H1N1 and H3N2 antigens respectively, for the 2011 cohort subjects born in those years in this study (Figure S6). Most striking, there are peaks for the B component in 1984-86 and 1990-91 in both studies, during seasons that were described as dominantly B seasons. These data suggest that subclinical infections not only boost vaccination responses in children in a contemporary setting, but may also contribute to vaccine responses in a similar manner later in life.

The shaping of the influenza antibody landscape will vary from individual to individual and there are many factors that will contribute to it. High levels of HI antibodies toward the A/California/7/2009 (pdmH1N1) strain in DoDSR serum collected from military service members in 2010, prior to vaccination, were reported indicating high rates of subclinical infection [22]. And it has been reported elsewhere that subclinical infections may influence the response to vaccination [61, 62]. From the data presented here and elsewhere [35], it is apparent that prior LAIV vaccination can also affect antibody titers after subsequent IIV vaccination. These data support the hypothesis that infections, including subclinical infections, early in life may also play a role in vaccine responses later in life. Recent prior vaccination with an IIV leads to lower vaccine response as measured by HI assay. Year of birth plays a significant role. While others have described differences in vaccine efficacy or mortality [20, 54], this study found differences in antibody generation after vaccination. Titers differed not only between birth cohorts, but also within the same birth cohorts receiving vaccinations at different ages. Differences in the intensity of prior influenza seasons people have been exposed to also appears to have an impact on later vaccine responses. It will be important to determine how these factors will affect responses in universal vaccines that are being developed [30]. Current influenza vaccines are designed to increase antibody titers toward contemporary strains but there is still variability in outcomes depending on many factors including age and prior vaccination status. Better understanding the effects of antigenic priming on influenza vaccination now will improve the interpretation of HI antibody responses toward universal vaccinations in the future.

Supplementary Materials: The following are available online at www.mdpi.com/link, Figure S1: Decline in HI Titer After Vaccination, Figure S2: Titers from Short and Long Service Subgroups, Figure S3: Amount of IgG from Short and Long Service Subgroups, Figure S4: HI Titers with Years of Service, Figure S5: HI Titers for subjects born after the emergence of the H1N1 strain, Figure S6: HI Titers for Different Birth Cohorts.

Acknowledgments: We would like to thank Dr. Xing Li for helping with the Octet measurements, and Vladimir Lugovtsev for critical review of the manuscript. Support (of funding) was provided by the Armed Forces Health Surveillance Branch of the Public Health Division, Combat Support Agency (CSA) Operations Directorate at the Defense Health Agency.

Author Contributions: E.P., A.E.-C., J.L.S., Z.Y., and M.C conceived and designed experiments; E.P., and Z.Y. performed HI assays; H.E. performed the modeling; E.P. performed IgG assays and analyzed data; E.P., A.E.-C., J.L.S., Z.Y., and M.C. wrote and edited the manuscript.

Conflicts of Interest: The authors declare no conflict of interest. The views expressed in this article are those of the authors and do not reflect the official policy or position of the Department of Defense or of the U.S. Government. The founding sponsors had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript, and in the decision to publish the results.

References

1. Hobson D, Curry RL, Beare AS, Ward-Gardner A. The role of serum haemagglutination-inhibiting antibody in protection against challenge infection with influenza A2 and B viruses. *J Hyg (Lond)*. 1972;70:767-77.
2. Coudeville L, Baillex F, Riche B, Megas F, Andre P, Ecochard R. Relationship between haemagglutination-inhibiting antibody titres and clinical protection against influenza: development and application of a bayesian random-effects model. *BMC Med Res Methodol*. 2010;10:18.

3. Black S, Nicolay U, Vesikari T, Knuf M, Del Giudice G, Della Cioppa G, et al. Hemagglutination inhibition antibody titers as a correlate of protection for inactivated influenza vaccines in children. *Pediatr Infect Dis J*. 2011;30:1081-5.
4. Hannoun C, Megas F, Piercy J. Immunogenicity and protective efficacy of influenza vaccination. *Virus Res*. 2004;103:133-8.
5. Petrie JG, Ohmit SE, Johnson E, Truscon R, Monto AS. Persistence of Antibodies to Influenza Hemagglutinin and Neuraminidase Following One or Two Years of Influenza Vaccination. *J Infect Dis*. 2015;212:1914-22.
6. Nabeshima S, Kashiwagi K, Murata M, Kanamoto Y, Furusyo N, Hayashi J. Antibody response to influenza vaccine in adults vaccinated with identical vaccine strains in consecutive years. *J Med Virol*. 2007;79:320-5.
7. Ng S, Ip DK, Fang VJ, Chan KH, Chiu SS, Leung GM, et al. The effect of age and recent influenza vaccination history on the immunogenicity and efficacy of 2009-10 seasonal trivalent inactivated influenza vaccination in children. *PLoS One*. 2013;8:e59077.
8. Thompson MG, Naleway A, Fry AM, Ball S, Spencer SM, Reynolds S, et al. Effects of Repeated Annual Inactivated Influenza Vaccination among Healthcare Personnel on Serum Hemagglutinin Inhibition Antibody Response to A/Perth/16/2009 (H3N2)-like virus during 2010-11. *Vaccine*. 2016;34:981-8.
9. Nachbagauer R, Choi A, Izikson R, Cox MM, Palese P, Krammer F. Age Dependence and Isotype Specificity of Influenza Virus Hemagglutinin Stalk-Reactive Antibodies in Humans. *MBio*. 2016;7:e01996-15.
10. Severson JJ, Richards KR, Moran JJ, Hayney MS. Persistence of influenza vaccine-induced antibody in lung transplant patients and healthy individuals beyond the season. *Hum Vaccin Immunother*. 2012;8:1850-3.
11. Kitphati R, Pooruk P, Lerdsamran H, Poosuwan S, Louisirirochanakul S, Auewarakul P, et al. Kinetics and longevity of antibody response to influenza A H5N1 virus infection in humans. *Clin Vaccine Immunol*. 2009;16:978-81.
12. Kositanont U, Assantachai P, Wasi C, Puthavathana P, Praditsuwan R. Kinetics of the antibody response to seasonal influenza vaccination among the elderly. *Viral Immunol*. 2012;25:471-6.
13. Mutsaerts E, Madhi SA, Cutland CL, Jones S, Hugo A, Trenor S, et al. Influenza vaccination of pregnant women protects them over two consecutive influenza seasons in a randomized controlled trial. *Expert Rev Vaccines*. 2016;15:1055-62.
14. Valero-Pacheco N, Perez-Toledo M, Villasis-Keever MA, Nunez-Valencia A, Bosco-Garate I, Lozano-Dubernard B, et al. Antibody Persistence in Adults Two Years after Vaccination with an H1N1 2009 Pandemic Influenza Virus-Like Particle Vaccine. *PLoS One*. 2016;11:e0150146.
15. Hsu JP, Zhao X, Chen MI, Cook AR, Lee V, Lim WY, et al. Rate of decline of antibody titers to pandemic influenza A (H1N1-2009) by hemagglutination inhibition and virus microneutralization assays in a cohort of seroconverting adults in Singapore. *BMC Infect Dis*. 2014;14:414.
16. Ruiz-Palacios GM, Leroux-Roels G, Beran J, Devaster JM, Esen M, Launay O, et al. Immunogenicity of AS03-adjuvanted and non-adjuvanted trivalent inactivated influenza vaccines in elderly adults: A Phase 3, randomized trial and post-hoc correlate of protection analysis. *Hum Vaccin Immunother*. 2016;12:3043-55.
17. Radin JM, Hawksworth AW, Myers CA, Ricketts MN, Hansen EA, Brice GT. Influenza vaccine effectiveness: Maintained protection throughout the duration of influenza seasons 2010-2011 through 2013-2014. *Vaccine*. 2016;34:3907-12.
18. Puig-Barbera J, Mira-Iglesias A, Tortajada-Girbes M, Lopez-Labrador FX, Librero-Lopez J, Diez-Domingo J, et al. Waning protection of influenza vaccination during four influenza seasons, 2011/2012 to 2014/2015. *Vaccine*. 2017;35:5799-807.
19. Young B, Sadarangani S, Jiang L, Wilder-Smith A, Chen MI. The Duration of Influenza Vaccine Effectiveness: A Systematic Review, Meta-analysis and Meta-regression of Test-Negative Design Case-control Studies. *J Infect Dis*. 2017.
20. Skowronski DM, Chambers C, Sabaiduc S, De Serres G, Winter AL, Dickinson JA, et al. Beyond Antigenic Match: Possible Agent-Host and Immuno-epidemiological Influences on Influenza Vaccine Effectiveness During the 2015-2016 Season in Canada. *J Infect Dis*. 2017;216:1487-500.
21. Giefing-Kroll C, Berger P, Lepperdinger G, Grubeck-Loebenstien B. How sex and age affect immune responses, susceptibility to infections, and response to vaccination. *Aging Cell*. 2015;14:309-21.
22. Plant EP, Fredell LJ, Hatcher BA, Li X, Chiang MJ, Kosikova M, et al. Different Repeat Annual Influenza Vaccinations Improve the Antibody Response to Drifted Influenza Strains. *Sci Rep*. 2017;7:5258.
23. Cobey S, Hensley SE. Immune history and influenza virus susceptibility. *Curr Opin Virol*. 2017;22:105-11.

24. Andrews SF, Kaur K, Pauli NT, Huang M, Huang Y, Wilson PC. High preexisting serological antibody levels correlate with diversification of the influenza vaccine response. *J Virol.* 2015;89:3308-17.
25. Fonville JM, Wilks SH, James SL, Fox A, Ventresca M, Aban M, et al. Antibody landscapes after influenza virus infection or vaccination. *Science.* 2014;346:996-1000.
26. Miller MS, Gardner TJ, Krammer F, Aguado LC, Tortorella D, Basler CF, et al. Neutralizing antibodies against previously encountered influenza virus strains increase over time: a longitudinal analysis. *Sci Transl Med.* 2013;5:198ra07.
27. Smith DJ, Forrest S, Ackley DH, Perelson AS. Variable efficacy of repeated annual influenza vaccination. *Proc Natl Acad Sci U S A.* 1999;96:14001-6.
28. Monto AS, Malosh RE, Petrie JG, Martin ET. The Doctrine of Original Antigenic Sin: Separating Good From Evil. *J Infect Dis.* 2017;215:1782-8.
29. Vatti A, Monsalve DM, Pacheco Y, Chang C, Anaya JM, Gershwin ME. Original antigenic sin: A comprehensive review. *J Autoimmun.* 2017;83:12-21.
30. Paules CI, Marston HD, Eisinger RW, Baltimore D, Fauci AS. The Pathway to a Universal Influenza Vaccine. *Immunity.* 2017;47:599-603.
31. Pinheiro J, Bates D, DebRoy S, Sarkar D. nlme: Linear and Nonlinear Mixed Effects Models. R package version 3.1-131.1 ed2018.
32. Lee J, Boutz DR, Chromikova V, Joyce MG, Vollmers C, Leung K, et al. Molecular-level analysis of the serum antibody repertoire in young adults before and after seasonal influenza vaccination. *Nat Med.* 2016;22:1456-64.
33. Christian LM, Beverly C, Mitchell AM, Karlsson E, Porter K, Schultz-Cherry S, et al. Effects of prior influenza virus vaccination on maternal antibody responses: Implications for achieving protection in the newborns. *Vaccine.* 2017;35:5283-90.
34. Sasaki S, He XS, Holmes TH, Dekker CL, Kemble GW, Arvin AM, et al. Influence of prior influenza vaccination on antibody and B-cell responses. *PLoS One.* 2008;3:e2975.
35. Levine MZ, Martin JM, Gross FL, Jefferson S, Cole KS, Archibald CA, et al. Neutralizing Antibody Responses to Antigenically Drifted Influenza A(H3N2) Viruses among Children and Adolescents following 2014-2015 Inactivated and Live Attenuated Influenza Vaccination. *Clin Vaccine Immunol.* 2016;23:831-9.
36. Nunez IA, Carlock MA, Allen JD, Owino SO, Moehling KK, Nowalk P, et al. Impact of age and pre-existing influenza immune responses in humans receiving split inactivated influenza vaccine on the induction of the breadth of antibodies to influenza A strains. *PLoS One.* 2017;12:e0185666.
37. Barker WH. Excess pneumonia and influenza associated hospitalization during influenza epidemics in the United States, 1970-78. *Am J Public Health.* 1986;76:761-5.
38. Thompson WW, Shay DK, Weintraub E, Brammer L, Cox N, Anderson LJ, et al. Mortality associated with influenza and respiratory syncytial virus in the United States. *JAMA.* 2003;289:179-86.
39. Alling DW, Blackwelder WC, Stuart-Harris CH. A study of excess mortality during influenza epidemics in the United States, 1968-1976. *Am J Epidemiol.* 1981;113:30-43.
40. Lui KJ, Kendal AP. Impact of influenza epidemics on mortality in the United States from October 1972 to May 1985. *Am J Public Health.* 1987;77:712-6.
41. Centers for Disease C, Prevention. Update: influenza activity--United States and worldwide, 2003-04 season, and composition of the 2004-05 influenza vaccine. *MMWR Morb Mortal Wkly Rep.* 2004;53:547-52.
42. Centers for Disease C, Prevention. Update: influenza activity - United States, 2011-12 season and composition of the 2012-13 influenza vaccine. *MMWR Morb Mortal Wkly Rep.* 2012;61:414-20.
43. Liu KA, Mager NA. Women's involvement in clinical trials: historical perspective and future implications. *Pharm Pract (Granada).* 2016;14:708.
44. Ingersoll MA. Sex differences shape the response to infectious diseases. *PLoS Pathog.* 2017;13:e1006688.
45. Sue K. The science behind "man flu". *BMJ.* 2017;359:j5560.
46. Engler RJ, Nelson MR, Klote MM, VanRaden MJ, Huang CY, Cox NJ, et al. Half- vs full-dose trivalent inactivated influenza vaccine (2004-2005): age, dose, and sex effects on immune responses. *Arch Intern Med.* 2008;168:2405-14.
47. Falsey AR, Treanor JJ, Tornieporth N, Capellan J, Gorse GJ. Randomized, double-blind controlled phase 3 trial comparing the immunogenicity of high-dose and standard-dose influenza vaccine in adults 65 years of age and older. *J Infect Dis.* 2009;200:172-80.

48. Khurana S, Verma N, Talaat KR, Karron RA, Golding H. Immune response following H1N1pdm09 vaccination: differences in antibody repertoire and avidity in young adults and elderly populations stratified by age and gender. *J Infect Dis.* 2012;205:610-20.
49. Simell B, Nurkka A, Lahdenkari M, Givon-Lavi N, Kayhty H, Dagan R, et al. Association of serotype-specific antibody concentrations and functional antibody titers with subsequent pneumococcal carriage in toddlers immunized with a 9-valent pneumococcal conjugate vaccine. *Clin Vaccine Immunol.* 2012;19:96-9.
50. Fink AL, Klein SL. Sex and Gender Impact Immune Responses to Vaccines Among the Elderly. *Physiology (Bethesda).* 2015;30:408-16.
51. Gergen PJ, McQuillan GM, Kiely M, Ezzati-Rice TM, Sutter RW, Virella G. A population-based serologic survey of immunity to tetanus in the United States. *N Engl J Med.* 1995;332:761-6.
52. Furman D, Hejblum BP, Simon N, Jojic V, Dekker CL, Thiebaut R, et al. Systems analysis of sex differences reveals an immunosuppressive role for testosterone in the response to influenza vaccination. *Proc Natl Acad Sci U S A.* 2014;111:869-74.
53. Worobey M, Han GZ, Rambaut A. Genesis and pathogenesis of the 1918 pandemic H1N1 influenza A virus. *Proc Natl Acad Sci U S A.* 2014;111:8107-12.
54. Gagnon A, Acosta E, Hallman S, Bourbeau R, Dillon LY, Ouellette N, et al. Pandemic Paradox: Early Life H2N2 Pandemic Influenza Infection Enhanced Susceptibility to Death during the 2009 H1N1 Pandemic. *MBio.* 2018;9.
55. Skowronski DM, Chambers C, De Serres G, Sabaiduc S, Winter AL, Dickinson JA, et al. Age-Related Differences in Influenza B Infection by Lineage in a Community-Based Sentinel System, 2010-2011 to 2015-2016, Canada. *J Infect Dis.* 2017;216:697-702.
56. Brodin P, Jojic V, Gao T, Bhattacharya S, Angel CJ, Furman D, et al. Variation in the human immune system is largely driven by non-heritable influences. *Cell.* 2015;160:37-47.
57. Belongia EA, Skowronski DM, McLean HQ, Chambers C, Sundaram ME, De Serres G. Repeated annual influenza vaccination and vaccine effectiveness: review of evidence. *Expert Rev Vaccines.* 2017;16:1-14.
58. Beyer WEP, Palache AM, Boulfich M, Osterhaus A. Rationale for two influenza B lineages in seasonal vaccines: A meta-regression study on immunogenicity and controlled field trials. *Vaccine.* 2017;35:4167-76.
59. Laurie KL, Horman W, Carolan LA, Chan KF, Layton D, Bean A, et al. Evidence for Viral Interference and Cross-reactive Protective Immunity Between Influenza B Virus Lineages. *J Infect Dis.* 2018.
60. Gross PA, Denning CR, Gaerlan PF, Bonelli J, Bernius M, Dran S, et al. Annual influenza vaccination: immune response in patients over 10 years. *Vaccine.* 1996;14:1280-4.
61. He XS, Holmes TH, Sanyal M, Albrecht RA, Garcia-Sastre A, Dekker CL, et al. Distinct patterns of B-cell activation and priming by natural influenza virus infection versus inactivated influenza vaccination. *J Infect Dis.* 2015;211:1051-9.
62. Kang EK, Eun BW, Kim NH, Lim JS, Lee JA, Kim DH. The priming effect of previous natural pandemic H1N1 infection on the immunogenicity to subsequent 2010-2011 influenza vaccination in children: a prospective cohort study. *BMC Infect Dis.* 2016;16:438.