

## SUPPORTING INFORMATION

**(+)-*epi*-Epoformin, a phytotoxic fungal  
cyclohexenepoxide: Structure activity relationships**

Antonio Cala<sup>1,†</sup>, Marco Masi<sup>2,†</sup>, Alessio Cimmino<sup>2</sup>, José M.G. Molinillo<sup>1</sup>, Francisco A. Macías<sup>1,\*</sup> and Antonio Evidente<sup>2</sup>

1 Allelopathy Group, Department of Organic Chemistry, School of Science, Institute of Biomolecules (INBIO), University of Cádiz, C/ República Saharaui 7, 11510-Puerto Real, Cádiz, Spain; antonio.cala@uca.es (A.C.); chema.gonzalez@uca.es (J.M.G.M.), famacias@uca.es (F.A.M.).


2 Department of Chemical Sciences, University of Naples "Federico II", Complesso Universitario Monte S. Angelo, Via Cintia 4, 80126 Napoli, Italy; marco.masi@unina.it (M.M.); alessio.cimmino@unina.it (A.C.), evidente@unina.it (A.E.).

<sup>†</sup> These authors contributed equally to this work.


\*Correspondence: Francisco A. Macías, famacias@uca.es

## Table of Contents


S3.	<sup>1</sup> H NMR Spectrum of 1
S4.	<sup>1</sup> H and <sup>13</sup> C NMR Spectra of 2
S5.	<sup>1</sup> H and <sup>13</sup> C NMR Spectra of 3
S6.	<sup>1</sup> H and <sup>13</sup> C NMR Spectra of 4
S7.	<sup>1</sup> H and <sup>13</sup> C NMR Spectra of 5
S8.	<sup>1</sup> H and <sup>13</sup> C NMR Spectra of 6
S9.	<sup>1</sup> H and <sup>13</sup> C NMR Spectra of 7
S10.	<sup>1</sup> H and <sup>13</sup> C NMR Spectra of 8
S11.	<sup>1</sup> H and <sup>13</sup> C NMR Spectra of 9



<sup>1</sup>H-NMR Spectrum of **1** (400MHz, CDCl<sub>3</sub>)




<sup>1</sup>H-NMR Spectrum of **2** (500MHz, CDCl<sub>3</sub>)




<sup>13</sup>C-NMR Spectrum of **2** (125MHz, CDCl<sub>3</sub>)

(+)-*epi*-Epoformin, a phytotoxic fungal cyclohexenepoxide: Structure activity relationships




<sup>1</sup>H-NMR Spectrum of **3** (500MHz, CDCl<sub>3</sub>)




<sup>13</sup>C-NMR Spectrum of **3** (125MHz, CDCl<sub>3</sub>)

(+)-*epi*-Epoformin, a phytotoxic fungal cyclohexenepoxide: Structure activity relationships




<sup>1</sup>H-NMR Spectrum of **4** (400MHz, CDCl<sub>3</sub>)




<sup>13</sup>C-NMR Spectrum of **4** (100MHz, CDCl<sub>3</sub>)

(+)-*epi*-Epoformin, a phytotoxic fungal cyclohexenepoxide: Structure activity relationships




<sup>1</sup>H-NMR Spectrum of 5 (500MHz, CDCl<sub>3</sub>)




<sup>13</sup>C-NMR Spectrum of 5 (125MHz, CDCl<sub>3</sub>)

(+)-*epi*-Epoformin, a phytotoxic fungal cyclohexenepoxide: Structure activity relationships





<sup>1</sup>H-NMR Spectrum of **6** (400MHz, CDCl<sub>3</sub>)




<sup>13</sup>C-NMR Spectrum of **6** (100MHz, CDCl<sub>3</sub>)

(+)-*epi*-Epoformin, a phytotoxic fungal cyclohexenepoxide: Structure activity relationships







$^1\text{H-NMR}$  Spectrum of 7 (400MHz,  $\text{CDCl}_3$ )




$^{13}\text{C-NMR}$  Spectrum of 7 (100MHz,  $\text{CDCl}_3$ )

(+)-*epi*-Epoformin, a phytotoxic fungal cyclohexenepoxide: Structure activity relationships





<sup>1</sup>H-NMR Spectrum of 8 (400MHz, CDCl<sub>3</sub>)




<sup>13</sup>C-NMR Spectrum of 8 (100MHz, CDCl<sub>3</sub>)

(+)-*epi*-Epoformin, a phytotoxic fungal cyclohexenepoxide: Structure activity relationships



<sup>1</sup>H-NMR Spectrum of **9** (400MHz, CDCl<sub>3</sub>)



<sup>13</sup>C-NMR Spectrum of **9** (100MHz, CDCl<sub>3</sub>)

(+)-*epi*-Epoformin, a phytotoxic fungal cyclohexenepoxide: Structure activity relationships