

Article

Micromorphological and chemical features of soils as evidence of Bronze Age ancient anthropogenic impact (Late Bronze Age Muradymovo settlement, Ural region, Russia)

Alexandra Golyeva¹, Olga Khokhlova², Marina Lebedeva³, Nickolay Shcherbakov⁴, Iia Shuteleva⁴

¹ Institute of Geography RAS, Moscow, Russian Federation, golyevaaa@yandex.ru

² Institute of Physical, Chemical and Biological Problems in Soil Science, Russian Academy of Sciences, ulitsa Institutskaya, 2, 142290, Pushchino, Moscow region, Russian Federation. E-mail: olga_004@rambler.ru

³ V.V. Dokuchaev Soil Science Institute, Moscow, Russian Federation. E-mail: m_verba@mail.ru

⁴ Archaeological Laboratory of Bashkir State Pedagogical University, Ufa, Russian Federation. E-mails: sherbakov@rambler.ru, shutelevai@gmail.com

* Correspondence: golyevaaa@yandex.ru; Tel.: +07-916-3294335

Abstract: In some cases, the human impact on ancient landscapes has been so profound that local soils still remain significantly affected even after hundreds and thousands of years after ending impact. We studied the Late Bronze Age Muradymovo settlement located in Urals, Russia, aiming to estimate the consequences of the ancient people activity for environment. Despite the present humid climate, the modern soils inside the cultural layer of the study site contain more than 27% of gypsum at a depth of just 10 cm from the surface, and a microrelief of the study site is typical of a gypsum desert. The nearby background Chernozems are gypsum-free to a depth of 2 meters. According to the archaeological data, the ancient people belonged to the 'Srubno-Alakul' archaeological culture (1750-1350 yrs BC cal) and had a tradition of building their houses from gypsum rocks. At the present time, this area is still unsuitable for human settlement. The properties of modern soils inside the cultural layer of the study site are directly affected by the Late Bronze Age human activities. It has been identified on soil morphology, micromorphology and chemical properties of soils developed inside the cultural layer of the settlement.

Keywords: Bronze Age settlement, modern soil, micromorphology, chemical properties, gypsum, anthropogenic impact

1. Introduction

In some cases, the human impact on ancient landscapes has been so profound that local soils still remain significantly affected even after hundreds and thousands of years after ending impact [1-9]. There are no natural soils left within such sites, being replaced by completely different anthropogenic soils with specific properties [10-17].

Studying the causes and implications of such negative influences of past human activities on soils and environment is necessary to prevent similar accidents in the future. The present article describes a case-study of the extremely severe and long-lasting impact of ancient people on soils and environment. The study site is the Late Bronze Age settlement of Muradymovo located in Bashkortostan Republic (Urals region, Russia). The site and its area have a peculiar hilly microrelief that doesn't occur anywhere else in Bashkortostan Republic. According to the archaeological data [18, 19], the Muradymovo people belonged to the 'Srubno-Alakul' culture of the Late Bronze Age and came here from an extra arid semidesert region of the southern Kazakhstan,

where they used to build their houses of gypsum rocks. There is a deposit of gypsum rocks just 5 km far from the Muradymovo settlement site [20].

The aim of our study was to investigate the micromorphological and chemical properties of the modern soils at the Bronze Age settlement Muradymovo site to estimate the consequences of the ancient people activity for environment.

2. Materials and Methods

The study site ($53^{\circ}58'44.8''$ N, $55^{\circ}30'58.8''$ E) is located 2.5 km north of the village of Muradymovo, in the Aurgazinskiy District of the Bashkortostan Republic of Russia (Figure 1). The site of Muradymovo ancient settlement having a total area of 6 ha is found on the first terrace at the right bank of the Urshak River, 0.2 km east of the mainstream, on a hill about 1.5-2 m high. The bedrock is composed of gypsum, anhydrite, dolomite and sandstone of the Kungur stage of the Permian Period. The bedrock is overlain by loess-like silty sediments (approximately 8-10 m thick) of the Quaternary Period that serve as parent rocks for soils.



Figure 1. Study site location.

Climate is continental, moderately cold. The mean annual air temperature is $+2.5^{\circ}\text{C}$, with the mean temperatures of January and July being -15°C and $+19.5^{\circ}\text{C}$, respectively. The mean annual precipitation is about 500 mm, with more than 300 mm falling during the growing season. The hygrothermal coefficient is about 1 [21].

The typical modern vegetation is represented by steppe communities.

The typical soils are Calcic Chernozems Loamic [22] that are naturally gypsum-free to a depth more than 3 meters [23, 24].

According to the archaeological data [25, 26], the settlement was built by ancient people of the 'Srubno-Alakul' archaeological culture of the Late Bronze Age (1750-1350 yrs BC cal), who lived here for no longer than 200-300 years. Later the settlement has remained abandoned. At present, the site is covered with sparse steppe vegetation, partially used as a pasture and bordered by a gully from the west and north.

We studied the archaeological excavation pit (no. IV) with the most representative morphology of cultural layers (Figures 2.a). The pit included several layers of house remains (no less than 5) and a hearth, i.e., this residential house was rebuilt several times [27].

The background soil outcropping from the bank of a small river at a short distance from the archeological excavation site was also studied. This soil was least affected by the ancient settlement construction, it was located on the opposite bank and had no traces of ancient human impact (Figure 2.b).

The modern microrelief consists of small hillocks separated by deep frost cracks, which covers the whole area of the study site and beyond, up to the banks of the river and gully. Frost cracks result from recent influence of continental climate. The hillocks are about 50 cm high and 1.5-2.2 m long. Such peculiar microrelief is absent at the other side of the gully.




Figure 2. Soil profiles: a) – archaeological pit; b) – background soil.

We described morphological characteristics of the pit within the settlement and the profile of background soil, collected samples in vertical columns in the field and conducted micromorphological and chemical analyses of the samples in the laboratory using conventional techniques [28-30]. The samples were prepared according to the requirements for each specific analysis.

Micromorphological observations were carried out on thin sections (3x4 cm square and 30 μ m thick), which were prepared from undisturbed soil samples from the upper part (10-20 cm), from the lower part (60-70 cm from the surface) of the cultural layer and from the humus horizon (10-20 cm) of the background soil. They were studied using an Olympus BX51 TL RL Pol microscope and an Olympus DP25 digital camera at the V.V. Dokuchaev Soil Science Institute, Moscow and using a polarizing microscope Carl Zeiss HBO 50 in the Chemical-Analytical Complex of the Institute of Physical, Chemical and Biological Problems in Soil Science, Russian Academy of Sciences, Pushchino.

The mineralogical composition of the dry residue of the water extract from the cultural layer samples taken at the depths of 10-20 and 60-70 cm was determined by X-ray diffraction in the Laboratory of Soil mineralogy and Micromorphology of the V.V. Dokuchaev Soil Science Institute by Dr. E. Varlamov. The analysis was conducted using an HZG-4A universal X-ray diffractometer with the following settings: Cu radiation, 30 kV tube voltages, 20 mA current and a scan speed of 2°-2 θ per minute. Calculations were performed using 'Difraktometr-Avto 2014' software produced by the LLC Iris.

Total phosphorus. The procedure included the sample combustion with concentrated sulfuric acid. Phosphate in the extract was determined calorimetrically using a SPECOL 211 spectrophotometer and a blue ammonium molybdate method with ascorbic acid as a reducing agent [30].

Gypsum was determined using 10% BaCl₂ solution [28]. Each sample was boiled for 3 minutes in 0.2N HCl, cooled for 30 minutes and passed through a filter. The filtrate was diluted with distilled water and passed through H-cationite. The resulting solution was titrated with BaCl₂ solution. The obtained values of SO₄ concentrations were recalculated for gypsum (CaSO₄·2H₂O).

pH_{H2O} was determined using a potentiometer, in suspension with soil to water ratio of 1:2.5, after a single shaking followed by settling for 30 min [28].

The organic carbon was determined by the Tyurin method, which included the wet combustion of organic substance in a mixture of 0.4 N K₂Cr₂O₇ and concentrated H₂SO₄ (1:1) at 150°C for 20 min. The measurements were performed by photometry on a SPECOL 211 spectrometer at 590 nm [28].

Calcium carbonate concentrations in the samples were determined by alkalimetry using the Kozlovskii procedure. A soil sample was treated with 2 M HCl; the released CO₂ was absorbed by a 0.4 M NaOH solution. Then a saturated BaCl₂ solution was added to the tube with NaOH, and the excess of alkali was titrated with 0.2 M HCl [30]. The obtained values of the carbonate ion concentrations were recalculated for calcium carbonates.

3. Results

3.1. Morphological description.

The soils studied were considerably different from each other (Figure 2).

3.1.1. Modern soil (background or reference soil)

Undisturbed background soils of the site were attributed to Calcic Chernozems Loamic and were characterized by the following features: a humus profile with the Ah + AhB horizons 50-(70) cm thick, followed by a carbonate-accumulative horizon Bk to a depth of 110 cm with the maximum occurrence of carbonate pedofeatures in the form of vertical chains of white soft spots (or soft carbonate nodules) at a depth of 90-110 cm, which, in turn, was followed by the soil-forming rock. The signs of burrowing animal's activity in the profiles are apparent: in the upper horizons the structure is coprogenic, the AhB horizons contain dark gray and yellowish-pale animal holes; in the Bk horizon a number of thin long passages filled with humus material were found; there are not fresh holes with a clear contour, but only indistinct, diffused ones.

The background (reference) soil had features of zonal Chernozem, with the organomineral horizon thicker than 50 cm, underlain by the calcic horizon.

3.1.2. The soil inside the cultural layer

The section was studied under forb-grassy vegetation. Soil of the excavation pit IV had a light-colored surface layer containing diffuse secondary calcium sulfate and abundant artefacts – pottery, bones and charcoal. According to archaeological data [26] it was the layer of habitation deposits. Under the habitation deposits, a well-preserved and morphologically distinct buried soil was found. We added the index “cul” to two surface horizons of the studied soil meaning that those horizons are developed inside the cultural layer.

Asod cul, 0-3 cm. Sod horizon.

A1 cul, 3-7(20) cm. Black, slightly dry, light loamy, with fine crumb structure; slightly compacted; siliceous powdering on ped faces; densely penetrated by roots; clear smooth boundary.

Cultural Layer 7(20)-80(86)cm. Light grayish pale, slightly dry, light loamy, structureless; powdery (ashlike); soft nodules and filaments of calcareous pseudomycelium with a characteristic size of 0.3-0.6 cm; roots; animal bones of 2-3 to 10-15 cm in size; fragments of broken pottery, clear boundary.

[A], 80(86)-106 cm. Black, slightly dry, heavy loam with coarse columnar-prismatic structure, calcareous powdering on ped faces; diffuse boundary.

[AB], 106-116 cm. Dark ochreous brown, with humus streaks, slightly wet heavy loam, fine columnar-prismatic structure; with ochreous-colored mottles.

3.2. Micromorphological description

The humus horizon of modern background soil has weakly separated granular microstructure, complex packing voids, small granules with great content of organic fine material, plant tissue residues and signs of mezofauna activity in voids (Fig. 3 a, b); the plant tissue residues have visible cell structure surrounded by brown and black small granules, signs of mites activity, speckled b-fabric (Fig 3 c, d) and no gypsum.

In the soil inside the cultural layer two samples, 10-20 and 60-70 cm depths, were compared. They have similarities and differences in their microfabric. The similarities include the absolute predominance of gypsum grains of various sizes and habits in the soil skeleton composition and the presence of granular peds consisting of isotropic amorphous organic fine material (Fig.4 a - h) [31].

The differences between the compared samples are expressed by quantitative rather than by the qualitative distinctions between prevailing forms of gypsum crystals and sizes of organic granular peds, as described below.




Figure 3. Micromorphological features of the humus horizon of modern reference soil at the depth 10-20 cm: a, b – weakly separated granular microstructure, complex packing voids, small granules with great content of organic fine material, tissue residues and signs of mezofauna activity in voids(a- PPL; b - XPL); c, d – tissue residue with visible cell structure surrounded by brown and black small granules, signs of mites activity, speckled b-fabric, complex packing voids (c - PPL; d -XPL).

Dominant features in the layer at the 10-20 cm depth are as follows: 1) gypsum crystals of two types: a) aggregates of corroded thin prismatic crystals (on average about 30 µm long) between humus granular peds, many of which are covered with humus coatings, and b) thick infillings of very dense and small (about 0.05 µm) tabular gypsum crystals in channel pores containing fine residues of plant tissues (Fig.4 a - d); 2) relatively large crumb peds (100-140 µm) that consist of agglomerated small granular peds (with an average size of about 40-50 µm); 3) bone fragments that are very small and rare; 4) rare micrite crystals [31].

At the depth of 60-70 cm, compared to the above sample, the number of humus aggregates is significantly lower and gypsum is a main soil-forming mineral. Among gypsum crystals, aggregates (clusters) of regularly shaped acicular gypsum crystals predominate. They form loose continuous infillings formed of lenticular and acicular gypsum crystals in pores between soil peds (Fig. 4 e, f). These infillings have only few microzones with microfeatures of re-crystallization resulting in formation of very small tabular gypsum crystals, similar to those prevailing within dense pore infillings containing plant tissues at the 10-20 cm depth. Humus microaggregates occurring in rarer groups have features of fragmentation due to the growth of acicular gypsum crystals within certain microzones and also contain long fragments of aragonite. Compared to the above sample, this layer has an increased number of large fragments of bones with microfeatures of their carbonatization and/or small micrite concentrations over small fragments of bones. Large fragments of bones have humus coatings on their surface (Fig. 4 g, h).




Figure 4. Micromorphological features of soil within the cultural at different depths: 10-20 cm (a - d); 60-70 cm (e -h). a-d – thick infillings of very dense and small (about 0.05 μm) tabular gypsum crystals in channel pores containing fine residues of plant tissues (a, c - XPL; b, d - PPL); e, f – loose continuous infillings composed of lenticular and acicular gypsum crystals; g, h – particles of bones with the carbonates (e, g - XPL; f, h - PPL).

3.3. The mineralogical investigation of dry residue of water extract

The analysis of X-ray diffraction patterns of powder samples from the compared samples in the soil within the cultural layer (10-20 and 60-70 cm depths) has revealed that their mineralogical composition is similar: only gypsum minerals with characteristic diffraction peaks are present in the dry residue of water extract (Fig.5). No other mineral salts have been identified.




Figure 5. Results of X-ray diffractions.

3.4. Chemical analyses.

The results of chemical analyses are presented in Table 1.

Table 1. Chemical properties of soils and cultural layers

Depth, cm	pH _{H2O}	Corg, %	P _{tot} , %	CaCO ₃ , %	CaSO ₄ , %
horizon					(gypsum)
Pit IV, inhabited house					
0-5 Asod cul	7.9	14.4	0.48	3.4	0.7
5-10 Ah cul	8.3	11.4	0.64	15.3	1.9
10-20 Ah+Cultural layer	8.3	3.8	0.26	9.5	49.8
20-30 Cultural layer	8.2	2.7	0.29	13.6	49.9
30-40 Cultural layer	8.2	2.9	0.42	15.3	39.2
40-50 Cultural layer	8.2	2.3	0.50	14.4	38.2
50-60 Cultural layer	8.0	2.7	0.48	19.4	8.0
60-70 Cultural layer	7.9	2.2	0.48	15.9	31.4
70-80 Cultural layer	7.9	1.5	0.45	18.4	25.6
80-86 [Ah]	8.0	2.1	0.38	24.6	3.9
86-96 [Ah]	8.4	2.4	0.21	33.1	1.3
96-106 [Ah]	8.6	1.8	0.19	38.3	2.7
106-116 [AhB]	9.1	1.8	0.19	42.2	2.3

Background reference soil

0-10 Ah	7.4	7.7	0.21	3.4	0
10-20 Ah	8.0	5.4	0.17	1.7	0
20-30 Ah	8.2	4.7	0.15	0	0
30-40 Ah	8.2	3.1	0.10	0	0
40-50 Ah	8.3	2.2	0.08	1.8	0
50-60 Ah	8.3	2.5	0.08	4.5	0
60-70 AhB	8.3	1.2	0.07	9.3	0
70-80 Bk	8.2	0.9	0.07	15.7	0
80-90 Bk	8.2	0.5	0.07	18.6	0.2
90-100 Bk	8.2	1.2	0.08	23.2	0.2
100-110 Bk	8.3	-	0.08	23.3	0.03

The background reference soil has a neutral reaction within its upper part and alkaline within its lower part. A similar pH distribution pattern is typical for native soils of the study region.

The organic carbon content and distribution are typical for the Chernozems [22], with the maximal organic carbon content within the upper 50 cm followed by a sharp decrease in deeper layers.

The content of P_{tot} in soil is low in comparison with the anthropogenic soils of excavation pit. The highest values (0.21%) occur within the litter horizon, while the mineral horizons have a uniform small concentration of phosphorus, which is typical for native soils.

Calcium carbonate content is low within the upper 40-cm-thick layer and significantly increases in deeper layers, which is also usual for the native soils.

The gypsum content is almost zero till the 70 cm depth. At the very bottom of the profile, there is a very small peak of gypsum and calcium carbonate concentration.

The excavation pit IV (residential house). The soil inside the cultural layer has a neutral reaction within its upper part and alkaline within its lower part. A similar pH distribution pattern is typical for native soils of the study region and reflects the general trend of downward migration of calcareous soil solutions. The organic matter content is high throughout the soil profile. The maximum amount of organic matter is concentrated within the upper 10-cm-thick layer, which is generally typical for Chernozems.

The P_{tot} distribution is irregular. The cultural layers occurring at 0-86 cm depth are characterized by a high content of total phosphorus (0.26-0.64%). The buried Chernozem (86-96 cm depth) has a lower P_{tot} content, virtually equal to that in the uppermost layer of the modern background soil.

The CaCO_3 content is high throughout the profile, being slightly higher in its lower part.

The gypsum content and distribution are very unusual. In the field, at a macromorphological scale, the cultural layer of the excavation pit appeared to be composed of whitish-gray ash-like material, relatively homogeneous, compacted, with inclusions of various artefacts. The laboratory analyses have revealed that the cultural layer is composed of a mixture of gypsum and organic matter, with small amount of calcium carbonate. The data obtained (Table 1) show that a high content of gypsum (more than 49%) is registered within the cultural layer at the depths from 10 to 86 cm, with a gradual downward decrease. The surface layer (0-10 cm) is relatively impoverished in gypsum (1.3%) as a result of leaching.

4. Discussion

The upper horizons of the pit IV is the ancient cultural layer according to their morphological characteristics and the archaeological data [26]. The modern background soil bears no traces of former human impact.

Background soil. By the content and distribution of P_{tot} the modern soil adequately reflects its natural background level, which can be used as a reference for separating the natural and anthropogenic layers: the former are poor in phosphorus, while the latter are characterized by the P_{tot} content above 0.22%.

A small amount of calcium carbonate within the uppermost horizon has a biogenic origin, resulting from the calcium carbonate uptake by plant roots and its return to soil upon the death and decay of those roots [32-34]. This is a general natural phenomenon for soils of the steppe regions of Russia.

The very limited amount of gypsum in the background soil is natural for soils developed in modern climatic conditions of the study region.

The excavation pit IV. There is an overall decrease of the organic matter content with depth, with occasional humus-rich lenses. These lenses in the upper layers apparently result from man-made depositions of the organic matter during the period of settlement building and exploitation and indicate the cultural layer. Similar lenses in the lower layers, deeper than 86 cm, are a part of the organic matter of the buried paleosol.

The “saw-tooth” pattern of the P_{tot} distribution within the total depth of the cultural layers reflects the stages of increase and decrease of anthropogenic pressure during the settlement functioning [35, 36, 12, 37]. There is no phosphorus depletion within the upper 10 cm, which is surprising after such a long period (more than 3000 years) following the abandonment of this site. Perhaps, presence of phosphorus within the surface layer could be its uptake by plants with subsequent decomposition of plant material. In addition, high amount of gypsum in the cultural layer may serve a “fixing factor” for phosphorus.

In this excavation pit, the content of calcium carbonate is significantly higher than that in background soil. The latter is calcareous only at depths more than 50 cm, while the excavation pit profile is calcareous from a 5 cm depth. Such large concentrations at a shallow depth are typical for the sites of ancient settlements [37]. They are residues of limestone that was used as a house building material. Calcium carbonate forms almost insoluble complexes with phosphates and organic matter [35]. That is why even under a percolative water regime, the cultural layers of settlements are calcareous.

Comparing the content of gypsum and calcium carbonates revealed a change in the use of bonding agents in the mud-bricks for house building [26]. The bonding agents were CaCO_3 -based at the beginning of the site occupation and later changed to gypsum. At the final stages of settlement existence gypsum was used with insignificant or no calcium carbonate content.

Infillings of gypsum are clearly pedogenic since they are accumulations in pores. Commonly, the presence of gypsum-saturated groundwater, under a non-percolating soil water regime, creates the conditions for precipitation of gypsum in macropores and the formation of these pedofeatures [38, 29]. The study of thin sections has revealed that dense gypsum infillings formed of tabular microcrystals (at the depth of 10-20 cm) and loose continuous infillings composed of lenticular and acicular gypsum crystals (at the depth of 60-70 cm) are modern gypsum pedofeatures. They can be formed by two different processes: first, crystallization from solutions saturated with Ca^{2+} and SO_4^{2-} that are drawn upwards by plant roots; second, presumably, re-crystallization of more ancient narrow prismatic crystals of gypsum (corroded forms of which are found within the upper part of the cultural layer) to form acicular crystals. Large fragments of bones have microfeatures of both their carbonatization and the development of collomorphic organic coatings on their surfaces. The

collomorphic micromorphotype of humus having a well-developed crumb or granular structure is formed under strongly hydromorphic conditions.

The analyses have shown that there is a high concentration of sulphates in the water extract of soil solution, which favors the migration and re-crystallization of gypsum throughout the ancient cultural layer.

Gypsum crystals are present even at the surface of the studied soil inside the cultural layer, which is dramatically different from the background soil which contains no gypsum at the whole profile.

The present day gradual dissolution of these salts results in the wider distribution of saline solutions beyond the settlement area. The high rainfall of the region causes calcium sulfate swelling, which leads to significant increases in the volume of the gypsum horizon and the further development of the characteristic hummocky microrelief. Such microrelief is often for gypsum deserts [39, 40] and absent in the study region, which is within the natural steppe zone.

5. Conclusions

On the basis of the data obtained it can be confidently concluded that the gypsum-bearing strata in the upper part of the excavation pit have an anthropogenic origin. In other words, people built their houses of mud-bricks made of a mixture of gypsum and organic matter (manure, for example), occasionally with addition of small amounts of calcareous rocks.

There is almost pure gypsum within the cultural layer and represents a consecutive series of houses built one after another at the same place. Accumulation of large amounts of gypsum rocks within the ancient settlement site resulted in contamination of the environment with gypsum.

During the period of abandonment of this site, the pedogenic processes homogenized the cultural layer of construction pits. The leaching of salts during the wet seasons was accompanied by the process of upward migration and precipitation of salts during occasional summer droughts as well as severe frosts. Still, the residual amount of gypsum is large after more than 3000 years after contamination. Therefore, it can be assumed that the initial man-made soil contamination by gypsum was extremely strong. The properties of the modern soil inside the cultural layer are directly affected by the Late Bronze Age human activities, with 3-3.5 thousand years being an insufficient timescale to restore the natural gypsum-free soils.

Acknowledgments: The archaeological excavation was supported by the Russian Foundation for Basic Research (project No. 16-11-02003). The field work and laboratory analyses of soils were supported by the Russian Science Foundation (project No. 16-17-10280) and by the State program of FANO RAS No. 0148-2016-0003. We also thank Inga Spiridonova and Michael Hayes for correcting the English language.

Author Contributions: Alexandra Golyeva and Olga Khokhlova performed the expedition with fieldwork and soil sampling, conceived and designed the experiments, and wrote the paper; Nikolay Shcherbakov and Iia Shuteleva organized and conducted the archaeological excavation, provided all the archaeological information; Marina Verba performed micromorphological analysis and determination of mineralogical composition of the water extract. All together we discussed the obtained data and corrected the text.

Conflicts of Interest: The authors declare no conflicts of interest. The funding sponsors did not take part in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript; or in the decision to publish the results.

References

1. Bettis, E.A. Pedogenesis in Late Prehistoric Indian Mounds, Upper Mississippi Valley. *Phys. Geogr.* **1988**, *9*(3), 263-279.
2. Weiss, T.J.; Courty, M.-A.; Wetterstrom, W.; Guichard, F.; Senior, L.; Meadow, R.; Curnow, A. The genesis and collapse of third millennium North Mesopotamia civilization. *Science* **1993**, *261*, 995-1004, DOI: 10.1126/science.261.5124.995

3. Lima, H.N.; Schaefer, C.E.R.; Mello, J.W.V.; Gilkes, R.J.; Ker, J.C. Pedogenesis and pre-Colombian land use of "Terra Preta Anthrosols" ("Indian black earth") of Western Amazonia. *Geoderma* **2002**, *110* (1-2), 1-17, DOI: 10.1016/S0016-7061(02)00141-6
4. Trofimov, S.Ya.; Yakimenko, O.S.; Sedov, S.N.; Dorofeeva, E.I.; Gorshkova, E.I.; Zazovskaya, E.P.; Oleinik, S.A.; Demin, V.V. Composition and properties of organic matter in the soils of ancient Slavic settlements in the forest zone. *Eurasian Soil Sci.* **2004**, *37*(9), 927-936.
5. Plekhanova, L.N.; Demkin, V.A. Ancient soil disturbances in river valleys within the steppe zone of the southeastern Urals. *Eurasian Soil Sci.* **2005**, *38*(9), 973-982.
6. Aleksandrovskii, A.L. Pyrogenic origin of carbonates: Evidence from pedoarchaeological investigations. *Eurasian Soil Sci.* **2007**, *40*(5), 471-477.
7. Khokhlova, O.S.; Khokhlov, A.A.; Kuznetsova, A.M.; Morgunova, N.L.; Chichagova, O.A. Paleosols buried under kurgans of the Pit-grave culture in the steppe zone of the Cis-Ural region. *Eurasian Soil Sci.* **2008**, *41*(5), 481-490.
8. Markiewicz, M.; Bednarek, R.; Jankowski, M.; Świtoniak, M. 'Paleotechnosols' of ancient settlements in Grodno and Kałdus. In : *Technogenic soils of Poland*; Charzyński, P., Hulisz, P., Bednarek, R., Eds.; Polish Society of Soil Science, Poland, **2013**, pp.111-121, ISBN 978-83-934096-1-7.
9. Sanchez-Perrez, S.; Solleiro-Rebolledo, E.; Sedov, S.; McClung de Tapia, E.; Golyeva, A.; Prado, B.; Ibarra-Morales, E. The Black San Pablo Paleosol of the Teotihuacan Valley, Mexico: Pedogenesis, Fertility, and Use in Ancient Agricultural and Urban Systems. *Geoarchaeology* **2013**, *28* (3), 249-267, DOI: 10.1002/gea.21439
10. Woods, W.I.; McCann, J.M. The anthropogenic origin and persistence of Amazonian dark earths. *Yearbook. Conference of Latin Americanist Geographers* **1999**, *25*, 7-14. <http://www.jstor.org/stable/25765871>
11. Nicosia, C.; Balista, C.; Cupitò, M.; Ertani, A.; Leonardi, G.; Nardi, S.; Vidale, M. Anthropogenic deposits from the Bronze Age site of Fondo Paviani (Verona, Italy): Pedochemical and micropedological characteristics. *Quat. Int.* **2011**, *243*, 280-292, DOI: 10.1016/j.quaint.2010.10.006
12. Engovatova, A.; Golyeva, A. Anthropogenic soils in Yaroslavl (Central Russia): history, development, and landscape reconstruction. *Quat. Int.* **2012**, *265*, 54-62, DOI: 10.1016/j.quaint.2012.02.039
13. Jankowski, M.; Kittel, P. Evidences of soil-forming processes in red colored Ochre soils (Rubic Arenosols) at the Szynekielew archaeological site, Central Poland. In: *Proceedings 14th International Working Meeting on Soil Micromorphology*; Poch, R.M., Casamitjana, M., Francis, L.M., Eds.; Lleida, Spain, **2012**, pp.40-42.
14. Antisari, L.V.; Cremonini, S.; Desantis, P.; Calastri, C.; Vianello, G. Chemical characterisation of anthro-technosols from Bronze to Middle Age in Bologna (Italy). *J. Archaeol. Sci.*, **2013**, *40*(10), 3660-3671, DOI: 10.1016/j.jas.2013.04.023
15. Pawłowski, D.; Milecka, K.; Kittel, P.; Woszczyk, M.; Spychalski, W. Palaeoecological record of natural changes and human impact in a small river valley in Central Poland. *Quat. Int.* **2015**, *370*, 12-28, DOI: 10.1016/j.quaint.2014.06.033
16. Thy, P.; Willcox, G.; Barfod, G.H.; Fuller, D.Q. Anthropogenic origin of siliceous scoria droplets from Pleistocene and Holocene archaeological sites in northern Syria. *J. Archaeol. Sci.* **2015**, *54*, 193-209, DOI: 10.1016/j.jas.2014.11.027
17. Golyeva, A.; Zazovskaia, E.; Turova, I. Properties of ancient deeply transformed man-made soils (cultural layers) and their advances to classification by the example of Early Iron Age sites in Moscow Region. *Catena* **2016**, *137*, 605-610, DOI:10.1016/j.catena.2014.12.030
18. Shcherbakov, N.B.; Shuteleva, I.A.; Leonova, T.A.; Goleva, A.A.; Lunkova, Yu.V.; Kraeva, L.A. The ceramic complex as reflection of intercultural exchange of the Late Bronze Age tribes of the Southern Transurals. In: *Creativity: an exploration through the Bronze Age and contemporary responses to the Bronze Age*; University of Cambridge, Great Britain, **2013**, pp. 32-33.
19. Shcherbakov, N.B.; Leonova, T.A.; Shuteleva, I.A.; Golyeva, A.A. Anthropogenetic factor of landscape change in the process of early urbanization on the territory of Southern Transurals. In: *Conference on the Environmental Archaeology of European Cities (CEAEC)*. Brussel: Museum Royal Belgian Institute of Natural Sciences, **2015**, pp. 80-81.
20. Abdrakthmanov, R.F.; Martin, V.I.; Popov, V.G.; Roshdestvenskij, A.P.; Smirnov, A.I.; Travkin, A.I. *Karst of Bashkortostan*; Ufa-press: Ufa, Russia, 2002. ISBN: 5-902279-07-0. (In Russian).
21. Yaparov, I.M. *Atlas of the Bashkortostan Republic*; Kitap-press: Ufa, Russia, 2005, 486 p. (In Russian).

22. IUSS Working Group WRB. World Reference Base for Soil Resources 2014. In *International Soil Classification System for Naming Soils and Creating Legends for Soil Maps*; World Soil Resources Reports No. 106; Food and Agriculture Organization (FAO): Rome, Italy, 2014.
23. Bogomolov, L.V. *Soils of Bashkiria ASSR*. USSR Academy of Science press: Moscow, Russia, 1954, pp.1-296.
24. Khaziev, F. Kh. *The soil of the Republic of Bashkortostan and the regulation of their fertility*; Gilem-press: Ufa, Russia, 2007, pp. 1-288, ISBN: 978-5-7501-0892-3. (In Russian).
25. Obydenova, G.T.; Shcherbakov, N.B.; Shuteleva, I.A.; Khokhlova, O.S.; Kovaljuh, N.N.; Skripkin, V.V. Some results of the application of a complex approach at studying a monument of the Late Bronze Age in Bashkir Priuraljy – settlements Muradymovo. In: *Proceedings of II (XVIII) All-Russian archaeological congress in Suzdal*; Derevyanko, A.P., Makarov, N.A., Eds.; Institute of Archaeology of RAS: Moscow, Russia, 2008, pp. 435-438. (In Russian).
26. Shuteleva, I.A.; Shcherbakov, N.B.; Balonova, M.G.; Khokhlova, O.S.; Goleva, A.A. Some results of the application of a complex approach to the research of the Late Bronze age settlement in the Volgo-Ural region. *Interdisciplinaria archaeologica: natural sciences in archaeology* **2010**, 1 (1), 29-36. On-line publication: <http://www.iansa.eu>
27. Golyeva, A.; Khokhlova, O.; Shcherbakov, N.; Shuteleva, I. Negative effects of Bronze age human activity on modern soils and landscapes, a case-study on the Muradymovo settlement, Urals, Russia. *Interdisciplinaria Archaeologica. Natural sciences in Archaeology* **2016**, 2, VII, 169-178. On-line publication: <http://www.iansa.eu>
28. Arinushkina, E.V. *Guide on the chemical analysis of soils*. Moscow State University Publishing House, Moscow, 1970. (In Russian).
29. Poch, R.M.; Artiedo, O.; Herrero, J.; Verba-Lebedeva, M. Gypsic features. In: *Interpretation of Micromorphological Features of Soils and Regoliths*; Stoops, G., Marcelino, V., Mees, F., Eds.; Chapter 10. Elsevier: Netherlands, 2010, pp. 195–216, ISBN: 978-0-444-53156-8.
30. Vorobieva, L.A. *Chemical analysis of soils*. Moscow: Lomonosov Moscow State University Press. 1998. (In Russian).
31. Bullock, P.; Fedoroff, N.; Jongerius, A.; Stoops, G.; Tursina, T. *Handbook for soil thin section description*. Waiane Research, 1985, 152 p.
32. Afanasyeva, E.A. *Chernozems of the Middle Russian Plain*. Nauka press: Moscow, 1966; pp. 1- 223 (In Russian).
33. Khokhlova, O.S.; Kovalevskaya, I.S.; Oleynik, S.A. Records of climatic changes in the carbonate profiles of Russian Chernozems. *Catena* **2001**, 43 (3), 203-215, DOI: 10.1016/S0341-8162(00)00164-8
34. Khokhlova, O.S.; Kouznetsova, A.M. Carbonate accumulation morphology in a soil chronosequence in the southern Pre-Ural, Russia: Significance for Holocene paleoenvironmental reconstruction. *Revista mexicana de ciencias geológicas* **2004**, 21(1), 185-194. <http://rmcg.geociencias.unam.mx/revista/index.html>
35. Hamond, F.W. Phosphate analysis of archaeological sediments. *Landscape Archeology in Ireland. BAR British Series* **1983**, 116, 47-80.
36. Holliday, V.T.; Gartner, W.G. Methods of soil P analysis in archaeology. *J. Archaeol. Sci.* **2007**, 34, 301-333, DOI: 10.1016/j.jas.2006.05.004
37. Golyeva, A. Soil researches on settlement Sosnovka IV (Moscow region). In: *Archaeology of Moscow Region*; Engovatova, A., Ed.; Institute of Archaeology RAS: Moscow, Russia, 2014; Volume 10, pp. 64-74. ISBN: 978-5-94375-161-5
38. Verba, M.P.; Yamnova, I.A. Gypsum neoformations in nonirrigated and irrigated soils of the serozem zone. In: *Proc. of the Int. Working Meeting on Soil Micromorphology, Moscow, Russia, July 8-13, 1996*; Shoba, S., Gerasimova, M., Miedema, R., Eds.; Printing Service Centre Van Gils B.V. Wageningen, The Netherlands. 1997. P. 187-195.
39. Nettleton, W.D.; Nelson, R.E.; Brasher, B.R.; Derr, P.S. Gypsiferous soils in the western United States. In: *Soil Science Society of America Special Publication 10, Acid Sulfate Weathering*; Kittrick, J.A., Fanning, D.S., Hossner, L.R., Eds.; 1982, pp. 147-168. DOI: 10.2136/sssaspecpub10.c9
40. Minashina, N.G.; Shishov, L.L. Gypsum-Containing Soils: Their Distribution, Genesis, and Classification. *Eurasian Soil Sci.* **2002**, 35(3), 240-249.