

Article

The (A)symmetry between the Exterior and Interior of a Schwarzschild Black Hole

Pawel Gusin ¹, Andy Augousti ², Filip Formalik¹ and Andrzej Radosz ^{1*}

¹ Wroclaw University of Science and Technology, Wroclaw, Poland ; Pawel.Gusin@pwr.edu.pl; Filip.Formalik@pwr.edu.pl; Andrzej.Radosz@pwr.edu.pl

² Kingston University, London UK; augousti@kingston.ac.uk;

* Correspondence: Andrzej.Radosz@pwr.edu.pl; Tel.: +48-660-112-122

Abstract: A black hole in a Schwarzschild spacetime is considered. A transformation is proposed that describes the relationship between the coordinate systems exterior and interior to an event horizon. Application of this transformation permits considerations of the (a)symmetry of a range of phenomena taking place on both sides of the event horizon. The paper investigates two distinct problems of a uniformly accelerated particle. In one of these, although the equations of motion are the same in the regions on both sides, the solutions turn out to be very different. This manifests the differences of the properties of these two ranges.

Keywords: black hole; event horizon; asymmetry

1. Introduction

Outside the horizon of a black hole the Killing vector is time-like and it becomes space-like inside the horizon –this means that energy conservation is broken. Such a conversion of the Killing vector results in a lot of interesting phenomena, Hawking radiation and the information paradox being the most prominent among them [1,2]. The status of these two outcomes arising from the presence of the event horizon is, in a sense, similar: both appeared mainly due to the work of Hawking and both of them are still debated [3]. More recent studies have considered a range of interesting questions. One of the latest was the supposition that black holes (BH) do indeed have hair [4]; another interesting outcome was the claim that the volume of a BH may be infinite [5]. In view of the argument that there is not enough space inside a black hole to store arbitrarily large amounts of information, this discovery turns out to be a contribution strictly associated with the problem of information stored and lost within an event horizon[6]. Following a proposal by Unruh [7] of development of analogues gravity, condensed, soft matter or photonic black hole-like systems that develop an event horizon have been experimentally realized [8]. A recent experiment has led to a Kerr-like BH including an ergo-sphere, not just an event horizon, opening up the possibility of experimental observations of the Penrose effect.

The interior of a black hole itself has been the subject of fewer but more diverse considerations. Hamilton et al. [9] presented a discussion on the question of (spectroscopic) vision inside the horizon of a Schwarzschild BH. In another approach Hamilton and Leslie [10] considered the exterior and interior of a BH within a so-called river model based on the use of a Painleve-Gullstrand co-moving frame. It is well-known that the static spacetime outside a BH's horizon turns out to be dynamic inside the outer horizon in the situation when two horizons are formed. Such a dynamic interior of a BH may be viewed as a cosmological model [11].

The aim of this paper is to study the properties of the interior of a Schwarzschild BH, i.e. which is isotropic and static outside the horizon. The investigations are presented from a special perspective. One finds a formal, specific symmetry between the exterior and interior ranges, even though the properties of these two regions are found to be completely different. We illustrate the differences using two problems related to the behaviour of a test particle uniformly accelerated outside and inside the horizon. In the first example the particle moves radially outside the horizon and its natural

extension is motion along a homogeneity axis (see below) inside the horizon. It is found then that although the equations of motion are the same in both regions the solutions differ significantly. In the other example a test particle follows a circular trajectory belonging to the photon sphere [12] and inside the horizon the test particle moves along a photon sphere analogy.

The paper is organized as follows. In Sec.2 we present a transformation between coordinate systems outside and inside horizon of the Schwarzschild black hole. Section 3 is devoted for thorough investigation of the problem of a uniformly accelerated motion of along a straight line: radially outside horizon along homogeneity direction inside horizon. In Sec. 4 we consider the case of uniform acceleration for motion along a circle on a photon sphere outside horizon and its analogue inside horizon. The discussion and final remarks are given in the last section.

2. Systems of co-ordinates in space-time with a horizon

Arbitrary co-ordinate system constitute a diffeomorphism Φ from an open set U of the space-time M into an open set U' of the real Euclidean flat space R^n , where n is the dimension of M :

$$\begin{aligned}\Phi: U \subset M &\rightarrow U' \subset R^n \\ \Phi(p) &= (x^1, \dots, x^n) \text{ where } p \in U.\end{aligned}\quad (1)$$

We will consider spherically symmetric space-time with a black hole defined by a horizon. The horizon H which is a global property of such a space-time, divides M into an exterior M_+ and an interior M_- :

$$M = M_- \cup H \cup M_+. \quad (2)$$

On M_+ (which is static) one can introduce the coordinate system which in the spherical coordinates (r, θ, φ) on R^3 takes the form:

$$\begin{aligned}\Phi_+: M_+ &\rightarrow R^4 \\ \Phi_+(p) &= (t, r, \theta, \varphi) = (x^\mu),\end{aligned}\quad (3)$$

where $r > r_g$ and r_g is a (gravitational) radius of the horizon H . The metric ds_+^2 on M_+ in this framework has the form

$$ds_+^2 = \left(1 - \frac{r_g}{r}\right) dt^2 - \left(1 - \frac{r_g}{r}\right)^{-1} dr^2 - r^2 d\Omega^2. \quad (4)$$

This is the well-known Schwarzschild space-time with $r_g = 2M$, and $d\Omega^2 = d\theta^2 + \sin^2\theta d\varphi^2$ is a unit metric on S^2 .

On M_- one can introduce the following coordinate system:

$$\begin{aligned}\Phi_-: M_- &\rightarrow R^4 \\ \Phi_-(p) &= (T, R, \theta, \varphi) = (X^\alpha),\end{aligned}\quad (5)$$

where $0 < T < r_g$. The metric ds_-^2 on M_- in this coordinate system has the following form (see [11])

$$ds_-^2 = \frac{dT^2}{-1 + \frac{r_g}{T}} - \left(\frac{r_g}{T} - 1\right) dR^2 - T^2 d\Omega^2. \quad (6)$$

Hence, on M there are two coordinate systems with two metrics $ds_+^2 = g_{\alpha\beta}(x)dx^\alpha dx^\beta$ and $ds_-^2 = h_{\alpha\beta}(X)dX^\alpha dX^\beta$. The transformation \mathbb{T} between these two systems of coordinates has the form:

$$\mathbb{T}: M_+ \rightarrow M_-, \quad (7)$$

where \mathbb{T} is given by the matrix:

$$\mathbb{T} = \begin{pmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} = \mathbb{T}^{-1}. \quad (8)$$

It acts as follows:

$$\begin{aligned}\mathbb{T}x &= X, \\ T = r, \quad R = t, \quad \theta = \theta, \quad \varphi = \varphi,\end{aligned}\quad (9)$$

and

$$h(X) = \mathbb{T}g(\mathbb{T}x)\mathbb{T}, \quad (10)$$

where $h(X) = (h_{\alpha\beta}(X))$ and $g(\mathbb{T}x) = (g_{\alpha\beta}(\mathbb{T}x))$.

There are two Killing vectors manifesting the symmetry properties of the exterior M_+ and interior M_- of the horizon. In M_+ these are a time-like one, ∂_t representing energy conservation due to the static character of the spacetime and a space-like one, ∂_φ reflecting angular momentum

conservation. In M_- there is a space-like one, ∂_R reflecting R -momentum conservation due to homogeneity along the R -axis and another space-like one, ∂_φ representing angular momentum conservation.

The exterior of the Schwarzschild black hole, M_+ is a static spacetime and its interior, M_- is a dynamic spacetime. One can introduce a class of static observers in the former and a class of resting observers in the latter case. The velocity vector of such observers has only a temporal non-vanishing coordinate $U = \alpha \partial_0$. It appears natural to label them as:

a) t-observers

$$U_t = \frac{1}{\sqrt{g_{tt}}} \partial_t \quad (11)$$

outside the horizon and

b) T-observers

$$U_T = -\frac{1}{\sqrt{h_{TT}}} \partial_T \quad (12)$$

inside the horizon.

3. Uniformly accelerated motion along straight line

In this section we will consider the problem of a uniformly accelerated test particle in the exterior and interior of the horizon. The motion of such a test particle will be confined to the x^0, x^1 and X^0, X^1 hyperplanes respectively. So one can discuss the case of a 1+1 dimensional spacetime

$$ds^2 = f(d\xi^0)^2 - f^{-1}(d\xi^1)^2 \quad (13)$$

such that

a) Outside the horizon, $\xi^\alpha \equiv x^\alpha$ and $f = g_{tt}$ is a function of spatial coordinate ξ^1

b) Inside the horizon, $\xi^\alpha \equiv X^\alpha$ and $f = h_{TT}$ is a function of temporal coordinate ξ^0 .

The components of a velocity vector u

$$u = u^\alpha \partial_\alpha \quad (14)$$

of the test particle, u^α will depend on ξ^1 in the case (a) and on ξ^0 in case (b).

An acceleration vector field a for u is:

$$a = \nabla_u u \quad (15)$$

and one obtains the following equations for its components:

a) outside the horizon

$$a^t = -g_{rr} u^r \frac{d}{dr} (g_{tt} u^t) \quad (16)$$

$$a^r = g_{tt} u^t \frac{d}{dr} (g_{tt} u^t) \quad (17)$$

and

b) inside the horizon

$$\bar{a}^T = h_{RR} u^R \frac{d}{dT} (h_{RR} u^R) \quad (18)$$

$$\bar{a}^R = -h_{TT} u^T \frac{d}{dT} (h_{RR} u^R) \quad (19)$$

Hence, one finds that these two pairs of equations are transformed into each other under the interchange of the temporal and spatial coordinates, $t \leftrightarrow R$ and $r \leftrightarrow T$ as emphasized in the former section (see Eq. (9)). Uniform acceleration is defined by the condition:

$$a^2 = f(a^0)^2 - f^{-1}(a^1)^2 = -\alpha^2 \quad (20)$$

where $\alpha = \text{const}$. The equations for the world line of the test particle are then derived as follows.

a) Outside the horizon

$$\frac{d}{dr} (g_{tt} u^t) = \pm \alpha. \quad (21)$$

Thus the world line $\gamma = \{t(\tau), r(\tau)\}$ of the uniformly accelerated particle is given in the case of a static metric by the integral curve of the vector field:

$$u^t = \frac{dt}{d\tau} = \frac{1}{g_{tt}} (E \pm \alpha \int dr) \quad (22)$$

$$u^r = \frac{dr}{d\tau} = \pm \sqrt{(E \pm \alpha \int dr)^2 - g_{tt}} \quad (23)$$

where E is an integration constant.

b) Inside the horizon

$$\frac{d}{dT}(h_{RR}u^R) = \pm\alpha. \quad (24)$$

In this case of a dynamic spacetime the world line of the test particle is given as follows:

$$u^R = \pm \frac{1}{h_{RR}} (E' \pm \alpha \int dT) \quad (25)$$

$$u^T = \sqrt{(E' \pm \alpha \int dT)^2 - h_{RR}}. \quad (26)$$

Applying the fact that inside the horizon

$$g_{tt} = 1 - \frac{r_g}{r} = h_{RR}(r) \quad (27)$$

one finds that the world line equations for the outer and inner horizon regions, Eqs. (22,23) and Eqs. (25, 26) take the same form:

$$\dot{t} = \frac{E+\alpha r}{1-\frac{r_g}{r}} \quad (28)$$

$$\dot{r} = \sqrt{(E + \alpha r)^2 - \left(1 - \frac{r_g}{r}\right)}. \quad (29)$$

Here we will examine the following point: how does the speed of a uniformly accelerated test particle change with respect to the static observers? In general a relative velocity four-vector w of u' as measured by another observer characterized by velocity u at the same event $p = (x^\alpha)$ is given by (see [13])

$$w = \frac{u'}{(u,u)} - u, \quad (30)$$

where (u,u') is the scalar product of the two velocity four-vectors, u and u' , $u^2 = u'^2 = 1$. Velocity w is orthogonal to u , $(w,u) = 0$, i.e. w is a space-like vector,

$$w^2 = \frac{1}{(u,u')^2} - 1 < 0. \quad (31)$$

Then the squared speed v^2 is given as:

$$v^2 = -w^2 > 0. \quad (32)$$

3.1. Black hole exterior

In the exterior of the Schwarzschild black hole the spacetime is static and static observers denoted as t -o are characterized by their velocity vector U_t (see Eq. (11)). Such an observer measures the speed v of a nearby passing uniformly accelerated (ua) test particle,

$$u_{ua} = \dot{t}\partial_t + \dot{r}\partial_r. \quad (33)$$

Hence, for $u = U_t$, and $u' = u_{ua}$ one applies the procedure described above to find:

$$w^2 = -\frac{\dot{r}^2}{(g_{tt}\dot{t})^2}. \quad (34)$$

Using Eqs. (28,29) we obtain

$$v^2 = \frac{(E+\alpha r)^2 - g_{tt}}{(E+\alpha r)^2}. \quad (35)$$

This case of a uniformly accelerated test particle initially at rest, $r = 2r_g$ is illustrated in Figure 1.




Figure 1. Squared speed v^2 (Eq. (35)) of a uniformly accelerated test particle initially at rest at $r_0 = 2r_g \equiv 4$ (in this case $r_g \equiv 2$) escaping radially from the gravitational field for different values of $\alpha=0.1$ (red), $\alpha=0.5$ (green), $\alpha=1$ (black)

3.2. Black hole interior

Inside the horizon, $h_{TT} = \left(\frac{r_g}{r} - 1\right)^{-1}$. A class of resting (or co-moving, see below) observers is determined by (12)

$$U_T = -\sqrt{\frac{r_g}{r} - 1} \partial_T. \quad (36)$$

Using Eqs. (26), (36), (32) one obtains the squared speed \tilde{v}^2 of the uniformly accelerated test particle inside the horizon:

$$\tilde{v}^2 = \frac{(E' + \alpha T)^2}{(E' + \alpha T)^2 - \left(1 - \frac{r_g}{r}\right)}. \quad (37)$$

Using the relation $T = r$ inside the horizon, one can express the squared speed \tilde{v}^2 as a function of a temporal coordinate, r . In Figure 2, the squared speed of the uniformly accelerated test particle initially at rest is illustrated by applying expression (37). The speed initially increases, reaches some (non-universal) maximal value and finally decreases to zero at the ultimate singularity, $r \rightarrow 0$.




Figure 2. Squared speed \tilde{v}^2 (Eq. (37)) a test particle initially, $T_0 = 0.9 r_g$ at rest (in this case $r_g \equiv 10$) uniformly accelerated along homogeneity axis $R(=t)$ for different values of $\alpha=0.05$ (blue), $\alpha=0.1$ (red), $\alpha=0.2$ (green), $\alpha=1$ (black).

4. Uniform acceleration on a photon sphere and on its analogue inside the horizon

A photon sphere outside the horizon, $r > r_g$, has a radius, $r_{ps} = \frac{3}{2}r_g$. It consists of null geodesic circles, $k = k^t \partial_t + k^\varphi \partial_\varphi$, $k^2 = 0$ ($r = r_{ps}$). Its analogy inside the horizon, $r < r_g$, consists of planar null geodesics having two non-vanishing components, $k = k^r \partial_r + k^\varphi \partial_\varphi$. However, in this case the radius, r of such a “sphere” diminishes from r_g to 0. We shall consider the problem of a uniformly accelerated test particle on a circle belonging to the photon sphere and on a “circle” belonging to the photon sphere analogue.

4.1. Black hole exterior

In a case of a test particle moving outside the horizon along a circle of radius $r_0 \geq r_{ps}$ the velocity vector is two-component

$$u = u^t \partial_t + u^\varphi \partial_\varphi, \quad (38)$$

where without the lost of generality, we have considered a circle in an equatorial plane, $\theta = \pi/2$. An acceleration vector field $a = \nabla_u u$ in the case of motion with constant speed is found to be $a = a^r \partial_r$ (see [12])

$$a^r = -\frac{1}{2} \left[(u^t)^2 \frac{dg_{tt}}{dr} + (u^\varphi)^2 \frac{dg_{\varphi\varphi}}{dr} \right]. \quad (39)$$

Applying a normalization condition, $u^2 = 1$, one finds that the acceleration of a photon sphere, $r_0 = r_{ps} = \frac{3}{2}r_g$, turns out to be speed independent [12]

$$a^r = -\frac{1}{2} \frac{1}{g_{tt}} \frac{dg_{tt}}{dr} = \frac{2}{3r_g}. \quad (40)$$

Hence in this case one can consider a generalization of the twin-paradox and other features such as specific non-geodesic motion [14].

4.2. Black hole interior

Motion along a photon sphere analogue in this case, $r < r_g$ is as follows.

The two-component velocity vector of a test particle takes the form:

$$u = u^r \partial_r + u^\varphi \partial_\varphi, \quad (41)$$

and its components will be r -dependent. An acceleration vector field $a = \nabla_u u$ is found to be two-component,

$$a = a^r \partial_r + a^\varphi \partial_\varphi, \quad (42)$$

and its components are:

$$a^r = -\frac{u^\varphi}{g_{rr}} \frac{d}{dr} (g_{\varphi\varphi} u^\varphi), \quad (43)$$

$$a^\varphi = \frac{u^r}{g_{\varphi\varphi}} \frac{d}{dr} (g_{\varphi\varphi} u^\varphi). \quad (44)$$

Uniform acceleration is defined by the condition (see Eq. (13))

$$a^2 = g_{rr} (a^r)^2 + g_{\varphi\varphi} (a^\varphi)^2 = -\beta^2 = \text{const.} \quad (45)$$

The equations for the world line of the test particle are then derived as follows. Applying Eqs. (45), (43), (44) one finds,

$$\frac{d}{dr} (g_{\varphi\varphi} u^\varphi) = \pm \beta \sqrt{-g_{rr} g_{\varphi\varphi}}. \quad (46)$$

Thus the world line $\gamma = \{r(\tau), \varphi(\tau)\}$ of the test particle uniformly accelerated on a sphere of ever diminishing radius, is given by an integral curve of the vector field u :

$$u^r = \frac{dr}{d\tau} = \pm \frac{1}{\sqrt{g_{rr}}} \left(1 - \frac{1}{g_{\varphi\varphi}} \left(C \pm \beta \int \sqrt{-g_{rr} g_{\varphi\varphi}} dr \right)^2 \right), \quad (47)$$

$$u^\varphi = \frac{d\varphi}{d\tau} = \frac{1}{g_{\varphi\varphi}} \left(C \pm \beta \int \sqrt{-g_{rr} g_{\varphi\varphi}} dr \right), \quad (48)$$

where C is an integration constant. Here we are interested in the following question (see above): how does the speed of a uniformly accelerated test particle change with respect to the static (resting) observers? Similarly, as in the former section, the speed is measured by (resting) T – observers, Eq. (36). In this case one finds:

$$\tilde{v}^2 = 1 - \frac{1}{g_{rr} (u^r)^2} = \frac{(C \pm \beta \int \sqrt{-g_{rr} g_{\varphi\varphi}} dr)^2}{(C \pm \beta \int \sqrt{-g_{rr} g_{\varphi\varphi}} dr)^2 - g_{\varphi\varphi}}. \quad (49)$$

The meaning of the result (49) is: the speed of the uniformly accelerated test particle, of total acceleration β tends to the speed of light as the ultimate singularity, $r \rightarrow 0$ approaches, as $g_{\varphi\varphi} = -r^2 \rightarrow 0$.

5. Discussion and final remarks

In order to investigate the properties of the interior of a Schwarzschild black hole one can apply a specific symmetry to the coordinate systems of its exterior and interior. The meaning of such a symmetry is that the radial and temporal coordinates interchange their roles. This symmetry may be regarded as a justification of the fact that one can apply a Schwarzschild system of coordinates, (t, r, θ, φ) both outside as well as inside the horizon. It is an almost trivial notion but it is important to remember the existence of a singularity in this system, which is the horizon itself. In this paper we

have considered the application of this interchange of roles – a symmetry which is particular to the case of a uniformly accelerated particle. In the case of radial motion, outside the horizon and motion along the direction of homogeneity inside the horizon, one finds that the equations of motion on both sides of the horizon are of the same form. In the case of circular motion on a photon sphere and “circular” motion on a photon sphere analogue, outside and inside the horizon, respectively, the equations of motion are found to be different. In both cases, with either the same or different equations of motion, one gets the very different solutions. Outside the horizon one reproduces more or less well-known outcomes; inside the horizon the outcomes are found to be rather unexpected. Outside the horizon a radially accelerated particle departs with ever increasing speed (if the acceleration is larger than some critical value); the speed of a test particle uniformly accelerated along the homogeneity axis inside the horizon, whose equation of motion is the same as the one outside the horizon, initially increases but then decreases to zero when approaching the ultimate singularity. The radial acceleration during circular motion of the test particle on the photon sphere is independent of its speed; the speed of a test particle following accelerated motion along a circle belonging to the photon sphere analogue (of ever decreasing radius) increases to the speed of light when approaching the ultimate singularity. Having in mind an (intimate) symmetry (7) between the exterior and interior of the Schwarzschild black hole leading in the simplest case to the same equation of motion, one may wonder how it arises that the properties of the solutions are so different. The answer is: it is because r and t , spatial (radial) and temporal coordinates, respectively, outside horizon interchange their roles, becoming temporal and spatial (in the direction of homogeneity) coordinates inside the horizon. Such an exchange of roles has a deeper consequence: outside the horizon spacetime is spherically symmetric and static and inside the horizon it is no longer static, it is dynamically changing (ie an “anisotropic cosmology”, see [11]) but homogeneous along one of its spatial directions.

References

1. S.W. Hawking, Black hole explosions? *Nature* **1974**, *248*, 30-31
2. S.W. Hawking, Breakdown of predictability in gravitational collapse. *Phys. Rev. D*, **1976** *14*, 2460-2473
3. A. Almheiri, D. Marolf, J. Polchinski, J. Sully, Black holes: complementarity or firewalls? *JHEP* **02** **2013** 062, 1-19
4. S.W. Hawking, M. Perry, A. Strominger, Soft Hair on Black Holes, **2016** *PRL* *116*, 231301 1-9
5. M. Christodoulou, C. Rovelli, How big is a black hole? *Phys. Rev. D* **2015**, *91*, 064046 1-7
6. S. Mathur, A model with no firewall. **2015** [arXiv:1506.04342](https://arxiv.org/abs/1506.04342)
7. W.G. Unruh, Experimental Black-Hole Evaporation? *PRL* **1981**, *46*, 1351-1353
8. H.S. Nguyen, et al., Acoustic Black Hole in a Stationary Hydrodynamic Flow of Microcavity Polaritons, *PRL* **2015**, *114*, 036402 1-5
9. A.J.S. Hamilton, and G. Polhemus, Stereoscopic visualization in curved spacetime: seeing deep inside a black hole. *New J. Phys.* **2010** *12*, 123027-123052
10. A. J. S. Hamilton and J. P. Lisle The river model of black holes, *Am.J.Phys.* **2008** *76*: 519-532
11. R. Doran, F.S.N. Lobo and P.Crawford, Interior of a Schwarzschild black hole revisited, *Foundations of Physics* **2008**, *38*, 160–187
12. M. A. Abramowicz, S. Bajtlik and W. Kluzniak: The twin paradox on the photon sphere, *Phys. Rev. A* **2007** *75*, 044101 1-2
13. V.J. Bolos, Intrinsic Definitions of "Relative Velocity" in General Relativity, *Comm. Math. Phys.* **2007** *273*, 217--236
14. M. A. Abramowicz and S. Bajtlik: Adding to the paradox: the accelerated twin is older, [arXiv:0905.2428v1](https://arxiv.org/abs/0905.2428v1) 2009