

Article

Land Use Planning and Wildlife-Inflicted Crop Damage in Zambia

Mitelo Subakanya¹, Gelson Tembo² and Robert B. Richardson^{3,*}

¹ Indaba Agricultural Policy Research Institute, Lusaka, Zambia; mitelo.subakanya@iapri.org.zm

² Department of Agricultural Economics and Extension, University of Zambia, Lusaka Zambia; tembogel@unza.zm

³ Department of Community Sustainability, Michigan State University, East Lansing, MI, USA; rbr@msu.edu

* Correspondence: rbr@msu.edu; Tel: +1 (517) 355-9533

Abstract: Damage to crops from wildlife interference is a common threat to food security among rural communities in or near Game Management Areas (GMAs) in Zambia. This study uses a two-stage model and cross-sectional data from a survey of 2,769 households to determine the impact of land use planning on the probability and extent of wildlife-inflicted crop damage. The results show that crop damage is higher in GMAs as compared to non-GMAs, and that land use planning could be an effective tool to significantly reduce the likelihood of such damage. These findings suggest that there is merit in the current drive to develop and implement land use plans as means to minimize human-wildlife conflict such as crop damage. This is especially critical as Zambian conservation policies do not have an explicit provision for compensation in the event of damage from wildlife.

Keywords: Land use planning; agriculture; crop damage; Game Management Areas; human-wildlife conflict; wildlife; Zambia

1. Introduction

In sub-Saharan Africa, the basic needs of most rural poor households—such as food, water, fuel, clothing and shelter—are usually met by natural resources and ecosystem services from the land around them. However, the increase in population growth means that the capacity of the land to support increasing demands continues to diminish [1,2]. In areas around game reserves, agriculture also has to compete with wildlife for resources. The ever-increasing demand for land among competing uses suggests a need for land use planning in order to maximize benefits, minimize losses, and avoid conflict.

The objective of a land use plan is to select and put into practice those land uses that will best meet the needs of the people while safeguarding resources for the future [1]. In Zambia, land use planning in Game Management Areas (GMAs) has been promoted and facilitated by the former Zambia Wildlife Authority (ZAWA) since around 1998 (in 2015, ZAWA was replaced by the Department of National Parks and Wildlife). Land use planning is a consultative process involving stakeholders including the local communities which shows how a particular GMA should be managed [3]. Land use planning has the benefit of increasing agricultural production [4] due to the decrease in wildlife-inflicted damage to the crops in the protected areas. Land use planning is a very important step that should be taken if wildlife conservation is to be effective, because a land use plan will identify appropriate land uses around the GMA [5].

Protected areas such as GMAs were formed to promote wildlife conservation [6] while giving benefits to the rural communities at the same time [7]. Yet, empirical evidence suggests that protected areas like the GMAs are associated with high incidence of wildlife-inflicted crop damage [8-16], arguably contributing to rural poverty. This problem is common to almost all problems where conservation is promoted alongside human habitats. In many several conservation areas of Nepal, for example, wildlife-inflicted crop damage is rampant [15,17]. Like in Zambia, most of the crop damage in developing countries is caused by elephants [10,15,17,18] and almost virtually uncompensated [8].

Alternative methods used by households living in GMAs to drive away wildlife from their agricultural fields such as drum beating, fire, and physically guarding the fields, have proved largely ineffective [15,19]. Electric fencing also proved ineffective in Zambia's Luangwa National Park and scouts could help in some instances but were very slow in responding to the problem of crop damage by wildlife [10]. To many planners, this identifies the need to develop and implement new and more effective ways of solving the problem [11]; ones that can enhance conservation without significantly jeopardizing human livelihoods. Some contend that land use planning should be an important member of such a solution set [20].

However, although conventional wisdom and anecdotal evidence suggest that land use planning could provide the much-needed long term solution to crop damage [9,21,22], evidence of its effectiveness is still very scanty. Like Zambia, many countries in Sub-Saharan Africa are either just turning to land use planning or already implementing some forms of land use plans; all of which could benefit immensely from research on the effectiveness of land use planning. Namibia and Zimbabwe, for example, are some of the examples of countries that are already actively implementing land use planning [23].¹

The study reported in this paper uses survey data from Zambia's GMAs to examine the impact of land use planning on wildlife-inflicted crop damage. To the best of our knowledge, no study has done this before. This report is arranged as follows. Section two provides background on wildlife conservation and GMAs in Zambia. Section three focuses on the conceptual framework while the methods are discussed in section four, which include data and data sources and the model specification. Results and discussion are presented in section five, and finally, the summary and conclusions are presented in section six.

2. Game Management Areas in Zambia

GMAs were declared in the National Parks and Wildlife Act in 1971 where protected areas were reduced to two categories: National Parks and GMAs. This Act was then replaced by the National Parks and Wildlife Act No. 10 of 1991 and later the Zambia Wildlife Act No.12 of 1998 which established ZAWA. It was also in this Act that the inclusion of local community participation in wildlife conservation was done. ZAWA was the agency responsible for the management of the National Parks and GMAs, with GMAs managed in partnership with the communities through the Community-Based Natural Resource Management (CBNRM) program. A share of revenue collected from hunting licenses is used by the rural communities to employ village scouts and to support development projects such as schools and clinics through the Community Resource Boards (CRBs) and the Village Action Groups (VAGs) [8,24].

Even with the positive benefits of the CBNRM program (such as creation of employment opportunities [25]), problems like crop damage can lead to negative impacts. Currently, policies do not give any form of compensation to rural farming households that live in the GMAs if wildlife causes damage to their crops [8].

At the time of this study, there were 36 GMAs and 19 national parks distributed around the country (see Figure 1). GMAs are defined as buffer zones around national parks in which licensed safari and subsistence hunting is permitted [24]. ZAWA used revenue collected from hunting as a basis for GMA categorization [8]. Four categories of GMAs were identified in this study: i) prime, ii) secondary, iii) specialized, and iv) under-stocked. In prime GMAs, there is an abundance and diversity of wildlife species while in the secondary GMAs wildlife species are less abundant. Sustained hunting can take place in both of these GMA types. Specialized GMAs have fewer numbers of wildlife species and are found in wetland areas. As the name suggests, the fourth category, under-

¹ In Zambia, preparation of land use plans involves the standard format determined by ZAWA and undergoes a strategic planning process, which addresses ecologically complex areas as influenced by ecological and socio-economic forces surrounding the protected areas [3].

stocked, refers to GMAs that have suffered losses to wildlife abundance and diversity, primarily due to uncontrolled hunting and poaching.




Figure 1. Map depicting Game Management Areas in Zambia [26]

3. Conceptual Framework

Crop damage from wildlife is a big challenge that farmers face in areas where wildlife and humans have to co-exist. Communities participate in wildlife conservation in Zambia through representation in CRBs and VAGs. Although the literature has demonstrated that communities do benefit from CBNRM programs (see [24]), losses from crop damage from wildlife can have deleterious effects in rural food security. A land use plan, by design, is expected to influence the way people use their land [1] and to respect land set aside for wildlife. In Zambia, preparation of land use plans includes a Strategic planning process that considers the ecological and socioeconomic dynamics of the protected areas.

Land use planning process involves participation from different relevant stakeholders including the rural community. The inclusion of the local community members in land use planning is important, as they better understand the local context and social-ecological dynamics. Ostrom [27] pointed out how community members can organize themselves to achieve certain goals such as land use planning. Thus, communities that have developed land use plans are expected to experience less human-wildlife conflict and to suffer less crop damage compared to those that have not developed any land use plan.

Even with the social and financial benefits of the CBNRM program, problems like crop damage from wildlife can lead to negative welfare impacts. Currently, policies do not allow for any form of compensation to rural farming households that live in the GMAs in the event of wildlife-inflicted crop damage [8], unlike in other countries such as Botswana where communities are compensated [28]. Land use planning is sporadic among VAGs and CRBs, as the practice has not been promoted widely. The existence of land use plans in GMAs allows for the examination of their impact on the probability and extent of losses from wildlife-inflicted crop damage.

4. Methods

4.1. Data

Nationally representative data covering 2,769 households from the Impact of Game Management Areas on Household Welfare survey (IGMAW) was used which covered areas adjacent to four national park systems - Bangweulu (including Isangano, Lavushi and Kasanka National Parks), Kafue (including Kafue, Blue Lagoon and Lochinvar National Parks), Lower Zambezi (Lower Zambezi National Park) and Luangwa (including North and South Luangwa National Parks). (For more details on the sampling procedure see [8,24]. The survey data were also supplemented with secondary data collected from various ZAWA documents.

4.2. Model specification

The Cragg [29] tobit alternative framework was used to model the probability and extent of crop damage. This method involves doing the estimation in two stages, where a probit model is used in the first stage to estimate the probability of crop damage, and a truncated regression is used in the second stage to estimate the extent of crop damage among those that have suffered the problem. The two-stage model can be represented as follows:

$$\Pr(c_i = 1 | \mathbf{x}_i) = \boldsymbol{\beta}'\mathbf{x}_i + \boldsymbol{\tau}'\mathbf{g}_i + \delta P_i + \mu_i \quad (1)$$

$$\ln y_i = \boldsymbol{\gamma}'\mathbf{x}_i + \boldsymbol{\psi}'\mathbf{g}_i + \lambda P_i + \varepsilon_i, \quad (2)$$

where c is a crop damage dummy variable taking a value of 1 if the household had suffered crop damage during the 2004/05 agricultural season, y_i is the extent of crop losses incurred, \mathbf{x} is a vector of covariates postulated to explain crop damage, P is a land use dummy variable equal to 1 if the community had already developed a land use plan, \mathbf{g} is a vector of GMA dummy variables equal to 1 for each type of GMA, δ and λ are coefficients to be estimated for the land use plan dummy variable, $\boldsymbol{\tau}$ and $\boldsymbol{\psi}$ are vectors of parameters to be estimated corresponding to the GMA dummy variables, $\boldsymbol{\beta}$ and $\boldsymbol{\gamma}$ are vectors of other parameters to be estimated, and $\mu_i \sim N(0, \sigma_\mu^2)$ and $\varepsilon_i \sim N(0, \sigma_\varepsilon^2)$ are mean-zero normally distributed random error terms.

Although there are four types of GMAs, the study did not include any depleted GMAs because hardly any wildlife can be found for hunting purposes. Furthermore, in our specification, GMA categories secondary and specialized were collapsed into one category. This is so because there were very few specialized GMAs in our data. Thus, in the end vector \mathbf{g} ended up with only two members, g_1 (for prime GMAs) and g_2 (for secondary and specialized GMAs).

Because extent of crop damage and, hence, the effectiveness of a land use plan are likely to be greatest in GMAs with large wildlife populations, we also model this differential effect of land use plans. This is done by introducing in both equations (1) and (2) interaction terms between the GMA effect and the effect of the land use plan:

$$\Pr(c_i = 1 | \mathbf{x}_i) = \boldsymbol{\beta}'\mathbf{x}_i + \boldsymbol{\tau}'\mathbf{g}_i + \delta P_i + \theta_1 (g_1 * P)_i + \theta_2 (g_2 * P)_i + \mu_i \quad (3)$$

$$\ln y_i = \boldsymbol{\gamma}'\mathbf{x}_i + \boldsymbol{\psi}'\mathbf{g}_i + \lambda P_i + \pi_1 (g_1 * P)_i + \pi_2 (g_2 * P)_i + \varepsilon_i, \quad (4)$$

where $\boldsymbol{\theta} = \{\theta_1, \theta_2\}$ and $\boldsymbol{\pi} = \{\pi_1, \pi_2\}$ are vectors of parameters to be estimated for the interaction terms.

In this latter specification (Equations 3 and 4), the effect of the land use plan is allowed to vary by wildlife numbers, represented here by the GMA dummy variables. Thus, the effect of land use planning on the probability of incurring crop damage could be derived from Equation (3) as:

$$\frac{\Delta \Pr(c_i = 1 | \mathbf{x}_i)}{\Delta P_i} = \begin{cases} \hat{\delta} + \hat{\theta}_1, & \text{if } g_1 = 1 \text{ and } g_2 = 0 \\ \hat{\delta} + \hat{\theta}_2, & \text{if } g_1 = 0 \text{ and } g_2 = 1 \\ \hat{\delta} & \text{if } g_1 = 0 \text{ and } g_2 = 0. \end{cases} \quad (5)$$

Similarly, the effect of land use planning on the extent of crop damage can be derived from Equation (4) as:

$$\frac{\Delta E(\ln y_i | \mathbf{x}_i)}{\Delta P_i^{0 \rightarrow 1}} = \begin{cases} \hat{\lambda} + \hat{\pi}_1, & \text{if } g1 = 1 \text{ and } g2 = 0 \\ \hat{\lambda} + \hat{\pi}_2, & \text{if } g1 = 0 \text{ and } g2 = 1 \\ \hat{\lambda} & \text{if } g1 = 0 \text{ and } g2 = 0. \end{cases} \quad (6)$$

All the models were tested for standard model specification problems such as heteroskedasticity and multicollinearity. Having detected heteroskedasticity at 5 percent level, we use heteroskedasticity-consistent standard errors in all the estimations. We include in vector \mathbf{x} all the variables postulated to affect crop damage [8]. Table 1 summarizes these variables and their postulated relationships with crop damage.

Table 1 presents the members of vector \mathbf{x} and proxies for ρ and the expected directions of their relationships with probability of crop damage. Several human capital variables are expected to be correlated with the likelihood of suffering and extent of wildlife-inflicted crop damage. Age, for example, was a proxy for experience. The more experienced the household head is, the more knowledgeable we expect them to be about wildlife and how to minimize human-wildlife conflict, including crop damage. However, younger heads may have the strength needed to thwart the actions of wildlife. Thus, we expect either sign on this variable. We also expect male-headed households to have greater ability to deal with wildlife as activities such as wildlife scaring (and snaring) are typically in the male domain in these communities.

Table 1. Covariates postulated to determine crop damage.

Variable	Expected sign
Age of household head in years	Positive/Negative
Sex of household head, 1=male	Negative
Maximum education (in years)	Negative
Household size	Negative
Distance to the nearest main road in kilometers	Positive
Total area cropped in hectares	Positive
Value of consumer assets in ZMK	Negative
Value of productive assets in ZMK	Negative/Positive
Infrastructure index	Negative
Population density per square km	Negative
Number of Scouts	Negative/Positive
Prime GMA, 1=yes (gma1)	Positive
Secondary/specialized GMA, 1=yes (gma2)	Positive
Value of harvest in ZMK	Positive
Land use plan, 1=yes (landuse)	Negative

Other human capital variables expected to be negatively correlated with crop damage include education of the head and labor supply as proxied by household size. More educated people are expected to have greater ability to analyze situations and to develop effective strategies for dealing with problematic situations such as wildlife attack, whereas larger households have a greater capacity to chase away wildlife from their fields. We also believe that the effectiveness of human capital endowments can be further enhanced by the household's wealth and physical capital endowments. Both consumer assets (such as television sets, radios, furniture, etc.) and productive assets (such as cars, trucks, farm equipment, etc.) are an indicator of wealth and could proxy for enhanced ability to respond to wildlife. However, a household with more valuable productive assets would be well positioned to expand total cultivated area, perhaps encroaching upon wildlife habitat and increasing crop damage. Therefore, wildlife-inflicted crop damage can be enhanced by the size of the farm operation. The larger the size of the farm, the less control the household will have per unit area, and the greater the likelihood and extent of wildlife attack. We use the size of cultivated land area and value of harvest as proxies for scale of operation.

The likelihood and extent of wildlife-inflicted crop damage is also expected to be inversely correlated by the level of human activity. The more remote a household is, for example, the greater

the likelihood of incurring wildlife-inflicted crop damage [8]. The further a household is from the road, the higher the probability of crop damage. Infrastructure such as the presence of clinics and schools was also used for concentration of human presence and activity. Households may also prefer working for wage employment due to the presence of infrastructure as a source of livelihood compared to farming. This could reduce the likelihood of crop damage by wildlife.

Closely related to this are variables such as human and wildlife population densities, which inhibit and enhance crop damage, respectively. Because data on wildlife populations were not readily available and GMAs are defined on the basis of wildlife populations, we use GMA classifications as proxies for wildlife populations. Prime GMAs have the largest populations of different kinds of wildlife. Thus, we expect greater crop damage in these areas compared to under-stocked GMAs [8]. Secondary and specialized GMAs have wildlife populations that are less than prime GMAs but more than under-stocked GMAs. This effect can, however, be moderated by presence of game scouts, although the scouts may also worsen crop damage if their primary focus is to help conserve wildlife.

5. Results and Discussion

5.1. Crop damage

Many different types of wildlife damage crops in the GMAs. However, as Osborn and Parker [10] show, elephants are the single most destructive species and that most of the wildlife-inflicted crop damage takes place during the harvest months of May, June and July. The IGMAW survey data showed that crop damage was mostly caused by monkeys and elephants with 31 and 20 percent of households reporting to have incurred crop damage from monkeys and elephants respectively.

5.2. Socioeconomic characteristics of households

Significance tests conducted using t-test showed that there were significant differences between households in GMAs and non-GMAs as can be seen in Table 2. Households in GMAs had lower average income levels, smaller household sizes, lower age of the household head and a lower average educational level compared with households in the non-GMAs. Households in GMAs also had fewer assets compared to households in non-GMAs. GMAs, as expected, are found in remote areas and are not densely populated compared to the non-GMAs.

Table 2. Socioeconomic characteristics of households in GMAs and control areas.

Variable	Full Sample	GMAs	Non-GMA control areas	Sig.
Number of sample households	2,717	1,574	1,143	
Total household income (ZMK)	4,235,762	3,591,253	5,123,301	*
Household size	5.28	5.08	5.57	***
Age of household head (in years)	42.46	41	44.48	***
Sex of household head (=1 if male)	0.74	0.73	0.76	**
Maximum education (in years)	6.78	6.42	7.27	***
Cropped area (hectares)	0.92	0.93	0.92	
Value of consumer assets (ZMK)	401,588	285,362	561,641	**
Value of productive assets (ZMK)	618,036	256,729	1,115,584	***
Distance to nearest main road (km)	5.09	6.08	3.8	***
Population density (per sq km)	35.2	41.41	26.97	***
Infrastructure index	3.62	3.64	3.59	
Prime GMA	0.17	0.3	<i>n.a.</i>	
Secondary or specialized GMA	0.2	0.35	<i>n.a.</i>	

n.a.= not applicable; *10% significance level, **5% significance level, ***1% significance level.

5.3. Characteristics of communities

Twenty-five (or 18.5 percent) of the 135 study communities had developed land use plans at the time of the survey. The t-test showed that characteristics differ in communities between those who don't have a land use plan, and those who have (Table 3). Communities that had a land use plan had older household heads, larger fields and were not remotely located compared to communities that did not have a land use plan. Communities with land use plans also had a higher population density as well as a higher presence of infrastructure compared to communities without land use plans. A higher proportion of communities with land use plans were found in prime GMAs while a higher proportion of communities without land use plans were found in secondary/specialized GMAs.

Table 3. Socioeconomic characteristics of households in communities with and without land use plans.

Categorical variable description	Communities without land use plans	Communities with land use plan	Sig.
Number of communities	110	25	
Household size	5.26	5.46	
Age of household head (in years)	42.25	43.48	*
Sex of household head (=1 if male)	0.75	0.75	
Maximum education (in years)	5.33	5.54	
Cropped area (hectares)	0.88	1.11	***
Value of consumer assets (Kwacha)	15.61	16.53	
Value of productive assets (Kwacha)	30.31	30.39	
Distance to nearest main road (km)	5.38	2.94	***
Population density (per sq km)	4.34	5.38	***
Infrastructure index	3.56	3.84	**
Prime GMA	0.12	0.39	***
Secondary /Specialized GMA	0.24	0.08	***

*10% significance level, **5% significance level, ***1% significance level.

5.4. Two-stage model results

The results of the two-stage model are shown in Table 4 where models 1 and 3 show the probability of crop damage while models 2 and 4 show the extent of damage. Further, the first two columns show the average partial effects (APEs) for the basic two-stage model without GMA-land use plan interaction terms, while the third and fourth columns show the APEs for the two-stage model with interaction terms. When land use planning is not interacted with the type of GMA, the results suggest that the land use plan does not affect crop damage in any statistically significant way. However, when interacted with the GMA effect, land use planning can lead to significant reductions in the likelihood of incurring crop damage for those households that are located in prime or secondary/specialized GMAs. Land use planning shows a greater impact in secondary/specialized GMAs compared to Prime GMAs. This could be that communities in GMAs that have abundant wildlife may not see the point of organizing themselves to formulate a land use plan [27]. Land use planning, however, does not have any such significant effects on the extent of crop damage for households that have incurred crop damage.

The results from the basic model show that increasing cultivated land area by 1 hectare increases the probability of crop damage by 2.7 percent. This finding and the fact that the probability of crop damage is significantly greater the larger the value of harvest suggest, as expected, that larger farms are more prone to crop damage. Also, increasing the value of harvest has a positive impact on the extent of crop damage. The results also suggest that households that are further away from the road are more likely to suffer from crop damage. An additional Km, on average, increases the probability of crop damage. This makes perfect sense as remote areas are more likely to be close to the national parks and to be further away from human activity.

Closely related to the remoteness variable is the presence of infrastructure, another proxy for concentration of human presence and activity. The results suggest, again consistent with expectations, that an additional physical infrastructure (such as a school, clinic, etc.) is associated with 1 percent reduction in the probability of incurring wildlife inflicted crop damage. Number of scouts has a positive and significant effect on the probability of crop damage showing that the presence of scouts could not be of an advantage to the communities but could only be there for the purpose of conservation of wildlife. The presence of a prime GMA has a positive and significant effect on crop damage showing that the presence of wildlife in the GMA increases the probability of farming households to suffer from crop damage.

The model with interaction terms in it shows some similar results to the basic model. The APEs on the probability of crop damage shows that increasing the size of the agricultural cropland by 1 hectare increases the probability of crop damage by 2.6 percent. The value of harvest variable shows a positive and significant effect on the probability and extent of crop damage. Increase in the distance from the main road increases the probability of crop damage and an additional infrastructure decreases the probability of crop damage.

A household in a prime GMA has a higher probability of crop damage (33.1 percent) compared with a household in a secondary/specialized GMA (5.7 percent). These results are consistent with the results found by Richardson *et al.* [8]. This shows that if a GMA has a high population of wildlife, the probability of crop damage is increased in such GMAs. Interestingly enough, when scouts are present in both prime GMA and secondary/specialized GMA they have no statistically significant effect but had a statistically significant effect on the extent of crop damage. The presence of village scouts in the GMA could be used for protecting farming households' fields from wildlife damage apart from protecting wildlife [8]. The joint significance of the district dummy variables shows that being in a different district has different effects on probability of crop damage and value of crop damage.

Table 4. Two-stage average partial effects for crop damage, 2005-06 agricultural season.

Variables	Basic Model		With Interactions	
	Stage 1 Model 1	Stage 2 Model 2	Stage 1 Model 3	Stage 2 Model 4
	(1)	(2)	(3)	(4)
Land use plan (dlandplan), 1=yes	-0.00673 (0.0203)	1.4020 (1.0530)	0.0324 (0.0311)	1.0230 (1.2540)
Prime GMA (g1), 1=yes	0.2240*** (0.0568)	-0.2080 (1.1580)	0.3310*** (0.0795)	0.0044 (1.6850)
Secondary/specialized GMA(g2), 1=yes	0.0393 (0.0260)	-0.5800 (0.8300)	0.0566* (0.0301)	0.1440 (0.9630)
dlandplan*g1			-0.0478* (0.0279)	1.0580 (1.8600)
dlandplan*g2			-0.0874*** (0.0228)	-1.0830 (2.2050)
Age of household in years (hage)	-8.19e-05 (0.0005)	0.0280 (0.0190)	-8.38e-05 (0.0005)	0.0295 (0.0186)
Sex of household head, 1= male (hmale)	-0.0150 (0.0171)	-0.2100 (0.8140)	-0.0140 (0.0168)	-0.3450 (0.8090)
Level of Education (hedu)	0.0009 (0.0022)	0.1430 (0.0915)	0.0008 (0.0022)	0.1420 (0.0915)
Household size (hhsz)	-0.0042 (0.0029)	0.1110 (0.1230)	-0.0045 (0.0028)	0.1050 (0.1240)
Distance to nearest main road in km (Kroad)	0.0015*** (0.0005)	-0.00889 (0.0180)	0.0012** (0.0005)	-0.0048 (0.0175)
Cropped area in hectares (care)	0.0266*** (0.0053)	-0.1390 (0.2290)	0.0263*** (0.0053)	-0.1370 (0.2240)
Value of consumer assets (vcasset2)	-0.1770 (0.5800)	12.3000 (22.8200)	-0.1100 (0.5750)	12.2200 (22.4800)
Value of productive assets (vpasset2)	-0.3930 (0.4220)	-22.4100 (17.4100)	-0.4010 (0.4200)	-19.3600 (18.0800)
Population density (Popdens)	0.0002 (0.0007)	-0.0099 (0.01480)	0.0004 (0.0007)	0.0031 (0.0162)
Infrastructure index (infras2)	-0.0102*** (0.0035)	-0.1580 (0.1350)	-0.0127*** (0.0037)	-0.1360 (0.1380)
Number of scouts (nscouts)	0.0069** (0.0030)	0.1130 (0.0706)	0.0045 (0.0070)	0.7950** (0.3580)
Value of harvest (harv2)	0.3050*** (0.0480)	8.8390*** (2.3230)	0.3000*** (0.0479)	9.0150*** (2.3150)
Nscouts*g1			-0.0118 (0.0117)	-1.2550** (0.5080)
Nscouts*g2			0.0066 (0.0079)	-0.7110* (0.3690)
Number of Observations	2,185	302	2,185	302
Goodness of Fit Chi ² -square statistic	246.23***		245.31***	
Log pseudo likelihood	-748.745	-866.880	-744.040	-862.678
Cragg & Uhler's R ²	0.202		0.209	
District dummy variable	104.80***	48.48***	95.36***	43.43***

Robust standard errors are presented in parentheses; *10% significance level, **5% significance level, ***1% significance level.

3. Conclusions

The main objective of this study was to estimate the impact of land use planning on wildlife inflicted crop damage in the GMAs. This objective was addressed by the use of a two-stage econometric model to identify the extent to which farmers cultivating crops in GMAs are more likely to be affected by crop damage caused by wildlife.

The findings showed that cultivated area, distance to the nearest road, infrastructure, value of harvest, number of scouts and the GMA effect are important factors affecting the probability of crop damage. Area cultivated, distance to the nearest main road, number of scouts, value of harvest, and living in a GMA are positively associated with crop damage. When number of scouts was interacted with GMA effect it was found not to have a significant effect on the probability of crop damage but had a significant negative effect on the extent of crop damage. Further, the presence of a land use plan in the GMAs and infrastructure were negatively associated with crop damage.

The empirical evidence has revealed that a land use plan has a significant effect in reducing the likelihood of crop damage caused by wildlife in the GMAs. Therefore, the implementation of land use plans should continue in all GMAs. This will help the rural communities that have to co-exist with wildlife have reduced impact of crop damage caused by wildlife therefore sustaining food security. The lack of policies that compensate the rural community when they suffer from crop damage makes land use planning even more of a valuable tool. In this study, population of wildlife was not controlled for, instead GMA type was used due to non-availability of an aerial count of wildlife in each GMA. Some communities might have a land use plan on paper but may not be implementing it. Further work should be done in this area to evaluate the effectiveness of various types of land use plans. There is also need for capacity building and extension for training in land use planning and best practices. This is especially important as all land use plans are different as they are made to suit the type of users.

Acknowledgments: The Impact of Game Management Areas on Household Welfare survey (IGMAW) was implemented by the Central Statistical Office (CSO) with funding from the World Bank (WB). The authors are responsible for all errors and omissions.

Author Contributions: M.S. led the writing of the manuscript and analysis of the data, and G.T. and R.B.R. contributed to the interpretation of the results. G.T. conceptualized the research design and led the collection of data. All authors contributed to the preparation of the manuscript.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Food and Agriculture Organisation (FAO). *Guidelines for Land-use Planning*; FAO. Rome, Italy, 1993.
2. Naab, F. Z.; Dinye, R. D.; Kasanga, R. K. Urbanisation and its impact on agricultural lands in growing cities in developing countries: a case study of Tamale, Ghana. *Modern Social Science Journal* **2013**, 2(2), 256-287.
3. Muleya, Z., *The Strategic Planning Process for Preparing General Management Plans for Protected Areas*; Unpublished manuscript, Zambia Wildlife Authority, Lusaka, Zambia, 2006.
4. Lamarque, F.; Anderson, J.; Fergusson, R.; Lagrange, M.; Osei-Owusu, Y.; Bakker, L. *Human-Wildlife Conflict in Africa. Causes, Consequences and Management Strategies*; Food and Agricultural Organization of the United Nations. Rome, Italy, 2009.
5. Kimeri-Mbote, P. *Land Tenure, Land Use and Sustainability in Kenya: Towards Innovative Use of Property Rights in Wildlife Management*; IELRC Working Paper 2005-4, International Environmental Law Research Centre, Geneva, Switzerland, 2005.
6. Dixon, J.A.; Sherman, P.B. Economics of protected areas. *Ambio* **1991**, 20(2): 68-74.
7. Ferraro, P.J. The local costs of establishing protected areas in low-income nations: Ranomafana National Park, Madagascar. *Ecological Economics* **2002**, 43: 261-275.
8. Richardson, R.B.; Fernandez, A.; Tschirley, D.; Tembo, G. Wildlife Conservation in Zambia: Impacts on Rural Household Welfare. *World Development* **2012**, 40(5): 1068-1081.
9. De Boer, W.F.; Baquete, D.S. Natural resource use, crop damage and attitudes of rural people in the vicinity of the Maputo Elephant Reserve, Mozambique. *Environmental Conservation* **1998**, 25(3): 208-218.
10. Osborn, F.V.; Parker, G.E. *Elephant/human conflict around the Luangwa National Parks, Zambia*; Consultancy Report for WWF/SARPO, Mid Zambezi Elephant Project, Harare, Zimbabwe, 2002.

11. Newmark, W. D.; Manyaza, D.N.; Gamassa, D.M.; Sariko, H.I. The conflict between wildlife and local people living adjacent to protected areas in Tanzania: Human density as a predictor. *Conservation Biology* **1994**, *8*(1): 249-255.
12. Hill, C.M. Conflict of interest between people and baboons: Crop raiding in Uganda. *International Journal of Primatology* **2000**, *21*(2):299-315.
13. Linkie, M.; Dinata, Y.; Nofrianto, A.; Leader-Williams, N. Patterns and perceptions of wildlife crop raiding in and around Kerinci Seblat National Park, Sumatra. *Animal Conservation* **2006**, *10*: 127-135.
14. Jones, T.B. *Human Wildlife Conflict Study, Namibian Case Study*; World Wide Fund, Windhoek, Namibia. 2006.
15. Shrestha, R. *A Case Study on Human-Conflict in Nepal (with particular reference to Human-Elephant conflict in Eastern and Western Terai regions)*; World Wide Fund, Nepal, 2007.
16. Naughton-Treves, L. Predicting patterns of crop damage by wildlife around Kibale National Park, Uganda. *Conservation Biology* **1998**, *12*(1): 156-168.
17. Bajracharya, S.B.; Furley, P.A.; Newton, A.C. Impacts of community-based conservation on local communities in the Annapurna Conservation Area, Nepal. *Biodiversity and Conservation* **2006**, *15*: 2765-2786.
18. Weladji, R. B.; Tchamba, M.N. Conflict between people and protected areas within the Bénoué Wildlife Conservation area, North Cameroon. *Oryx* **2003**, *37*: 72-79.
19. Otuoma, J. *The effects of wildlife-livestock-Human interactions on Habitat in the Meru conservation Area, Kenya*; LUCID Working paper 39, Land Use Change Impacts and Dynamics, International Livestock Research Institute, Nairobi, Kenya, 2004.
20. Muruthi, P. *Human Wildlife Conflict: Lessons Learned From AWF's African Heartlands*; AWF Working Papers, African Wildlife Foundation, Nairobi, Kenya, 2005.
21. Granados, A.; Weladji, R.B. Human-elephant conflict around Bénoué National Park, Cameroon: Influence on local attitudes and implications for conservation. *Human Dimensions of Wildlife: An International Journal* **2012**, *17*:77-90.
22. O'Connell-Rodwell, C. E.; Rodwell, T.; Rice, M.; Hart, L.A. Living with the modern conservation paradigm: Can agricultural communities co-exist with elephants? A five-year case study in East Caprivi, Namibia. *Biological Conservation* **2000**, *93*: 381-391.
23. Taylor, R.D. *A Review of Problem Elephant Policies and Management Options in Southern Africa*; Human Elephant Task Force, IUCN, African Elephant Specialist Group, Nairobi, Kenya, 1999.
24. Bandyopadhyay, S.; G. Tembo, G. Household consumption and natural resource management around national parks in Zambia. *Journal of Natural Resources Policy Research* **2010**, *2*(1): 39-55.
25. Larson, D.; Pienaar, E.; Jarvis, L. Wildlife conservation, labor supply and time values in rural Botswana. *Environment and Development Economics* **2016**, *21*(2): 135-157.
26. Zambia Wildlife Authority (ZAWA). *Annual Report*; Unpublished manuscript, Directorate of Conservation and Management, Zambia Wildlife Authority, Lusaka, Zambia, 2012.
27. Ostrom, E. A general framework for analyzing sustainability of social-ecological systems. *Science* **2009**, *325*: 419-422.
28. DeMotts, R.; Hoon, P. Whose elephants? Conserving, compensating, and competing in northern Botswana. *Society & Natural Resources* **2012**, *25*(9): 837-851.
29. Cragg, J. Some statistical models for limited dependent variables with application to the demand for durable goods. *Econometrica* **1971**, *39*: 829-44.