

Supplementary Materials for the article

Nitrogen Fertilization Reduces the Capacity of Soils to take up Atmospheric Carbonyl Sulphide

Aurore Kaisermann^{1*}, Sam P. Jones¹, Steven Wohl¹, Jérôme Ogée¹ and Lisa Wingate^{1,*}

¹ INRA ISPA UMR 1391, 33130 Villenave d'Ornon, France; aurore.kaisermann@inra.fr

* Correspondence: aurore.kaisermann@inra.fr; lisa.wingate@inra.fr; Tel.: +33 557 12 25 96






Figure S1. Variations in the (a) net COS flux; (b) the COS production rate (P) and (c) COS hydrolysis rate (k) with soil inorganic N content (NO_3^- and NH_4^+ contents, expressed in log) across the 27 soils in Europe. Each point represents the mean flux \pm standard deviation for each soil ($n=3$). Best statistical models were found comparing the adjusted R^2 and the AIC of each models with the easynls package (R).




Figure S2. Variations in the (a) net COS flux; (b) the COS production rate (P) and (c) COS hydrolysis rate (k) with soil nitrate (NO_3) and ammonium (NH_4^+) contents across 27 soils in Europe. The two levels of NO_3^- and NH_4^+ content are below and above 10 $\mu\text{g.g}^{-1}$ of N- NO_3^- and N- NH_4^+ respectively. Each black point represents the replicate means ($n=3$) for one soil. Letters represents results of TukeyHSD test when anova was significant.

Table S1. Characteristics of sites sampled for soil gas exchange. The two names in bold represent the two fertilized parcels in manipulation field experiment. The 9 underlined names represent the 9 soils used in the ammonium-nitrate addition experiment. Altitude, mean annual temperature (MAT) and precipitation (MAP) are estimated from the literature. VW is the volumetric water content, MBC and MBN are the microbial biomass C and N, DOC and DN are the dissolved organic C and N.

| SiteID | Country | Site | Landuse | Latitude | Longitude | Altitude (m) | MAT (°C) | MAP (mm) | pH | Bulk Density (g.cm-3) | VW (cm ³ .cm ⁻³) | MBC (ug.g ⁻¹) | MBN (ug.g ⁻¹) | DOC (ug.g ⁻¹) | DN (ug.g ⁻¹) | NO3 (ug.g ⁻¹) | NH4 (ug.g ⁻¹) |
|---------|------------------|------------------------|------------|----------|-----------|--------------|----------|----------|-----|-----------------------|---|---------------------------|---------------------------|---------------------------|--------------------------|---------------------------|---------------------------|
| ES-Amo | Spain (ES) | Amoladeras | Grassland | 36.8 | -2.3 | 54 | 18.5 | 200 | 8.5 | 0.93 | 0.12 | 1027 | 112 | 90 | 31 | 27 | 4 |
| ES-Bal | Spain (ES) | Balsablanca | Grassland | 36.9 | -2.0 | 200 | 18.5 | 220 | 8.5 | 0.89 | 0.10 | 650 | 57 | 83 | 23 | 20 | 3 |
| CH-Cha | Switzerland (CH) | Chamau | Grassland | 47.2 | 8.4 | 393 | 9.5 | 1136 | 5.4 | 0.69 | 0.15 | 963 | 84 | 71 | 69 | 76 | 12 |
| PT-Cor | Portugal (PT) | Coruche | Forest | 39.1 | -8.3 | 156 | 17 | 656 | 5.8 | 1.13 | 0.11 | 154 | 14 | 64 | 30 | 1 | 47 |
| CH-Dav | Switzerland (CH) | Davos | Forest | 46.8 | 9.9 | 1639 | 2.8 | 1062 | 4.2 | 0.33 | 0.21 | 1447 | 145 | 276 | 37 | 3 | 31 |
| CH-Fru | Switzerland (CH) | Fruebuell | Grassland | 47.1 | 8.5 | 982 | 7.2 | 1651 | 4.8 | 0.74 | 0.18 | 903 | 98 | 182 | 167 | 160 | 48 |
| DE-Hai | Germany (DE) | <u>Hainich</u> | Forest | 51.1 | 10.5 | 463 | 7.9 | 646 | 5.7 | 0.88 | 0.25 | 640 | NA | 146 | 173 | 125 | 58 |
| FR-Hes | France (FR) | <u>Hesse</u> | Forest | 48.7 | 7.1 | 313 | 9.7 | 650 | 4.8 | 0.87 | 0.18 | 509 | 53 | 127 | 45 | 5 | 42 |
| SE-Hy1 | Sweden (SE) | <u>Hyltemossa</u> | Forest | 56.1 | 13.4 | 111 | 7 | 850 | 3.7 | 0.52 | 0.12 | 854 | 110 | 157 | 53 | 1 | 66 |
| FI-Hyy | Finland (FI) | <u>Hyytiala</u> | Forest | 61.8 | 24.3 | 187 | 3.8 | 709 | 4.3 | 0.51 | 0.23 | 1058 | 120 | 273 | 29 | 3 | 12 |
| CH-Lae | Switzerland (CH) | <u>Laegern</u> | Forest | 47.1 | 8.5 | 866 | 8.3 | 1100 | 6.1 | 0.78 | 0.20 | 813 | 22 | 77 | 95 | 94 | 8 |
| FR-Lqu2 | France (FR) | Laqueuille_Cont | Grassland | 45.6 | 2.7 | 1041 | 7.9 | 897 | 4.9 | 0.55 | 0.22 | 2304 | 190 | 267 | 101 | 115 | 18 |
| FR-Lqu1 | France (FR) | Laqueuille_Fert | Grassland | 45.6 | 2.7 | 1041 | 7.9 | 897 | 4.2 | 0.54 | 0.16 | 1015 | 7 | 305 | 262 | 278 | 57 |
| DE-Lei | Germany (DE) | Leinefelde | Forest | 51.3 | 10.4 | 474 | 7.6 | 775 | 5.0 | 0.88 | 0.15 | 542 | 55 | 102 | 56 | 52 | 29 |
| PT-Mit | Portugal (PT) | Mitra | Forest | 38.5 | -8.0 | 240 | 15.6 | 656 | 6.0 | 0.80 | 0.11 | 479 | 52 | 75 | 18 | 9 | 10 |
| SE-Nor | Sweden (SE) | Norunda | Forest | 60.1 | 17.5 | 71 | 5.6 | 544 | 4.4 | 0.55 | 0.20 | 1258 | 135 | 94 | 25 | 1 | 36 |
| CH-Oe2 | Switzerland (CH) | Oensingen | Agrosystem | 47.3 | 7.7 | 452 | 9.2 | 1100 | 6.9 | 0.89 | 0.17 | 552 | 43 | 42 | 19 | 18 | 3 |
| IL-Reh | Israel (IL) | Rehovot | Agrosystem | 31.9 | 34.8 | 50 | 19.7 | 185 | 7.9 | 1.01 | 0.15 | 362 | 31 | 71 | 27 | 25 | 3 |
| SE-Ros2 | Sweden (SE) | Rosinedal_Cont | Forest | 64.2 | 19.7 | 145 | 1.8 | 614 | 4.6 | 0.86 | 0.12 | 99 | 5 | 132 | 16 | 3 | 15 |
| SE-Ros1 | Sweden (SE) | Rosinedal_Fert | Forest | 64.2 | 19.7 | 145 | 1.8 | 614 | 4.3 | 0.81 | 0.12 | 151 | 13 | 211 | 52 | 39 | 41 |
| DK-Sor | Denmark (DK) | Soro | Forest | 55.5 | 11.6 | 45 | 8.5 | 564 | 4.2 | 0.63 | 0.17 | 569 | 58 | 95 | 42 | 34 | 29 |
| SE-Sva | Sweden (SE) | Skartberget | Forest | 64.2 | 19.8 | 161 | 1.8 | 614 | 4.0 | 0.55 | 0.22 | 919 | 112 | 153 | 19 | 1 | 9 |
| ES-Ube2 | Spain (ES) | Ubeda_noVeg | Agrosystem | 37.9 | -3.2 | 366 | 14.8 | 20.4 | 8.6 | 0.86 | 0.12 | 766 | 54 | 110 | 15 | 4 | 4 |
| ES-Ube1 | Spain (ES) | Ubeda_Veg | Agrosystem | 37.9 | -3.2 | 366 | 14.8 | 20.4 | 8.4 | 0.99 | 0.12 | 795 | 63 | 105 | 40 | 40 | 3 |
| FI-Var1 | Finland (FI) | Varrtio1 | Forest | 67.8 | 29.6 | 400 | -1 | 660 | 5.2 | 0.76 | 0.11 | 212 | 19 | 150 | 14 | 1 | 5 |
| FI-Var2 | Finland (FI) | Varrtio2 | Forest | 67.8 | 29.6 | 400 | -1 | 660 | 4.9 | 0.76 | 0.15 | 546 | 73 | 104 | 17 | 1 | 14 |
| IL-Yat | Israel (IL) | Yatir | Forest | 31.3 | 35.1 | 661 | 17.9 | 292 | 8.6 | 0.87 | 0.16 | 1240 | 144 | 151 | 13 | 2 | 4 |

Table S2. Explanatory power of each soil properties for the three COS fluxes across the 27 European soils measured with the marginal R^2 value of the linear mixed model with a single soil properties. The significance of correlations has been tested with an analysis of variance. Abbreviations are: FCOS, net COS flux; k, COS hydrolysis rate; P, COS production rate; MBC, microbial biomass C; MBN, microbial biomass N; Ninor, inorganic N content (i.e. nitrate + ammonium); NO₃, nitrate; NH₄, ammonium; Redox, potential redox; BD, bulk density; VW, volumetric water content.

| Predictors | FCOS | | P | | k | |
|-----------------|----------------|---------|----------------|---------|----------------|---------|
| | R ² | P value | R ² | P value | R ² | P value |
| MBC | 0.20 | <0.0001 | 0.05 | 0.04 | 0.16 | <0.0001 |
| MBN | 0.37 | <0.0001 | 0.00 | 0.70 | 0.34 | <0.0001 |
| DOC | 0.00 | 0.10 | 0.10 | 0.01 | 0.00 | 0.73 |
| DN | 0.18 | <0.0001 | 0.54 | <0.0001 | 0.14 | <0.0001 |
| Ninor | 0.21 | <0.0001 | 0.41 | <0.0001 | 0.18 | <0.0001 |
| NO ₃ | 0.22 | <0.0001 | 0.43 | <0.0001 | 0.16 | <0.0001 |
| NH ₄ | 0.1 | 0.005 | 0.28 | <0.0001 | 0.14 | <0.0001 |
| pH | 0.00 | 0.70 | 0.16 | <0.0001 | 0.08 | 0.01 |
| Redox | 0.00 | 0.87 | 0.21 | <0.0001 | 0.05 | 0.06 |
| BD | 0.12 | 0.00 | 0.05 | 0.05 | 0.01 | 0.46 |
| VW | 0.03 | 0.11 | 0.13 | <0.002 | 0.00 | 0.75 |