

# S1 Subunit of Spike Protein from a Current Highly Virulent Porcine Epidemic Diarrhea Virus is an Important Determinant of Virulence in Piglets

Tohru Suzuki <sup>1,\*</sup>, Yutaka Terada <sup>2</sup>, Luis Enjuanes <sup>3</sup>, Seiichi Ohashi <sup>1</sup>, and Wataru Kamitani <sup>2</sup>

**Affiliations:** <sup>1</sup> Division of Viral Disease and Epidemiology, National Institute of Animal Health, National Agricultural Research Organization, Tsukuba, 3050856 Ibaraki, Japan; S.T.; suzuki106@affrc.go.jp, O.S.; ohashis@affrc.go.jp

<sup>2</sup> Research Institute for Microbial Diseases, Osaka University, Suita, 5650871 Osaka, Japan; Y.T.; yterada@biken.osaka-u.ac.jp; K.W.; wakamita@biken.osaka-u.ac.jp

<sup>3</sup> Department of Molecular and Cell Biology, Centro Nacional de Biotecnología, Campus Universidad Autónoma de Madrid, Madrid 28049, Spain; l.enjuanes@cnb.csic.es

**\*Correspondence:** Tohru Suzuki, Ph. D.

Division of Viral Disease and Epidemiology, National Institute of Animal Health, National Agricultural Research Organization, 3-1-5 Kannondai, Tsukuba, 3050856 Ibaraki, Japan.

Phone: +81-29-838-7914.

E-mail: suzuki106@affrc.go.jp.

## Abstract

Base on the sequence of S genes, which encode spike proteins, we previously identified three different types (North American, S INDEL and S large-DEL types) of porcine epidemic diarrhea virus (PEDV) that have re-emerged in Japan since 2013. Base on experimental infections with the North American and S large-DEL types, we also hypothesized that PEDV virulence may be linked to the S1 subunit of the S protein. To test this hypothesis, we have now assayed in gnotobiotic piglets various recombinant PEDVs generated by reverse genetics. Piglets inoculated with CV777 maintained in National Institute of Animal Health, along with piglets infected with a recombinant form of the same virus, developed subclinical to mild diarrhea. In contrast, severe watery diarrhea, dehydration, weight loss, astasia, and high mortality were observed in piglets inoculated with recombinant strains in which the S gene was partially or fully replaced with corresponding sequences from the highly virulent Japanese PEDV isolate OKN-1/JPN/2013. Indeed, symptoms resembled those in piglets inoculated with the OKN-1/JPN/2013, and were especially pronounced in younger piglets. Collectively, the data demonstrate that the S1 subunit of the S protein is an important determinant of PEDV virulence, and advance development of new vaccine candidate.

**Keywords:** porcine epidemic diarrhea virus; coronavirus; spike protein; virulence; gnotobiotic piglets; reverse genetics

## 1. Introduction

Porcine epidemic diarrhea is characterized by severe watery diarrhea which causes dehydration and high mortality among piglets. The disease is due to porcine epidemic diarrhea virus (PEDV), which belongs to the family *Coronaviridae* and genus *Alphacoronavirus*, and has an envelope surrounding a single-stranded positive-sense RNA genome [1]. The disease was first described in the United Kingdoms in 1971 [2], and has since widely spread not only to other European countries, but also to Asian countries including China, Korea, and Japan [3-8]. Notably, severe outbreaks caused by a strain genetically distinct from the prototype strain CV777 based on analysis of S genes, which encode spike proteins, have been reported in China since late 2010, with considerable morbidity and mortality among suckling piglets [9-12]. In April 2013, the disease was detected for the first time in the US, and was rapidly dispersed over 30 states in just one year [13, 14]. There were mainly two different types of PEDV strains in the US based on analysis of the S genes: the original highly virulent strain (North American type), which is genetically closely related to strains that emerged in China since 2010, and a mildly virulent strain (S INDEL type) with insertions and deletions at the 5' end of the S gene [15, 16]. US-like epidemics have also occurred in Canada and Mexico in North America, Korea and Taiwan in Asia, and Germany and Belgium in Europe [16-22].

In Japan, porcine epidemic diarrhea was first reported in 1982 [4]. In 1996, outbreaks occurred in 80,000 pigs at over 100 farms in nine prefectures, of which approximately half died. Although no cases were reported beginning in 2006, the disease re-emerged in October 2013. Thereafter, Ministry of Agriculture, Forestry and Fisheries (<http://www.maff.go.jp>) has reported over 1,000 outbreaks across almost all (39/47) prefectures, killing approximately 500,000 pigs until July 2018. To investigate the cause of re-emergence, we analyzed the S genes in 38 PEDV strains collected from 18 prefectures between 2013 and 2014, and detected the North American type (n = 33), the S INDEL type (n = 4), and the S large-DEL type (n = 1), a novel variant with a large deletion

of 582 nucleotides (194 amino acids) at the 5' end [23, 24]. In addition, this analysis revealed that the strains are more similar to global PEDV strains detected in recent years than to classical strains detected in Japan decades ago. Moreover, a phylogenetic dendrogram constructed from complete genomes indicated that the strains are genetically closely related to strains widespread in the US and Korea in 2013-2014, and that the S large-DEL variant may have spontaneously arisen from strains already circulating in domestic pigs [24, 25]. On experimental infections with the North American or the S large-DEL type, colostrum-deprived piglets, which are highly susceptible to pathogens, developed acute severe or moderate watery diarrhea, although the former was lethal while the other was not [26]. Moreover, the strains propagated in varying fashion to different tissues and formed different intestinal lesions.

The coronavirus spike protein consists of S1 and S2 subunits [27], of which the former binds putative cellular receptors such as aminopeptidase N and sialic acid, whereas the latter mediates virus-cell membrane fusion during entry [28]. Strikingly, porcine respiratory coronavirus has a large deletion of 200-230 amino acids in the S1 subunit, and is believed to be a naturally occurring mutant of, but with different tropism and pathogenicity as transmissible gastroenteritis virus, also of the genus *Alphacoronavirus* [29-31]. Therefore, these observations imply that the S protein, especially the S1 subunit, may play an important role in virulence and tissue tropism.

Reverse genetics, a powerful approach to analyze the function and role of a single gene, has been also used recently in coronaviruses which have large genomes of approximately 30 Kb, including for PEDV, severe acute respiratory syndrome and middle east respiratory syndrome coronaviruses [32-35]. To evaluate whether the S gene, especially the S1 subunit is an important determinant of PEDV virulence, we have now infected gnotobiotic piglets with several recombinant PEDVs that were generated by reverse genetics to fully or partially replace S genes but otherwise retain common genetic backbones.

## 2. Material and methods

### 2.1 Cells and viruses

Huh7 (human liver carcinoma) and Vero (African green monkey kidney cells) cells were maintained in Dulbecco's modified Eagle's medium (DMEM) (Nacalai Tesque, Kyoto, Japan) supplemented with 10 % fetal bovine serum (FBS), 100 U/mL penicillin, and 100 µg/mL streptomycin (Nacalai Tesque, Kyoto, Japan). All cells were cultured at 37 °C in a humidified atmosphere with 5 % CO<sub>2</sub>. The prototype PEDV strain, CV777 was kindly provided by Dr. MB Pensaert, and thereafter passaged and maintained at several times in our institute (The cells maintained in our institute were designated as CV777-niah strain). The complete genome of the CV777-niah strain had eleven nucleotide changes including eight non-synonymous substitutions, and three deletion (1-52 nucleotide in length) as compared with that of the reference CV777 strain (GenBank accession no. AF353511) (Table 1). The highly virulent PEDV strain, OKN-1/JPN/2013 was isolated and maintained in Vero cells as described in our previous study [24].

### 2.2 Construction of bacterial artificial chromosomes (BAC)

A BAC clone carrying full-length genome of the CV777-niah strain, pBAC-PEDV-CV777-FL, was generated on the backbone of SARS-CoV-Rep [36] using a Red/ET Recombination System Counter-Selection BAC Modification Kit (Gene Bridges, Heidelberg, Germany). However, the S gene of pBAC-PEDV-CV777-FL was modified to encode the entire spike protein based on CV777 reference sequence (GenBank accession no. AF353511). Using the same kit, ORF3 gene was replaced with green fluorescent protein (GFP) gene driven by the ORF3 transcription regulatory sequence, yielding pBAC-PEDVGFP-CV777, an infectious clone expressing GFP from subgenomic RNA. Finally, this clone was modified using the same kit to generate chimeric clones in which the S gene is fully or partially (S1 subunit) replaced with that of a high virulent Japanese

PEDV strain, OKN-1/JPN/2013 (GenBank accession no. LC063836), yielding pBAC-PEDVGFP-S-OKN1 and pBAC-PEDVGFP-S1-OKN1, respectively.

### 2.3 Recovery of recombinant PEDVs

Huh7 cells were grown to approximately 80 % confluence in a 6-well plate (VIOLAMO, Osaka, Japan), and transfected with 8 µg BAC DNA using X-tremeGENE 9 DNA Transfection Reagent (Sigma-Aldrich, St. Louis, MO). After 6 h, media were replaced with fresh DMEM. Transfected cells were then cultured at 37 °C for 3 days, washed three times with DMEM without FBS, and incubated with DMEM containing 5 µg/ml of trypsin (Sigma-Aldrich, St. Louis, MO). Viruses were harvested 6 h thereafter by freeze-thawing transfected cells and culture supernatant three times, and stored at -80 °C as P0 virus. To propagate the virus, Vero cells were seeded onto a 35 mm dish (VIOLAMO, Osaka, Japan), cultured overnight, washed three times with DMEM without FBS, and inoculated with 2 mL of P0 virus. After 2-3 h, wells were washed once with DMEM without FBS, and filled with 2 mL of DMEM containing 10 µg/mL of trypsin and 0.3% tryptose phosphate broth (Sigma-Aldrich, St. Louis, MO). Cells were then incubated at 37 °C for 3 or 4 days until cytopathic effects (CPEs) were observed. Harvested P1 viruses were passaged one more time on Vero cells to generate the P2 viruses, which were stored at -80 °C until use.

### 2.4 Animal studies

Gnotobiotic newborn piglets, which are highly susceptible to infectious agents, were obtained from four specific pathogen-free sows, following guidelines for the proper conduct of animal experiment at National Institute of Animal Health (17-061,5 December, 2017, NIAH Animal experiment committee).

#### *Experimental infection in 7-day-old gnotobiotic piglets.*

Twelve 7-day-old piglets were randomly assigned to orally receive CV777-niah ( $10^{6.3}$  TCID<sub>50</sub>/head, n = 3), its recombinant form, rPEDVGFP-CV777 ( $10^{5.6}$  TCID<sub>50</sub>/head, n = 5),

and the S-chimeric strain, rPEDVGFP-S-OKN1 ( $10^{4.7}$  TCID<sub>50</sub>/head, n = 4). Fecal samples were collected from each piglet every day from 0 to 5 days post-inoculation (DPI) and every 2–4 days thereafter. Sera were collected from each piglet every 2–3 days from 0 to 7 DPI and every 3–4 days thereafter. All animals were monitored for clinical signs and virus shedding until 28 DPI. Viral RNA was extracted from 10 % fecal suspensions and sera using QIAmp Viral RNA Mini Kit (Qiagen, Hilden, Germany), and titers were quantified by real-time reverse-transcription quantitative PCR using Takara One Step PrimeScript RT-PCR Kit (Takara Bio, Shiga, Japan) as described previously [26].

#### *Experimental infection in 5-day-old gnotobiotic piglets.*

Eighteen 5-days-old piglets were randomly assigned to orally receive the S-chimeric strain, rPEDVGFP-S-OKN1 ( $10^{5.8}$  TCID<sub>50</sub>/head, n = 6), the S1-chimeric strain, rPEDVGFP-S1-OKN1 ( $10^{5.8}$  TCID<sub>50</sub>/head, n = 6), or DMEM with no anything as mock control (n = 6). Feces and sera were collected as described above. Three piglets from each group were euthanized at 2 DPI, and remaining piglets were monitored for clinical signs and virus shedding until 28 DPI. On necropsy, small and large intestines were collected, along with heart, lung, kidney, liver, spleen, tonsil, trachea, muscle, stomach, and mesenteric lymph node. Viral RNA was extracted from 10 % fecal suspensions, sera, and 10 % tissue homogenates as described above. Virus shedding in feces and sera, and virus distributions in tissues at 2 DPI were quantified by real-time reverse-transcription quantitative PCR of the N gene as described above.

#### *2.5 Virus isolation and sequencing*

To isolate viruses from PEDV-inoculated piglets, feces collected at peaks virus shedding were suspended at 10 %, and inoculated into Vero cells. The inoculum was removed after 1 h at 37 °C in a humidified atmosphere with 5 % CO<sub>2</sub>. Cells were then washed three times with DMEM without FBS, and filled with DMEM containing 10 µg/mL trypsin and 0.3% tryptose phosphate broth. After 2 or 3 days at 37 °C, CPEs such

as cell fusion and syncytium formation were assessed by microscopy, along with GFP expressions.

The genomes of viruses isolated from PEDV-inoculated piglets were sequenced on an Ion Torrent PGM (Thermo Fisher Scientific, Carlsbad, CA, USA) as described previously [24].

### *2.6 Antibody detection by immunofluorescence assay*

Confluent monolayers of Vero cells grown in 96-well plates (Sumitomo Bakelite, Tokyo, Japan) were inoculated with  $10^{2.5}$  TCID<sub>50</sub> OKN-1/JPN/2013 in 100  $\mu$ L, and incubated at 37 °C in a humidified atmosphere with 5 % CO<sub>2</sub>. Inocula were removed after 18 to 24 h. Subsequently, cells were washed with PBS once, and fixed with cold 70 % acetone (FUJIFILM Wako Pure Chemical, Osaka, Japan) for 10 min. Plates were then air dried, sealed and stored at -20°C until probed with 50  $\mu$ L sera serially diluted two-fold from 1:20 to 1:1280 in PBS. Sera were removed after 1 h at 37 °C, and plates were washed 5 times with PBS. Wells were then reacted for 1h at 37 °C with 50  $\mu$ L/well of 1:50 goat anti-swine IgG (H+L) conjugated to fluorescein isothiocyanate (Jackson ImmunoResearch, West Grove, PA), washed 5 times with PBS, and examined for PEDV-specific cytoplasmic staining under a fluorescent microscope. Antibody titers are reported as the reciprocal of the highest dilution that produced clear, specific cytoplasmic staining.

## **3. Results**

### *3.1 Recovery of recombinant PEDVs derived from CV777*

To evaluate virulence due to the S gene, we first attempted to establish an infectious clone (pBAC-PEDV-CV777-FL) carrying a whole genome of the CV777-niah strain except for 3' end of the S gene by using a BAC system. The CV777 cDNA was assembled in the BAC under the control of the cytomegalovirus promoter and flanked at the 3' by 25 bp of poly(A), a hepatitis D virus (HDV) ribozyme, and a bovine growth hormone (BGH)



sequence (Fig. 1). The ORF3 gene was then replaced with GFP gene driven by the ORF3 transcription regulatory sequence to generate pBAC-PEDVGFP-CV777. Two chimeric clones, pBAC-PEDVGFP-S-OKN1 and pBAC-PEDVGFP-S1-OKN1, were subsequently generated by replacing the S and S1 genes in pBAC-PEDVGFP-CV777, respectively, with those of the OKN-1/JPN/2013 strain. The P2 viruses isolated in Vero cells were confirmed the CPEs, the same as GFP expressions by microscopy (data not shown). Moreover, genomic sequences of the P2 viruses were confirmed to be identical to those of the sources.

### *3.2 Experimental infection in 7-day-old gnotobiotic piglets.*

Newborn piglets inoculated with the CV777-niah and its recombinant form, rPEDVGFP-CV777 developed subclinical to mild diarrhea and anorexia 3–7 DPI (Table 2). In contrast, the piglets inoculated with the S-chimeric strain, rPEDVGFP-S-OKN1 developed severe watery diarrhea and eventually anorexia, lethargy, severe dehydration, weight loss, and astasia 2–7 DPI. However, these animals recovered beginning at 14 DPI and ultimately survived to the end of the experiment. Immunofluorescent assay for antibodies confirmed that all PEDV-inoculated piglets were infected (Table 2).

Piglets inoculated with the S-chimeric strain exhibited peak viral shedding of  $10^8$ – $10^9$  genome equivalents (GE)/mL in feces at 4–7 DPI. Shedding was sustained at high levels for approximately two weeks (Fig. 2). Piglets inoculated with the CV777-niah and the recombinant strain reached transient peak shedding of  $10^4$ – $10^6$  and  $10^4$ – $10^7$  GE/mL in feces at 5–7 DPI. Viral RNAs was detected at  $10^5$ – $10^6$  GE/mL in the sera 2–4 days after inoculation with the S-chimeric strain (Fig. 2). Viral RNA also peaked at  $10^3$ – $10^5$  GE/mL 4 days after infection with the CV777-niah and the recombinant strain. The recombinant and chimeric viruses collected from feces of infected piglets at the peak virus shedding elicited GFP expressions and CPEs in Vero cells, as assessed by microscopy, confirmed that the viruses are maintained and propagated in piglets, as suggested by sequencing (Fig. 3).

### 3.3 Experimental infection in 5-day-old gnotobiotic piglets.

Piglets inoculated with the rPEDVGFP-S-OKN1 and rPEDVGFP-S1-OKN1 strains exhibited acute severe watery diarrhea from 1 DPI. At 2–4 DPI, these animals additionally developed anorexia, lethargy, severe dehydration, weight loss, and astasia (Table 3). Finally, all animals died by 7 DPI. On the other hand, piglets infected with culture medium did not develop clinical signs, and survived to the end of the experiment.

Viral RNA titers in feces and sera are plotted in Fig. 4. Fecal samples collected 1–2 DPI with the S- and S1-chimeric strains tested positive for PEDV by PCR, with peak viral shedding at  $10^7$ – $10^9$  GE/mL within 3–5 DPI. Those titers were sustained at high levels until death. Viruses collected from feces at the peak shedding of piglets infected with the S- and S1-chimeric strains were confirmed by virus isolation and sequencing to be derived from inocula (Fig. 3). Viral RNAs were also detected at  $10^4$ – $10^6$  and  $10^5$ – $10^7$  GE/mL in sera 2 DPI with the S- and S1-chimeric strains, respectively. In piglets infected with the S-chimera viruses were mainly detected at  $10^7$ – $10^{10}$  GE/mL in intestinal tissues from the stomach to the rectum, as well as in mesenteric lymph node (Fig. 5). In animals infected with the S1-chimera, viruses at  $10^7$ – $10^{12}$  GE/mL were widely distributed in the small and large intestines and in mesenteric lymph node. In contrast, fecal, sera, and tissue samples from mock-infected piglets tested negative on PEDV-specific PCR (data not shown).

## 4. Discussion

In comparison of whole genomes between the CV777-niah and the reference CV777 (GenBank accession no. AF353511), we found 11 nucleotide difference in 5' UTR, ORF1a/1b, spike, and nucleocapsid, two deletions (1 or 4 nucleotides) in 5' UTR and 52 nucleotides deletion in Spike/ORF3 junction (Table 1). Especially, 52 nucleotides deletion in Spike/ORF3 junction in the CV777-niah can be predicted to result in the removal of 7 amino acids from C-terminus of spike protein and loss of ORF3 protein expression as

reported in a previous study [37]. In addition, we generated the recombinant and chimeric PEDV strains which replaced their ORF3 genes with GFP genes, and hence these strains can also be expected to be complete loss of their ORF3 functions. Piglets inoculated with the CV777-niah and its recombinant form showed similar symptoms and patterns of fecal virus shedding, and survived throughout the experiment. In contrast, Pensaert et al. reported that pathogenicity of piglets inoculated with the prototype strain CV777 is identical to those observed by the North American PEDV strains [38]. The difference of pathogenicity between the CV777-niah and the prototype CV777 might be involved in nucleotide change by cell adaptation, especially loss of ORF3 function, because there are previous reports that ORF3 differentiation through cell adaptation is associated with reduced virulence [39-41].

In the first experiment, all 7-day-old gnotobiotic piglets inoculated with the S-chimeric PEDV strain, rPEDVGFP-S-OKN1 ( $10^{4.7}$  TCID<sub>50</sub>/head), exhibited extreme debilitation at 2–7 DPI, including severe watery diarrhea, anorexia, lethargy, severe dehydration, weight loss, and astasia, but ultimately survived until 28 DPI (Table 2). In contrast, in the second experiment, three 5-day-old gnotobiotic piglets inoculated with the same S-chimeric strain ( $10^{5.8}$  TCID<sub>50</sub>/head) died at 7 DPI after exhibiting anorexia, lethargy, severe dehydration, weight loss, and astasia in addition to severe watery diarrhea (Table 3). Moreover, two of three 5-day-old gnotobiotic piglets inoculated with the S1-chimeric strain, rPEDVGFP-S1-OKN1 ( $10^{5.8}$  TCID<sub>50</sub>/head) died at 5DPI, and the other died at 7DPI with similar clinical signs as piglets inoculated with the S-chimeric strain. The discrepancy in mortality of piglets with the S-chimeric strain suggests to be associated with the difference of viral titers inoculated with piglets or the difference of age in onset of two experiment.

Piglets inoculated with the S- and S1-chimeric PEDVs showed similar patterns of acute virus shedding in feces and sera from 1–2 DPI (Fig. 2 and Fig. 4). Moreover, necropsy at 2 DPI indicated that viruses were predominantly restricted to the intestinal tract from jejunum to rectum, and to mesenteric lymph node (Fig. 5). Previously, we reported that 6-day-old colostrum-deprived piglets inoculated with OKN-1/JPN/2013,

which is the source of S and S1 fragments inserted into chimeric PEDVs, showed severe clinical signs such as dehydration, weight loss, acute viral shedding in feces and sera from 1–2 DPI, and high virus titers in small and large intestines and mesenteric lymph node, and ultimately died by 4 days post-infection [26]. Thus, the chimeric strains with S and S1 genes from OKN-1/JPN/2013 appear to reproduce its virulence. Consequently, our data strongly suggests that the S gene, especially the S1 subunit, is an important factor of severe symptoms and mortality in piglets. The difference of pathogenicity in piglets using these recombinant viruses might be caused by the function of spike protein in virus entry or process of budding. In future, we would perform *in vitro* assay using these viruses to elucidate the mechanism.

Also using reverse genetics, Hou et al. found that deletion of 197 amino acids from the N-terminal domain of the S1 subunit attenuated the pathogenicity of a highly virulent PEDV [32]. This results strongly supports the hypothesis that the S1 subunit is an important determinant of virulence, especially since we now demonstrate the opposite effect, in which virulence is conferred via S1 from a highly virulent strain upon an avirulent strain CV777-niah. In contrast, Wang et al. reported that the S gene is necessary but not sufficient for the virulence, based on experimental infections with various chimeric PEDVs [33]. However, these experiments were based on conventional piglets, while we and Hou et al. used gnotobiotic piglets, so results may not be directly comparable, especially since there were clear differences in pathogen susceptibility between conventional and gnotobiotic piglets. In any case, further studies based on reverse genetics are needed to clarify the function and role of other PEDV genes.

In summary, we generated recombinant PEDVs derived from the CV777-niah maintained in cell culture, as well chimeric PEDVs in which the original S gene is replaced with a virulent form by reverse genetics. The resulting S- and S1-chimeric strains caused severe clinical signs or high mortality in highly susceptible gnotobiotic piglets of different ages, reproducing the pathogenicity of OKN-1/JPN2013 as we described previously. Collectively, the data indicate that the S1 subunit is an important determinant

of PEDV virulence. We will thus attempt to further map the sequences in the S1 subunit that are critical to virulence.

**Author contributions:** Conceptualization, T.S.; Methodology, W.K. & T.S.; Validation, W.K. & T.S.; Investigation, S.O. & T.S.; Resources, Y.T. & E.L.; Data Curation, S.O. & T.S.; Writing – Original Draft Preparation, T.S.; Writing – Review & Editing, W.K. & T.S. All authors read and approved the manuscript.

**Funding:** TS was partially supported by a grant from National Agricultural Research Organization. WK was supported by grants from The Ito Foundation and The Mishima Kaiun Memorial Foundation. YT was supported by a grant from The Futaba Research Grant Program of the Futaba Foundation

**Acknowledgements:** We thank Mrs. K. Yukawa for secretarial assistance, Ms. K. Yoshizawa and N. Shimada for technical assistance, Drs. M. Sugiyama and N. Ito (Gifu University) for providing advice and discussion.

**Conflict of interest:** We have no conflict of interest to declare.

## References

1. Pensaert, M.B.; de Bouck, P. A new coronavirus-like particle associated with diarrhea in swine. *Arch Virol* **1978**, *58*, 243–247
2. Wood, E.N. An apparently new syndrome of porcine epidemic diarrhea. *Vet Rec* **1977**; *100*, 243–244.
3. Chasey, D.; Cartwright, S.F. Virus-like particles associated with porcine epidemic diarrhoea. *Res Vet Sci* **1978**, *25*, 255–256.
4. Takahashi, K.; Okada, K.; Ohshima, K. An outbreak of swine diarrhea of a new-type associated with coronavirus-like particles in Japan. *Jpn J Vet Sci* **1983**, *45*, 829–832.
5. Kweon, C.H.; Kwon, B.J.; Jung, T.S.; Kee, Y.J.; Hur, D.H.; Hwang, E.K.; Rhee, J.C.; An, S.H. Isolation of porcine epidemic diarrhea virus (PEDV) in Korea. *Korean J Vet Res* **1993**, *33*, 249–254
6. van Reeth, K.; Pensaert, M.B. Prevalence of infectious with enzootic respiratory and enteric viruses in feeder pigs entering fattening herds. *Vet Rec* **1994**, *135*, 594–597.
7. Nagy, B.; Nagy, G.; Meder, M.; Mocsari, E. Enterotoxigenic *Escherichia coli*, rotavirus, porcine epidemic diarrhoea virus, adenovirus and calici-like virus in porcine postweaning diarrhoea in Hungary. *Acta Vet Hung* **1996**, *44*, 9–19.
8. Song, D.; Park, B. Porcine epidemic diarrhoea virus: a comprehensive review of molecular epidemiology, diagnosis and vaccine. *Virus Genes* **2012**, *44*, 167–175.
9. Chen, J.; Wang, C.; Shi, H.; Qiu, H.; Liu, S.; Chen, X.; Zhang, Z.; Feng, L. Molecular epidemiology of porcine epidemic diarrhea virus in China. *Arch Virol* **2010**, *155*, 1471–1476.
10. Chen, X.; Yang, J.; Yu, F.; Ge, J.; Lin, T.; Song, T. Molecular characterization and phylogenetic analysis of porcine epidemic diarrhea virus (PEDV) samples from field cases in Fujian, China. *Virus Genes* **2012**, *45*, 499–507.
11. Li, W.; Li, H.; Liu, Y.; Pan, Y.; Deng, F.; Song, Y.; Tang, X.; He, Q. New variants of porcine epidemic diarrhea virus, China, 2011. *Emerg Infect Dis* **2012**, *18*, 1350–1353.

12. Li, Z.L.; Zhu, L.; Ma, J.Y.; Zhou, Q.F.; Song, Y.H.; Sun, B.L.; Chen, R.A.; Xie, Q.M.; Bee, Y.Z. Molecular characterization and phylogenetic analysis of porcine epidemic diarrhea virus (PEDV) field cases in south China. *Virus Genes* **2012**, *45*, 181–185.
13. Huang, Y.W.; Dickerman, A.W.; Pineyro, P.; Li, L.; Fang, L.; Kiehne, R.; Opriessnig, T.; Meng, X.J. Origin, evolution, and genotyping of emergent porcine epidemic diarrhea virus strains in the United States. *mBio* 2013, 4:e00737–00713.
14. Stevenson, G.W.; Hoang, H.; Schwartz, K.J.; Burrough, E.R.; Sun, D.; Madson, D.; Cooper, V.L.; Pillatzki, A.; Gauger, P.; Schmitt, B.J.; Koster, L.G.; Killian, M.L.; Yoon, K.J. Emergence of porcine epidemic diarrhea virus in the United States: clinical signs, lesions, and viral genomic sequences. *J Vet Diagn Invest* **2013**, *25*, 649–654.
15. Wang, L.; Byrum, B.; Zhang, Y. New variant of porcine epidemic diarrhea virus, United States, 2014. *Emerg Infect Dis* **2014**, *20*, 917–919.
16. Vlasova, A.N.; Marthaler, D.; Wang, Q.; Culhane, M.R.; Rossow, K.D.; Rovira, A.R.; Collins, J.; Saif, L.J. Distinct characteristics and complex evolution of PEDV strains, North America, May 2013-February 2014. *Emerg Infect Dis* **2014**, *20*, 1620–1628.
17. Pasick, J.; Berhane, Y.; Ojkic, D.; Maxie, G.; Embury-Hyatt, C.; Swekla, K.; Handel, K.; Fairles, J.; Alexandersen, S. Investigation into the role of potentially contaminated feed as a source of the first-detected outbreaks of porcine epidemic diarrhea in Canada. *Transbound Emerg Dis* **2014**, *61*, 397–410.
18. Lee, S.; Lee, C. Outbreak-related porcine epidemic diarrhea virus strains similar to US strains, South Korea, 2013. *Emerg Infect Dis* **2014**, *20*, 1223–1226.
19. Choi, J.; Lee, K.; Pi, J.H.; Park, S.; Song, C.; Choi, I.; Lee, J.; Lee, D.; Lee, S. Comparative genome analysis and molecular epidemiology of the reemerging porcine epidemic diarrhea virus strains isolated in Korea. *Infect Genet Evol* **2014**, *26*, 348–351.

20. Chiou, M.; Yu, C.; Chang, C.; Chung, W.; Wu, H.; Lin, C.; Lin, C. Molecular characterization of the porcine epidemic diarrhea virus TW4/2014 in Taiwan. *Austin Virol and Retrovirology* **2014**, *1*, id1007.
21. Hanke, D.; Jenckel, M.; Petrov, A.; Pitzmann, M.; Stadler, J.; Akimkin, V.; Blome, S.; Pohlmann, A.; Schirrmeyer, H.; Beer, M.; Hoper, D. Comparison of porcine epidemic diarrhea viruses from Germany and the United States, 2014. *Emerg Infect Dis* **2015**, *21*, 4893–4896.
22. Theuns, S.; Conceição-Neto, N.; Christiaens, I.; Zeller, M.; Desmarests, L.M.; Roukaerts, I.D.; Acar, D.D.; Heylen, E.; Matthijnsens, J.; Nauwynck, H.J. Complete genome sequence of a porcine epidemic diarrhea virus from a novel outbreak in Belgium, January 2015. *Genome Announc* **2015**, *3*, e00506-15.
23. Murakami, S.; Miyazaki, A.; Takahashi, O.; Hashizume, O.; Hase, Y.; Ohashi, S.; Suzuki, T. Complete genome sequence of the porcine epidemic diarrhoea virus variant Tottori2/JPN/2014. *Genome Announc* **2015**, *3*, e00877-15.
24. Suzuki, T.; Murakami, S.; Takahashi, O.; Kodera, A.; Masuda, T.; Itou, S.; Miyazaki, A.; Ohashi, S.; Tsutsui, T. Molecular characterization of pig epidemic diarrhoea viruses isolated in Japan from 2013 to 2014. *Infect Genet Evol* **2015**, *36*, 363-368.
25. Masuda, T.; Murakami, S.; Takahashi, O.; Miyazaki, A.; Ohashi, S.; Yamasato, H.; Suzuki, T. Discovery of a new PEDV variant with a large deletion in the spike gene identified in domestic pigs. *Arch Virol* **2015**, *160*, 2565-2568.
26. Suzuki, T.; Shibahara, T.; Yamaguchi R, Nakade, K.; Yamamoto, T.; Miyazaki, A.; Ohashi, S. Pig epidemic diarrhoea virus S gene variant with a large deletion non-lethal to colostrum-deprived newborn piglet. *J Gen Virol* **2016**, *97*, 1823-1826.
27. de Haan, C.A.M.; Haijema, B.J.; Schellen, P.; Wichgers Schreur, P.; te Lintelo, E.; Vennema, H.; Rottier, P.J.M. Cleavage of group 1 coronavirus spike proteins: how furin cleavage is traded off against heparan sulfate binding upon cell culture adaptation. *J Virol* **2008**, *82*, 6078-6083.



28. Liu, C.; Tang, J.; Ma, Y.; Liang, X.; Yang, Y.; Peng, G.; Jiang, S.; Li, J.; Du, L.; Li, F. Receptor usage and cell entry of porcine epidemic diarrhea coronavirus. *J Virol* **2015**, *89*, 6121-6125.
29. Callebaut, P.; Correa, I.; Pensaert, M.; Jiménez, G.; Enjuanes, L. Antigenic differentiation between transmissible gastroenteritis virus of swine and a related porcine respiratory coronavirus. *J Gen Virol* **1988**, *69*, 1725-1730.
30. Gallagher, T.M.; Buchmeier, M.J. Coronavirus spike proteins in viral entry and pathogenesis. *Virology* **2001**, *279*, 371-374.
31. Hulswit, R.J.G.; de Haan, C.A.M.; Bosch, B.J. Coronavirus spike protein and tropism changes. *Adv Virus Res* **2016**, *96*, 29-57.
32. Hou Y, Lin C, Yokoyama M, Yount, B.L.; Marthaler, D.; Douglas, A.L.; Ghimire, S.; Qin, Y.; Baric, R.S.; Saif, L.J.; Wang, Q. Deletion of a 197-amino-acid region in the N-terminal domain of spike protein attenuates porcine epidemic diarrhea virus in piglets. *J Virol* **2017**, *91*, e00227-17.
33. Wang, D.; Ge, X.; Chen, D.; Li, J.; Cal, Y.; Deng, J.; Zhou, L.; Guo, X.; Han, J.; Yang, H. The S is necessary but not sufficient for the virulence of porcine epidemic diarrhea virus novel variant strain BC2011C. *J Virol* **2018**, *92*, e00603-18.
34. Almazán, F.; Sola, I.; Zuñiga, S.; Marquez-Jurado, S.; Morales, L.; Becares, M.; Enjuanes, L. Coronavirus reverse genetic systems; infectious clones and replicons. *Virus Res* **2014**, *189*, 262-270.
35. Scobey, T.; Yount, B.L.; Sims, A.C.; Donaldson, E.F.; Agnihotharm, S.S.; Menachey, V.D.; Graham, R.L.; Swanstrom, J.; Bove, P.F.; Kim, J.D.; Grego, S.; Randell, S.H.; Baric, R.S. Reverse genetics with a full-length infectious cDNA of the Middle East respiratory syndrome coronavirus. *Proc Natl Acad Sci U S A* **2013**, *110*, 16157-16162.
36. Almazán, F., Dediego, M.L., Galán, C., Escors, D.; Alvarez, E.; Ortego, J.; Sola, I.; Zuñiga, S.; Alonso, S.; Moreno, J.L.; Nogales, A.; Capiscol, C.; Enjuanes, L. Construction of a severe acute respiratory syndrome coronavirus infectious cDNA

- clone and a replicon to study coronavirus RNA synthesis. *J Virol* **2006**, *80*, 10900-10906.
37. Rasmussen, T.B.; Boniotti, M.B.; Papetti, A.; Papetti, A.; Grasland, B.; Frossard, J.; Dastjerdi, A.; Hulst, M.; Hanke, D.; Pohlmann, A.; Blome, S.; van der Poel, W.H.M.; Steinbach, F.; Blanchard, Y.; Lavazza, A.; Bøtner, A.; Belsham, G.J. Full-length genome sequences of porcine epidemic diarrhoea virus strain CV777; Use of NGS to analyse genomic and sub-genomic RNAs. *PLoS One* 2018, *13*, e0193682.
38. Pensaert, M.B.; Martelli, P. Porcine epidemic diarrhea: A retrospect from Europe and matters of debate. *Virus Res* **2016**, *226*, 1-6.
39. Wood, R.D. Efficacy of a transmissible gastroenteritis coronavirus with an altered ORF-3 gene. *Can J Vet Res* **2001**, *65*, 28–32.
40. Song, D.S.; Yang, J.S.; Oh, J.S.; Han J.H.; Park, B.K. Differentiation of a Vero cell adapted porcine epidemic diarrhea virus from Korean field strains by restriction fragment length polymorphism analysis of ORF3. *Vaccine* **2003**, *21*, 1833–1842.
41. Park, S.J.; Moon, H.J.; Luo, Y.; Kim, H.K.; Kim, E.M.; Yang, J.S. Song, D.S.; Kang, B.K.; Lee, C.S. Park, B.K. Cloning and further sequence analysis of the ORF3 gene of wild- and attenuated-type porcine epidemic diarrhea viruses. *Virus Genes* **2008**, *36*, 95–104.