

The Status of Cardiovascular Health in Rural and Urban Areas of Janów Lubelski District in Eastern Poland: a Population-Based Study

Grzegorz Józef Nowicki*, Barbara Ślusarska, Honorata Piasecka, Agnieszka Bartoszek, Katarzyna Kocka and Alina Deluga

Table S1. The definition of cardiovascular health in our research (according to AHA criteria).

Metric	Definition
Behavioural component	
Smoking	
Poor	Currently smoke (at least 1 cigarette a day or the last cigarette was burned in the last month)
Intermediate	Smoked in the past and quit smoking 1–12 months ago
Ideal	Never smoked or quit smoking > 12 months ago
Body mass index	
Poor	≥ 30 kg/m ²
Intermediate	25–29.9 kg/m ²
Ideal	18.5–24.9 kg/m ²
Physical activity	
Poor	< 150 minutes of physical activity / week
Intermediate [#]	-
Ideal	≥ 150 minutes of physical activity / week
Healthy diet score	
Poor	≥ 12 points in the diet test
Intermediate	20–13 points in the diet test
Ideal	30–21 points in the diet test
Biological component	
Total cholesterol	
Poor	≥ 240 mg/dl
Intermediate	200–239 mg/dl or < 200 mg/dl and intake of cholesterol-lowering medications
Ideal	< 200 mg/dl and not taking cholesterol-lowering medications
Blood pressure	
Poor	SDP ≥ 140 mm Hg and DBP ≥ 90 mm Hg
Intermediate	SBP 121–139 and DBP 80–89 or intake of hypertensive
Ideal	SBP ≤ 120 mm Hg and DBP ≤ 80 mm Hg without hypotensive treatment

Table S1. *Cont.*

Metric	Definition
Biological component	
Fasting plasma glucose	
Poor	≥ 126 mg/dl
Intermediate	100–125 mg/dl or > 100 mg/dl and intake of hypoglycaemic medications
Ideal	< 100 mg/dl and not taking hypoglycaemic medications

^aValue not taken into consideration

Table S2. Characteristics of the researched group according to their gender.

Characteristics of population	Female		Male		P-value
	n	%	n	%	
	2298	58.9	1603	41.1	
Smoking					
Poor	262	11.4	358	22.3	< 0.001 [#]
Intermediate	369	16.1	451	28.1	< 0.001 ^{##}
Ideal	1667	72.5	794	49.5	
Body Mass Index					
Poor	814	35.4	589	36.7	< 0.001 [#]
Intermediate	828	36	734	45.8	< 0.001 ^{##}
Ideal	656	28.5	280	17.5	
Physical Activity					
Poor	1383	60.2	851	53.1	< 0.001 [#]
Intermediate	-	-	-	-	
Ideal	915	39.8	752	46.9	
Healthy Diet					
Poor	1946	84.7	1438	89.7	< 0.001 [#]
Intermediate	166	7.2	97	6.1	< 0.001 ^{##}
Ideal	186	8.1	68	4.2	
Fasting Total Cholesterol					
Poor	701	30.5	453	28.3	0.2 [#]
Intermediate	799	34.8	554	34.6	0.12 ^{##}
Ideal	798	34.7	596	37.2	
Blood Pressure					
Poor	1010	44	1016	63.4	< 0.001 [#]
Intermediate	955	41.6	518	32.3	< 0.001 ^{##}
Ideal	333	14.5	69	4.3	
Fasting Serum Glucose					
Poor	192	8.4	154	9.6	0.06 [#]
Intermediate	658	28.6	497	31	0.02 ^{##}
Ideal	1448	63	952	59.4	
Cardiovascular health metrics					
Poor cardiovascular health (0–2 ideal metrics)	1148	50	986	61.5	<0.001 [#]
Intermediate cardiovascular health (3–4 ideal metrics)	975	42.4	581	36.2	< 0.001 ^{##}
Ideal cardiovascular health (5–7 ideal metrics)	175	7.6	36	2.2	

Table S2. *Cont.*

Characteristics of population	Female		Male		P-value
	n	%	n	%	
	2298	58.9	1603	41.1	
No. of ideal cardiovascular health metrics					
0	59	2.6	102	6.4	<0.001
1	383	16.7	349	21.8	
2	706	30.7	535	33.4	
3	611	26.6	417	26	
4	364	15.8	164	10.2	
5	136	5.9	31	1.9	
6	34	1.5	5	0.3	
7	5	0.2	0	0	

*Ideal vs. poor vs. intermediate; **Ideal vs. poor + intermediate