

1 Type of the Paper (Article, Review, Communication, etc.)

2 Bromatological and mineralogical studies in fruit 3 pulps cultivated in the Northern Amazon

4
5 Ismael Montero Fernández ^{1*}, Edvan Alves Chagas ^{1,2,3}, Pollyana Cardoso Chagas ^{1,8}, Selvin
6 Antonio Saravia Maldonado ^{3,5}, Ricardo Carvalho dos Santos ^{1,6}, Pedro Rômulo Estevam Ribeiro ⁶,
7 Ana Cristina Gonçalves Reis de Melo ⁶, Antonio Alves de Melo Filho ^{1,7}

8
9 1 Post Graduate Program in Biodiversity and Biotechnology, Bionorte, State Coordination of Roraima,
10 UFRR, Campus Paricarana, CEP 69304000, Boa Vista-RR-Brazil; ismamontero@hotmail.com
11 2 Embrapa, Rodovia 174, Km 8, Industrial District, CEP 69301970, Boa Vista-RR-Brazil; CNPq Research
12 Productivity Scholarship; edvan.chagas@embrapa.br
13 3 Post Graduate Program in Agronomy, CCA/UFRR, Campus Cauamé, Boa Vista-RR-Brazil
14 4 University National of Agriculture, Catacamas, Olancho, Honduras saraviaselvin@yahoo.com
15 5 Insikiran Institute, Universidade Federal de Roraima (UFRR), Campus Paricarana, CEP 69304000, Boa
16 Vista-RR-Brazil; ricardo.carvalho@ufrr.br
17 6 Post-Graduate Program in Natural Resources, PRONAT, UFRR. Paricarana Campus, CEP 69304-000, Boa
18 Vista-RR-Brazil; pedro.ribeiro@ufrr.br
19 7 Department of Chemistry, Center of Science and Technology, UFRR, Campus Paricarana, CEP 69304000,
20 Boa Vista-RR-Brazil; antonio.alves@ufrr.br
21 8 Department of Agriculture, Center of Horticultural Science, UFRR, Campus Cauamé, CEP 69304000, Boa
22 Vista-RR-Brazil; pollyana.chagas@ufrr.br

23
24
25 * Correspondence: ismamontero@hotmail.com; Tel.: +55-95-988040640
26

27 **Abstract:** In this work, nine fruits cultivated in the northern Amazon were studied: abiu, acerola,
28 araçá, bacupari, biribá, caçari, fruta-do-conde, graviola and taperebá, with the objective of carrying
29 out a bromatological and nutritional study of the pulps of fruits studied. Of all of them, are the pulps
30 of graviola (76.83 ± 0.02 kcal 100 g⁻¹) bacupari (76.83 ± 0.02 kcal 100 g⁻¹) and fruta-do-conde ($46.66 \pm$
31 0.02 kcal 100 g⁻¹). Among the macronutrients, the high concentration of potassium stands out,
32 especially in the graviola (541.16 ± 0.24 mg 100 g⁻¹) and the biribá (468.21 ± 0.13 mg 100 g⁻¹). Among
33 the micronutrients, iron concentrations are representative for araçá pulp (3.04 ± 0.02 mg 100g⁻¹),
34 abiu is rich in zinc (3.71 ± 0.02 mg 100 g⁻¹) and manganese (6.61 ± 0.11 mg 100 g⁻¹). The presence of
35 cobalt at the level of traces in some of the pulps studied stands out. The Pearson correlation
36 coefficient was evaluated, as well as the statistical treatment by multivariate analysis PCA to
37 establish the correlation between the variables studied.

38 **Keywords:** Amazonian fruit, functional food, PCA, Person.

39 1. Introduction

40 The Amazon region along with the freshwater biome, presents the largest biodiversity on the
41 planet with more than 5000 species [1]. The fruits of this biodiversity, native and exotic, present an
42 expressive potential of bioactive compounds, which can be a source of bioproducts for the
43 development of humanity [2].

44 The Amazonian fruits arouse great study interest due to their great biodiversity, which,
45 according to [3-4] present outstanding results in quality and attractive attributes such as appearance
46 in large sizes, different shapes, colors, textures and different flavors.

47 In the human diet, fruits are considered the main sources of necessary minerals, playing a vital
48 role in the peculiar development and good health of the human body [5] because they participate in
49 many biochemical reactions, being divided according to [6] in macronutrients (minerals required for
50 humans in amounts greater than or equal to 100 mg day⁻¹ as calcium, magnesium, potassium,
51 phosphorus, sulfur, chlorine and sodium) and micronutrients (minerals required for humans in
52 amounts less than 100 mg day⁻¹, such as copper, iron, zinc, manganese, selenium, molybdenum and
53 fluorine.

54 Thus, many of the fruits cultivated, Amazonian or introduced in the region, do not present
55 information on the nutritional and mineralogical composition. For this reason, the objective of this
56 work is to analyze the composition of minerals and nutrients in the pulp of nine fruits grown in the
57 Northern Amazon (Figure 1) and to study the correlation between existing data using the Pearson
58 test as well as to use multivariate analysis methods such as Principal Component Analysis (PCA).

59

60



Fruta-do-conde
(*A. squamosa*)



Biribá
(*R. mucosa*)



Graviola
(*A. muricata*)



Bacupari
(*R. gardneriana*)



Abiu
(*P. caimito*)



Acerola
(*M. emarginata*)



Taperebá
(*S. mombin*)



Caçari
(*M. dubia*)



Araçá
(*P. cattleianum*)

61

62 **Figure 1.** Fruits grown in the North Amazon under study (Pictures by Selvin Antonio Saravia Maldonado and
63 Ismael Montero Fernández).

64 2. Materials and Methods

65 2.1. Preparation of samples

66 Samples (Table 1) were collected from fruit markets and producers in Roraima state, Brazil.
67 Then, the collected fruits were taken to the Laboratory of the Agronomic Research Center, at the
68 Agricultural Sciences Center, Cauamé campus, Federal University of Roraima, fruits with good
69 appearance were selected, washed previously with distilled water and then with hypochlorite
70 solution of sodium chloride and finally with distilled water again according to the procedure
71 described [7].

72

73 **Table 1** - Names and families of fruits cultivated in the Northern Amazon studied in this work.

Scientific name	Family	Name in Brazil
<i>Pouteria caimito</i>	Sapotaceae	Abiu
<i>Malpighia emarginata</i>	Malpighiaceae	Acerola
<i>Psidium cattleianum</i>	Myrtaceae	Araçá
<i>Rheedia gardneriana</i>	Clusiaceae	Bacuparí
<i>Rollinia mucosa</i>	Annonaceae	Biribá
<i>Myrciaria dúbia</i> (Krunth) Mc Vaugh, Myrtaceae	Myrtaceae	Caçari
<i>Annona squamosa</i>	Annonaceae	Fruta-do-conde
<i>Annona muricata</i>	Annonaceae	Graviola
<i>Spondias mombin</i> L.	Anacardiaceae	Taperebá

74

75 The fruits were pulped, weighed and frozen in an ultra-freezer at -80 °C for further lyophilization
76 in Liotop L101 lyophilizer for 48 hours, until complete drying. After drying, the samples were ground
77 in a knife mill and sieved between 30-40 Mesh, and stored in hermetically sealed sachets and
78 protected from light to perform nutritional and mineralogical analysis.

79

80

81 2.2. *Nutritional analysis*

82 The physical parameters evaluated to determine the nutritional composition were the
83 percentage of moisture and ash. The other nutritional parameters evaluated were the determination
84 of total proteins, lipids and carbohydrates, to determine the total energy content.

85

86 2.2.1. *Determination of Humidity*

87 To determine moisture, 5 g of fresh samples were placed in porcelain capsules for 6 hours at 105
88 °C to constant mass, and then cooled in desiccator to room temperature [8].

$$89 \text{ Humidity (g } 100\text{g}^{-1}) = ((P' - P'') / (P' - P)) \cdot 100$$

90 being:

91 P = weight of porcelain capsule (g)

92 P' = weight of the porcelain capsule + fresh sample (g)

93 P'' = weight of the capsule + sample after the oven (g)

94

95 2.2.2. *Determination of ashes*

96 To determine the ash in the samples, the methodology proposed for the food analysis [8] with
97 modifications was used, where 5 grams of the lyophilized samples were weighed. These were placed
98 in preheated porcelain crucibles in an oven at 110 °C for one hour, to remove moisture, and cool them
99 in a desiccator to room temperature. The samples were incinerated at 600 °C in a FDG 3P-S EDG
100 muffle for 16 hours, after which the samples were left in the desiccator until reaching room
101 temperature.

102

$$103 \% \text{ ashes} = ((N \cdot 100) / M)$$

104

105 N = mass in grams of ash and M = mass of the sample in grams.

106

107 2.2.3. *Determination of total protein*

108 Protein determination is performed from the total nitrogen analysis by Kjeldahl distillation, in
109 which the existing organic matter is transformed into ammonia. The nitrogen content of the different
110 proteins is approximately 16%, which introduces the empirical factor of 5.75 (conversion factor for
111 vegetable protein), this will transform the number of grams of nitrogen, found with the number of
112 grams of protein [8].

113

$$\% \text{ proteins} = \% N \cdot 5.75$$

114 2.2.4. *Determination of lipids*

115 To determine the total amount of lipids, 20 g of each sample was weighed, and placed in the
116 Soxhlet extractor apparatus with hexane as the solvent for six hours. The solvent was recovered in a
117 rotary evaporator [8].

118

$$119 \% \text{ lipids} = ((N \cdot 100) / m)$$

120

121 Where: N = mass in grams of lipids and M = mass of the sample in grams.

122 2.2.5. Determination of Carbohydrates

123 The carbohydrate content is achieved by the difference of the value 100 subtracted from the sum
124 of the already obtained values of moisture, ashes, lipids and proteins.

125

$$126 \text{ Carbohydrates} = 100 - (\% \text{ moisture} + \% \text{ ash} + \% \text{ lipids} + \% \text{ proteins})$$

127 2.2.5. Energetic value

128 In order to quantify the energy value, it was necessary to use the protein (P), lipid (L) and
129 carbohydrate (C) contents of each sample. The result should be expressed in kcal 100g⁻¹ [9].

130

$$131 \text{ Energy value (kcal 100g}^{-1}\text{)} = (P * 4) + (L * 9) + (C * 4)$$

132

133 P = value of protein (%), L = lipid value (%), C = carbohydrate value (%), 4 = conversion factor in kcal
134 determined in calorimetric pump for proteins and carbohydrates and 9 = conversion factor in kcal
135 determined in a calorimetric pump for lipids.

136

137 2.2. Mineralogical analysis

138

139 The extraction of the minerals into the pulps was done according to the methodology described
140 by [10] in which the perchloric nitric digestion (3:1) was used in TECNAL model TE 0079 digester
141 block, washed with distilled water up to 25 mL for subsequent analysis.

142 Calcium (Ca), magnesium (Mg), iron (Fe), copper (Cu), zinc (Zn), manganese (Mn) and
143 aluminum (Al) were determined by Flame Atomic Absorption Spectrophotometry (FAAS) Shimadzu
144 AA-7000, coupled with ASC-7000 auto sample. Calibration was performed with standard solutions
145 prepared from commercial standards of 1000 mg L⁻¹ Qhemis High Purity PACU 1000-0125, according
146 to the specific conditions of each element (Table 2).

147

148 **Table 2.** Analytical Parameters.

Element	Technique	(λ) nm	Calibration Line
Ca	AAS	422.70	y= 0.0092 x - 0.0005 r ² = 0.999
Mg	AAS	285.21	y= 0.2353 x - 0.0658 r ² = 0.997
P	UV-Vis Spectroscopy	660.00	y= 0.2181 x - 0.0005 r ² = 0.999
K	Flame Photometry	766.50	y= 0.1231 x - 0.0013 r ² = 0.993
S	UV-Vis Spectroscopy	420.00	y= 0.0213 x - 0.0012 r ² = 0.998
Fe	AAS	248.33	y= 0.0399 x + 0.0067 r ² = 0.996
Zn	AAS	213.80	y= 0.0600 x - 0.0171 r ² = 0.991
Mn	AAS	279.48	y= 0.0282 x + 0.0041 r ² = 0.999
Cu	AAS	324.75	y= 0.0512 x - 0.0099 r ² = 0.997
Na	EAS	589.00	y= 1.0000 x + 0.0005 r ² = 0.999
Al	AAS	309.30	y= 0.0088 x + 0.0005 r ² = 0.998
B	UV-Vis Spectroscopy	460.00	y= 0.0537 x + 0.0002 r ² = 0.999
Co	AAS	240.73	y= 0.0286 x - 0.0066 r ² = 0.997

149 FAAS = Flame Atomic Absorption Spectroscopy. FAES = Flame Atomic Emission Spectroscopy.
150 Gas flow rate 266 cm s⁻¹. Burner thickness 10 cm. Air-acetylene flame temperature 2300 ° C

151

152 As the ionization suppressor for the Ca and Mg elements, 0.1% of the lithium oxide solution
153 (Li₂O) was used. In the case of sodium (Na), it was determined in the same equipment, but in atomic
154 emission mode. As for potassium (K), it was determined by means of flame photometry on the
155 Digimed Flame Photometer DH-62, calibrated using a Digimed standard solution whose
156 concentration range was 2 - 100 mg L⁻¹.

157 For the determination of the phosphorus (P), boron (B) and sulfur (S) elements, the ultraviolet
158 molecular absorption spectrophotometry technique was used using a SHIMADZU UV-1800 model,
159 according to the [10], by formation of the colorimetric reaction with ammonium molybdate
160 ((NH₄)₂MoO₄). In the case of P, blue complex formed, where the readings were made at λ = 660 nm;
161 in the case of B complex was formed with azomidine-H of yellow color and absorbs light at λ = 460
162 nm; and for the sulfur was precipitated with BaCl₂, calibrating with potassium sulphate, at λ = 420
163 nm.

164 Nitrogen determination was carried out by the distillation method followed by titration
165 (Kjeldahl), where the ammonium ion produced in the digestion with sulfuric acid (H₂SO₄) is distilled
166 in strongly alkaline medium in the Kjeldahl distiller model TECNAL TE-036/1, collected (0.01%) and
167 methyl red (0.04%) and titrated with 0.01 mol L⁻¹ HCl solution were added in 2% boric acid solution
168 with a mixture of green bromocresol (0.01%) and methyl [10].

169

$$170 \quad \% \text{ N total} = (V \cdot 0.028) / m$$

171

172 Where V = difference in the titration volume of the sample blank; m = mass of the sample in
173 grams; and the value 0.028 = milliequivalents grams of nitrogen multiplied by the concentration.

174

175 2.3. Statistical analysis

176 Correlations between the amounts of the different minerals in the different parts of the fruit were
177 evaluated using the Pearson statistical test using INFOSAT [11] for significance levels of 5%, 1% and
178 0.1% respectively, as well as the principal component analyzes (PCA).

179 3. Results and Discussion

180 3.1 Bromatological analysis from Amazonian fruits

181 Table 3 presents the nutritional analysis values for the pulps of the different Amazonian fruits
182 studied. The readings were done in triplicate. The standard deviation was calculated for a level of significance
183 of the t-student of 95%

184

185

186

187

188

189

190

191

192

193 Table 3 - Nutritional composition in Amazonian fruit pulps.

Fruits	Nutritional Contribution					
	Moisture	Ashes	Lipids	Carbohydrates	Proteins	Energetic Value
	%					Kcal 100 g ⁻¹
Abiu	92.43 ± 0.02	0.13 ± 0.07	0.12 ± 0.01	6.48 ± 0.02	0.84 ± 0.01	30.36 ± 0.03
Acerola	94.21 ± 0.01	0.16 ± 0.03	0.06 ± 0.00	4.45 ± 0.02	1.12 ± 0.01	22.82 ± 0.02
Araçá	59.21 ± 0.08	0.21 ± 0.07	0.17 ± 0.09	30.87 ± 0.03	3.93 ± 0.07	18.09 ± 0.02
Bacupari	86.61 ± 0.11	0.19 ± 0.01	0.07 ± 0.00	12.36 ± 0.01	0.77 ± 0.01	53.15 ± 0.02
Biribá	91.34 ± 0.01	0.31 ± 0.02	0.22 ± 0.02	6.95 ± 0.03	1.18 ± 0.04	34.5 ± 0.01
Caçari	95.21 ± 0.14	0.25 ± 0.11	0.05 ± 0.00	3.14 ± 0.02	1.35 ± 0.04	18.41 ± 0.02
Fruta-do-conde	88.27 ± 0.05	0.29 ± 0.07	0.18 ± 0.07	10.08 ± 0.01	1.18 ± 0.04	46.66 ± 0.02
Graviola	80.77 ± 0.07	0.31 ± 0.11	0.23 ± 0.08	17.34 ± 0.01	1.35 ± 0.02	76.83 ± 0.02
Taperebá	88.23 ± 0,10	0.15 ± 0.01	0.05 ± 0.01	10.01 ± 0.03	1.56 ± 0.01	46.73 ± 0.03

195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238

The first parameter analyzed is moisture, which according to [12] the moisture content is used as a factor indicative of propensity for food spoilage, and may think that the greater stability of the food is in the control of the minimum humidity.

The amount of moisture in the pulps varies from 64.22-95.21%, where the highest moisture values are for caçari ($95.21 \pm 0.14\%$) and acerola ($94.21 \pm 0.11\%$); and the lowest value for araçá with $64.22 \pm 0.12\%$. Regarding the values of humidity for abiu, acerola and graviola, are slightly lower, but close to those presented by [13] and in the case of araçá lower than those presented by the same author.

The presented values of humidity are within the percentage humidity range given by the [8] that establishes the values in fruits between 65-90%. In the case of caçari, [14] determined the percentage of moisture in the camu-camu pulp of 92.65%, close to that found in the present study.

The ash content reflects the amount of minerals present in the food. On the other hand, the [8] establishes the values of percentage of fruit ashes between 0.3 - 2.1%, being the values of ashes studied in this work according to this range.

For proteins, it is those of animal origin that have higher biological value compared to proteins of plant origin [15] and the identification of plant species with a certain content of proteins, are important to satisfy the nutritional deficiencies of people with different dietary habits and diets [16] plus many native Amazonian species have not yet been studied to evaluate their protein potential.

The protein content in the fruit pulps studied in this study ranged from 0.77% for the bacupari to 3.93% for the araçá pulp. Caçari has a protein content of 1.35% as does graviola with the same value.

The three groups of primary metabolites in fruits, carbohydrates are the major ones, presenting values between 3.14-30.87% for pulps, being the one with the lowest value is the potato pulp and the highest is the araçá pulp.

The lipid content observed in the fruits cultivated in the Northern Amazon was observed in the caçari pulps, 0.05%, taperebá, 0.05%, and biribá, 0.22%, with lipid values being relatively low in fruit pulps. Among the fruits of the Annonaceae family, the species that had the highest lipid content was graviola with 0.23%, followed by the biribá with 0.22% and the fruit of the count with 0.18%. The values obtained for the biribá are close to those determined [17]. The oil of other fruits of the Annonaceae family, as is the case of *A. squamosa* and *A. atemoya*, presents natural bioinsecticidal properties [18], as well as other biological properties such as *A. hypoglauca* on inhibition of fungi and bacteria and the acetylcholinesterase enzyme [19].

3.2 Mineral analysis

In Tables 4 and 5, the values of macronutrients and micronutrients are presented for the different pulps studied. The readings were done in triplicate. The standard deviation was calculated for a level of significance of the t-student of 95%

239

240 Table 4 - Macronutrients analyzed in fruits grown in the northern Amazon.

Fruit	Macronutrients					
	Calcium (Ca)	Magnesium (Mg)	Phosphorous (P)	Potassium (K)	Sulfur (S)	Nitrogen (N)
	mg 100 g ⁻¹					%
Abiu (<i>P. caimito</i>)	4.51 ± 0.02	1.71 ± 0,07	8.21 ± 0,04	255.21 ± 0.03	11.11 ± 0.04	0.15 ± 0.01
Acerola (<i>M. emarginata</i>)	11.23 ± 0.12	18.41 ± 0.21	11.93 ± 0.04	154.34 ± 0.18	34.13 ± 0.14	0.19 ± 0.01
Araçá (<i>P. cattleianum</i>)	24.13 ± 0.03	12.21 ± 0.08	6.32 ± 0.04	137.11 ± 0.08	9.02 ± 0.01	0.68 ± 0.07
Bacupari (<i>R. gardneriana</i>)	32.41 ± 0.02	14.21 ± 0.08	12.31 ± 0.14	329.12 ± 0.04	5.21 ± 0.04	0.13 ± 0.01
Biribá (<i>R. mucosa</i>)	32.11 ± 0.08	112.32 ± 0.12	23.41 ± 0.01	468.21 ± 0.13	21.31 ± 0.12	0.21 ± 0.04
Caçari (<i>M. dubia</i>)	9.51 ± 0.02	8.49 ± 0.04	6.21 ± 0.04	124.13 ± 0.12	7.21 ± 0.04	0.23 ± 0.04
Fruta-do-conde (<i>A. squamosa</i>)	52.21 ± 0.13	32.12 ± 0.09	17.30 ± 0.12	431.21 ± 0.17	27.78 ± 0.13	0.21 ± 0.04
Graviola (<i>A. muricata</i>)	39.21 ± 0.13	27.11 ± 0.15	19.24 ± 0.16	541.16 ± 0.24	29.31 ± 0.08	0.23 ± 0.02
Taperebá (<i>S. mombin</i>)	38.12 ± 0.12	16.32 ± 0.09	24.12 ± 0.11	149.13 ± 0.23	4.38 ± 0.08	0.27 ± 0.01

241 The readings were done in triplicate. The standard deviation was calculated for a level of significance of the t-student of 95%

242

243

244

245

246

247

248 Table 5 - Micronutrients analyzed in fruits grown in the northern Amazon.

Fruit	Micronutrients							
	Iron (Fe)	Zinc (Zn)	Manganese (Mn)	Copper (Cu)	Sodium (Na)	Aluminum (Al)	Boron (B)	Cobalt (Co)
	mg 100 g⁻¹							
Abiu (<i>P. caimito</i>)	0.18 ± 0.04	3.71 ± 0.22	6.61 ± 0.11	0.12 ± 0.02	0.22 ± 0.01	0.17 ± 0.02	0.27 ± 0.07	N.D.
Acerola (<i>M. emarginata</i>)	0.80 ± 0.12	0.08 ± 0.01	0.24 ± 0.05	0.17 ± 0.01	35.13 ± 0.12	0.93 ± 0.04	0.11 ± 0.03	N.D.
Araçá (<i>P. cattleianum</i>)	3.04 ± 0.02	1.14 ± 0.02	1.25 ± 0.07	1.73 ± 0.02	1.93 ± 0.02	0.14 ± 0.06	0.10 ± 0.02	0.012 ± 0,003
Bacupari (<i>R. gardneriana</i>)	0.71 ± 0.02	3.46 ± 0.02	0.24 ± 0.01	0.15 ± 0.00	0.09 ± 0.01	0.12 ± 0,01	0.14 ± 0.01	0.021 ± 0.000
Biribá (<i>R. mucosa</i>)	1.82 ± 0.11	1.23 ± 0.04	0.33 ± 0.04	1.14 ± 0,13	18.44 ± 0.21	0.06 ± 0.01	0.51 ± 0.05	0.006 ± 0,001
Caçari (<i>M. dubia</i>)	0.29 ± 0.03	0.13 ± 0.04	2.39 ± 0.02	0.17 ± 0.08	1.91 ± 0.04	0.09 ± 0.01	0.11 ± 0.06	0.067 ± 0.001
Fruta-do-conde (<i>A. squamosa</i>)	0.91 ± 0.09	0.22 ± 0.03	0.12 ± 0.02	0.31 ± 0.08	4.24 ± 0.31	0.04 ± 0.01	0.12 ± 0.03	0.018 ± 0.001
Graviola (<i>A. muricata</i>)	0.87 ± 0.12	0.39 ± 0.02	0.09 ± 0.00	0.19 ± 0.04	8,76 ± 0,31	0.07 ± 0.01	0.17 ± 0.02	0.012 ± 0.001
Taperebá (<i>S. mombin</i>)	1.13 ± 0.05	0.19 ± 0.03	0.04 ± 0.00	0.07 ± 0.00	3.24 ± 0.83	0.02 ± 0.00	0.19 ± 0.01	N.D.

249 N.D. not detected. The readings were done in triplicate. The standard deviation was calculated for a level of significance of the t-student of 95%

250

251

252

253

254

255

256

257

258

259 Among the macroelements, the high values of potassium in the fruits studied were the lowest,
260 with the lowest value presented for the caçari pulp, with $124.13 \pm 0.12 \text{ mg } 100 \text{ g}^{-1}$ and 541.16 ± 0.24
261 $\text{mg } 100 \text{ g}^{-1}$. The levels of potassium daily are 4700 mg day^{-1} [20], with consumption of potassium-rich
262 foods beneficial for controlling blood pressure, type II diabetes and bone health.

263 Phosphorus is an essential element that, besides appearing in fruits, its main contribution to the
264 organism is the source of animal origin, mainly in red, white and viscera meats [21], in fruits, the
265 phosphorus levels oscillate between $20\text{-}100 \text{ mg } 100 \text{ g}^{-1}$ [22]. The fruits in the study presented low
266 values of phosphorus, being the caçari pulp, which presents a lower value $6.21 \pm 0.04 \text{ mg } 100 \text{ g}^{-1}$ and
267 for taperebá $24.12 \pm 0.11 \text{ mg } 100 \text{ g}^{-1}$, the recommended dose of phosphorus per day is 800 mg [22].

268

269 The fruits studied, it is the abiu that presents a lower concentration of calcium $4.51 \pm 0.02 \text{ mg } 100 \text{ g}^{-1}$, 52.21
270 $\pm 0.13 \text{ mg } 100 \text{ g}^{-1}$. The nutritional contribution of Ca in adults is $1000\text{-}1200 \text{ mg day}^{-1}$ [23].

271 Magnesium is other important macroelement in fruits, and it appears within a very variable range in the
272 fruits studied. In the abiu pulp, it is in low concentrations $1.71 \pm 0.07 \text{ mg } 100 \text{ g}^{-1}$, presenting the highest
273 concentration of magnesium for pulp of the biribá with $112.32 \pm 0.12 \text{ mg } 100 \text{ g}^{-1}$. The main function of
274 magnesium in the body is to stabilize the structure of ATP in enzymatic reactions, as cofactor in enzymatic
275 reactions, in neuromuscular transmission [24].

276 As for sulfur, it is required in small concentrations, being an element that forms part of the structure of
277 essential amino acids such as cysteine and methionine and enzymatic activator [25]. Of the fruits studied, it is
278 the acerola that has the highest concentrations of sulfur, $34.13 \pm 0.14 \text{ mg day}^{-1}$, being the pulp of taperebá,
279 which presents lower concentrations.

280 Nitrogen is not one of the most studied micronutrients in fruits, being more studied associated with the
281 proteins of the fruit. Its fruit pulp quantity is low, presenting the lowest value for the bacuparí with $0.13 \pm$
282 0.01% , being the fruit that presents a higher value the pulp of araçá with $0.68 \pm 0.07\%$.

283 In the case of graviola, the values obtained for the case of Ca, K and Mg are close to those obtained [26] in
284 the sodium and potassium are lower than those obtained by the same author and in the case of sulfur the value
285 obtained is bigger.

286 In the fruit of the earl, the value obtained from Ca, K and S, are similar to those obtained by the same author
287 for the Mg and Na are many lower than those presented [26].

288 Among the micronutrients, iron is very important in the human diet, because its deficiency can cause anemia,
289 fatigue and impairment in neurological growth and development [27]. The highest values of iron presented in
290 this work are for araçá with concentrations of $3.04 \pm 0.02 \text{ mg } 100 \text{ g}^{-1}$, the fruit having the lowest amounts of
291 iron $0.18 \pm 0.04 \text{ mg } 100 \text{ g}^{-1}$. Another of the rich fruits in iron is the pulp of biribá with $1.82 \pm 0.11 \text{ mg } 100 \text{ g}^{-1}$.

292 As for zinc, it is important in the organism at the physiological level as an antioxidant [28], as well as
293 developing a fundamental role in the polymer organization of macromolecules such as DNA and RNA, as well
294 as their synthesis [29]. According to Food and Nutrition Board [30]. The zinc recommendations for the
295 population are 8 mg day^{-1} for women and 11 mg day^{-1} for men. In the fruits studied, the highest pulp
296 concentration was pulp pulp $3.71 \pm 0.22 \text{ mg } 100 \text{ g}^{-1}$, with the acerola pulp being $0.08 \pm 0.01 \text{ mg } 100 \text{ g}^{-1}$.

297 Several Amazonian fruits, among them the araçá and acerola, presenting values of zinc concentration in the
298 araçá of $0.17 \text{ mg } 100 \text{ g}^{-1}$ e for acerola of $0.19 \text{ mg } 100 \text{ g}^{-1}$ [26], for the edible fraction of fruit, being for acerola
299 lesser than that obtained in this work (table 5) and in the case of araçá is smaller than the value presented in
300 Table 5.

301 Other important microelement in enzymatic metabolic reactions is manganese is part of two
302 metalloenzymes, carboxylase pyruvate and Mn-superoxide dismutase [31].

303 Among the studied fruits, the abiu presents high concentrations for the pulp with $6.61 \pm 0.11 \text{ mg } 100 \text{ g}^{-1}$.
304 Other fruits with considerable concentrations of manganese are the caçari pulp with $2.39 \pm 0.02 \text{ mg } 100 \text{ g}^{-1}$,
305 with the lowest values of manganese in the graviola pulp $0.09 \pm 0.00 \text{ mg } 100 \text{ g}^{-1}$ and the taperebá pulp $0.04 \pm$
306 $0.00 \text{ mg } 100 \text{ g}^{-1}$.

307 Minerals studies in tropical fruits and found values of manganese for the graviola of $0.07 \pm 0.02 \text{ mg } 100$
308 g^{-1} low value in relation to the value found in the present work for the pulp $0.39 \pm 0.02 \text{ mg } 100 \text{ g}^{-1}$ and for the
309 fruit of the count $0.16 \pm 0.00 \text{ mg } 100 \text{ g}^{-1}$, a value close to the finding in this work for the pulp $0.12 \pm 0.02 \text{ mg}$
310 100 g^{-1} [32]. On the other hand, evaluated fruits cultivated in Colombia and found manganese values for the
311 *araçá* $0.08 \text{ mg } 100 \text{ g}^{-1}$ [26], the value found in this work for the pulp of $1.25 \pm 0.07 \text{ mg } 100 \text{ g}^{-1}$ slightly higher
312 than the value described by the previous author and for acerola $0.09 \text{ mg } 100 \text{ g}^{-1}$, being lower than that found in
313 the present study with a value of $0.24 \pm 0.05 \text{ mg } 100 \text{ g}^{-1}$.

314 Copper is a trace element that may exhibit various oxidation states and within the cell predominates the
315 cuprous ion [33]. Copper levels, compared to the other elements, are low, with the exception of *araçá* that
316 presents copper concentrations of $1.73 \pm 0.02 \text{ mg } 100 \text{ g}^{-1}$ for the pulp, and the taperebá is the one with the lowest
317 concentration of copper, with only $0.07 \pm 0.00 \text{ mg } 100 \text{ g}^{-1}$.

318 The need for copper is $1\text{-}2 \text{ mg dia}^{-1}$, and 10 mg dia^{-1} is tolerated according to DRIs [34] for the maintenance
319 of the human organism, the above fruits being above tolerable levels for the organism [32] determined copper
320 concentrations in the soursop of $0.15 \pm 0.03 \text{ mg } 100 \text{ g}^{-1}$, being close to that found in the present study with 0.19
321 $\pm 0.04 \text{ mg } 100 \text{ g}^{-1}$ and for $0.22 \pm 0.03 \text{ mg } 100 \text{ g}^{-1}$, the concentration of copper for the pulp in $0.31 \pm 0.08 \text{ mg}$
322 100 g^{-1} , slightly higher to the value found in the literature.

323 An important trace element is boron, being related to the cerebral metabolism [35] among other functions.
324 In the case of fruits, boron has an important function of stimulating the germination and generation of pollen
325 and pollen tube growth, being a fundamental factor for the adequate formation of fruits [36]. The highest
326 concentration of boron was found in the pulp of the biribá $0.51 \pm 0.05 \text{ mg } 100 \text{ g}^{-1}$ and the lowest concentration
327 was the aracá $0.10 \pm 0.02 \text{ mg } 100 \text{ g}^{-1}$.

328 The aluminum is a toxic metal, whose concentration in food is low, of the order of 5 mg Kg^{-1} [37]. The
329 consumption of foods contaminated by this metal may be related to Alzheimer's disease [38]. Thus, the fruits
330 analyzed had relatively low concentrations, varying between $0.02 - 0.17 \text{ mg } 100 \text{ g}^{-1}$ in pulps, being within the
331 recommended levels.

332 Among all the evaluated minor elements, cobalt is the lowest concentration in relation to the
333 microconstituents, only present in some of the studied fruits and the highest values show the crab with $0.067 \pm$
334 $0.001 \text{ mg } 100 \text{ g}^{-1}$. The estimated cobalt doses are between $0.5\text{-}1.4 \text{ mg dia}^{-1}$, therefore, the levels found in the
335 fruits studied would be below the recommended levels [39].

336 A present a study of determination of seven minerals in different Amazonian fruits in the different parts of
337 the same, among them the biribá, being the values obtained for this fruit, very close to those presented with the
338 exception of the sodium concentration obtained for the pulp of the biribá that presents a much lower value [17].

339 A study of mineral determination in acerola *in natura* and juice, presenting very close values for Mg, P, K
340 and Zn, and for the Fe, Na and Cu elements we obtained larger values compared to the author [40].

341 The results obtained in this study were similar to those and the values for Ca, Mg and Ca were higher for P,
342 K, Na, Fe, Mn and Cu [41].

343

344 3.3. Statistic analysis

345

346 3.3.1. Pearson correlation coefficient

347

348 Table 6 presents the Pearson correlation matrix between the different elements for the pulps of the
349 different fruits.

350

351

352

353

354

355

356

357

358

359

360

361

362

363

364

365

366

367

368

369

370 **Table 6** - Pearson correlation matrix between the different elements for the pulps of Amazonian fruits.

	Ca	Mg	P	K	S	N	Fe	Zn	Mn	Cu	Na	Al	B	Co
Ca	1													
Mg	0,76*	1												
P	0,40ns	0,13ns	1											
K	0,69*	0,43ns	0,36ns	1										
S	0,22ns	0,23ns	-0,29ns	0,42ns	1									
N	0,02ns	-0,08ns	0,60ns	0,12ns	-0,14ns	1								
Fe	0,28ns	0,22ns	0,76*	0,31ns	-0,04ns	0,92**	1							
Zn	-0,25ns	-0,23ns	0,37ns	0,11ns	-0,47ns	0,02ns	0,10ns	1						
Mn	-0,66*	-0,41ns	-0,43ns	-0,28ns	-0,32ns	-0,10ns	-0,26ns	0,56ns	1					
Cu	0,10ns	0,08ns	0,68*	0,27ns	-0,09ns	0,94**	0,97**	0,22ns	-0,06ns	1				
Na	-0,03ns	0,28ns	0,01ns	-0,02ns	0,66*	0,21ns	0,34ns	-0,34ns	-0,34ns	0,27ns	1			
Al	-0,48ns	-0,21ns	-0,25ns	-0,40ns	0,52ns	-0,11ns	-0,10ns	-0,17ns	-0,05ns	-0,10ns	0,80**	1		
B	0,42ns	0,80**	0,17ns	0,40ns	-0,03ns	0,09ns	0,35ns	0,26ns	0,13ns	0,32ns	0,15ns	-0,27ns	1	
Co	0,21ns	0,45ns	-0,14ns	0,08ns	-0,15ns	-0,09ns	-0,04ns	-0,18ns	-0,08ns	0,02ns	-0,11ns	-0,33ns	0,34ns	1

371 Subtitle: ns (not significant) $p > 0.05$, * $p < 0.05$, ** $p < 0.01$

372

373

374

375

376

377

378 In Table 6, Pearson correlation coefficient showed highly significant correlation values at 1% significance
 379 for the following elements: copper with iron (0.97), iron with nitrogen (0.92), copper with nitrogen (0.94) and
 380 aluminum with sodium (0.80). On the other hand, there are significant interactions at the significance level of
 381 5%, calcium with magnesium (0.76), calcium with potassium (0.69), and at the same time, calcium with
 382 manganese (0.66). Phosphorus also has a significant interaction with iron and copper. In the case of the
 383 interaction of iron with phosphorus presents with Pearson correlation coefficient (0.76) and for the interaction
 384 of phosphorus with copper (0.68). Sodium with sulfur also has a significant interaction of (0.66). The other
 385 elements do not present significant interactions between them.

386


387 3.3.2. Principal component analysis (PCA)

388 The analyzes of main components were carried out jointly for the evaluated systems (abiu, bacupari,
 389 acerola, graviola, caçari, araçá, biribá and taperebá), independently for each part of the fruit, in order to (minerals
 390 present in different parts of the fruit), in order to find a new set of variables (main components), uncorrelated,
 391 that explain the structure of the variation, being represented the weight of each variable analyzed in each
 392 component (axes).

393

394 **Figure 2.** Distribution of the original variables between the different fruits for the pulp on the first and second
 395 main component (CP1 and CP2).

396



397

398 In the biplot (Figure 2), the results of the analysis of the main components (PCA) for the pulps of the
 399 different fruits are presented, explaining the 52.3% of the original variability of the data retained in these
 400 components.

401 The arrangement of the sequence in Figure 2 of the supplementary material shows that the systems can be
402 grouped into two sets, the first major component (CP1), contributed with 30.8% of the total variance explained,
403 however most of the minerals that were strongly affected, between (S), sodium (Na), magnesium (Mg), calcium
404 (Ca), cobalt (Co), potassium (K) and boron (B) contributing positively to CP1 and inverse with aluminum
405 elements) and manganese (Mn).

406 These results indicate that CP1 allowed to distinguish the fruits that are associated to the minerals, being
407 the fruit araçá, biribá, fruit of the count and graviola that have been associated, being the last three of the same
408 family.

409 The second main component (CP2) explained 21.5% of the total data, relating the elements aluminum
410 (Al) and manganese (Mn). The analysis of this component also showed that these attributes negatively projected
411 on the elements S, sodium (Na), magnesium (Mg), calcium (Ca), cobalt (Co), potassium (K) and boron (B), being
412 the fruits acerola, taperebá, abiu, caçari and bacuparí who were associated.

413

414 4. Conclusions

415 The present work establishes the nutritional importance of Amazonian fruits, which present a great
416 richness in minerals especially in micronutrients, establishing a correlation between the different constituents,
417 as well as establishing methods of multivariate analysis to establish the relationship between the different
418 studied variables.

419 Due to the nutritional importance of fruits, they could be used to develop bioproducts with interest being at the
420 same time a part with high energetic potential.

421

422 **Acknowledgments:** To CAPES for the scholarship and agronomy research laboratory (NUPAGRI) of the Federal
423 University of Roraima (Brazil).

424

425 **Conflicts of Interest:** The authors declare no conflict of interest.

426

427 **References**

- 428 1. Hubbell, S.P.; He, F.; Condit, R.; Borda-de-água, L.; Kellner, J.; Ter, S.H. How many tree species are there
429 in the Amazon and how many of them will go extinct? *Proceedings of National Academy of Sciences*. **2008**, *105*,
430 11498-11504.
- 431 2. Mariutti, L.R.B.; Rodrigues, E.; Chisté, R.C.; Fernandes, E.; Mercadate, A.Z.; Freitas, S. P. The Amazonian
432 fruit *Byrsonima crassifolia* effectively scavenges reactive oxygen and nitrogen species and protects human
433 erythrocytes against oxidative damage. *Food Research International*. **2014**, *64*, 618-625.
- 434 3. Drewnowski, A.; Fulgoni, V. Nutrient profiling of foods: creating a nutrient-rich. *Food Index*. **2014**, *66*, 23-
435 29.
- 436 4. Darmon, N.; Briend, A.; Drewnowski, A. Energy-dense are associated with lower diet costs; a community
437 study of french adults. *Public Health Nutrition*. **2004**, *7*, 21-27.
- 438 5. Hardisson, A.; Rubio, C.; Baez, A.; Martin, M.; Alvarez, R.; Diaz, E. Mineral composition of the banana
439 (*Musa acuminata*) from the island of Tenerife. *Food Chemistry*. **2001**, *73*, 153-161.
- 440 6. Krause, M.V.; Mahan, L.K. *Minerals*. In: food, nutrition and diet therapy. 11 ed. São Paulo, 2005.
- 441 7. Avila, O.V.; Fernández, I.M.; da Costa, H.N.R.; Melo Filho, A.A.; Ribeiro, P.R.E. Bromatological Analysis,
442 Chemical Composition and Bioassays from the Genipa americana L. (Rubiaceae). *Journal of Agricultural
443 Science*. **2018**, *10*, 244-252.
- 444 8. Instituto Adolfo Lutz (IAL). Physicochemical methods for food analysis (IV ed.) São Paulo, 2008.
- 445 9. Mendes-Filho, N.E.; Carvalho, M.P.; de Souza, J.M.T. Determination of macronutrients and minerals
446 nutriente of the mango pulp (*Mangifera indica* L.). *Perspectivas da Ciência e Tecnologia*. **2014**, *6*, 22-36
- 447 10. Empresa Brasileira de Pesquisa Agropecuária- EMBRAPA. Manual of chemical analyzes of soils, plants
448 and fertilizers. 2nd edition revised and extended, Brasília, DF, 627 p, 2009.
- 449 11. Rienzo, J.A.di.; Casanoves F.; Balzarini, M.G.; Gonzales, L.; Tablada, M.; Robledo, C.W. InfoStat Release
450 2016. InfoStat Group FCA, Universidad Nacional de Córdoba, Argentina, 2016.
- 451 12. Welti, J.; Vergara, F. Water activity: concept and application in foods with high moisture content. In:
452 AGUILERA, J.M. *Topics in food technology*. **1997**, *1*, 11-26.
- 453 13. Canuto, G.A.B.; Xavier, A.A.O.; Neves, L.C.; Benassi, M.T. Physical chemical characterization of fruit pulps
454 from Amazonia and their correlation with free radical anti-free radicals. *Rer. Bras. Frutic. Jaboticabal*. **2010**,
455 *32*, 1196-1205.
- 456 14. Maeda, R.N.; Pantoja, L.; Yuyama, L.K.O.; Chaar, J.M. Determination of the formulation and
457 characterization of camu-camu nectar (*Myrciaria dub* McVaugh). *Ciência e Tecnologia de alimentos*. **2006**, *26*,
458 70-74.
- 459 15. Kinupp, V.F.; Barros, I.B.I.; Protein and mineral contents of native species, potential vegetable and fruits.
460 *Ciência e Tecnologia de alimentos*. **2006**, *28*, 846-857.
- 461 16. Aletor, V.A.; Adeogun, O.A. Nutrient and antinutrient constituents of some tropical leafy vegetables. *Food
462 Chemistry*. **1995**, *53*, 3775-3779.
- 463 17. Berto, A.; Da Silva, A.F.; Visentainer, M.M.; de Souza, N.E. Proximate compositions, mineral contents and
464 fatty acid composition of native Amazon fruits. *Food Research International*. **2015**, *77*, 441-449.
- 465 18. Seffrin, R.C.; Shikano, I.; Akhtar, Y.; Isman, M.B. Effects of crude seed extracts of *Annona atemoya* and
466 *Annona squamosa* L. against the cabbage looper, *Trichoplusia* in the laboratory and greenhouse. *Crop
467 Protection*. **2010**, *29*, 20-24.
- 468 19. Santos, R. C.; Melo Filho, A. A.; Chagas, E. A.; Takahashi, J. A.; Ferraz, V. P.; Costa, A. K. P.; Melo, A. C. G.
469 R.; Montero, I. F.; Ribeiro, P. R. E. Fatty acid profile and bioactivity from *Annona hypoglauca* seeds oil. *African
470 Journal of Biotechnology*. **2015**, *14*, 2377-2382.
- 471 20. Food and nutrition board. Dietary reference intakes for vitamin a, vitamin k, arsenic, boron, chromium,
472 copper, iodine, iron, manganese, molybdenum, nickel, silicon, vanadium and zinc. *National Academy of
473 Sciences, Washington*, 2001.
- 474 21. Cozzolino, S.M.F. Bioavailability of nutrients. SP: Manile, 2007.
- 475 22. Tomassi, G. Phosphorus- an essential nutrient for human diet. *Imphos*. **2002**, *16*, 1-3.
- 476 23. Pereira, G.A.P.; Genaro, P.S.; Pinheiro, M.M.; Szenjfeld, V.L.; Martini, L.A. Diet Calcium-Strategies to
477 Optimize Consumption. *Revista Brasileira Reumatol*. **2009**, *49*, 164-180.
- 478 24. Iseri, L.T.; French, J.M. Magnesium-nature's physiologic calcium blocker. *American Heart Journal*. **1984**, *108*.

- 479 25. Silva, D.J.; Pereira, J.R.; do Carmo, M.A.; de Albuquerque, J.A.S.; Van Raji, B.; Silva, C.A. Mineral nutrition
480 and fertilization of the hose under irrigated conditions. *Technical Circular*, **2004**, 77. Ministry of Agriculture,
481 Livestock and Food Supply. EMBRAPA.
- 482 26. Leterme, P.; Buldgen, A.; Estrada, F.; Londoño, A.M. Mineral content of tropical fruits and unconventional
483 foods of the Andes and the rain forest of Colombia. *Food Chemistry*. **2006**, 95, 644-652.
- 484 27. Carvalho, M.C.; Baracat, E.C.E.; Sgarbieri, V.C. Iron deficiency anemia and chronic disease anemia:
485 disturbances of iron metabolism. *Revista SAN*. **2010**, 13, 54-56.
- 486 28. Powell, S.R. The antioxidante properties of zinc. *J. Nut.* **2000**, 130, 1447-1454.
- 487 29. Vallee, B.L.; Falchuk, K.H. The biochemical basis of zinc physiology. *Physiol Rev.* **1993**, 73, 1.
- 488 30. Food and Nutrition Board of the Institute of Medicine, 2013.
- 489 31. Panziera, F.B.; Dorneles, M.M.; Durgante, P.C.; da Silva, V.L. Evaluation of antioxidant minerals intake in
490 elderly. *Ver. Bras. Gerontol.* **2011**, 14, 49-58.
- 491 32. Almeida, M.M.A.; de Souza, P.H.M.; Fonseca, M.L.; Magalhães, C.E.C.; Lopes, M.F.G.; de Lemos, T.L.G.
492 Evaluation of macro and micro-mineral content in fruits cultivated in the northeast of Brazil. *Ciência e*
493 *Tecnologia de Alimentos*. **2009**, 23, 581-589.
- 494 33. Bairele, M.; Valentini, J.; Paniz, G.; Moro, A.; Junior, F.B.; Garcia, S.C. Possible effects of blood copper on
495 hematological parameters in elderly. *Journal Bras Patol Med Lab.* **2010**, 46, 463-470.
- 496 34. Dietary Reference Intakes (DRI). <https://www.nal.usda.gov/fnic/dietary-reference-intakes>
- 497 35. Penland, J.G. Dietary boron, brain function and cognitive performace. *Environ Health Perspect.* **1994**, 102, 65-
498 72.
- 499 36. SangHyun, L.; WolSoo, K.; TaeHo, H. Effects of post-harvest foliar boron and calcium applications on
500 subsequent season's pollen germination and pollen tube growth of pear (*Pyrus pyrifolia*). *Scientie*
501 *Horticulturae*. **2009**, 122, 77-82.
- 502 37. Dantas, S.T.; Saron, E.S.; Dantas, F.B.H.; Yamashita, D.M.; Kiyataka, P.H.M. Determining aluminuim
503 dissolution when cooking food in aluminuim cans. *Ciência e Tecnologia de Alimentos*. **2007**, 27, 291-297.
- 504 38. Martyn, C.N., Coggan, D., Inskip, H., Lacey, R.F., & Young, W.F. (1997). Aluminium concentrations in
505 drinking water and risk of Alzheimer's disease. *Epidemiology*, 8, 281-286.
- 506 39. Martyn, C.N., Coggan, D., Inskip, H., Lacey, R.F., & Young, W.F. (1997). Aluminium concentrations in
507 drinking water and risk of Alzheimer's disease. *Epidemiology*, 8, 281-286.
- 508 40. Vaitsman, D.S.; Alonso, J.C.; Dutra, P.B. What are the chemical elements for? *Editora Interciência*, 2011.
- 509 41. Freitas, C.A.S.; Maia, G.A.; da Costa, J.M.C.; Figueiredo, R.W.; de Sousa, H.M. Acerola: production,
510 composition nutritional aspects and products. *Revista Brasileira Agrociência*. **2006**, 12, 395-400.
- 511 42. Caldeira, S.D.; Hiane, P.A.; Ramos, M.I.L.; Filho, M.M.R. (2004). Physical-chemical characterization of araçá
512 (*Psidium guineense* SW) and tucumã (*Vitez cymosa* Bert.) Of the State of Mato Grosso do Sul. *Ceppa*. **2006**,
513 22, 145-154.