

Running head: ORGANOGENESIS: A NEW THEME

# Organogenesis: A New Theme to Study Genetic Expression in Adult Liver Regeneration After Partial Hepatectomy

Haydar Witwit

National University

## Abstract

Researchers in different disciplines studied liver's genetic expression of organogenesis in embryogenesis; however, organogenesis has not been studied as an independent and complementary process during adult liver regeneration. This paper reviewed studies and extracted information related to organogenesis in adult liver regeneration because of organogenesis' significant role in cancer and tissue regeneration. The paper discussed the accumulated and current status of knowledge of organogenesis after partial hepatectomy, it also dived into the gaps associated with this knowledge and compare it to cellular proliferation process of hepatocytes in liver regeneration. A new theme proposed to signify the role of organogenesis as a separate and parallel process to cellular proliferation in liver regeneration. In addition; the paper reviewed evidence-based contributions that explored the organogenesis and described the associated changes with the targeted factors or genes.

**Key words:** organogenesis, signaling, adult, regeneration, gene, expression, hepatectomy, cancer, hepatocytes, liver

## I. Introduction

The liver has a unique ability to regenerate and to retain a specific three-dimensional shape after partial hepatectomy; researchers investigated the liver's abilities because of the multiple factors involved and the complexities of the processes. The focus of this review is organogenesis, and because it is part of the regeneration process, and it overlaps and intermingles with cellular proliferation, both processes were reviewed. While organogenesis defined as the formation of organs<sup>1,2</sup>, but this definition formulated in the context of embryogenesis<sup>3</sup> and these definitions do not specify which process in the regeneration referred to (three-dimensional constitution, cellular proliferation or regeneration as a whole). We propose a definition to define the term organogenesis as following: [Organogenesis is the process in which the cells constitute the three-dimensional shape, and this in turn may contribute to the function, of an organ in the normal spatiotemporal parameters

of that organ]. With this definition, we generalize the process to both embryonic organogenesis and adult organogenesis and specialize and differentiate the term organogenesis apart from cellular proliferation and liver regeneration. By utilizing this definition, we look for how and why cells move and organize to form the destined three-dimensional shape. Most previous studies focused on the proliferation and differentiation of liver cells (for citations, please refer to Table 1, 2, and 3). In addition, this paper will discuss a theme in which the organogenesis process would be recognized as a separate yet parallel and complementary process in all phases of liver regeneration to help identify factors associated with each process.

Table 1. *Summary of Phosphorylation Processes in Liver Regeneration and Organogenesis*

Process	Observation during proliferation	Observation during organogenesis
Phosphorylation of Lysine-Rich Histone (Balhorn et al., 1971) <sup>31</sup>	High rate	not stated
Nuclear proteins phosphorylation pattern (Ballal et al., 1975) <sup>32</sup>	High rate, except for two protein (All and A24) were markedly decreased.	not stated
Acidic nonhiston proteins phosphorylation of nuclear matrix (Allen et al., 1977) <sup>33</sup>	High rate	not stated
Phosphorylation of H4 histidine (Chen et al., 1977) <sup>34</sup>	High rate	not stated
Polyadenylated mRNA formation (Atryzek & Fausto, 1979) <sup>35</sup>	High level	not stated
Mitochondrial oxidative phosphorylation and mitochondrial glutathione concentration (Vendemiale et al., 1995) <sup>36</sup>	Low level in prereplective phase, but recovered in second phase	not stated
STAT3 phosphorylation (Shu et al., 2009) <sup>11</sup>	Fat excess in hepatocyte harm regeneration process	Steatosis correlates with a defective regeneration. Same observation in adiponectin-deficiency
Tyrosine phosphorylation (Pagano et al., 2012) <sup>26</sup>	Signaling initiation	not stated

Table 2. *Hormonal Effect on Liver Regeneration and Organogenesis*

Hormone	Effect on proliferation	Effect on organogenesis
Parathormone (Rixon et al., 1971) <sup>37</sup>	It is important to maintain parenchymal cells competency to proliferate	not stated
Testosterone (Vic et al., 1982) <sup>38</sup>	Enhancer	not stated
Estrogen (Biondo-Simões Mde et al., 2009) <sup>39</sup>	Enhancer	not stated
Growth hormone (Asakawa et al., 1989 <sup>40</sup> ; Pennisi et al., 2004 <sup>41</sup> )	Enhancer	not stated
Steroid (Glanemann et al., 2004) <sup>42</sup>	no effect	not stated
Prolactin (Moreno-Carranza et al., 2011) <sup>43</sup>	Enhancer	not stated
Thyroid hormone (Alisi et al., 2004) <sup>44</sup>	Enhancer	not stated

Table 3. *Factors Associated with Liver Regeneration and Organogenesis*

Factors	Observation during proliferation	Observation during organogenesis
DNA polym+A4:D13 (Fry et al., 1985) <sup>45</sup>	There was an increase of 13 folds of DNA polymerase alpha stimulating protein during liver regeneration	not stated
Albumin and alpha feto-protein (Panduro et al., 1986) <sup>46</sup>	A reduction in Albumin level and increase level of Fetoprotein was observed during liver regeneration	not stated
Acidic nonhiston proteins phosphorylation of nuclear matrix (Allen et al., 1977) <sup>33</sup>	High rate	not stated
Hepatocytes' number and size, surface and volume fractions, their nuclei, cytoplasm, and their relation to the sinusoidal bed (Vizzotto et al., 1989) <sup>22</sup>	An increase in the number of nuclei and the nucleus/cytoplasm ratio associated with regenerating liver	not stated
Transcription factors (c-jun, jun B, c-fos fos B, fra 1, zif268, nurl77) (Dubois, 1990) <sup>47</sup>	High level of transcription factors is associated with liver regeneration	not stated
Hepatocyte Growth Factor (HGF) mRNA and DNA synthesis (Zarnegart et al., 1991) <sup>48</sup>	High level of Hepatocyte Growth Factor (HGF) and an increase of DNA synthesis associated with liver regeneration process, researchers direct toward that HGF is a signaling factor	not stated
DNA Polymerase delta, alpha and proliferating cell nuclear antigen activities (Yang et al., 1991) <sup>49</sup>	High levels of DNA Polymerase delta, alpha and proliferating cell nuclear antigen activities observed in association with regenerating rat liver	not stated

## ORGANOGENESIS: A NEW THEME

3

LDH, uric acid, albumin, DNA, laminin and collagen IV (Parsons-Wingenter & Saltzman, 1993) <sup>24</sup>	High levels of uric acids and albumin associated with proliferation compared to low level in hepatocytes monolayer, also high level of laminin and collagen IV associated with proliferation compared to low level with mature cells	not stated
STAT3 (signal transducer and activator of transcription 3), NF- $\kappa$ B (nuclear factor kappa B), OVA1, OVA2 and OVA3 (Sanchez et al., 2004) <sup>50</sup>	OV1 used as a marker for identification and sorting of oval cells, researchers established that both NF- $\kappa$ B and STAT3 were highly activated in the OV1+ cell population.	not stated
Tissue inhibitors of metalloproteinases (TIMP-1) (Mohammed et al., 2005) <sup>51</sup>	TIMP-1 inhibits proliferation process.	not stated
Hepatocyte nuclear factor (vHnf1) (Lokmane et al., 2008)	vHnf1 deficiency associated with abnormal growth	Absence of vHNF1 forms defective hepatic bud and leads abnormal gut regionalization
Bone morphogenic protein-7 (BMP-7)	Please refer to organogenesis observation	Liver growth defect was observed and Was associated with Bone morphogenic protein-7 (BMP-7) deficiency
Myeloid differentiation factor (MyD) 88 (Seki et al., 2005) <sup>52</sup>	It is important in signaling pathway for proliferation	not stated
Fibroblast Growth Factor (FGF) (Wang et al., 2015) <sup>5</sup>	Absence of FGF signaling associated with halting of anterior murine liver bud	FGF signaling is required for anterior but not posterior specification of the murine liver bud
Lymphatic vessel endothelial hyaluronan receptor 1 (LYVE1) and prospero homeobox protein 1 (PROX1) (Meng, 2017) <sup>12</sup>	PROX1 and LYVE1 correlate with regeneration process and can be used as biomarkers to identify formation of new liver sinusoidal endothelial cells.	not stated
Tumor Necrosis Factor-alpha (TNF- $\alpha$ ) and Inteleukin-6 (IL-6), Transforming Growth Factor Beta (TGF- $\beta$ ) and related TGF- $\beta$ family members (Tao et al., 2017) <sup>53</sup>	TNF- $\alpha$ and IL-6 act as complete mitogen, (TGF- $\beta$ ) and related TGF- $\beta$ family members associated with termination of liver regeneration	not stated

Studies investigated adult liver organogenesis, and factors associated with it were scarce (Lokmane et al., 2008<sup>4</sup>; Wang et al., 2015<sup>5</sup>). Signaling and genes expression that control three-dimensional modeling of liver are important for many reasons, including the identification of termination factors that may help in controlling both proliferation and organogenesis in tumor tissue, and the identification of signaling factors that may help in initiating organogenesis of regenerative tissue for organ donation. Researchers also proposed phases to describe these processes (Xu et al., 2005<sup>6</sup>; Zimmermann, 2002<sup>7</sup>). There is gap in current knowledge because few studies identified causality of specific factors in the processes, especially morphogens. Furthermore, some factors associated with triggering and termination of organogenesis are still unknown (see Table 3), and most importantly, there is a lack of controlled studies to identify factors that control how the liver achieves its final three-dimensional shape. Some studies experimented with models of three-dimensional cultures to study a variety of effects (Li et al., 2018<sup>8</sup>; Tegge et al., 2018<sup>9</sup>). In addition, similarities (or a lack thereof) between adults and embryonic genetic expression of organogenesis

process need to be investigated (Weiss et al., 2016<sup>10</sup>). The paper reviewed and critiqued the early measures of regeneration activities, the hormonal role, the factors role, the genes involved, the three-dimensional modeling of adult liver organogenesis, and the stages and phases of adult liver organogenesis. Then, the paper discussed and proposed a new theme to consider adult liver organogenesis as a separate and parallel process to cell proliferation in liver regeneration. The information was tabulated for readability.

## **II. Early Methodologies and Measurements of Regeneration Activities**

Emergence of new discoveries related to phosphorylation and mRNA directed the focus of researchers toward the application of these concepts to explore their association with liver regeneration. Studies found that high phosphorylation activities are associated with cell proliferation; however, these studies did not state whether or not an association with organogenesis exists (see Table 1). Table 1 reveals important studies in regeneration and their lack of indication to organogenesis in regeneration process. An exception to these studies is Shu et al.'s (2009)<sup>11</sup> contribution of an important observation associated with STAT3 phosphorylation (see orange highlight in Table 1): their team observed that defective regeneration correlates with steatosis, this observation extended to the group with adiponectin-deficiency, but this study did not describe the pattern in which the defective liver regenerates.

## **III. Hormones**

Hormonal effects were studied to explore whether hormones have an enhancing or inhibitory effect on the regeneration process, but studies were not specific as to whether or not hormones have any effect on organogenesis (Table 2). Proliferation could give rise to mass, but organogenesis is the process that determines how the liver forms its three-dimensional figure, lobar shapes, and chirality.

## **IV. Factors Associated with Regeneration Process Pace**

While many studies investigated different factors involved in liver regeneration, only a few researched factors involved in organogenesis (Table 3). Wang et al. (2015)<sup>5</sup> observed that the absence of FGF signaling associated with halting of only anterior murine liver bud; FGF signaling was specific for anterior hepatic bud and it has no effect on the posterior portion. The posterior hepatic bud might be controlled by another signaling pathway. Lokmane et al. (2008)<sup>4</sup> investigated hepatocyte nuclear factor (vHnf1) gene and its effect on regeneration process and concluded that a

lack of vHNF1 leads to formation of a defective hepatic bud and an abnormal gut regionalization. Meng (2017)<sup>12</sup> found that PROX1 and LYVE1 correlate with liver regeneration and can be used as biomarkers to identify formation of new liver sinusoidal endothelial cells. However, Meng (2017)<sup>12</sup> did not indicate how the endothelial cells constitute sinusoidal final shape. The abovementioned studies found factors correlated with organogenesis (see orange highlight in Table 3). In addition, these studies demonstrate the importance of organogenesis in liver regeneration. The rest of the studies in Table 3 showed association of factors with regeneration process; however, they did not indicate the association with organogenesis.

### V. Genetics

Insufficient studies identified genes involved in organogenesis (including signaling factors or inhibitors associated with gene activities). Arai et al. (2003)<sup>13</sup> conducted extensive gene expression analysis during liver regeneration of rats and found 496 out of 2,304 genes showed a measurable expression with up- and down-regulation, including 317 genes showed at least 2-fold change. Nevertheless, Arai et al. (2003)<sup>13</sup> did not indicate how these genes are associated with the organogenesis of the liver after partial hepatectomy. Youn et al. (2003)<sup>14</sup> reported high expression of *abdB* *hox* genes during liver regeneration of murine rats; however, *hoxa1* and *b* had the same level pre- and post-organogenesis process, indicating both *hoxa1* and *b* are not involved in this organogenesis process. Group IX and X—including *hoxad10* and *10*—of *hox* genes are regeneration-specific. Matz-Soja (2016)<sup>15</sup> reviewed signaling pathways of morphogenesis in adult liver mathematical model (e.g., Hedgehog and *Wnt/β-Catenin* pathways). These signaling pathways were lagging yet had similarities with those that occur in embryogenesis. Matz-Soja (2016)<sup>15</sup> further emphasized the importance of signaling pathways of morphogenesis in initiating and orchestrating liver regeneration. *Homeobox containing 1* could prevent liver tumor progression; its action might be attributed to different mechanisms, including promoting autophagy, inhibiting CSC phenotype, or increasing tumor cells' sensitivity to natural killer cell cytolysis (Zhao et al., 2018)<sup>16</sup>. The Wingless/Integrated “*Wnts*” signaling pathway was confirmed to have an important role in hepatic progenitor cells' fate (Boulter et al., 2012)<sup>17</sup>; however, it was not demonstrated how its deficiency would affect organogenesis. Godoy et al. (2013)<sup>18</sup> reviewed morphogens—particularly Wnts (p. 1327) and their team identified canonical Wnts' signaling pathway as a major determinant in

zonation of the parenchyma. *β-Catenin* is important in the expression of enzymes that metabolize drugs and in localizing liver reactions to xenobiotic inducers (Godoy et al., 2013)<sup>18</sup>. Aging is associated with a decrease in liver ability to regenerate and an increase in transcription of inflammatory and immune response; this aging effect on liver regeneration remains ambiguous, but it might be contributed to a depressed level of extracellular factors and hormones (Pibiri et al., 2015)<sup>19</sup>. Lokmane et al. (2008)<sup>4</sup> studied hepatocyte nuclear factor (vHnf1) gene and concluded that a lack of vHNF1 leads to formation of a defective hepatic bud and an abnormal gut regionalization. Based on their experiment on HNF4A, Taniguchi et al. (2018)<sup>20</sup> indicated that a mutation that reduced HNF4A transcription led to tumor growth; and noted that it acts as both a major regulator of organogenesis and a tumor suppressor to hepatocyte growth. Mizuta et al. (1996)<sup>21</sup> used competitive reverse-transcriptase polymerase chain reaction to clone and sequence *homebox* genes at different phases after partial hepatectomy of rat liver, high levels of caudal related gene (RCdx-3) at one hour, while low levels of *hox* gene (RHoxB5) was noticed; however, other identified *hox* genes [RHoxA1, B2, B3, A4, and A5] that were expressed before and after hepatectomy. The focus of the majority of previous studies was liver regeneration, but the study of *Homeobox containing 1* demonstrated the importance of organogenesis factors in halting tumors; in addition, some of *hox* genes are associated with regeneration; however, their role in organogenesis needs to be determined. Furthermore, there is lack of studies that compare genes and their expression of organogenesis in adults versus embryos.

### VI. Three-Dimensional Modeling of Adult Liver Organogenesis

Multiple studies were conducted to investigate three-dimensional cultures of the liver. A study found that hepatocytes' size increases dramatically 18 hours after hepatectomy; it was followed by a noticeable increment in volume which happens 24 hours after partial hepatectomy (Vizzotto et al., 1989)<sup>22</sup>. Hepatocyte stellate cells play a mediator role in communication between hepatocytes and endothelial cells in terms of endothelial cells morphogenesis (Kasuya et al., 2011)<sup>23</sup>. Parsons-Wingerter & Saltzman (1993)<sup>24</sup> reported that aggregate cultures retained two folds of functional parameter compared to singlet cultures; in addition, functional recovery starts when growth is completed. However, the abovementioned studies did not indicate how the volume cells orient, regionalize, or take specific shape. Three-dimensional organotypic liver culture was used to understand liver sinusoidal endothelial cells' (LSECs) and Kupfer cells' (KCs) behavior in three-

dimensional cultures and found that LSECs, accompanied by KCs, demonstrate a decrement in focal adhesion kinase's (FAK) signaling expression, which is a pathway linked to liver sinusoidal endothelial cells dedifferentiation (Tegge et al., 2018)<sup>9</sup>. A functional three-dimensional spheroid was produced, and a hepatic bud was developed from naive mesenchymal stem cells (MSCs), MSC-derived hepatocytes, hepatic stellate cells (HSCs), HSC- and LSEC-like cells (Li et al., 2018)<sup>8</sup>; however, specific mechanisms and pathways of organogenesis in this study is still unraveled and needs to be determined.

### VII. Three or Six Phases and Two Processes

Most of previous research conducted on liver regeneration concentrated on the tissue's competency to proliferate and the proliferation process. Three distinct and sequential stages were described during liver regeneration: (1) initiation stage involves replicating competency; (2) proliferation stage includes the expansion of cells population; and (3) termination stage: here the cellular growth is halted to terminate regeneration at a set point; however, organogenesis is still vague process (Zimmermann, 2002)<sup>7</sup>. In their review, Pahlavan et al. (2006)<sup>25</sup> used staging to discuss factors associated with regeneration; the team also stated the process is ambiguous, but they predicted the presence of inhibitors to control the size of regeneration. A three-stages model was diagrammed with factors and pathways associated with these factors in the regeneration (e.g., TNF-alpha and IL6 activate TNFR, STAT3, NF-kB and AP1 in priming [initiation] stage) (Pagano et al., 2012)<sup>26</sup>. Xu et al. (2005)<sup>6</sup> recognized six temporal phases based on gene expression: 1) immediate early; 2) early; 3) intermediate; 4) early-late; 5) late; and 6) terminal.

A networks triad was suggested to explain the mechanism of liver regeneration: cytokine, growth factor, and metabolic (Fausto et al., 2006)<sup>27</sup>. There was no indication on how are these networks are correlated with organogenesis. Furchtgott et al. (2009)<sup>28</sup> proposed a model of liver regeneration based on interaction of multiple factors, the model divided regeneration process into three phases: 1) quiescent; 2) primed; and 3) replicating. This model lacks a termination phase as a separate phase, and the model did not discuss organogenesis in liver regeneration.


There was a difference in the staging timelines and phase counts between liver regeneration models in the previously mentioned studies mainly because the phases were established according to the timing of the samples' retrieval, and these models did not reflect the actual timeline of



regeneration events. Another factor was the technology that was used in analyzing the data and the kind of data were analyzed. However, these staging models will aid in the classification and assigning of different factors to their temporal window in future studies.

Stimulators of signaling pathways', enhancers', and inhibitors' roles in organogenesis are not well identified. But, the independency of organogenesis process in liver regeneration is supported by ongoing evidences. A study found that liver growth defect associated with Bone morphogenic protein-7 (BMP-7) deficiency (Sugimoto et al., 2006)<sup>29</sup>, BMP-7 included in table 3 due to lack of consensus on the description of its role -whether or not it is a hormone—. Adult hepatocytes play a major role in organogenesis (Utoh et al., 2016)<sup>30</sup>. Lokmane et al. (2008)<sup>4</sup> reported that the lack of vHNF1 leads to formation of a defective hepatic bud and an abnormal gut regionalization. Organogenesis requires Fibroblast Growth Factor (FGF) signaling for anterior but not posterior specification of the murine liver bud (Wang et al., 2015)<sup>5</sup>. These studies demonstrated that some areas and regions in the liver are controlled by different factors during organogenesis; which makes the process more unique.

The paper proposes a theme (Figure 1) in which organogenesis process would be identified as independent, yet parallel and complementary process to proliferation in all phases of liver regeneration, this theme will help identify factors associated with each process. In this theme, organogenesis starts at prime stage, and continue to direct the expansion of three-dimensional growth and ends the process in termination stage. Factors that affect organogenesis are different from those associated with proliferation because of specificity of these factors to three-dimensional spatial control other than random mass expansion (mass can be achieved by proliferation alone).



### VIII. Discussion

Organogenesis is a distinct process that encompasses multiple factors that need to be studied during proliferation. Signaling pathway, cell-cell interactions, extracellular matrix and multiple factors are associated and required to achieve the organogenesis process. This paper discussed a theme to recognize organogenesis in adult's liver as separate, yet parallel and complementary process to proliferation in order to achieve liver regeneration. The theme will help explain organogenesis's factors rather than proliferation alone in regeneration.

Based on retrieved studies, four important observations were concluded after review stratification: (a) Organogenesis is an overlooked process in adult liver regeneration. (b) The factors associated with organogenesis lack causality relationship and mechanisms. (c) Multiple factors are probably associated with organogenesis; most are unknown. (d) Structural architecture and heterogeneity of cells and tissues (including neovascularization and sinusoidal endothelial

integration) to achieve final liver function are controlled by organogenesis. Further studies are warranted to determine each factor and its relationship to its pathway. Three conclusions were reached: (a) Phases of liver regeneration fall under two parallel processes: organogenesis and cell proliferation. (b) Current regeneration staging classification depends on timing of tissue retrieval from experimental groups for testing and does not reflect actual system biology of liver regeneration. And (c) Time recording is important to identify factors in each phase; however, causality of factors in their phase is more important. Further studies are warranted to determine each factor and its relationship to its pathway.

**Acknowledgement:** I'd like to thank Dr. Veronica Ardi for her help and National University library team.

**Conflict of Interest:** All authors declare no conflicts of interest in this paper.

#### References

1. King RC, Stansfield WD, Mulligan PK. *A Dictionary of Genetics*. Oxford University Press; 2006. Seventh Edition. 314
2. Cammack R. *Biochemistry and Molecular Biology*. Oxford University Press, USA; 2006. Second Edition. 483
3. Daintith J, Martin E. *Oxford Dictionary of Science*. Sixth. (Daintith J, Martin E, eds.). Oxford University Press Inc., New York; 2010. Sixth Edition. 587
4. Lokmane L, Haumaitre C, Garcia-Villalba P, Anselme I, Schneider-Maunoury S, Cereghini S. Crucial role of vHNF1 in vertebrate hepatic specification. *Development*. 2008;135(16):2777-2786. doi:10.1242/dev.023010
5. Wang J, Rhee S, Palaria A, Tremblay KD. FGF signaling is required for anterior but not posterior specification of the murine liver bud. *Dev Dyn*. 2015;244(3):431-443. doi:10.1002/dvdy.24215
6. Xu C-S, Chang C-F, Yuan J-Y, et al. Expressed genes in regenerating rat liver after partial hepatectomy. *World J Gastroenterol*. 2005;11(19):2932-2940. <http://www.ncbi.nlm.nih.gov/pubmed/15902731><http://www.pubmedcentral.nih.gov/arti>

- clerender.fcgi?artid=PMC4305662.
7. Zimmermann A. Liver regeneration: the emergence of new pathways. *Med Sci Monit.* 2002;8(3):RA53-63. doi:2366 [pii]
  8. Li J, Xing F, Chen F, et al. Functional 3D Human Liver Bud Assembled from MSC-Derived Multiple Liver Cell Lineages. *Cell Transplant.* 2018;(29):096368971878033. doi:10.1177/0963689718780332
  9. Tegge AN, Rodrigues RR, Larkin AL, Vu L, Murali TM, Rajagopalan P. Transcriptomic Analysis of Hepatic Cells in Multicellular Organotypic Liver Models. *Sci Rep.* 2018;8(1):1-14. doi:10.1038/s41598-018-29455-x
  10. Weiss MC, Le Garrec J-F, Coqueran S, Strick-Marchand H, Buckingham M. Progressive developmental restriction, acquisition of left-right identity and cell growth behavior during lobe formation in mouse liver development. *Development.* 2016;143(7):1149-1159. doi:10.1242/dev.132886
  11. Shu RZ, Zhang F, Wang F, et al. Adiponectin deficiency impairs liver regeneration through attenuating STAT3 phosphorylation in mice. *Lab Invest.* 2009;89(9):1043-1052. doi:10.1038/labinvest.2009.63
  12. Meng F. LYVE1 and PROX1 in the reconstruction of hepatic sinusoids after partial hepatectomy in mice. *Folia Morphol.* 2017;76(2):239-245. doi:10.5603/FM.a2016.0074
  13. Arai M, Yokosuka O, Chiba T, et al. Gene expression profiling reveals the mechanism and pathophysiology of mouse liver regeneration. *J Biol Chem.* 2003;278(32):29813-29818. doi:10.1074/jbc.M212648200
  14. Youn A study of Hox Gene expression Profile During Murine Liver Regeneration.pdf.
  15. Matz-Soja M. Computational modelling of Hedgehog signalling in liver regeneration. *Drug Discov Today Dis Model.* 2016;22(2016):45-50. doi:10.1016/j.ddmod.2017.06.001
  16. Zhao Homeobox containing 1 inhibits liver cancer progression by promoting autophagy as well as inhibiting stemness and immune escape.pdf.
  17. Boulter L, Govaere O, Bird T. Macrophage derived Wnt signalling opposes Notch signalling in a Numb mediated manner to specify HPC fate in chronic liver disease in human and mouse. *Nat ....* 2012;18(4):572-579. doi:10.1038/nm.2667.Macrophage

18. Godoy P, Hewitt NJ, Albrecht U, et al. *Recent Advances in 2D and 3D in Vitro Systems Using Primary Hepatocytes, Alternative Hepatocyte Sources and Non-Parenchymal Liver Cells and Their Use in Investigating Mechanisms of Hepatotoxicity, Cell Signaling and ADME*. Vol 87.; 2013. doi:10.1007/s00204-013-1078-5
19. Pibiri M, Sulas P, Leoni VP, et al. Global gene expression profile of normal and regenerating liver in young and old mice. *Age (Omaha)*. 2015;37(3):1-14. doi:10.1007/s11357-015-9796-7
20. Taniguchi H, Fujimoto A, Kono H, Furuta M, Fujita M, Nakagawa H. Loss-of-function mutations in Zn-finger DNA-binding domain of HNF4A cause aberrant transcriptional regulation in liver cancer. *Oncotarget*. 2018;9(40):26144-26156. doi:10.18632/oncotarget.25456
21. Mizuta I. RCriIEF Identification of Homeobox Genes Expressed During. 1996;34:1-2.
22. Vizzotto L, Romani F, Ferrario VF, Degna CT, Aseni P. Characterization by morphometric model of liver regeneration in the rat. *Am J Anat*. 1989;185(4):444-454. doi:10.1002/aja.1001850407
23. Kasuya J, Sudo R, Mitaka T, Ikeda M, Tanishita K. Hepatic Stellate Cell-Mediated Three-Dimensional Hepatocyte and Endothelial Cell Triculture Model. *Tissue Eng Part A*. 2011;17(3-4):361-370. doi:10.1089/ten.tea.2010.0033
24. Parsons-Wingerter PA, Saltzman WM. Growth versus Function in the Three-Dimensional Culture of Single and Aggregated Hepatocytes within Collagen Gels. *Biotechnol Prog*. 1993;9(6):600-607. doi:10.1021/bp00024a006
25. Pahlavan PS, Feldmann RE, Zavos C, Kountouras J. Prometheus' Challenge: Molecular, Cellular and Systemic Aspects of Liver Regeneration. *J Surg Res*. 2006;134(2):238-251. doi:10.1016/j.jss.2005.12.011
26. Pagano MA, Tibaldi E, Gringeri E, Brunati AM. Tyrosine phosphorylation and liver regeneration: A glance at intracellular transducers. *IUBMB Life*. 2012;64(1):27-35. doi:10.1002/iub.576
27. Fausto N, Campbell JS, Riehle KJ. Liver regeneration. *Hepatology*. 2006;43(2 SUPPL. 1):45-53. doi:10.1002/hep.20969
28. Furchtgott LA, Chow CC, Periwal V. A model of liver regeneration. *Biophys J*.

- 2009;96(10):3926-3935. doi:10.1016/j.bpj.2009.01.061
29. Sugimoto H, Yang C, LeBleu VS, et al. BMP-7 functions as a novel hormone to facilitate liver regeneration. *FASEB J*. 2006;21(1):256-264. doi:10.1096/fj.06-6837com
  30. Utoh R, Komori J, Kuge H, et al. Adult hepatocytes direct liver organogenesis through non-parenchymal cell recruitment in the kidney. *J Hepatol*. 2018;68(4):744-753. doi:10.1016/j.jhep.2017.12.016
  31. Balhorn R, Rieke W, Chalkley R. Rapid Electrophoretic Analysis for Histone Phosphorylation. 1971;(21):3952-3959.
  32. Raghuveera Ballal N, Kang YJ, Olson MOJ, Busch H. Changes in nucleolar proteins and their phosphorylation patterns during liver regeneration. *J Biol Chem*. 1975;250(15):5921-5925.
  33. Allen L. No Title. 1977;75(1).
  34. Chen CC, Bruegger BB, Kern CW, Lin YC, Halpern RM, Smith R a. Phosphorylation of nuclear proteins in rat regenerating liver. *Biochemistry*. 1977;16(22):4852-4855. doi:10.1021/bi00641a016
  35. Biol RRJM, Biophys AB, Mech- HGM, Atryzek V, Fausto N. Accumulation of Polyadenylated mRNA during Liver Regeneration '. 1979;18(7):1281-1287.
  36. Vendemiale G, Guerrieri F, Grattagliano I, Didonna D. Rat Liver Regeneration. 1995:1450-1454.
  37. Rixon Parathormone Proceedings of the Society for Experimental Biology and Medicine.pdf.
  38. Vic P, Saint-Aubert B, Astre C, et al. Complete Liver Regeneration in One-Stage 90% Hepatectomized Rats Treated with Testosterone. *Hepatology*. 1982;2(2):247S-248S. doi:10.1002/hep.1840020210
  39. de Lourdes Pessole Biondo-Simões MI, Rolf Erdmann TI, Ossamu Ioshii III S, et al. The influence of estrogen on liver regeneration. An experimental study in rats. *Acta Cirúrgica Bras*. 2009;24(1):3-6. doi:10.1590/S0102-86502009000100002
  40. Asakawa K, Hizuka N, Takano K, et al. Human growth hormone stimulates liver regeneration in rats. *J Endocrinol Invest*. 1989;12(5):343-347. doi:10.1007/BF03350004
  41. Pennisi PA, Kopchick JJ, Thorgeirsson S, Leroith D, Yakar S. Role of Growth Hormone (GH) in liver regeneration. *Endocrinology*. 2004;145(10):4748-4755. doi:10.1210/en.2004-0655

42. Glanemann Steroid administration before partial hepatectomy.pdf.
43. Moreno Carranza prolactine liver regeneration.pdf.
44. Alisi A, Demori I, Spagnuolo S, Pierantozzi E, Fugassa E, Leoni S. Cellular Physiology Biochemistry and Biochemistr y Partial Hepatectomy by Regulating Cell Cycle and Apoptosis Proteins. 2005:69-76.
45. Natl BP, Sci A, Fry M, Lapidot J, Weisman-shomer P. A DNA Template Recognition Protein : Partial Purification from Mouse Liver and Stimulation of DNA Polymerase at. 1985;(1974):7549-7556.
46. Panduro A, Shalaby F, Weiner FR, Biempica L, Zern MA, Shafritz DA. Transcriptional Switch from Albumin to  $\alpha$ -Fetoprotein and Changes in Transcription of Other Genes during Carbon Tetrachloride Induced Liver Regeneration. *Biochemistry*. 1986;25(6):1414-1420. doi:10.1021/bi00354a034
47. Mean T. Editorial Early Changes in Gene Expression during Liver Regeneration : What do They Mean ? :1079-1082.
48. Zarnegar R, DeFrances MC, Kost DP, Lindroos P, Michalopoulos GK. Expression of hepatocyte growth factor mRNA in regenerating rat liver after partial hepatectomy. *Biochem Biophys Res Commun*. 1991;177(1):559-565. doi:10.1016/0006-291X(91)92020-K
49. Yang C, Zhang S, Toomey NL, Palmer TN, Lee MYWT. Induction of DNA Polymerase Activities in the Regenerating Rat Liver ? 1991;(1990):7534-7541.
50. Sánchez A, Factor VM, Schroeder IS, Nagy P, Thorgeirsson SS. Activation of NF- $\kappa$ B and STAT3 in Rat Oval Cells during 2-Acetylaminofluorene/Partial Hepatectomy-Induced Liver Regeneration. *Hepatology*. 2004;39(2):376-385. doi:10.1002/hep.20040
51. Mohammed FF, Pennington CJ, Kassiri Z, et al. Metalloproteinase inhibitor TIMP-1 affects hepatocyte cell cycle via HGF activation in murine liver regeneration. *Hepatology*. 2005;41(4):857-867. doi:10.1002/hep.20618
52. Seki E, Tsutsui H, Iimuro Y, et al. Contribution of Toll-like receptor/myeloid differentiation factor 88 signaling to murine liver regeneration. *Hepatology*. 2005;41(3):443-450. doi:10.1002/hep.20603
53. Tao Y, Wang M, Chen E, Tang H. Liver Regeneration: Analysis of the Main Relevant

Signaling Molecules. *Mediators Inflamm.* 2017;2017. doi:10.1155/2017/4256352