

Review

Safety Evaluation of Absorbent Hygiene Pads: a Review on Assessment Framework and Test Methods

Jihyun Bae ¹, Hoonjeong Kwon ^{2,3} and Jooyoun Kim ^{3,4,*}

¹ Department of Clothing & Textiles, Hanyang University; jbae2@hanyang.ac.kr

² Department of Food & Nutrition, Seoul National University; hjkwon@snu.ac.kr

³ Research Institute of Human Ecology, Seoul National University

⁴ Department of Textiles, Merchandising and Fashion Design, Seoul National University; jkim256@snu.ac.kr

* Correspondence: jkim256@snu.ac.kr; Tel.: +82-2-880-6846

Abstract: Disposable absorbent hygiene products have evolved for superior performance, enhancing the convenience of daily lives. Yet the use of disposable hygiene pads has brought safety concerns on chemical exposure, and significant efforts have been made to assess the potential risks associated with use of hygiene pads. This article intends to overview the safety assessment framework of diapers and feminine pads, which includes hazard identification, hazard characterization, exposure assessment, risk characterization, and postmarket risk management. Risk assessment of various constituents are reviewed for quantification methods and conservative estimation of exposure parameters. By reviewing the up-to-date considerations in risk assessment, we aim to provide insightful discussion on safety evaluation of current versions of disposable absorbent products. More clinical testing and postmarket surveillance are needed for continuous monitoring of potential health impacts of advanced products and constituents.

Keywords: absorbent hygiene pad; diaper; feminine pad; risk assessment; safety; hazard; exposure; test method

1. Introduction

Disposable absorbent hygiene products such as diapers and feminine pads have evolved with availability of the advanced materials, and their superior functions over cloth diapers have improved the quality of daily lives. Though feminine pads and baby/adult diapers are used for different purposes, main constituents of the absorbent hygiene product share the similar functions and materials. The absorption capacity of disposable absorbent products had significantly improved with the implementation of superabsorbent polymer (SAP) in the core absorbent layer [1-4]. The safety of SAP material has been evaluated rather thoroughly including its potential risks involving ingestion and dermal contact [2-4]. Also, potential hazards associated with process aids and solvents were investigated for human health risks. For example, phthalates [5] that are used as plasticizer of polymers and chlorine compounds [1,6,7] that are used as bleaching agent were identified as potential hazards; and as the absorbent pads are worn in a prolonged skin contact, the safety assurance of products has been regarded significant.

Regulatory classifications of absorbent products including baby diapers, adult diapers and feminine hygiene pads are varied for countries, and are regulated by different disciplines [8-14]. For example, in US, menstrual pads are regulated under the guidelines (not mandatory) by the Food and Drug Administration (FDA) [10,11]; in EU, they are regulated as broadly applicable consumer products [14-16]; in Korea, baby and adult diapers are regulated by the Korean Ministry of Food and Drug Safety (MFDS) under 'Sanitary Products,' and menstrual pads are regulated under 'Quasi-Drugs' [9]. While the specific criteria may vary among the regulatory bodies, human health risk of products needs to be assured with due diligence from any regulatory frameworks. The commonly adopted is the exposure-based quantitative risk assessment (QRA), of which approach includes hazard identification, hazard characterization, exposure assessment, and risk characterization

[4,5,14,17-27]. Many studies have followed such a procedural framework in assessing the human exposure risks involved with absorbent products. Within such a framework, a wide range of designs have been tested for the toxicological aspects of raw materials, the controlled clinical trials, and postmarket surveillance [28,29].

Among the absorbent pads, many risk assessment was done dominantly for baby diapers [2-5,18,24,30-40], while there would exist similar potential risks for feminine hygiene pads and other absorbent products. This paper aims to review the risk assessment methodology that had been applied to disposable absorbent products including baby/adult diapers and feminine hygiene pads. The constituents of absorbent pads and the potential risks associated with dermal exposure are reviewed with specific focus on assessment framework, quantification methods and exposure profiles. By reviewing the up-to-date considerations in risk assessment framework, it is intended to bring up insightful discussion on safety evaluation of current versions of disposable absorbent products.

2. Disposable absorbent hygiene pads

2.1. Types of disposable absorbent hygiene pads

Disposable absorbent hygiene pads consist of several different product categories including baby diapers, feminine hygiene pads, and adult urinary incontinence diapers. Table 1 shows the user and usage profiles for different absorbent products. Among the product types, baby diapers have been studied rather thoroughly in the perspective of safety and risk assessment [1-5,18,21,24,30,33-41]. Baby diapers are generally used for babies in 0-36 months old to absorb urine and feces [3]. In assessing the risks associated with baby diapers, biometric profiles and usage characteristics have been surveyed; which includes body weight [3,42-45], surface area [46], frequency of use [3,24], length of wear, and exposure routes [2,3,47].

Feminine pads are used for menstrual hygiene by women in the age of approximately 14-49 years old. Unlike baby diapers, the pads are worn periodically for about a week per month [14]. The risk assessment frame for feminine hygiene pads has a lot in common with that of baby diapers, but with different user and application characteristics [14,17,29,41,48-52].

Adult incontinence diapers are used for protection from urine leakage, and little studies are available for this product category [53]. However, as the disposable absorbent pads share the common aspects of product constituents and usage habit, a similar assessment framework can be used, if relevant user and usage profiles are employed.

Table 1. Types of disposable absorbent hygiene pads.

Pad type		Baby diaper	Feminine hygiene pad	Adult incontinence diaper
User profile	Age	0-36 months ¹	14-49 years ³	NA
	Body weight	TWA body weight 10.2 kg ¹	50-60 kg ³	60 kg ⁶
Typical application	Frequency	4.7 diapers/day ²	5-7.5 pads/day ⁴ 7 days/month ⁵	2-5 pads/day ⁶
	Length of use	36 months	35 years	NA

¹ TWA body weight is reference value from US EPA 2011 [44]. Values for mean body weight, 10th percentile and 95th percentile body weights for different age groups are reported in references [2,3].

² A 4.7 pads/day is a mean value for all size diapers from the US diary data from Dey et al.'s study [3].

³ 50 kg is referenced from Woeller & Hochwalt's study [14] is noted while it has a wide range.

⁴ 5 pads/day is referred from Woeller & Hochwalt's study [14]; 7.5 pads/day is referred from Korean MFDS [54].

⁵ Data based on Korean MFDS [54].

⁶ Data based on Proctor & Gamble consumer research referenced in Rai et al.'s study [53]. USA institutional and home care uses 3-5 pads or briefs/day; Japan uses 3.5-4 pads/day or 1.5-1.75 pull-on diapers/day.

2.2. Layer construction of absorbent hygiene pads

Figure 1 shows a typical layer construction of a disposable absorbent hygiene pad, which consists of a fluid permeable top sheet, an optional acquisition layer, a superabsorbent core, and a fluid impermeable back sheet with and without adhesives [2,4,14,55]. The layer structure of baby diapers, feminine hygiene pads, and adult diapers are very similar; whereas the design of fastening systems (such as fastening cuffs/wings and adhesives) are different between baby diapers and feminine pads.

For feminine pads, main layer design has changed little since its earliest commercial product by Johnson & Johnson in 1896 [29], while the constituent materials have been evolved for higher level of protection and comfort. One of the main improvements was made in 1970s when the fastening adhesive was added to the back sheet of the pads, replacing pins and belts of the fastening system. Further improvement was made with incorporation of superabsorbent gel particles in the absorbent core, which allowed a higher capacity and quick absorption of fluids [1-4,29,30,40].

For all type of absorbent pads, the topsheet commonly consists of a thin layer of perforated polypropylene/or polyethylene nonwoven and this layer is in direct contact with skin [56]. The main function of this layer is to transfer fluids and feces quickly to the layer beneath, thus a high wicking ability is desirable. Mostly, emollient or lotion is applied onto the topsheet material to grant protection from irritation and skin softness [2,14,49,50,52].

Between the topsheet and the absorbent core, an optional acquisition layer (also called as distribution layer) may exist for modern design of pads [4]. This layer is composed of a cellulose patch and a polyester nonwoven, and its function is to facilitate the spread of fluid evenly across the entire area. Also, this layer helps prevent fluid reflux by retaining the fluid, and helps fluid transfer to the next absorbent layer.

The absorbent core beneath the acquisition layer functions for fluid storage. This layer is composed of a blend of superabsorbent polymer granules and fluff cellulose, which are encapsulated by cellulose or polypropylene nonwoven. The superabsorbent polymer (SAP) is typically made from sodium polyacrylate granules [4,14,33,34], and it transforms into a gel-like substance once it is wet, absorbing up to 30 times its weight liquid. The superabsorbent material alone has been thoroughly evaluated in many studies for skin irritation, sensitization, systemic toxic effects, and concluded to be safe for human use [4,30,40]. The cellulose fluff in the absorbent core helps to absorb liquid quickly and transfer to SAP [4]. Then fluid is locked and stored within this core layer even under applied pressure. For the core constituents to be transferred to skin, the constituents need to be first solubilized in the fluid vehicle then released from the layer upon the applied pressure [2,3,47]. As SAP is not water soluble, it is hardly released from the pad by wetting or reflux.

The backsheet is typically made of a water-proof polyethylene or polypropylene film laminated with polypropylene nonwoven [2,14,55]. This layer prevents fluid from leaking. For feminine pads, the backsheet has adhesives to fix on cloth. Baby diapers employ different fastening system, thus do not have such adhesives on the backsheet. Constituents of the backsheet and adhesives are reported to have negligible skin exposure through reflux [14].




Figure 1. Layer construction of disposable sanitary pads and diapers.

2.3. Regulatory classifications

Regulatory classifications of absorbent hygiene pads vary globally; it is subject to either mandatory legal enforcement or voluntary manufacturers' control depending on countries. In EU,

the sanitary pads are subject to consumer product regulations, which are less stringent [14-16]. In US, menstrual pads and adult incontinence diapers are regulated by FDA under Class I Medical Device [10,11]. This FDA regulation is not legally enforceable and regarded as guidance or recommendations for manufacturing control. For example, FDA recommends that chemicals and their quantities of all constituent materials be provided. From 510(k) premarket submissions guideline, a health risk of adverse tissue reaction is suggested to be identified for menstrual pads, and manufacturer's voluntary testing for biocompatibility is recommended in accordance with ISO-10993 (Biological Evaluation of Medical Devices Part-1: Evaluation and Testing). Tampons, as comparison to pads, are noted for more identified risks, which include adverse tissue reaction, vaginal injury, vaginal infection, and toxic shock syndrome [10,11]. Also, 510(k) suggests to identify the bleaching process of pads, if any, such as "Elemental Chlorine-Free (ECF)." For new formulations that are dissimilar from the previously marketed, it is suggested to conduct voluntary risk analysis and provide clinical data [8,10,11].

In China, safety of baby diapers and sanitary pads are regulated under GB 15979-2002: Hygienic Standard for Disposable Sanitary Products (HS Code of 9619001000 Pull-up baby diapers, diapers/nappies; and HS Code of 9619002000 Sanitary napkins-panty liners- and tampons), which is a mandatory regulation [12]. Sanitary absorbent pads are also regulated under GB/T 8939-2008 [13], which is not the mandatory control. In accordance with those regulations, microbial indicators need to be reported for bacterial and fungal colonies, pathogenic purulent bacteria, and coliforms as well as other test results such as pH value, water absorbency, and infiltration capacity [12,13]. For the imported products into China, toxicological indicators need to be reported, such as skin irritation (for diaper), vaginal mucosa irritation (for menstrual pad), and dermal allergic reaction (for both menstrual pad and diapers).

In Korea, Korean Ministry of Food and Drug Safety (MFDS) regulates baby and adult diapers under 'Sanitary Products,' and menstrual pads under 'Quasi-Drugs' [9]. The regulation has restrictions on substances such as fluorescent whitening agent, formaldehyde, chlorinated phenols (pentachlorophenol, tetrachlorophenol), azoic dyes, phthalates, and metal contaminants (antimony, barium, cadmium, chrome, lead, mercury, arsenic, selenium, etc). However, a test protocol simulating exposure assessment has not been confirmed as of September 2018, and it could possibly create argument on risk management decisions for feminine pads [54,57]. Thus, an urgent need is brought about on revisiting and developing exposure-based risk assessment methodology for absorbent hygiene pads.

Table 2. Safety regulations for diapers and sanitary pads.

	Product	Regulation	Restriction
US [10,11]	Baby diaper	Not available	Manufacturers are recommended to identify new formulations and bleaching process.
	Menstrual pad and adult diaper	FDA Class I Medical Device (guidance)	
China [12,13]	Baby diapers	GB 15979-2002 (mandatory)	Manufacturers should provide microbial and toxicological indicators.
	Sanitary pads	GB 15979-2002 (mandatory) GB/T 8939-2008 (guidance)	
Korea [9]	Baby and adult diapers	MFDS Sanitary Products (mandatory)	There are restrictions on chemicals and heavy metals for baby ¹ and adult diapers. ²
	Menstrual pad	MFDS Quasi-Drugs (mandatory)	

¹ For baby diapers, chemicals and heavy metals restricted include: fluorescent whitening agent, formaldehyde, chlorinated phenols (pentachlorophenol, tetrachlorophenol), azoic dyes, phthalates, antimony, barium, cadmium, chrome, lead, mercury, arsenic, and selenium [9].

² For adult diapers, chemicals restricted include: fluorescent whitening agent, formaldehyde, pentachlorophenol, tetrachlorophenol, azoic dyes [9].

3. Safety assessment framework of absorbent hygiene pads

In general, product safety assessment follows the process established by scientific committees [58,59], regulatory agencies [60,61], and other authoritative bodies [62]. Farage [28,29] suggested a

step-wise process in assessing potential toxicological effects of raw materials, including hazard identification, hazard characterization, exposure assessment, and risk characterization. Additionally, clinical evaluation of product use and postmarket surveillance can be conducted for thorough risk management of products. An overview of assessment approach is shown in Figure 2. Each step of assessment is summarized in the following section, with an emphasis on the exposure studies associated with diapers and feminine pads.




Figure 2. Safety assessment approach.

3.1. Hazard identification and characterization

Hazard identification is generally the first step of risk assessment and is the process to identify the specific chemical hazard and to determine whether exposure to this chemical has the potential to harm human health, such as acute, cumulative and mechanical skin irritation, contact sensitization, and the potential for acute or subchronic effects [63].

Generally, the list of chemicals used in the industrial process as well as the associated safety data can be easily identified by the supplier's chemical safety data sheet. The identification may also include whether mixture of chemicals form other hazardous chemicals or behave differently. If the identity of the chemical is unknown, the possible types of chemicals should be inferred from the descriptions of sources, production processes, or chemicals of other commercial industrial operations (available from emission scenario documents) published by Organization for Economic Cooperation and Development (OECD) [64]. Those documents are useful in quantifying the emission of chemicals into water, air, soil or solid waste, and such information supports the hazard identification process.

The potential hazard of a chemical can be determined from the available scientific database from toxicological or epidemiological studies provided by various international and national agencies [65,66]. For example, the International Agency for Research on Cancer (IARC) [65] coordinates epidemiological and laboratory research for the causes of human cancer, providing five categories based on the strength of evidence for carcinogenicity in humans. World Health Organization (WHO) works to establish the scientific basis for the sound management of chemicals for the safety of human health and the environment through the International Programme on Chemical Safety (IPCS) [67]. The Agency for Toxic Substances and Disease Registry (ATSDR) [68] also prepares toxicological profiles for hazardous substances found at National Priorities List sites. The Integrated Risk Information System (IRIS) [69] maintained by the U.S. Environmental Protection Agency (EPA) also provides an electronic database containing information on human health effects that may result from exposure to various chemicals in the environment. The European Commission (EC) in the framework of the Regulatory Fitness Programme (REFIT) [70] conducts studies that establish the links between the chemical substances and their impacts on human health and environment, to influence the legislation to reduce such impacts. The European Chemicals Agency (ECHA) also provides the list of restricted substances under the framework of Registration, Evaluation, Authorization and Restriction of Chemicals (REACH) [66].

For feminine pads, potential hazardous chemicals and descriptions of chemical sources have been published in several case studies [71-73]. From those studies, volatile organic compounds including styrene, chloromethane, chloroethane, chloroform, and acetone were detected from the commercial products [71-73], where those chemicals are identified as carcinogens, reproductive and developmental toxins according to the National Toxicology Program ATSDR [68,74] and the State of California Environmental Protection Agency [75]. In another study [76], methylidibromo glutaronitrile (MDBGN) was identified from the adhesive used in the pad, which is known to cause dermatitis [76-80]. Cinnamaldehyde and cinnamic alcohol were found from the scented pad, and those are reported to be causative agents of allergic skin reaction and rash by European Chemical agency (ECHA) [81].

Dioxins were detected from the fluff pulp of the absorbent core in pad products [82-84], where dioxins and dioxin-like compounds (DLCs) are highly toxic and environmental persistent organic pollutants. DLCs such as polychlorinated biphenyls can be produced from bleaching process of wood pulps, and such a compound is identified as Group 1 carcinogen by IARC. Acrylic acid, a residual monomer of SAP in the diaper core, is linked to skin irritations and toxic shock syndrome [85].

Hazard characterization is the qualitative or quantitative description of the inherent properties of chemicals that cause health risks, in terms of chemical toxicity, dose-response relationship, exposure route and time [23,24]. Based on the hazard characterization, health based references or point of departure (POD) are determined; where those reference values include no observed adverse effect level (NOAEL), lowest observed adverse effect level (LOAEL), oral reference dose (RfD), inhalation reference concentration (RfC), benchmark dose limit (BMDL), acceptable daily intake (ADI), tolerable daily intake (TDI).

As the disposable absorbent hygiene pads including baby/adult diapers, feminine pads share the main constituents of materials, analyses for hazard identification and characterization on those products can be shared to some extent. With the information on hazard identification and hazard characterization, policies and regulations may be established for risk management associated with exposure to diapers and feminine pads. For example, phthalates, common plasticizers for polymeric materials, can be found from absorbent pads, and those chemicals are known to have potential carcinogenicity, kidney and liver damage; thus, certain phthalates in a concentration > 0.1% [86] are restricted for childcare articles in European REACH [87] and Japan [88].

3.2. Exposure assessment and risk characterization

In Ishii's study [5,21], seven different phthalates were quantified from the topsheet of diapers using artificial sweat and urine as eluents. The eluted liquid was extracted using dichloromethane and analyzed by gas chromatography-mass spectrometry (GC-MS). The quantified contents were processed to produce daily exposure per body weight, taking an estimate of transdermal absorption rate of phthalates [5]. After the exposure assessment, risk characterization was conducted by calculating the margin of exposure (MOE), the ratio of a daily exposure to a No Observed Adverse Effect Level (NOAEL); as MOEs were greater than 1,000, it was concluded that the tested diapers were safe to phthalate exposure [5]. The test was performed only on the topsheet, thus the phthalate contents could be underestimated, because the phthalates beneath the topsheet were not counted in the study.

Organic derivatives of tin such as dibutyltin (DBT), monobutyltin (MBT), tributyltin (TBT) and dioctyltin (DOT) are identified as potential hazard that can be present in diapers and pads as tin is used as a catalyst in manufacturing processes [89]. With the known harmful effect such as neurotoxicity, reproductive and developmental toxicity, immunotoxicity and endocrine disruption [89], organotin compounds were quantified from baby diapers by the energy dispersive x-ray fluorescence with ²⁴¹Am radioactive source [39]. Assuming the daily use of 5 diapers, daily organotin exposure was estimated to be 0.021 µg tin/kg weight·day. However, the dermal absorption rate of organotin compounds was not accounted in this study. Based on such information, risk characterization was conducted reporting that this content was 20 to 100% of the tolerable daily intake (TDI), and the risk of tin exposure is present [39].

Dioxins are another subject of attention regarding the potential risk of diaper wearing, as they can be generated from wood pulp bleaching with elemental chlorine, if total chlorine free (TCF) or elemental chlorine free (ECF) bleaching process is not used [1,6,7,21,82,90]. Ishii [21] performed the risk assessment of sanitary pads for the levels of dioxins, by comparing the daily exposure of dioxins from the tolerable daily intake (TDI). In this study [21], the amounts of polychlorinated dibenzo-p-dioxins (PCDDs), polychlorinated dibenzofurans (PCDFs), and dioxin-like polychlorinated biphenyls (DL-PCBs) in fluff pulp samples of sanitary pads were determined by toluene extraction followed by GC-MS analysis. The quantified contents were converted to the toxic equivalency quantity (TEQ) using the World Health Organization's toxic equivalency factors (TEF) [26]. The TEQ concentration was then used to calculate the daily exposure [21] by taking into consideration of frequency of use, body weight, and reference absorption rate of dioxins (3%) [71,91]. From the result, the daily exposure was lower than 1/9,500 times TDI, thus the risk of exposure was concluded to be negligible. This study did not consider the reflux of dioxins by fluids; instead, total amount of dioxins extracted by toluene was quantified. As the fluff cellulose exists in the absorbent core, the actual exposure of dioxins in the fluff cellulose may occur only through fluid reflux, if ever; thus, actual daily exposure in real situation would be negligible, or be much lower than the values obtained from this study [21]. The bleaching of pulp can also leave chlorinated compounds in pulp products. Later on, Wada et al. [7] updated the test protocol in assessing the exposure risk of total amount chlorine in pulp products, using an artificial sweat as an eluent, extracting at 40 °C for 60 min. The eluted liquid was analyzed for chloroform, chloroketones, chlorophenols, chloroacetic acids and organically bound chlorine [7]. From this study, little residual chlorine was quantified in products, posing little health risk.

Recently, the emission of 74 types VOCs (including pesticides) from feminine products was investigated by Korean MFDS [54]. Test protocol was designed to measure the maximum amount of residual VOCs in products; in this method, pads were frozen at -196 °C to avoid evaporation of VOCs before detection, grounded at room temperature, then the VOCs were analyzed at 120 °C by GC-MS. The systemic exposure was calculated based on the following usage assumptions: 7.5 pads/day, 7 menstruation days/month, lifelong use. Risk characterization was performed by the margin of safety (MOS), calculated by the NOAEL values obtainable from US EPA-IRIS, ATSDR, or WHO IPCS; where MOS values for all tested pads were found to be greater than 5. It is probable that the VOCs measured by this method are overestimation of the actual exposure, as the total amount of VOCs in the whole pad was counted (assuming 100% dermal absorption rate). A protocol that simulates the actual wearing environment may provide more accurate representation of exposure.

3.2.1. Exposure model

The systemic and dermal exposure rates for the absorbent hygiene products can be estimated as following equation [14]:

$$\text{Systemic exposure } (\mu\text{g}/\text{kg}/\text{d}) = \frac{M \times C \times F \times D \times T_{\text{direct or T}_{\text{indirect}}} \times A}{B} \quad (1)$$

$$\text{Dermal exposure } (\mu\text{g}/\text{cm}^2/\text{d}) = \frac{M \times C \times F \times D \times T_{\text{direct or T}_{\text{indirect}}} \times A}{S} \quad (2)$$

where M is the raw material mass in product (g), C is the constituent concentration in the raw material (%), F is the frequency of the use (numbers/day), D is the exposure duration (% of day), T_{direct} (%) is the constituent transfer rate from materials to skin (in direct skin contact), or T_{indirect} (%) is the constituent transfer rate from materials to skin (indirect skin contact), A is the dermal absorption rate (%), B is the body weight (kg), and S is the exposed skin surface area (cm²). Systemic exposure is calculated based on the body weight, and dermal exposure is expressed by dose per unit area of skin exposed.

Important parameters for exposure-based assessment include; the routes (oral, inhalation, and dermal exposure) and pathways of exposure (i.e., direct and indirect skin contact), exposure concentration, duration and frequency of exposure, and transfer media (air, water, soil). While the main exposure route of hygiene pads is dermal absorption, the pathways of exposure can be varied. The pathway of exposure refers to the physical course of chemical transfer as the chemical moves from a source to a point of contact. Common pathways of chemical transfer for pad products are direct skin contact and indirect skin contact by fluid reflux. Exposure assessments for different pathways are further discussed in the later section.

The user biometric parameters like body weight and skin contact are critical parameters for the robust exposure-based risk assessment. Especially in diaper/pad product use, the body weight also accounts for the pressure applied onto the materials. Because infant body weight changes every day, the body weight should be specified during the given time period; thus, the US EPA Exposure Factors Handbook [44] and the Centers for Disease Control (CDC) [45] recommend using time-weighted average (TWA) body weights for the diapering period of 0-36 months. For assessment of skin sensitization, the surface area of the skin contact, instead of body weight, may be a relevant parameter for exposure-based assessment [92].

Frequency of use and length of wear are other key parameters in assessing the exposure. This information can be obtained through survey statistics for the affected individuals, demographic data, behavioral observation, and activity diaries/models [62]. In applying human exposure models, WHO provides general guidance and suggestion of uncertainty factors [92,93].

3.2.2. Exposure by the direct skin contact

Among the constituents of diapers and pads, polymeric materials are of little concern because large molecular weight polymers can be hardly transferred to skin. Of particular concern is small molecular weight materials such as aesthetic ingredients (scent, dyes), solvents and process additives (bleaching agent). In a simulation study by Farage [49,50,52], emollient transfer to skin was estimated to be lower than 20%, which was a conservative estimate. In Odio et al's study [36,37], a lotion was spread on the topsheet of a diaper, then the transferred amount of lotion to the collection tape (Tegaderm™, 3M, St. Paul, MN) was measured during the children's diaper-wearing [37,38]. From this experiment, the lotion transfer to skin was estimated to be 3.0-4.3% of the initial amount of lotion per diaper [3,37,38,47]; this is a half the amount that Rai et al. [24] used (7%) as the skin transfer rate by the direct contact.

3.2.3. Indirect exposure by reflux

Material transfer by the indirect skin contact involves rewetting of skin by fluid reflux under body weight, where rewetting was defined as the fraction of fluid that was resurfaced back from the absorbed state. Components of absorbent pads that are not in direct skin contact can be solubilized in aqueous carrier like urine or menstruation, and can be resurfaced to the topsheet under body weight [3]. Rai et al. [24] explained that modern disposable diapers that contain SAP in the core has reflux value of merely about 0.25% as the resurfacing of liquid is minimized by SAP.

In Dey et al's study [2,3], rewetting test was performed employing a controlled gravimetric approach where collagen is used as a model skin. In this method, the rewetting by the prolonged exposure was simulated by applying a saline solution as a model urine under 0.41 psi, which was noted as the simulated pressure from an infant of 18-24 months. A 60 ml of 0.9% saline solution was loaded onto a diaper at an interval of 82 min, and the liquid transferred to the collagen sheet was measured after a predetermined time of wear [3]. Assuming that the applied pressure from a baby will not be on 100% of diaper area, rewetting fraction was adjusted to count 50% diaper area; the fraction of constituents transferred to skin by the reflux was estimated to be 0.32-0.66% (mean value of 0.46%). This value is higher than 0.25% that was suggested by Rai [24]. This conservative value of 0.46% was suggested by Dey et al. as a default rewet as the worst scenario [3].

The developed reflux protocol was applied to a target chemical, acrylic acid which can exist as a residual monomer in the SAP [2,3,47]. The predetermined amount of acrylic acid was added to the

SAP material, then the resurfaced acrylic acid was collected on the collagen sheet. Quantification of acrylic acid from the extract of collagen sheet resulted in rewet value of 0.15-0.21%, which is lower than the suggested default value of 0.46% [3,47]. The authors discussed that using a default rewet value is a practical alternative to conducting a series of routine rewet testing for every constituent [3,47]. The concept of 'default value' or 'lotion transfer factor' was mentioned also from Kosemund et al. [4] as a practical approach to estimate the transfer rate of the chemicals by the indirect skin contact. When the added quantities of constituents are known, transferrable amounts of all constituents can be calculated using a conservative, default rewet value. Figure 3 overviews hazard characterization methods applied for different parts of products; (i) VOCs from the whole product construction were characterized; (ii) chemicals from the topsheet were analyzed based on the direct contact assumption; (iii) refluxed chemicals were analyzed with simulated rewet testing.




Figure 3. Exposure assessment for different pathways.

3.2.4. Risk characterization

Risk characterization is a comparison between measured (or estimated) exposure and the reference values, such as NOAEL, LOAEL, RfD and RfC [94-96]. In this step, information obtained from previous steps of hazard identification, hazard characterization and exposure assessment is integrated, and the margin of exposure (MOE) or margin of safety (MOS) is calculated as a final evaluation for safety assurance.

MOE is the ratio of a point of departure (POD) dose such as NOAEL, LOAEL, and BMDL to an estimated human exposure, without consideration of uncertainty factor. NOAEL and LOAEL are obtained from animal toxicity studies and are noted in a unit of mg/kg body weight/day or ppm. BMDL, which incorporates the quantitative dose-response aspects, is also used for risk assessment associated with hazardous compounds.

MOS takes into account uncertainty factor, and is determined by the ratio of a derived reference dose with the no or minimal adverse effects- which includes RfD, RfC, ADI, etc.- to an estimated human exposure level [22,26,97]. With uncertainty factor implemented, $MOS \geq 1$ is considered to be acceptable; that is, if the product is used in its intended way or in foreseeable misuse, the product is safe with adequate margin of safety [4,24]. When MOS is lower than the desired, it is recommended to refine the conservative default assumptions. It is noteworthy that the reference values such as RfD, RfC, ADI, etc., are not the definite values, but rather evolve as new information or new scientific approach becomes available. While this fact does not invalidate the current reference doses, the reference values and MOS need to be regarded as the currently agreed ones.

3.3. Clinical evaluation of product safety in-use

Woeller et al. [14] conducted clinical studies on sanitary pads by the patch test to assess the overall mean erythema score (OMES) as a measure of cumulative irritation. During 21-day study, the

OMES of a finished sanitary product (with emollient) was significantly less irritating than that of product without emollient and that of absorbent core only [14]. On the other side, when an absorbent core is used in a finished product, hygiene benefits including reduced dermatitis [34-36,98-100] and reduced bacterial contamination [26,101-103] were demonstrated. The Behind-the-Knee (BTK) clinical test [29,104] was performed by applying a test pad to the popliteal fossa of one leg under an elastic bandage for 6 h per day for 4 consecutive days. From BTK test, skin irritation and dryness were scored after 30-60 min after removal of products, and the score of erythema and dryness for the test product was compared with those of reference product that had safe use history [48,49,52].

The clinical researches before market introduction are desirable to assure product safety especially for such cases as: change of materials in use, introduction of the new process, modification of product design. Such clinical studies are commonly conducted under practical or exaggerated conditions of use. The clinical study generally follows the guidelines from the International Committee on harmonization/Good Clinical Practice (ICH/GCH) and gets approvals by the Institutional Review Board or ethical committee of the pertinent institution [29]. From the development of protocols to the investigation, independent academic physicians and medical experts in obstetrics and gynecology, dermatology, and microbiology are involved. Consumers habits and practices, as well as different geographic effect, are considered in the protocol design and investigation [105-107]. In US, though it is not legally enforceable, manufacturers are recommended to provide premarket notification submissions under FDA 510(k) for menstrual tampons and pads [8]. Then a critical review on safety assurance is conducted by a panel of independent experts with the most up-to-date information and best scientific judgement.

3.4. Postmarket surveillance

Risk management in the stage of consumer usage in market can be conducted through postmarket surveillance by monitoring consumer's experience and satisfaction. Electronic media such as bulletin boards, social media, and manufacturer's website can facilitate the communication between consumers and manufactures. Any unexpected risk or health-related questions/comments would be alerted through such communications [71,108]. Especially for chemicals that has cumulative long-term effects, decades of after-market record would be very useful as critical references in various types and levels of comments.

While there are studies that revealed the symptom of skin irritation and allergic rash caused by the use of hygiene pads [109,110], in practice, it is not obvious for users and medical care providers to identify the problematic ingredients unless the whole product formulations are disclosed. In fact, it is not a mandatory requirement in many countries to disclose the formulations of hygiene pads. Recently in Korea, users of hygiene pads have claimed through social communities for the negative health signals after using specific brand pads, which include reduced menstrual bleeding, skin rashes and painful cramps. In 2017, the Korea MFDS collected all feminine pads sold in Korea, either manufactured or imported for the past three years, and investigated the levels of volatile organic compounds in products [54,55,57]. Such postmarket surveillance and continuous monitoring would provide important cases that would explain unknown risks.

4. Conclusions

This article described a systematic approach for risk assessment of disposable absorbent hygiene products such as feminine pads and diapers. The approach consists of: (i) hazard identification and hazard characterization; (ii) exposure assessment and risk characterization; (iii) evaluation of safety-in-use by means of prospective clinical trials; (iv) postmarket surveillance. More specifically, the following contents related to the absorbent pad products were reviewed [62]: identification of the concerned chemicals and their potential harms to human; characteristics of chemicals that can adversely affect health; chemical reference values that can be used with minimal harmful effects; relevancy of assumptions made in exposure assessment; pathways of chemical exposure in use; quantity, duration, frequency of exposure; comparison of the estimated exposure with the reference values. In exposure assessment of disposable hygiene pads, a protocol that simulates the actual

wearing environment would provide more accurate representation of realistic impacts. For exposure models, it is important to use appropriate and conservative exposure parameters. Then the risk characterization is conducted by comparing the estimated exposure with the appropriate reference values.

Consumers expect safe use of disposable hygiene products in the extended period of time and frequent use, yet the expectation is challenged at times. A study by the non-profit organization for women safety and academic research have reported [57,72,81,111,112] that feminine pads includes several chemicals of concern which are not identified by manufacturers. However, very limited data is available to confirm this claim or to demonstrate the effect of cumulative chemical exposure to health. More research and testing are clearly needed to better characterize and understand the potential health impact by the exposure [75,111,113]. In addition, it is critical to establish the rigorous and effective postmarket surveillance to monitor the product defects and health effects. Extensive review of collected consumers' feedback by the independent experts may provide unknown risks involved with long-term and cumulative exposure.

Author Contributions: Conceptualization, J. Kim, J. Bae, and H. Kwon; Investigation and Analysis, J. Kim, J. Bae and H. Kwon; Writing-Original Draft Preparation, J. Kim and J. Bae; Writing-Review & Editing, J. Bae, J. Kim, and H. Kwon.

Funding: This research was supported by Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education (2018R1D1A1B07040340).

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Counts, J.; Weisbrod, A.; Yin, S. Common diaper ingredient questions: Modern disposable diaper materials are safe and extensively tested. *Clinical Pediatrics* **2017**, *56*, 23S-27S, doi: 10.1177/0009922817706998.
2. Dey, S.; Kenneally, D.; Odio, M.; Hatzopoulos, I. Modern diaper performance: Construction, materials, and safety review. *Int J Dermatol* **2016**, *55 Suppl 1*, 18-20, doi: 10.1111/ijd.13333.
3. Dey, S.; Purdon, M.; Kirsch, T.; Helbich, H.; Kerr, K.; Li, L.; Zhou, S. Exposure factor considerations for safety evaluation of modern disposable diapers. *Regul Toxicol Pharmacol* **2016**, *81*, 183-193, doi: 10.1016/j.yrtph.2016.08.017.
4. Kosemund, K.; Schlatter, H.; Ochsenhirt, J.L.; Krause, E.L.; Marsman, D.S.; Erasala, G.N. Safety evaluation of superabsorbent baby diapers. *Regul Toxicol Pharmacol* **2009**, *53*, 81-89, doi: 10.1016/j.yrtph.2008.10.005.
5. Ishii, S.; Katagiri, R.; Minobe, Y.; Kuribara, I.; Wada, T.; Wada, M.; Imai, S. Investigation of the amount of transdermal exposure of newborn babies to phthalates in paper diapers and certification of the safety of paper diapers. *Regul Toxicol Pharmacol* **2015**, *73*, 85-92, doi: 10.1016/j.yrtph.2015.06.010.
6. Koda, K.; Goto, H.; Shintani, H.; Matsumoto, Y.; Meshitsuka, G. Oxidative cleavage of lignin aromatics during chlorine bleaching of kraft pulp. *Journal of Wood Science* **2001**, *47*, 362-367, doi: 10.1007/BF00766786.
7. Wada, T.; Iwasaki, K.; Minobe, Y.; Wada, M.; Imai, S.; Ishii, S. Proposal of a flow scheme for the chemical-form-based quantitative analysis of chlorine compounds in pulp for sanitary products and verification of safety. *Regul Toxicol Pharmacol* **2017**, *91*, 109-116, doi: 10.1016/j.yrtph.2017.10.009.
8. US FDA. Guidance for industry and FDA staff- menstrual tampons and pads: information for premarket notification submissions 510(k). 2017. Available online: <https://www.fda.gov/MedicalDevices/ucm071781.htm> (accessed on 08. 06. 2018).
9. Korean Ministry of Food and Drug Safety (MFDS). Available online: http://www.mfds.go.kr/brd/m_543/list.do (accessed on 08. 06. 2018).
10. Code of Federal Regulations. CFR 884.5425: Scented or scented deodorized menstrual pad. Available online: https://www.ecfr.gov/cgi-bin/retrieveECFR?gp=1&SID=df43bf980a5782408f10c44a86b616bf&ty=HTML&h=L&mc=true&r=SECTION&n=se21.8.884_15425 (accessed on 08. 06. 2018).
11. Code of Federal Regulations. CFR 884.5435: Unscented menstrual pad. Available online: https://www.ecfr.gov/cgi-bin/retrieveECFR?gp=1&SID=df43bf980a5782408f10c44a86b616bf&ty=HTML&h=L&mc=true&r=SECTION&n=se21.8.884_15435 (accessed on 08. 06. 2018).
12. Standardization Administration of China. GB 15979-2002: Hygienic standard for disposable sanitary products. 2002.
13. Standardization Administration of China. GB/T 8939-2008: Sanitary absorbent pads (including pantiliner). 2008.
14. Woeller, K.E.; Hochwalt, A.E. Safety assessment of sanitary pads with a polymeric foam absorbent core. *Regul Toxicol Pharmacol* **2015**, *73*, 419-424, doi: 10.1016/j.yrtph.2015.07.028.
15. European Disposables and Nonwovens Association (EDANA). Nonwovens standard procedures 2015. Available online: <https://www.edana.org/docs/default-source/default-document-library/toc-cover-preamble.pdf?sfvrsn=1> (accessed on 08. 06. 2018).
16. European Disposables and Nonwovens Association (EDANA). Fact sheet health & hygiene benefits of absorbent hygiene products & personal care wet wipes. Available online: <https://www.edana.org/docs/default-source/default-document-library/fact-sheet---health-hygiene-benefits-of-absorbent-hygiene-products-personal-care-wet-wipes.pdf?sfvrsn=2> (accessed on 08. 06. 2018).
17. Farage, M.A.; Bjerke, D.L.; Mahony, C.; Blackburn, K.L.; Gerberick, G.F. Quantitative risk assessment for the induction of allergic contact dermatitis: Uncertainty factors for mucosal exposures. *Contact Dermatitis* **2003**, *49*, 140-147, doi: 10.1111/j.0105-1873.2003.00192.x
18. Felter, S.P.; Carr, A.N.; Zhu, T.; Kirsch, T.; Niu, G. Safety evaluation for ingredients used in baby care products: Consideration of diaper rash. *Regul Toxicol Pharmacol* **2017**, *90*, 214-221, doi: 10.1016/j.yrtph.2017.09.011.
19. Felter, S.P.; Robinson, M.K.; Basketter, D.A.; Gerberick, G.F. A review of the scientific basis for uncertainty factors for use in quantitative risk assessment for the induction of allergic contact dermatitis. *Contact Dermatitis* **2002**, *47*, 257-266, doi: 10.1034/j.1600-0536.2002.470502.x
20. Gerberick, G.F.; Robinson, M.K.; Felter, S.P.; White, I.R.; Basketter, D.A. Understanding fragrance allergy using an exposure-based risk assessment approach. *Contact Dermatitis* **2001**, *45*, 333-340, doi: 10.1034/j.1600-0536.2001.450603.x.

21. Ishii, S.; Katagiri, R.; Kataoka, T.; Wada, M.; Imai, S.; Yamasaki, K. Risk assessment study of dioxins in sanitary napkins produced in Japan. *Regul Toxicol Pharmacol* **2014**, *70*, 357-362, doi: 10.1016/j.yrtph.2014.07.020.
22. Kroes, R.; Kleiner, J.; Renwick, A. The threshold of toxicological concern concept in risk assessment. *Toxicol Sci* **2005**, *86*, 226-230, doi: 10.1093/toxsci/kfi169.
23. Meek, M. E.; Boobis, A. R.; Crofton, K. M.; Heinemeyer, G.; Raaij, M. V.; Vickers, C. Risk assessment of combined exposure to multiple chemicals: A WHO/IPCS framework. *Regul Toxicol Pharmacol* **2011**, *60*, S1-S14, doi: 10.1016/j.yrtph.2011.03.010.
24. Rai, P.; Lee, B.M.; Liu, T.Y.; Yuhui, Q.; Krause, E.; Marsman, D.S.; Felter, S. Safety evaluation of disposable baby diapers using principles of quantitative risk assessment. *J Toxicol Environ Health A* **2009**, *72*, 1262-1271, doi: 10.1080/15287390903212246.
25. Robinson, M.K.; Gerberick, G.F.; Ryan, C.A.; McNamee, P.; White, I.R.; Basketter, D.A. The importance of exposure estimation in the assessment of skin sensitization risk. *Contact Dermatitis* **2000**, *42*, 251-259, doi: 10.1034/j.1600-0536.2000.042005251.x.
26. Van den Berg, M.; Birnbaum, L. S.; Denison, M.; De Vito, M.; Farland, W.; Feeley, M.; Fiedler, H.; Hakansson, H.; Hanberg, A.; Haws, L.; Rose, M.; Safe, S.; Schrenk, D.; Tohyama, C.; Tritscher, A.; Tuomisto, J.; Tysklind, M.; Walker, N.; Peterson, R. E. The 2005 World Health Organization reevaluation of human and Mammalian toxic equivalency factors for dioxins and dioxin-like compounds. *Toxicological Sciences* **2006**, *93*, 223-241, doi: 10.1093/toxsci/kfl055.
27. Repon, M.R.; Islam, M.T.; Mamun, M.A.A. Ecological risk assessment and health safety speculation during color fastness properties enhancement of natural dyed cotton through metallic mordants. *Fashion and Textiles* **2017**, *4*, 24, doi: 10.1186/s40691-017-0109-x
28. Farage, M.A.; Stadler, A.; Elsner, P.; Maibach, H.I. Safety evaluation of modern feminine hygiene pads: Two decades of use. *The female patient* **2004**, *29*, 23-30.
29. Farage, M.A. A behind-the-scenes look at the safety assessment of feminine hygiene pads. *Ann NY Acad Sci* **2006**, *1092*, 66-77.
30. Dey, S.; Helmes, C.T.; White, J.C.; Zhou, S. Safety of disposable diaper materials: Extensive evaluations validate use. *Clin Pediatr* **2014**, *53*, 17s-19s, doi: 10.1177/0009922814540378.
31. Akin, F.; Spraker, M.; Aly, R.; Leyden, J.; Raynor, W.; Landin, W. Effects of breathable disposable diapers: Reduced prevalence of candida and common diaper dermatitis. *Pediatr Dermatol* **2001**, *18*, 282-290.
32. Alberta, L.; Sweeney, S.M.; Wiss, K. Diaper dye dermatitis. *Pediatrics* **2005**, *116*, e450-e452, doi: 10.1542/peds.2004-2066.33. Campbell, R.L. Clinical tests with improved disposable diapers. *Pediatrician* **1987**, *14 Suppl 1*, 34-38.
34. Campbell, R.L.; Seymour, J.L.; Stone, L.C.; Milligan, M.C. Clinical studies with disposable diapers containing absorbent gelling materials: Evaluation of effects on infant skin condition. *J Am Acad Dermatol* **1987**, *17*, 978-987, doi: 10.1016/S0190-9622(87)70287-4.
35. Davis, J.A.; Leyden, J.J.; Grove, G.L.; Raynor, W.J. Comparison of disposable diapers with fluff absorbent and fluff plus absorbent polymers: Effects on skin hydration, skin ph, and diaper dermatitis. *Pediatr Dermatol* **1989**, *6*, 102-108, doi: 10.1111/j.1525-1470.1989.tb01005.x.
36. Odio, M.; Friedlander, S.F. Diaper dermatitis and advances in diaper technology. *Curr Opin Pediatr* **2000**, *12*, 342-346.
37. Odio, M.R.; O'Connor, R.J.; Sarbaugh, F.; Baldwin, S. Continuous topical administration of a petrolatum formulation by a novel disposable diaper. 2. Effect on skin condition. *Dermatology* **2000**, *200*, 238-243, doi: 10.1159/000018366.
38. Odio, M.R.; O'Connor, R.J.; Sarbaugh, F.; Baldwin, S. Continuous topical administration of a petrolatum formulation by a novel disposable diaper. 1. Effect on skin surface microtopography. *Dermatology* **2000**, *200*, 232-237, doi: 10.1159/000018365.
39. Šmajgl, D.; Obhodaš, J. Occurrence of tin in disposable baby diapers. *X-Ray Spectrometry* **2015**, *44*, 230-232, doi: 10.1002/xrs.2609.
40. Wilson, P.A.; Dallas, M.J. Diaper performance: Maintenance of healthy skin. *Pediatr Dermatol* **1990**, *7*, 179-184, doi: 10.1111/j.1525-1470.1990.tb00277.x.
41. Farage, M.A.; Gilpin, D.A.; Enane, N.A.; Baldwin, S. Development of a new test for mechanical irritation: Behind the knee as a test site. *Skin Res Technol* **2001**, *7*, 193-203, doi: 10.1034/j.1600-0846.2001.70309.x
42. US Environmental Protection Agency. Guidance on selecting age groups for monitoring and assessing childhood exposures to environmental contaminants. 630-P-03-003F, November 2005. Available online: <https://www.epa.gov/risk/guidance-selecting-age-groups-monitoring-and-assessing-childhood-exposures-environmental> (accessed on 08. 07. 2018).

43. US Environmental Protection Agency. Child-specific exposure scenarios examples (final report). EPA/600/R-14-217F, 2014. Available online: <https://cfpub.epa.gov/ncea/risk/recordisplay.cfm?deid=262211> (accessed on 08. 07. 2018).
44. US Environmental Protection Agency. Exposure factors handbook 2011 edition (final report). EPA/600/R-09/052F, 2011. Available online: <https://cfpub.epa.gov/ncea/risk/recordisplay.cfm?deid=236252> (accessed on 08. 07. 2018).
45. Centers for Disease Control and Prevention. WHO growth standards are recommended for use in the U.S. for infants and children 0 to 2 years of age. 2010. Available online: https://www.cdc.gov/growthcharts/who_charts.htm#The%20WHO%20growth%20charts (accessed on 08. 07. 2018).
46. Boniol, M.; Verriest, J.P.; Pedoux, R.; Dore, J.F. Proportion of skin surface area of children and young adults from 2 to 18 years old. *J Invest Dermatol* **2008**, *128*, 461-464, doi: 10.1038/sj.jid.5701032.
47. Dey, S.; Rothe, H.; Page, L.; O'Connor, R.; Farahmand, S.; Toner, F.; Marsh, R.; Wehmeyer, K.; Zhou, S. An in vitro skin penetration model for compromised skin: Estimating penetration of polyethylene glycol 14c-peg-7 phosphate. *Skin Pharmacology and Physiology* **2015**, *28*, 12-21, doi: 10.1159/000362284.
48. Farage, M.A. Enhancement of visual scoring of skin irritant reactions using cross-polarized light and parallel-polarized light. *Contact Dermatitis* **2008**, *58*, 147-155, doi: 10.1111/j.1600-0536.2007.01284.x.
49. Farage, M.A.; Segarra, V.S.; Bramante, M.; Elsner, P.; Maibach, H.I. Dermatological testing of an emollient-treated menstrual pad with a novel foam absorbent core. *Cutan Ocul Toxicol* **2008**, *27*, 333-343, doi: 10.1080/15569520802337749.50. Farage, M.A. Evaluating lotion transfer from products to skin using the behind-the-knee test. *Skin Res Technol* **2010**, *16*, 243-252.
51. Hanke-Baier, P.; Johannigmann, J.; Levin, R.J.; Wagner, G. Evaluation of vaginal and perineal area during the use of external sanitary protection throughout the menstrual cycle. *Acta Obstetrica et Gynecologica Scandinavica* **1994**, *73*, 486-491, doi: 10.3109/00016349409013437.
52. Farage, M.A.; Berardesca, E.; Maibach, H. Skin moisturization and frictional effects of an emollient-treated menstrual pad with a foam core. *Cutaneous and Ocular Toxicology* **2009**, *28*, 25-31, doi: 10.1080/15569520902738408.
53. Rai, P.; Marsman, D.S.; Felter, S.P. Dermal safety evaluation: Use of disposable diaper products in the elderly. In *Textbook of aging skin*, Farage, M.A.; Miller, K.W.; Maibach, H.I., Eds. Springer: Berlin, 2010; pp 857-868; ISBN 978-3-540-89656-2.
54. Korean Ministry of Food and Drug Safety (MFDS). 생리대. 팬티라이너 74종 VOCs 인체 위해 우려 없어. 12. 28. 2017. Available online: http://www.mfds.go.kr/brd/m_99/view.do?seq=40034 (accessed on 08. 06. 2018).
55. Helmes, C.T.; O'Connor, R.; Sawyer, L.; Young, S. Disposable diaper absorbency: Improvements via advanced designs. *Clinical Pediatrics* **2014**, *53*, 14S-16S, doi: 10.1177/0009922814540377.
56. Ajmeri, J.R.; Ajmeri, C.J. Nonwoven personal hygiene materials and products. In *Applications of nonwovens in technical textiles*, Chapman, R.A., Ed. Woodhead Publishing: 2010; pp 85-102.
57. Kim, S. Reviewing the Korean episodes of environmental chemicals in summer 2017. *The Korean Journal of Public Health* **2017**, *54*, 3-12, doi: 10.17262/KJPH.2017.12.54.2.3.
58. National Academy of Science. Risk assessment in the federal government: Managing the process. National Academy Press: Washington, DC, 1983.
59. Scientific Committee on Consumer Safety. *The SCCS's notes of guidance for the testing of cosmetic ingredients and their safety evaluation*. 9th ed.; 2016.
60. European Chemicals Agency. E.C. Guidance on informatin requirements and chemical safety assessment. Helsinki, Finland, 2013.
61. US Environmental Protection Agency (EPA). *Human health risk assessment*; Agency (EPA): 2012 <https://www.epa.gov/risk/human-health-risk-assessment> (accessed on 08.07.2018).
62. World Health Organization. WHO human health risk assessment toolkit: chemical hazards. 2010. Available online: http://www.who.int/ipcs/methods/harmonization/areas/ra_toolkit/en/ (accessed on 08. 08. 2018).
63. Czerwinski B.S. Adult feminine hygiene practices. *Applied Nursing Research* 1996, *9*, 93123-93129, doi 10.1016/S0897-1897(96)80218-7.
64. OECD, Emission scenario document on textile finishing industry, Available online: <http://www.oecd.org/chemicalsafety/risk-assessment/emissionscenariodocuments.htm> (accessed on 07.24. 2018).
65. International Agency for Research on Cancer, IARC Monographs on the evaluation of carcinogenic risks to humans. Available online: <https://monographs.iarc.fr/> (accessed on 08.27.2018).

66. ECHA, Substances restricted under REACH. Available online: <https://echa.europa.eu/substances-restricted-under-reach> (accessed on 08.27.2018).
67. International Programme on Chemical Safety. Public health impact of chemicals: knowns and unknowns, 2016. Available online: <http://www.who.int/ipcs/publications/chemicals-public-health-impact/en/> (accessed on 08.27.2018).
68. Agency for Toxic Substances and Disease Registry, Substance priority list. Available online: <https://www.atsdr.cdc.gov/spl/resources/index.html> (accessed on 08.27.2018).
69. US Environmental Protection Agency (EPA), Integrated Risk Information System (IRIS) advanced search. Available online: <https://cfpub.epa.gov/ncea/iris/search/index.cfm> (accessed on 08.27.2018).
70. European Commission, Study on the calculation of the benefits of chemicals legislation on human health and the environment: Development of a system of indicators. Available online: http://ec.europa.eu/environment/chemicals/reach/pdf/study_final_report.pdf (accessed on 08.27.2018).
71. Klasmeyer, J.; McLachlan, M.S. PCDD/Fs in textiles- part 1: A screening method for detection of octachlorodibenzo-p-dioxin and octachlorodibenzofuran. *Chemosphere* **1998**, *36*, 1627-1635, doi: 10.1016/S0045-6535(97)10093-5.
72. Eason, E.L.; Feldman, P. Contact dermatitis associated with the use of always sanitary napkins. *Canadian Medical Association Journal* **1996**, *154*, 1173-1176.
73. Nicole, W. A question for women's health: Chemicals in feminine hygiene products and personal lubricants. *Environmental Health Perspectives* **2014**, *122*, A70-A75 doi:10.1289/ehp.122-A70.
74. National Toxicology Program, Substances Listed in the Fourteenth Report on Carcinogens. Available online: <https://ntp.niehs.nih.gov/pubhealth/roc/index-1.html#toc1> (accessed on 08.27.2018).
75. Office of Environmental Health Hazard Assessment, Proposition 65 list. Available online: <https://oehha.ca.gov/proposition-65/about-proposition-65> (accessed on 08.27.2018).
76. Christina L. B. Methylidibromo Glutaronitrile, *Int J Toxicol.* **2017**, *36*, 395-415, doi: 10.1177/1091581817716647
77. Sterry, W.; Schmoll, M. Contact urticaria and dermatitis from self-adhesive pads. *Contact Dermatitis* **1985**, *13*, 284-285, doi:10.1111/j.1600-0536.1985.tb02575.x
78. Wilhelm, K.P.; Maibach, H.I. Factors predisposing to cutaneous irritation. *Dermatologic Clinics* **1990**, *8*, 17-22, doi.org/10.1016/S0733-8635 (18) 30514-X.
79. Williams, J.D.; Frowen, K.E.; Nixon, R.L. Allergic contact dermatitis from methylidibromo glutaronitrile in a sanitary pad and review of Australian clinic data. *Contact Dermatitis* **2007**, *56*, 164-167.
80. Schaudé, C.; Mohr, G.J. Indicator washcloth for detecting alkaline washing solutions to prevent dermatitis patients and babies from skin irritation. *Fashion and Textiles* **2017**, *4*, 7, doi: 10.1186/s40691-017-0092-2.
81. Walter, G.; Larsen, W.G. Sanitary napkin dermatitis due to the perfume. *Archives of Dermatology* **1979**, *115*, 363.
82. DeVito, M.J.; Schecter, A. Exposure assessment to dioxins from the use of tampons and diapers. *Environ Health Perspect* **2002**, *110*, 23-28.
83. Schecter, A.J.; Papke, O.; Marquardt, S. Dioxins and dibenzofurans in American sanitary products: Tampons sanitary napkins, disposable and cloth diapers, and incontinence pads. In *Organohalogen Compounds*, Swedish Environmental Protection Agency: Stockholm, Sweden, 1998; Vol. 36, pp 281-284.
84. Jeong, H.S.; Yun, G.A. Analysis of polychlorinated dibenzo-p-dioxins and dibenzo-furans in sanitary products of women. *Textile Research Journal* **2007**, *77*, 597-603, doi.org/10.1177/0040517507078786.
85. Chemistry Explained. Disposable diapers. Available online: <http://www.chemistryexplained.com/Di-Fa/Disposable-Diapers.html> (accessed on July 24, 2018).
86. Earls, A.O.; Axford, I.P.; Braybrook, J.H. Gas chromatography-mass spectrometry determination of the migration of phthalate plasticisers from polyvinyl chloride toys and childcare articles. *J. Chromatogr A* **2003**, *983*, 237-246, doi: 10.1016/S0021-9673(02)01736-3.
87. EU Scientific Committee on Toxicity, Ecotoxicity and the Environment. Phthalate migration from soft PVC toys and child-care articles. Opinion expressed at the CSTE thrid plenary meeting, Brussels, April 24, 1998. Available online: http://ec.europa.eu/health/ph_risk/committees/sct/documents/out12_en.pdf (accessed on 08. 07. 2018).
88. Japan Ministry of Health, Labour and Welfare. Phthalate esters, review of amendment by public comment, etc. 2015. Available online: <https://www.mhlw.go.jp/shingi/2010/02/dl/s0222-6g.pdf> (accessed on 08. 07. 2018).
89. Risk & Policy Analysts Limited (RPA). Risk assessment studies on targeted consumer applications of certain organotin compounds. September 2005. Available online: file:///C:/Users/%EC%82%AC%EC%9A%A9%EC%9E%90/Downloads/organotins_3rd_report_16_sept_20_05_en.pdf (accessed on 08. 07. 2018).

90. Hise, R.G.; Wright, B.T.; Swanson, S.E. Formation of chlorinated dioxins and furans from lignin and lignin model compounds. *Chemosphere* **1990**, *20*, 1723-1730, doi: 10.1016/0045-6535(90)90335-Q.
91. Klasmeier, J.; Mühlebach, A.; McLachlan, M.S. PCDD/Fs in textiles- Part II: Transfer from clothing to human skin. *Chemosphere* **1999**, *38*, 97-108, doi: 10.1016/S0045-6535(98)00176-3.
92. Kimber, I.; Dearman, R.J.; Basketter, D.A.; Ryan, C.A.; Gerberick, G.F.; McNamee, P.M.; Lalko, J.; Api, A.M. Dose metrics in the acquisition of skin sensitization: Thresholds and importance of dose per unit area. *Regul Toxicol Pharmacol* **2008**, *52*, 39-45.
93. World Health Organization. Part 1: Guidance document on characterizing and communicating uncertainty in exposure assessment. 2008. Available online: <http://www.who.int/ipcs/methods/harmonization/areas/exposure/en/> (accessed on 08. 08. 2018).
94. US Environmental Protection Agency. Guidelines for carcinogen risk assessment. March 2005. Available online: https://www.epa.gov/sites/production/files/2013-09/documents/cancer_guidelines_final_3-25-05.pdf (accessed on 08. 06. 2018).
95. US Environmental Protection Agency. A review of the reference dose and reference concentration processes. December 2002. Available online: <https://www.epa.gov/sites/production/files/2014-12/documents/rfd-final.pdf> (accessed on 08 06 2018).
96. US Environmental Protection Agency. Guidelines for exposure assessment, Risk assessment forum. EPA/600/Z-92/001, 1992. Available online: <https://cfpub.epa.gov/ncea/risk/recordisplay.cfm?deid=15263> (accessed on 08. 07. 2018).
97. Eriksson, L.; Jaworska, J.; Worth, A.P.; Cronin, M.T.; McDowell, R.M.; Gramatica, P. Methods for reliability and uncertainty assessment and for applicability evaluations of classification- and regression-based qsars. *Environ Health Perspect* **2003**, *111*, 1361-1375.
98. Lane, A.T.; Rehder, P.A.; Helm, K. Evaluations of diapers containing absorbent gelling material with conventional disposable diapers in newborn infants. *American Journal of Diseases of Children* **1990**, *144*, 315-318, doi: 10.1001/archpedi.1990.02150270065027.
99. Seymour, J.L.; Keswick, B.H.; Hanifin, J.M.; Jordan, W.P.; Milligan, M.C. Clinical effects of diaper types on the skin of normal infants and infants with atopic dermatitis. *Journal of the American Academy of Dermatology* **1987**, *17*, 988-997, doi: 10.1016/S0190-9622(87)70288-6.
100. Seymour, J.L.; Keswick, B.H.; Milligan, M.C.; Jordan, W.P.; Hanifin, J.M. Clinical and microbial effects of cloth, cellulose core, and cellulose core/absorbent gel diapers in atopic dermatitis. *Pediatrician* **1987**, *14 Suppl 1*, 39-43.
101. Campbell, R.L.; Bartlett, A.V.; Sarbaugh, F.C.; Pickering, L.K. Effects of diaper types on diaper dermatitis associated with diarrhea and antibiotic use in children in day-care centers. *Pediatric Dermatology* **1988**, *5*, 83-87, doi: 10.1111/j.1525-1470.1988.tb01143.x.
102. Kubiak, M.; Kressner, B.; Raynor, W.; Davis, J.; Syverson, R.E. Comparison of stool containment in cloth and single-use diapers using a simulated infant feces. *Pediatrics* **1993**, *91*, 632-636.
103. Van, R.; Wun, C.; Morrow, A. L.; Pickering, L. K. The effect of diaper type and overclothing on fecal contamination in day-care centers. *JAMA* **1991**, *265*, 1840-1844, doi: 10.1001/jama.1991.03460140068028.
104. ASTM. F2808-17: Standard test method for performing behind-the-knee (BTK) test for evaluating skin irritation response to products and materials that come into repeated or extended contact with skin. 2010. Available online: https://compass.astm.org/EDIT/html_annot.cgi?F2808+17 (accessed on 08. 06. 2018).
105. Xuemin W.; Sato N.; Chao Y.; Na L.; Fujimura T.; Takagi Y.; Nojiri H.; Kitahara T.; Takema Y. Cutaneous and sensory effects of two types of sanitary pads with different surfaces in the Shanghai, Chinese population. *Cutan Ocul Toxicol*. **2011**, *30*, 212-216, doi: 10.3109/15569527.2010.547542.
106. Seenivasan, P.; Priya, K.C. A cross sectional study on awareness about menstrual hygiene among rural women. *Stanley Medical Journal* **2015**, *2*, 17-21.
107. Xuemin, W.; Sato, N.; Chao, Y.; Na, L.; Fujimura, T.; Takagi, Y.; Nojiri, H.; Kitahara, T.; Takema, Y. Cutaneous and sensory effects of two types of sanitary pads with different surfaces in the shanghai, chinese population. *Cutan Ocul Toxicol* **2011**, *30*, 212-216, doi: 10.3109/15569527.2010.547542.
108. South Korea to risk assess hundreds of sanitary pads. Available online: <https://chemicalwatch.com/58514/south-korea-to-risk-assess-hundreds-of-sanitary-pads> (accessed on 08.29. 2018).
109. Runeman, B. Skin interaction with absorbent hygiene products. *Clinics in Dermatology* **2008**, *26*, 45-51.
110. Rademaker, M. Allergic contact dermatitis to a sanitary pad. *Australasian Journal of Dermatology* **2004**, *45*, 234-235, doi:10.1111/j.1440-0960.2004.00105.x.
111. Scranton, A. Chem Fatale: potential health effects of toxic chemicals in feminine care products. Missoula, MT: Women's Voices for the Earth (November 2013). Available online: <http://goo.gl/Bglwdu> (Accessed 08.10 2018).

112. Women's Voices for the Earth. 'Product testing results: Always pads. 2014. Available online: <http://www.womensvoices.org/wp-content/uploads/2014/10/Always-fact-sheet-FINAL.pdf> (accessed on 08. 06. 2018).
113. H.R. 2332 (112th Congress): Robin Danielson Act. Available online: <https://www.congress.gov/bill/115th-congress/house-bill/2379?q=%7B%22search%22%3A%5B%22robin+danielson%22%5D%7D&r=1> (accessed on 08. 22. 2018).