

Vibriocidal activity of leaf and rhizome extracts of *Maranta arundinacea* L.

**Priyadarshani Samal¹, Priyaranjan Tripathy¹, Ritarani Das¹, Santi Lata Sahoo¹,
Chinmay Pradhan¹, Bijaya Kumar Padhi² and Jyoti Ranjan Rout^{2*}**

¹Biochemistry and Molecular Biology Laboratory, Post Graduate Department of
Botany, Utkal University, Vani Vihar, Bhubaneswar-751004, Odisha, India.

²School of Biological Sciences, AIPH University, Pahala, Bhubaneswar-752101,
Odisha, India.

9

10

11

12

13

14

15

16

17

18

*Corresponding author

Jyoti Ranjan Rout, PhD, FSPP ([✉](mailto:routjr@gmail.com))

School of Biological Sciences, AIPH University,

Pahala, Bhubaneswar – 752101, Odisha, India.

E-mail: routjr@gmail.com; routjr@aiph.ac.in

Contact No: +91 9438047975

Fax: +91 674 2433556

26

27

28 Co-author 1:

29 **Priyadarshani Samal**

30 Biochemistry and Molecular Biology Laboratory,

31 Post Graduate Department of Botany, Utkal University,

32 Vani Vihar, Bhubaneswar-751004, Odisha, India.

33 E-mail: priyadarshanisamal@gmail.com

34

35 Co-author 2:

36 **Priyaranjan Tripathy**

37 Biochemistry and Molecular Biology Laboratory,

38 Post Graduate Department of Botany, Utkal University,

39 Vani Vihar, Bhubaneswar-751004, Odisha, India.

40 E-mail: priyaranjan5@gmail.com

41

42 Co-author 3:

43 **Ritarani Das**

44 Biochemistry and Molecular Biology Laboratory,

45 Post Graduate Department of Botany, Utkal University,

46 Vani Vihar, Bhubaneswar-751004, Odisha, India.

47 E-mail: ritarani.das@gmail.com

48

49 Co-author 4:

50 **Santi Lata Sahoo**

51 Biochemistry and Molecular Biology Laboratory,

52 Post Graduate Department of Botany, Utkal University,

53 Vani Vihar, Bhubaneswar-751004, Odisha, India.

54 E-mail: santi_bot_uu@yahoo.co.in

55

56 Co-author 5:

57 **Chinmay Pradhan**

58 Biochemistry and Molecular Biology Laboratory,

59 Post Graduate Department of Botany, Utkal University,

60 Vani Vihar, Bhubaneswar-751004, Odisha, India.

61 E-mail: chinmay.uubot@gmail.com

62

63 Co-author 6:

64 **Bijaya Kumar Padhi**

65 School of Biological Sciences, Asian Institute of Public Health,

66 Pahala, Bhubaneswar – 752101, Odisha, India.

67 E-mail: bkpadhi@aiph.ac.in

68

69

70

Abstract: Globally, the gastroenteritis or diarrhoea has become a more significant problem today due to infection caused by foodborne/ waterborne pathogen *Vibrio cholera*. In this concern, an investigation was carried out to evaluate the vibriocidal potential of the different solvent extracts of leaf and rhizome of *Maranta arundinacea* under *in vitro* condition. For this, aqueous, methanolic, ethanolic and hexane extracts of both leaf and rhizome of *M. arundinacea* were tested against the pre-isolated strains of *Vibrio cholerae* such as SPAB1, SPAB4 and SPAB5 by agar well diffusion and minimum inhibitory concentration (MIC) method. All the solvent extracts of both leaf and rhizome were found to be active against the tried strains of *V. cholera* however, ethanolic extract showed maximum inhibitory effect against SPAB1 strain with an inhibition zone of 26.23 ± 0.53 mm (MIC of 80.00 ± 10.06 μ g/ ml) and 24.27 ± 0.12 mm (MIC of 100.00 ± 12.82 μ g/ ml) in rhizome and leaf samples, respectively. Then, the effectiveness was followed in SPAB4 and SPAB5 however, it was not much more significant to that of SPAB1. Therefore, it was suggested that the rhizome and leaf extracts which proved to be potentially effective can be used as the natural alternative for the treatment of diarrhoea caused by *Vibrio* infection.

Keywords: Cholera; *Maranta arundinacea* L.; Phytochemical; *Vibrio cholerae*; Vibriocidal

1. Introduction

In the modern time, human beings are facing a serious threat from various water-or foodborne bacterial diseases. Amongst them, diarrhoea (caused by the intestinal pathogen) is one of the major concerns, as it is responsible for primary causes of infant mortality especially in developing countries [1]. More specifically, the cholera which is a severe form of acute secretory diarrhoea is the second leading

cause of mortality worldwide among children under 5 years and also does cause of morbidity in adults [2,3]. According to World Health Organization, it is estimated that the more than one million people are reported by acute diarrhoeal cases annually due to *Vibrio cholerae* infection [4]. Cholera caused by *Vibrio cholerae* is one of the most notorious enteric pathogens responsible for many cholera outbreaks, which are motile, Gram negative, comma shape belonging to the family Vibrionaceae [5]. There are approximately 200 recognized serogroups, of which serogroup O1 and O139 are associated with cholera epidemics in humans [6]. The cholera toxin (also known as cholera toxin) is a protein complex which is secreted by *V. cholerae* mainly responsible for causing copious, painful, watery diarrhoea, leading to vomiting, severe dehydration and even death (if the treatment is not so prompt or even if too late) [7,8].

The emergence and spread of multidrug resistance pathogenic bacteria have created the need for the development of novel therapeutic agents. For which, plants are considered as the best sources of natural products as it cures various diseases from an ancient era [9,10]. Moreover, the World Health Organization certified that the more than 80 % of the World's population relies on traditional medicine for their primary health care needs [11]. During the second half of the 20th century, the acceptance of traditional medicine is increased terrifically and become an alternative option in modern health care as it has no/ less side effects [12,13]. Different plant species containing a wide range of natural products are screened for vibriocidal activities and suggested that the plants are the best potential sources to treat cholera caused by some selective strains of *Vibrio* spp. [14]. It has been also reported that the plants like *Morinda citrifolia*, *Ganophyllum falcatum*, *Leea aequata*, *Lawsonia inermis*, *Saraca indica*, *Syzygium cumini*, *Terminalia belerica*, *Allium sativum*, and

120 *Datura stramonium* are having better effectiveness against vibriocidal activity with
121 different solvent extracts [15,16].

122 *Maranta arundinacea* L. (commonly known as arrowroot plant) belongs to the
123 family Marantaceae is a medicinally as well as economically important plant. Due to
124 its starchy rhizome, it is widely cultivated in India which yields an easily digestible
125 starch. Rhizome parts are used in confectionery for making of biscuits and weaned
126 foods. Medicinally, the rhizome is utilized for various digestive disorders like
127 abdominal pain, indigestion, acidity and irritation on gastrointestinal system [17,18].
128 The plant also possesses anti-inflammatory, antiseptic and antioxidant activity [19].
129 Castor oil-induced antidiarrheal activity was successfully studied in rat the model [20]
130 however, no evidence is found regarding its vibriocidal activity. Hence, the present
131 study is undertaken to evaluate the vibriocidal activity of leaf and rhizome extracts of
132 *M. arundinacea* against *V. cholerae*.

133 2. Materials and Methods

134 2.1. Collection of Plant Material

135 Rhizomes of *M. arundinacea* were collected from the coastal fertile belt of
136 Cuttack, Odisha during the month of February 2013 and planted in the garden of Post
137 Graduate Department of Botany, Utkal University. After identification by Dr. P.C.
138 Panda, Principal Scientist, Regional Plant Resource Centre (RPRC), Bhubaneswar
139 and deposition of the specimen with accession number BOTU10573, the further work
140 was carried out within the laboratory.

141 2.2. Preparation of Various Solvent Extracts of *M. arundinacea*

142 The healthy leaves and rhizomes were harvested, washed under running tap
143 water, cut into small pieces and allowed to air dry up to achieving a constant weight.
144 After drying the plant materials were subjected to extraction by using different

solvents systems (aqueous, ethanol, hexane and methanol) with the help of Soxhlet apparatus for 48 hours. The obtained crude extracts were concentrated by rotary evaporator at reduced pressure and stored at 4 °C for further analysis.

2.3. Microorganisms

The microorganisms used in this investigation included 3 bacterial strains of *V. cholerae* like SPAB1, SPAB4 and SPAB5. The isolated bacteria were identified based on morphological and biochemical characters. Identification and confirmation of *Vibrio cholerae* at the molecular level was done by 16s r RNA sequencing and submitted to NCBI GenBank having accession number KT985959.1, KT985960.1 and KT985961.1 against SPAB1, SPAB4 and SPAB5, respectively.

2.4. Evaluation of Vibriocidal Activity of Plant Extracts

2.4.1. Agar Well Diffusion Assay

Agar well diffusion method initially tried for screening of vibriocidal activity and was carried out as per the method of Pervez *et al.*, 1990 [21]. Mueller Hinton Agar (MHA) plates were swabbed with sterile cotton swabs by taking broth culture of *V. cholerae*. This procedure was repeated by twice and rotated the plates approximately 60° each time for even distribution of the inoculums. Wells (6 mm diameter) were made in each of these plates using cork borer. Each extract was checked for vibriocidal activity by introducing 250 µl of 100 mg/ ml concentration into the wells by using sterile micropipettes. The plates were incubated at 37 °C for 24 hours. After incubation, the zone of inhibition was measured and expressed in millimetres (mm). Ofloxacin (30 µg) was used as the positive control.

2.4.2. Determination of Minimum Inhibitory Concentration

The evaluation of MIC was determined by applying the method of Akinpelu and Kolawole, 2004 [22]. The different concentrations of extracts (20 µg/ ml to 200

µg/ ml) were used with the simple dilution processes. Ofloxacin is taken as a standard drug for the vibrocidal activity. MIC was recorded as lowest extracts of concentration demonstrating no visible growth in the broth.

2.5. Statistical Analysis

All results are the mean of three independent experimental replicates (n = 6) and data is reported as mean \pm standard error.

3. Results and Discussion

3.1. Well Diffusion Analysis

The various solvent (aqueous, ethanol, hexane and methanol) extracts from the leaf and rhizome samples of *M. arundinacea* showed various degrees of the inhibition against three strains of *V. cholerae* using the agar well diffusion method (Table 1). The results are presented by assessing in terms of inhibition of bacterial growth and compared with control (Ofloxacin). The growth inhibition zone measured ranged from 10.04 ± 0.03 - 26.23 ± 0.19 mm for all the strains of *V. cholerae* and for all extracts. However, the maximum inhibitory zone of diameter (24.27 ± 0.12 , 26.23 ± 0.53 mm) were observed in ethanolic extracts of both leaf and rhizome, respectively against SPAB1 and then followed in SPAB4 and SPAB5 (Figure 1). Among all tested extracts, ethanol is highly sensitive and then in methanol, aqueous and hexane. Moreover, the efficacy of rhizome is better than the leaf with respect to both solvents as well as strains (Table 1). The above study significantly points out that both leaf and rhizome of *M. arundinacea* possesses a toxicological impact against Cholera.

Presently there is an increasing demand in herbal medicines by the screening of pharmaceutical or bioactive compounds and their ability to treat various diseases [18,23,24]. Marantaceae plants have been received much more attention since they can produce many complex compounds that are useful for preparation of herbal

medicine [25]. To achieve the effectiveness of vibriocidal activity, various extracts such as aqueous, ethanol, hexane and methanol of leaf and rhizome were tried and significant activeness results were obtained against ethanol extracts which indicates that the ethanol extracts may possess more bioactive compounds and chemical constituents which are responsible for strong vibriocidal activity [26,27]. Phytoconstituents mainly phenols, flavonoids, tannins, alkaloids, steroids and terpenoids are responsible for antimicrobial activity which was confirmed from our previous study on *M. arundinacea* with ethanolic extracts [18,20].

3.2. MIC Analysis

The MIC values of different solvent extracts of leaves and rhizomes are represented in Table 2, indicating that the variations are dependent not only as per the solvent system but also depend on the strains of *V. cholerae*. Among the various solvent extracts studied in this assay, ethanolic extract of rhizome exhibited the smallest inhibitory concentration (80.00 ± 10.06 $\mu\text{g/ml}$, 90.00 ± 10.82 , 100.00 ± 16.11) which indicates the maximum prevention power against SPAB1, SPAB5, SPAB4, respectively. However in case of leaf samples, no such significant variations were observed.

To support the vibriocidal activity of *M. arundinacea*, the MIC values were determined to know the efficacy of concentration against different strains of *V. cholerae* (SPAB1, SPAB4 and SPAB5). The presence of possible constituents was observed even in a low concentration of solvent extracts of leaf and rhizome. Here the same ethanol yielding extracts were more powerful than others which indicate also might be due to dissolving or diffusing nature of bioactive compounds present within plant extracts [24]. Moreover, the hydrophobic and hydrogen bonding nature of phenolic compounds from plant extracts attach to membrane proteins, followed by

involvement in the lipid bilayer do inhibition of spreading of tested bacteria as it is a Gram negative in nature and hence acted as a potent bacteriostatic agent to *V. cholerae* [28,29]. *M. arundinacea* leaf and rhizome ethanol extracts have both growth inhibition as well as minimum inhibition activity against different strains of *V. cholerae* which confirms that the tested plant has efficacy towards antidiarrheal activity.

4. Conclusion

The present study concludes that the ethanol extract of rhizome and leaf of *M. arundinacea* is a better option to inhibit the spreading of *V. cholerae*, which is a very challenging pathogen that can cause cholera to humans. For example, is being used for decades as a medicinal plant against various gastrointestinal disturbances, but to our knowledge, this is the first ever report of its efficacy in controlling growth of *V. cholerae*. Moreover, the study indicates that the experimental plant contains some natural anti-vibrio compounds which may need further isolation and characterization and that may be used in future for human health and disease prevention perspective. For that, additional work is essential at the cellular and subcellular level to understand the mechanisms of action of the active plant extracts.

Author Contributions: Conceived, designed and supervised the experiments, J.R.R., S.L.S. and R.D.; performed the experiments and writing-original draft preparation, P.S. and P.T.; analysed the data and wrote the manuscript, P.S., J.R.R. and R.D.; made critical revisions, C.P. and B.K.P.; all authors read, reviewed and approved the final manuscript.

Funding: This research received no external funding.

Acknowledgements: The authors are grateful to the Head of the Department, Post Graduate Department of Botany for providing all the experimental facilities.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Kosek, M.; Bern, C.; Guerrant R.L. The global burden of diarrhoeal disease, as estimated from studies published between 1992 and 2000. *Bull. World Health. Organ.* 2003, 81(3), 197–204.
2. Bryce, J.; Boschi-Pinto, C.; Shibuya, K.; Black, R.E. WHO estimates of the causes of the death in children. *Lancet* 2005, 365, 1147–1152.
3. Harris, J.B.; Larocque, R.C.; Qadri, F.; Ryan, E.T.; Calderwood, S.B. Cholera. *Lancet* 2012, 379, 2466–2476.
4. World Health Organization. Cholera 2012. *Wkly. Epidemiol. Rec.* 2013, 88, 321–336.
5. Nishibori, T.; de Vries, G.C.; Rahardjo, D. *et al.* Phenotypic and genotypic characterization of *Vibrio cholerae* clinically isolated in Surabaya, Indonesia. *Jpn. J. Infect. Dis.* 2011, 64(1), 7–12.
6. Ried, J.; Klose, K.E. *Vibrio cholerae* and cholera: out of the water into the host. *FEMS Microbiol. Rev.* 2002, 26, 125–139.
7. Diallo, D.; Hveem, B.; Mahmoud, M.A.; Betge, G.; Paulsen, B.S.; Maiga, A. An international survey of herbal drugs of Gourma district. *J. Pharm. Biol.* 1999, 37(1), 80–91.
8. Bharati, K.; Ganguly, N.K. Cholera toxin: a paradigm of a multifunctional protein. *Indian J. Med. Res.* 2011, 133(2), 179–187.
9. Ji, L.; Larregieu, C.A.; Benet, L.Z. Classification of natural products as sources of drugs according to the biopharmaceutics drug disposition classification system (BDDCS). *Chin. J. Nat. Med.* 2016, 14(12), 0888–0897.

- 269 10. Dias, D.A.; Urban, S.; Roessner, U. A historical overview of natural products
270 in drug discovery. *Metabolites* 2012, 2(2), 303–336.
- 271 11. Ekor, M. The growing use of herbal medicines: issues relating to adverse
272 reactions and challenges in monitoring safety. *Front. Pharmacol.* 2013,
273 4(177), 1–10.
- 274 12. Panda, S.; Rout, J.R.; Pati, P.; Ranjit, M.R.; Sahoo, S.L. Antimalarial activity
275 of *Artemisia nilagirica* against *Plasmodium falciparum*. *J. Parasit. Dis.* 2018,
276 42(1), 22–27.
- 277 13. Ozkan, G.; Kamiloglu, S.; Ozdal, T.; Boyacioglu, D.; Capanoglu, E. Potential
278 use of turkish medicinal plants in the treatment of various diseases. *Molecules*
279 2016, 21(257), 1–32.
- 280 14. Aminzare, M.; Hashemi, M.; Abbasi, Z.; Mohseni, M.; Amiri, E. Vibriosis
281 phytotherapy: a review on the most important world medicinal plants effective
282 on *Vibrio* spp. *J. Appl. Pharm. Sci.* 2018, 8(1), 170–177.
- 283 15. Chander, M.P.; Vijayachari, P. Vibriocidal activity of selected medicinal
284 plants used by Nicobarese tribe of Andaman and Nicobar Islands, India. *J.*
285 *Pharm. Pharmacol.* 2017, 5, 164–168.
- 286 16. Sharma, A.; Patel, V.K.; Chaturvedi, A.N. Vibriocidal activity of certain
287 medicinal plants used in Indian folklore medicine by tribals of Mahakoshal
288 region of central India. *Indian J. Pharmacol.* 2009, 41(3), 129–133.
- 289 17. Jayakumar, A.; Suganthi, A. Biochemical and phytochemical analysis of
290 *Maranta arundinacea* (L.) rhizome. *Int. J. Pharm. Pharm. Sci.* 2017, 2(3), 26–
291 30.

- 292 18. Samal, P.; Rout, J.R.; Das, R.; Sahoo, S.L.; Padhi, B.K. Screening and
 293 evaluation of phytochemicals from *Maranta arundinacea* L. *Int. J. Biol. Med.*
 294 *Res.* 2018, 9(1), 6212–6217.
- 295 19. Ruba, A.A.; Mohan, V.R. Evaluation of total phenolic and flavonoid contents
 296 and *in vitro* antioxidant activity of rhizome of *Maranta arundinacea* L. *Int J.*
 297 *Pharm. Sci.* 2013, 4(2), 3914–3928.
- 298 20. Rahman, M.K.; Chowdhury, M.A.U.; Islam, M.T.; Chowdhury, M.A.; Uddin,
 299 M.E.; Sumi, C.D. Evaluation of antidiarrhoeal activity of methanolic extract of
 300 *Maranta arundinacea* Linn. Leaves. *Adv. Pharmacol. Sci.* 2015, 2015, 1–6.
- 301 21. Pervez, C.; Pauli, M.; Bazerque, P. An antibiotic assay by agar well diffusion
 302 method. *Acta Biol. Med. Exp.* 1990, 15, 113–115.
- 303 22. Akinpelu, D.A.; Kolawole, D.O. Phytochemical and antimicrobial activity of
 304 leaf extract of *Piliostima thonningii* (Schum.). *Science Focus* 2004, 7, 64–70.
- 305 23. Lobo, R.; Prabhu, K.S.; Shirwaikar, A.; Shirwaikar, A. *Curcuma zedoaria*
 306 Rosc. (white turmeric): a review of its chemical, pharmacological and
 307 ethnomedicinal properties. *J. Pharm. Pharmacol.* 2009, 61, 13–21.
- 308 24. Choudhury, S.; Sharan, L.; Sinha, M.P. Screening of some commonly used
 309 medicinal plants against enteric human pathogen *Vibrio cholera*. *European J.*
 310 *Med. Plants* 2015, 9(3), 1–6.
- 311 25. Hattori, S. Utilization of Marantaceae plants by the Baka hunter-gatherers in
 312 southeastern Cameroon. *Afr. Stud. Monogr.* 2006, 33, 29–48.
- 313 26. Koffi, A.E.; Francis, Y.A.; Felix, Y.H.; Calixte, B.; Joseph, D.A. Comparative
 314 study of *in vitro* susceptibility of clinical and environmental strains of *Vibrio*
 315 *cholerae* O1 strains to ethanol extracts of *Morinda morindoides*. *Eur. J. Sci.*
 316 *Res.* 2011, 66(2), 284–292.

27. Payne, A.; Mukhopadhyay, A.K.; Deka, S.; Saikia, L.; Nandi, S.P. Anti-vibrio and antioxidant properties of two weeds: *Euphorbia serpens* and *Amaranthus viridis*. *Res. J. Med. Plants* 2015, 9(4), 170–178.
28. Kamruzzaman, M.; Bari, S.M.N.; Faruque, S.M. *In vitro* and *in vivo* bactericidal activity of *Vitex negundo* leaf extract against diverse multidrug resistant enteric bacterial pathogens. *Asian Pac. J. Trop. Med.* 2013, 6(5), 352–359.
29. Akrayi, H.F.S. Antibacterial effect of aqueous extracts of spices and herbs against bacteria isolated from frozen meat. *Med. J. Islamic World Acad. Sci.* 2014, 22(1), 30–35.




Figure 1. Plate showing zone of inhibition against leaf and rhizome ethanolic extracts of *M. arundinacea* against different strains of *Vibrio cholerae*. **A:** *Vibrio cholerae* strain SPAB1; **B:** *Vibrio cholerae* strain SPAB4; **C:** *Vibrio cholerae* strain SPAB5.

Table 1: The inhibition zone diameter (mm) of leaf and rhizome extracts of *M. arundinacea* against different strains of *Vibrio cholerae*.

Extracts	Parts used	Strains of <i>Vibrio cholerae</i>		
		SPAB1	SPAB4	SPAB5
AQ	Le	21.08 ± 0.16	20.10 ± 0.28	21.08 ± 0.21
	Rh	22.09 ± 0.78	22.14 ± 0.71	23.16 ± 0.48
ET	Le	24.27 ± 0.12	23.20 ± 0.29	22.15 ± 0.37
	Rh	26.23 ± 0.53	25.20 ± 0.22	24.20 ± 0.12
HE	Le	16.06 ± 0.51	10.04 ± 0.18	18.06 ± 0.54
	Rh	16.18 ± 0.48	15.05 ± 0.46	20.07 ± 0.59
ME	Le	23.20 ± 0.94	24.13 ± 0.38	23.18 ± 0.18
	Rh	25.31 ± 0.41	22.16 ± 0.72	22.18 ± 0.75
Ofloxacin (30 µg)		13.01 ± 0.68	12.06 ± 0.56	11.00 ± 0.33

AQ: Aqueous extract; ET: Ethanolic extract; HE: Hexane extract; ME: Methanolic extract; Le: Leaf; Rh: Rhizome. The data represents mean ± SE of replicates (n = 6).

381 **Table 2:** The minimum inhibitory concentration ($\mu\text{g}/\text{ml}$) of the ethanolic leaf
 382 and rhizome extracts of *M. arundinacea* against different strains of *Vibrio cholerae*.

Solvent extracts	Parts used	Different strains of <i>Vibrio cholerae</i>		
		SPAB1	SPAB4	SPAB5
AQ	Le	110.00 \pm 12.61	120.00 \pm 11.55	120.00 \pm 12.23
	Rh	100.00 \pm 20.17	120.00 \pm 13.46	110.00 \pm 11.02
ET	Le	100.00 \pm 12.82	100.00 \pm 14.23	110.00 \pm 12.78
	Rh	80.00 \pm 10.06	100.00 \pm 16.11	90.00 \pm 10.82
HE	Le	120.00 \pm 12.90	120.00 \pm 13.32	120.00 \pm 13.22
	Rh	110.00 \pm 09.15	120.00 \pm 12.26	110.00 \pm 12.55
ME	Le	110.00 \pm 10.56	120.00 \pm 12.44	110.00 \pm 12.54
	Rh	90.00 \pm 11.15	110.00 \pm 09.18	100.00 \pm 11.03
Ofloxacin (30 μg)		140.00 \pm 11.72	160.00 \pm 16.60	140.00 \pm 15.91

383 AQ: Aqueous extract; ET: Ethanolic extract; HE: Hexane extract; ME: Methanolic
 384 extract; Le: Leaf; Rh: Rhizome. The data represents mean \pm SE of replicates (n = 6).

385