

On the Eternal Role of Planck Scale in Machian Cosmology

U. V. S. Seshavatharam¹ and S. Lakshminarayana²

¹Honorary Faculty, I-SERVE, S. No-42, Hitex Road, Hitech City, Hyderabad-84, Telangana, India.

²Department of Nuclear Physics, Andhra University, Visakhapatnam-03, AP, India.

Email(s): seshavatharam.uvs@gmail.com (and) sln@auvsp.edu.in

Abstract: Considering Planck mass as a representation of baby universe and evolving universe as a growing Planck ball, a heuristic model of quantum cosmology can be developed. By considering the famous relation $GM = c^2 R$, a Machian model of evolving quantum cosmology can also be developed with increasing matter content and decreasing matter density. In all directions, ‘from and about’ the baby universe, cosmic expansion velocity can be shown to be increasing with decreasing matter density. Estimated current cosmic mass, radius, matter density, expansion velocity and age are: 4.3352×10^{53} kg, 3.207×10^{26} m, 3.138×10^{-27} kg.m⁻³, $2.43c$ and 19.78 Billion years respectively. As a peculiar case, considering the equality of current Hubble parameter and current angular velocity, current cosmic rotational kinetic energy can be shown to be 0.667 times the current critical energy! Proceeding further, with reference to Planck scale, cosmic temperature seems to be redshifted by a factor $(Z_T)_t \cong \sqrt{1 + \ln(H_{pl}/H_t)} - 1$ where (H_{pl}, H_t) represent Planck scale and time dependent Hubble parameters respectively.

Keywords: Planck scale; Mach’s relation; quantum cosmology; critical density; ordinary matter; dark matter; thermal redshift; expansion velocity; rotational kinetic energy; dark energy; Hubble’s law;

1. INTRODUCTION

Replacing ‘big bang’ (Gamow 1948a, 1948b, Abhas Mitra 2011, 2014) with a growing Planck ball and considering ‘Mach’s relation’ (Patrick Grahn et al 2018; Annala 2012; Berman 2008a, 2009b; Arbab 2001a, 2004b; Miyazaki 2001;) as a deep cosmic probe, in a hypothetical approach, an evolving model of quantum cosmology can be developed (Bojowald 2015, Arbab 2004, Padmanabhan 2006, Sivaram 2000). To make the model sensible, we assume that, with reference to Planck scale, cosmic temperature is redshifted by a factor, $(Z_T)_t \cong \sqrt{\gamma_t} - 1 \cong \sqrt{1 + \ln(H_{pl}/H_t)} - 1$ where (H_{pl}, H_t) represent Planck scale and time dependent Hubble parameters respectively and $(Z_T)_t$ is the redshift associated with temperature T_t about the baby universe. Considering the characteristic γ_t , current cosmic temperature and current Hubble parameter can be fitted and we make an attempt to estimate the current cosmic mass and radius. Proceeding further, in analogy with ‘cosmic acceleration’, in this paper, using γ_t , we proposed simple relations for a decreasing trend of total matter density with increasing trend of expansion velocity referred to ‘from and about’ the baby universe in all directions.

1.1 Motivating Points

- 1) Mach’s principle (Einstein 1918) is one of the iconic principles underlying general theory of relativity and can be given a priority in developing a workable or unified model of cosmology.
- 2) Cosmic expansion, Lambda term, dark matter, cosmic temperature, inflation, cosmic acceleration and dark energy and vacuum energy are different concepts, by using which alternative models of GTR are emerging and are being extended in many ways. In this sequence, quantum cosmology can also be given some consideration.
- 3) Quantum cosmology is a wide range physical model intended for understanding the in-built cosmological quantum phenomena on small scale as well as large scale distances. So far, progress in this direction is very nominal and ‘GTR’ needs a serious review with reference to ‘quantum cosmology’.
- 4) When universe is able to give birth to atoms, elementary particles and photons that show quantum behaviour, universe can certainly be considered as a quantum gravitational object for ever.
- 5) What to quantize? How to quantize? When to quantize? and What to measure? etc. are some interesting questions in current quantum cosmology and need a special focus. In this context, cosmic temperature can be considered as a characteristic feature of quantum cosmology.
- 6) According to current notion, if the known laws of physics are extrapolated to the highest density regime, the result is a singularity which is typically associated with the Big Bang. In this context, Planck mass can be considered as a

characteristic seed of the evolving universe and big bang can be replaced with an evolving Planck ball. Planck mass can be called as the ‘baby Planck ball’ or ‘baby universe’.

- 7) With reference to particle physics, current technological limits on particle colliding energy, unidentified/unseen particles, unknown particle interactions and incomplete final unification scheme - to some extent, one can hopefully believe in the existence of dark matter (Rennan Barkana, 2018).
- 8) Basically, ‘dark energy’ was proposed for understanding cosmic acceleration. Careful analysis of improved supernovae data suggests that universe is coasting at constant velocity and evidence for acceleration is only marginal (Neilsen et al. 2016, Lawrence H. Dam et al., 2017). In this context, now a days, a great debate has been initiated among mainstream cosmologists on the existence of dark energy (Balakrishna et al 2017, G’abor R’acz et al 2017, Lasha Berezhiani et al, 2017, Joel Smoller et al, 2017). According to a new study (Gong-Bo Zhao et al, 2017), the nature of dark energy is ‘dynamic’ and conceptually seems to deviate from the famous cosmological constant or vacuum energy. According to another new study (Deng Wang and Xin-He Meng, 2017), evidence for dynamical dark energy is very poor.
- 9) Density perturbations (Joel Smoller et al 2017) and interaction between dark matter and baryons (Lasha Berezhiani et al 2017) seem to play a crucial role in understanding observed cosmic acceleration and need of introducing dark energy seems to be ad-hoc.
- 10) Even though redshift is an index of cosmic expansion, without knowing the actual galactic distances and actual galactic receding speeds, with 100% confidence level, it may not be possible to decide the absolute nature of cosmic expansion rate.

2. ASSUMPTIONS, CONCEPTS AND BASIC RELATIONS

2.1 Nomenclatures

- 1) $(\Omega_{OM})_t$ = Ratio of ordinary matter density to critical density.
- 2) $(\Omega_{DM})_t$ = Ratio of dark matter density to critical density.
- 3) H_t = Hubble parameter and H_{pl} = Planck scale Hubble parameter.
- 4) $(V_{exp})_t$ = Cosmic expansion velocity from and about the baby universe or baby Planck ball.
- 5) $(M_{OM})_t$ = Cosmic ordinary mass content, $(M_{DM})_t$ = Cosmic dark matter content.
- 6) $(M_{OM} + M_{DM})_t \cong M_t$ = Total matter content = Total mass of evolving Planck ball.
- 7) R_t = Cosmic radius associated with M_t = Radius of evolving Planck ball.
- 8) T_t = Cosmic temperature and $(Z_T)_t$ = Redshift associated with T_t about the baby universe.
- 9) $(T_c)_t = \left(\frac{3H_t^2 c^2}{8\pi G a} \right)^{\frac{1}{4}}$ = Critical temperature.
- 10) $\gamma_t \cong \left[1 + \ln \left(\frac{H_{pl}}{H_t} \right) \right] \cong \sqrt{\frac{3H_t^2 c^2}{8\pi G (aT_t^4)}} = \text{Root of ratio of critical energy density and thermal energy density.}$
- 11) (d_g) = Galactic distance from and about the baby universe or baby Planck ball.
- 12) (v_g) = Galactic receding speed from and about the baby universe or baby Planck ball.

Note-1: For the above symbols, subscript 0 denotes current value and subscript pl denotes Planck scale value.

2.2 Proposed Assumptions

With respect to our earlier publications (Tatum et al 2015, Seshavatharam & Lakshminarayana 2017a, 2017b, 2018c, 2018d) in this paper we review the basic assumptions.

- 1) Planck scale and Mach’s relation play a crucial role in entire cosmic evolution.
- 2) With reference to Planck scale, cosmic temperature is redshifted by a factor $(Z_T)_t \cong \sqrt{\gamma_t} - 1 \cong \sqrt{1 + \ln(H_{pl}/H_t)} - 1$.
- 3) Ordinary matter and dark matter, both, play a crucial role in estimating cosmic expansion velocity and there is no need to consider dark energy for the purpose of cosmic expansion rate.

Note-2: If there is no redshift in cosmic temperature, at any stage of cosmic evolution, thermal energy density and critical energy density will always be equal in magnitude. It may also be noted that, proposed thermal redshift is with reference to the baby universe and currently believed notion of redshift is with reference to distance between earth and any receding galaxy.

Note-3: Based on these assumptions, from the beginning of Planck scale, a) With a ‘decreasing’ trend of total matter density ratio, cosmic expansion velocity can be shown to be increasing. b) With an ‘increasing’ trend of total matter density ratio, cosmic expansion velocity can be shown to be decreasing. c) With a constant trend of total matter density ratio, cosmic expansion velocity can be shown to be constant.

2.3 Choosing the Magnitude of H_0

- 1) As per the 2015 Planck data (Planck Collaboration 2015) : $H_0 \cong (67.31 \pm 0.96)$ km/sec/Mpc and the present temperature of the CMB radiation is, $T_0 \cong (2.722 \pm 0.027)$ K.
- 2) According to the advanced observational data analysis by Riess et al.(2016) current best value of $H_0 \cong (73.24 \pm 1.74)$ km/sec/Mpc.
- 3) With reference to $T_0 \cong 2.722$ K and our proposed set of concepts, in this paper, we choose,
 $H_0 \cong 70 \text{ km/sec/Mpc} \cong 2.26853 \times 10^{-18} \text{ sec}^{-1}$. This value seems to lie in between (67.31 and 73.24) km/sec/Mpc.

2.4 The Planck Scale in Entire Cosmic Evolution

So far no mainstream cosmological model implemented Planck scale in current cosmic evolution. In this complicated situation, in a positive approach, we make an attempt to implement the ‘Planck scale’ in the entire cosmic evolution. With further study, our approach can be developed for a better understanding. Based on quantum gravity, we define the

Planck scale Hubble parameter, $H_{pl} \cong \sqrt{\frac{c^5}{G\hbar}} \cong 1.854921 \times 10^{43} \text{ sec}^{-1}$. To proceed further, we define that,

$$\left. \begin{aligned} \sqrt{\left(\frac{3H_t^2 c^2}{8\pi G (aT_t^4)} \right)} &\cong \gamma_t \cong \left[1 + \ln \left(\frac{H_{pl}}{H_t} \right) \right] \\ \left(\frac{3H_t^2 c^2}{8\pi G (aT_t^4)} \right)^{\frac{1}{4}} &\cong \sqrt{\gamma_t} \cong \sqrt{1 + \ln \left(\frac{H_{pl}}{H_t} \right)} \cong ((Z_T)_t + 1) \end{aligned} \right\} \quad (1)$$

Based on this relation, if defined $H_{pl} \cong 1.854921 \times 10^{43} \text{ sec}^{-1}$, one can choose different values of γ in between $\gamma_{pl} \cong 1$ and $\gamma_0 \cong 141.2564$. For each assumed value of H , one can get a corresponding γ and all other physical parameters can

be estimated. For the Planck scale, $\sqrt{\left(\frac{3H_{pl}^2 c^2}{8\pi G (aT_{pl}^4)} \right)} \cong \gamma_{pl} \cong 1$ and for the current case,

$\sqrt{\left(\frac{3H_0^2 c^2}{8\pi G (aT_0^4)} \right)} \cong \gamma_0 \cong \left[1 + \ln \left(\frac{H_{pl}}{H_0} \right) \right] \cong 141.2564$. In a simplified form, cosmic temperature can be expressed as,

$$T_t \cong \frac{1}{\sqrt{\gamma_t}} \left(\frac{3H_t^2 c^2}{8\pi G a} \right)^{\frac{1}{4}} \cong \frac{(T_c)_t}{\sqrt{\gamma_t}} \cong \frac{1}{\sqrt{\gamma_t}} \left\{ \frac{0.652632 \hbar \sqrt{H_{pl} H_t}}{k_B} \right\} \quad (2)$$

3. TO ESTIMATE THE COSMIC MASS, RADIUS AND EXPANSION VELOCITY

$$\text{Let, } (M_{OM} + M_{DM})_t \cong M_t \cong \left[(\Omega_{OM})_t + (\Omega_{DM})_t \right] \left(\frac{3H_t^2}{8\pi G} \right) \left(\frac{4\pi}{3} R_t^3 \right) \quad (3)$$

$$G(M_{OM} + M_{DM})_t \cong GM_t \cong R_t c^2 \quad (4)$$

$$R_t \cong \frac{G(M_{OM} + M_{DM})_t}{c^2} \cong \frac{GM_t}{c^2} \quad (5)$$

Based on these relations, it is possible to show that,

$$R_t \cong \frac{G(M_{OM} + M_{DM})_t}{c^2} \cong \sqrt{\frac{2}{(\Omega_{OM} + \Omega_{DM})_t}} \left(\frac{c}{H_t} \right) \quad (6)$$

$$(V_{\text{exp}})_t \cong R_t H_t \cong \sqrt{\frac{2}{(\Omega_{OM} + \Omega_{DM})_t}} (c) \quad (7)$$

$$(M_{OM} + M_{DM})_t \cong M_t \cong \sqrt{\frac{2}{(\Omega_{OM} + \Omega_{DM})_t}} \left(\frac{c^3}{GH_t} \right) \cong \left(\frac{c^2 (V_{\text{exp}})_t}{GH_t} \right) \quad (8)$$

Based on relations (6 to 8) and in terms of $(\Omega_{OM})_t$ and $(\Omega_{DM})_t$,

$$(M_{OM})_t \cong \left[\frac{(\Omega_{OM})_t}{(\Omega_{OM} + \Omega_{DM})_t} \right] \left(\frac{c^2 (V_{\text{exp}})_t}{GH_t} \right) \quad (9)$$

$$(M_{DM})_t \cong \left[\frac{(\Omega_{DM})_t}{(\Omega_{OM} + \Omega_{DM})_t} \right] \left(\frac{c^2 (V_{\text{exp}})_t}{GH_t} \right) \quad (10)$$

4. UNDERSTANDING THE TREND OF ORDINARY AND DARK MATTER DENSITY RATIOS

With the help of defined $\gamma_t \equiv \left[1 + \ln \left(\frac{H_{pl}}{H_t} \right) \right]$ and $\sqrt{\gamma_t} \equiv \sqrt{1 + \ln \left(\frac{H_{pl}}{H_t} \right)}$, on ad-hoc basis and with reference to current observed values of $(\Omega_{OM})_0$ and $(\Omega_{DM})_0$, we are making an attempt to estimate the past values of $(\Omega_{OM})_t$ and $(\Omega_{DM})_t$. In this context, the basic question to be answered is: Is there any scope for the existence for dark matter at Planck scale? It needs further study.

4.1 Current ordinary matter density ratio

If it is assumed that,

$$\left\{ [(M_{OM})_0 c^2] / \left[\frac{4\pi}{3} R_0^3 \right] \right\} \propto \sqrt{\left(\frac{3H_0^2 c^2}{8\pi G} \right)} (aT_0^4) \quad (11)$$

As per relation (1),

$$\left\{ [(M_{OM})_0 c^2] / \left[\frac{4\pi}{3} R_0^3 \right] \right\} \propto \frac{1}{\gamma_0} \left(\frac{3H_0^2 c^2}{8\pi G} \right) \quad (12)$$

By considering an ad-hoc proportionality coefficient of the form, $\beta_0 \equiv \left(\frac{1 + \sqrt{\gamma_0}}{2} \right) \equiv \left(1 + \frac{(Z_T)_0}{2} \right)$,

$$\left\{ \left[(M_{OM})_0 c^2 \right] / \left[\frac{4\pi}{3} R_0^3 \right] \right\} \cong \frac{\beta_0}{\gamma_0} \left(\frac{3H_0^2 c^2}{8\pi G} \right)$$

$$\text{and } (\Omega_{OM})_0 \cong \left(\frac{1}{\gamma_0} \right) \left(\frac{1 + \sqrt{\gamma_0}}{2} \right) \cong \frac{\beta_0}{\gamma_0} \cong 0.04561$$
(13)

where $\sqrt{\gamma_0} \cong [1 + (Z_T)_0] \cong (1 + Z_{2.722K}) \cong 11.885$ and $Z_{2.722K} \cong 10.885$.

4.2 Current dark matter density ratio

Like the case of ordinary matter density,

$$\left\{ \left[(M_{DM})_0 c^2 \right] / \left[\frac{4\pi}{3} R_0^3 \right] \right\} \propto \sqrt{\left(\frac{3H_0^2 c^2}{8\pi G} \right) (aT_0^4)}$$
(14)

As per relation (1),

$$\left\{ \left[(M_{DM})_0 c^2 \right] / \left[\frac{4\pi}{3} R_0^3 \right] \right\} \propto \frac{1}{\gamma_0} \left(\frac{3H_0^2 c^2}{8\pi G} \right)$$
(15)

By considering an ad-hoc proportionality coefficient of the form, $\beta_0^2 \cong \left(\frac{1 + \sqrt{\gamma_0}}{2} \right)^2 \cong \left(1 + \frac{(Z_T)_0}{2} \right)^2$,

$$\left\{ \left[(M_{DM})_0 c^2 \right] / \left[\frac{4\pi}{3} R_0^3 \right] \right\} \cong \frac{\beta_0^2}{\gamma_0} \left(\frac{3H_0^2 c^2}{8\pi G} \right)$$

$$\text{and } (\Omega_{DM})_0 \cong \left(\frac{1}{\gamma_0} \right) \left(\frac{1 + \sqrt{\gamma_0}}{2} \right)^2 \cong \frac{\beta_0^2}{\gamma_0} \cong 0.29384$$
(16)

4.3 General expressions for understanding the trend of ordinary and dark matter density ratios

By extrapolating the coefficient to past and future as,

$$\beta_t \cong \left(\frac{1 + \sqrt{\gamma_t}}{2} \right) \text{ and } \beta_t^2 \cong \left(\frac{1 + \sqrt{\gamma_t}}{2} \right)^2$$
(17)

$$\left. \begin{array}{l} \text{A) } (\Omega_{OM})_t \cong \left(\frac{1}{\gamma_t} \right) \left(\frac{1 + \sqrt{\gamma_t}}{2} \right) \cong \frac{\beta_t}{\gamma_t}; \\ \text{B) } (\Omega_{DM})_t \cong \left(\frac{1}{\gamma_t} \right) \left(\frac{1 + \sqrt{\gamma_t}}{2} \right)^2 \cong \frac{\beta_t^2}{\gamma_t}; \\ \text{C) } \frac{(\Omega_{DM})_t}{(\Omega_{OM})_t} \cong \left(\frac{1 + \sqrt{\gamma_t}}{2} \right) \cong \beta_t; \\ \text{D) } (\Omega_{OM} + \Omega_{DM})_t \cong \frac{\beta_t + \beta_t^2}{\gamma_t} \end{array} \right\}$$
(18)

See the following picture-1. With reference to critical density, dashed blue curve represents the trend of ordinary matter density ratio and black curve represents the trend of dark matter density ratio.

Qualitatively, we noticed that, $\frac{(Z_T)_t}{2} \approx \ln(\gamma_t)$ and $\beta_t \approx 1 + \ln(\gamma_t)$. With reference to the 'status of existence' of ordinary matter and dark matter and with further research, various relations for β_t can be developed using by which, in near

future, trend of ordinary and dark matter density ratios can be understood right from the beginning of Planck scale. See figure -2 for the decreasing trend of total matter density ratio.

Figure 1: Decreasing trend of ordinary and dark matter density ratios





Figure 2: Decreasing trend of total matter density ratio



5. CURRENT AND PLANCK SCALE COSMIC PHYSICAL PARAMETERS

See table-1 for various cosmic physical parameters associated with current and Planck scales.

Table-1: Current and Planck scale cosmic physical parameters

Current scale	Planck scale
$H_0 \cong 70\text{km/sec/Mpc} \cong 2.26853 \times 10^{-18} \text{ sec}^{-1}$	$H_{pl} \cong \sqrt{\frac{c^5}{G\hbar}} \cong 1.855 \times 10^{43} \text{ sec}^{-1}$
$\gamma_0 \cong \left[1 + \ln \left(\frac{H_{pl}}{H_0} \right) \right] \cong 141.2564$	$\gamma_{pl} \cong \left[1 + \ln \left(\frac{H_{pl}}{H_{pl}} \right) \right] \cong 1$
$(\Omega_{OM})_0 \cong 0.04561$	$(\Omega_{OM})_{pl} \cong 1$

$(\Omega_{DM})_0 \cong 0.29384$	$(\Omega_{DM})_{pl} \cong 1$
$T_0 \cong \left(\frac{1}{\sqrt{\gamma_0}}\right) \left(\frac{3H_0^2 c^2}{8\pi G a}\right)^{\frac{1}{4}} \cong 2.721 \text{ K}$	$T_{pl} \cong \left(\frac{1}{\sqrt{\gamma_{pl}}}\right) \left(\frac{3H_{pl}^2 c^2}{8\pi G a}\right)^{\frac{1}{4}} \cong 9.247 \times 10^{31} \text{ K}$
$R_0 \cong \sqrt{\frac{2}{[(\Omega_{OM})_0 + (\Omega_{DM})_0]}} \frac{c}{H_0}$ $\cong 3.207 \times 10^{26} \text{ m} \cong 10.40 \text{ Gpc}$	$R_{pl} \cong \sqrt{\frac{2}{[(\Omega_{OM})_{pl} + (\Omega_{DM})_{pl}]}} \left(\frac{c}{H_{pl}}\right)$ $\cong 1.616 \times 10^{-35} \text{ m}$
$(V_{\text{exp}})_0 \cong R_0 H_0 \cong 2.42654c$	$(V_{\text{exp}})_{pl} \cong R_{pl} H_{pl} \cong c$
$(M_{OM})_0 \cong 5.836 \times 10^{52} \text{ kg}$	$(M_{OM})_{pl} \cong 1.0882 \times 10^{-8} \text{ kg}$
$(M_{DM})_0 \cong 3.752 \times 10^{53} \text{ kg}$	$(M_{DM})_{pl} \cong 1.0882 \times 10^{-8} \text{ kg}$
$[(M_{OM})_0 + (M_{DM})_0] \cong M_0 \cong 4.3352 \times 10^{53} \text{ kg}$	$[(M_{OM})_{pl} + (M_{DM})_{pl}] \cong M_{pl} \cong 2.176 \times 10^{-8} \text{ kg}$
$t_0 \cong \frac{2R_0}{[(V_{\text{exp}})_0 + (V_{\text{exp}})_{pl}]} \cong 19.78 \text{ BillionYears}$	$t_{pl} \cong 0$

6. COSMIC SCALE FACTOR AND REDSHIFT

With reference to the proposed relations (1) and (2) and with reference to the current definitions of cosmic redshift and scale factor, it is possible to show that,

$$\begin{aligned} \left(\frac{1}{a} \cong (z+1) \cong \frac{T_t}{T_0}\right) &\cong \sqrt{\frac{\gamma_0 H_t}{\gamma_t H_0}} \cong \sqrt{\frac{\gamma_0}{\gamma_t}} \sqrt{\frac{H_t}{H_0}} \\ &\cong \sqrt{\frac{\gamma_0}{\gamma_t}} \left\{ \exp\left(\frac{\gamma_0 - \gamma_t}{2}\right) \right\} \cong \sqrt{\left(\frac{\gamma_0}{\gamma_t}\right)} \exp(\gamma_0 - \gamma_t) \end{aligned} \quad (19)$$

Redshift can be expressed in the following form.

$$z \cong \sqrt{\left(\frac{\gamma_0}{\gamma_t}\right)} \exp(\gamma_0 - \gamma_t) - 1 \quad (20)$$

We are working on interpreting this relation and it needs further study.

7. TO ESTIMATE THE CURRENT COSMIC AGE

From the beginning of cosmic evolution, based on the proposed cosmic expansion velocities, cosmic age can be approximated with the following relation.

$$t \cong \frac{(R_t - R_{pl})}{\left[\left((V_{\text{exp}})_t + (V_{\text{exp}})_{pl}\right)/2\right]} \quad (21)$$

where $\left[\left((V_{\text{exp}})_t + (V_{\text{exp}})_{pl}\right)/2\right]$ can be considered as average expansion velocity.

For the current case,

$$t_0 \cong \frac{(R_0 - R_{pl})}{\left[\left((V_{\text{exp}})_0 + (V_{\text{exp}})_{pl}\right)/2\right]} \cong \frac{2R_0}{\left((V_{\text{exp}})_0 + (V_{\text{exp}})_{pl}\right)} \cong 19.78 \text{ BillionYears} \quad (22)$$

For a temperature of 3000 K, it is possible to show that,

$$\left. \begin{aligned} H_{3000K} &\cong 2.49 \times 10^{-12} \text{ sec}^{-1} \\ \gamma_{3000K} &\cong 1 + \ln \left(\frac{H_{pl}}{H_{3000K}} \right) \cong 127.34774 \\ Z_{3000K} &\cong \sqrt{\gamma_{3000K}} - 1 \cong 10.285 \\ z_{3000K} &\cong \sqrt{\frac{\gamma_0}{\gamma_{3000K}} \exp(\gamma_0 - \gamma_{3000K})} - 1 \cong 1102.407 \end{aligned} \right\} \quad (23)$$

Cosmic age corresponding to a temperature of $T=3000K$ can be estimated to be,

$$t_{3000K} \cong \frac{(R_{3000K} - R_{pl})}{\left[\left((V_{\text{exp}})_{3000K} + (V_{\text{exp}})_{pl} \right) / 2 \right]} \cong 17987.07 \text{ Years} \quad (24)$$

This estimation is 21.13 times less than the current estimations and needs further study.

8. VELOCITY-DISTANCE RELATION

From and about the hypothetical baby Planck ball or baby universe, current galactic receding speeds can be approximated with,

$$(v_g)_0 \cong \left(\frac{(d_g)_0}{R_0} \right) (V_{\text{exp}})_0 \cong \left(\frac{(V_{\text{exp}})_0}{R_0} \right) (d_g)_0 \cong H_0 (d_g)_0 \quad (25)$$

When $(d_g)_0 \rightarrow R_0$, $(v_g)_0 \cong H_0 R_0$. This can be compared with currently believed Hubble's law for the current expanding universe.

9. DISCUSSION

9.1 Cosmological Constant Problem

With reference to proposed concepts, ratio of the Planck scale critical density to the current critical density is,

$$\left(\frac{3H_{pl}^2 c^2}{8\pi G} \right) \div \left(\frac{3H_0^2 c^2}{8\pi G} \right) \cong \left(\frac{H_{pl}}{H_0} \right)^2 \cong 6.685 \times 10^{121} \quad (26)$$

We wish to appeal that, this idea can be considered as a characteristic tool for constructing a model of 'quantum gravity' with cosmic evolution.

9.2 Horizon Problem

The 'horizon problem' is a problem with the standard cosmological model of the Big Bang. It points out that different regions of the universe have not 'contacted' each other because of the large distances between them, but nevertheless they have the same temperature and other physical properties. If one is willing to consider the concept of 'matter causes the space-time to curve', 'horizon problem' can be understood. According to hot big bang model, during its evolution, as the universe is expanding, thermal radiation temperature decreases and matter content increases. As matter content increases, based on Mach's principle, at any stage of evolution, it is possible to have an increasing radius of curvature,

$R_t \equiv \frac{G}{c^2} [(M_{OM})_t + (M_{DM})_t]$. For the current case, $R_0 \equiv \frac{G}{c^2} [(M_{OM})_0 + (M_{DM})_0] \cong 10.40 \text{ Gpc}$ and there is no scope for 'causal disconnection' of distant visible matter.

9.3 Cosmic Inflation

Basically, 'inflation' (Guth 1981, Linde 1982) is a period of super cooled cosmic expansion, where temperature seems to be dropped by a factor of 100,000 and is mostly model dependent. Mainstream cosmologists believe that just after the hot big bang, 'inflation' seems to be started at 10^{-37} seconds and ended by 10^{-32} seconds. During this time span, cosmic temperature seems to be dropped from 10^{27} K to 10^{22} K. Since that time, expansion initially decelerated and then, after about 6 billion years, began very slowly to accelerate. Most puzzling point to be noted is that, the detailed mechanism responsible for inflation is still unknown. Many cosmologists proposed different starting mechanisms for initiating and fine tuning the believed 'inflation'. Since its introduction by Alan Guth in 1980, the inflationary paradigm has become widely accepted. Currently, a great debate is going on its existence (Steinhardt 2011, Ijjas et al 2014). In this proposed model, when $H_t \approx \frac{1}{t} \approx 10^{37} \text{ sec}^{-1}$, estimated ($\gamma_t \cong 15.43$, $T_t \cong 1.73 \times 10^{28} \text{ K}$). Similarly, when $H_t \approx \frac{1}{t} \approx 10^{32} \text{ sec}^{-1}$, estimated ($\gamma_t \cong 26.946$, $T_t \cong 4.14 \times 10^{25} \text{ K}$). Thus, for the currently believed inflationary time span of (10^{-37} to 10^{-32}) sec, estimated ratio of initial and final temperatures seems to be around 400 and needs further study with respect to the believed ratio of 100000. We are working in this direction.

9.4 Cosmic Acceleration and Expansion Velocity

Before 1990s, most of the cosmologists thought that the cosmic expansion would be decelerating due to the gravitational attraction of the matter in the universe. In between 1995 and 1998, as space technology was advancing, based on the available supernovae redshift data, cosmologists noticed that, rather than slowing down, universe was expanding with increasing rate (Riess et al. 1998, Perlmutter et al. 1999). Even though it was a shocking news for most of the science community, confirmatory evidence has been found in baryon acoustic oscillations and other cosmological observations (Weinberg DH. et al, 2013). Considering the updated supernovae redshift data, in 2016, cosmologists noticed that, universe is coasting at constant speed rather than acceleration (Neilsen et al, 2016). In this context, now a days, a great debate has been initiated among various groups of cosmologists on 'cosmic acceleration' (Zhengxiang Li et al, 2011; D. Rubin and B. Hayden, 2016; Tutusaus et al, 2018; Lawrence H. Dam et al, 2017; Christos G Tsagas, 2011)

Now a days, main stream cosmologists are seriously working on 'eternal light speed expansion' (John 2016, Jun-Jie Wei, et al 2015, Melia 2012, Melia et al 2016). In this context, in our earlier published papers, based on ordinary matter density and Hubble's law, we come across different magnitudes of cosmic expansion velocities ranging from $2c$ to $12c$. We would like to suggest that, by considering a decreasing trend of ordinary matter and dark matter density, starting from the Planck scale, it is possible to get an expression for cosmic expansion velocity comparable to speed of light. It can be expressed as follows.

$$\frac{(V_{\text{exp}})_t}{c} \cong \sqrt{\frac{2}{[(\Omega_{OM})_t + (\Omega_{DM})_t]}} \quad (27)$$


Based on this expression, for the Planck scale, $(V_{\text{exp}})_{pl} \cong c$ and for the current scale, $(V_{\text{exp}})_0 \cong 2.427c$. Interesting point to be noted is that, after 20 billion years of cosmic expansion, increment in expansion velocity seems to be only $[(V_{\text{exp}})_0 - (V_{\text{exp}})_{pl}] \cong 1.427c$. See the following picture-3.

From figure-3, it is very clear that, right from the beginning of cosmic evolution, cosmic expansion velocity seems have an increasing trend. To some extent, this can be compared with currently believed cosmic acceleration concept.

Interesting point to be noted is that, expansion velocity seems to depend on $\sqrt{\frac{2}{[(\Omega_{OM})_t + (\Omega_{DM})_t]}}$. In near future, if decrease in $[(\Omega_{OM})_t + (\Omega_{DM})_t]$ is found to be significant, one can expect 'acceleration' and if decrease in

$\left[(\Omega_{OM})_t + (\Omega_{DM})_t \right]$ is found to be insignificant, one can expect cosmic ‘constant rate of expansion’. It is for further study.

Figure 3: Increasing trend of cosmic expansion velocity



9.5 Cosmic Rotation and Mysterious Dark Energy

With reference to our estimates of current cosmic mass of 4.3352×10^{53} kg and current radius of 3.207×10^{26} m and considering the equality of current Hubble parameter and current angular velocity as a peculiar case, (Serkan Zorba 2012, Sivaram et al 2012), current cosmic rotational kinetic energy can be estimated to be 0.667 times the current critical energy. It can be understood as:

$$(KE_{rot})_0 \cong \frac{1}{3} M_0 R_0^2 \omega_0^2 \cong \frac{1}{3} M_0 R_0^2 H_0^2 \cong 7.95244 \times 10^{70} \text{ J} \quad (28)$$

$$\left\{ \begin{aligned} & \left[(KE_{rot})_0 / \frac{4\pi}{3} R_0^3 \right] \cong 5.756 \times 10^{-10} \text{ Jm}^{-3} \\ & \text{and } \left[(KE_{rot})_0 / \frac{4\pi}{3} R_0^3 \right] \div \left[\frac{3H_0^2 c^2}{8\pi G} \right] \cong \frac{2}{3} \end{aligned} \right\} \quad (29)$$

Whether it is speculative or real, shall be decided with further study and observations (Korotky V.A and Obukhov 1995, Li-Xin Li 1998, Obukhov 2000, Saulo Carneiro 2002, Godlowski W and Szydlowski Marek 2005, Godlowski 2011, Longo 2011, Rong-Gen Cai et al 2012, Chechin 2016, 2017). We would like to suggest that, currently believed (unseen, unproven and unidentified) mysterious dark energy can be identified with current cosmic rotational kinetic energy. Point to be noted is that, dark energy can be considered as a characteristic energy output pertaining to cosmic rotational phenomenon and in no way it seems responsible for cosmic acceleration or expansion velocity.

10. CONCLUSIONS

As current observational status of ‘quantum cosmology’ is very poor (Bojowald 2015), we would like to suggest that:

- 1) Universe can be considered as a quantum gravitational object.
- 2) Planck mass can be considered as a characteristic representation of the mass of baby universe.
- 3) Mach’s relation can be given some consideration in developing a workable model of cosmology.

- 4) Proposed temperature redshift $(Z_T + 1)$ assumed to be associated with $\sqrt{1 + \ln\left(\frac{H_{pl}}{H_t}\right)}$ can be recommended for further research. For the current case, $(Z_{2.722K} + 1) \cong 11.885$ and $Z_{2.722K} \cong 10.885$.
- 5) Even though it is very interesting, it is also true that the proposed relation $\gamma_t \equiv \left[1 + \ln\left(\frac{H_{pl}}{H_t}\right)\right] \cong \sqrt{\frac{3H_t^2 c^2}{8\pi G(aT_t^4)}}$ is new and peculiar. We are confident that, its role, scope and validity can be verified with further study.
- 6) Estimated current cosmic Planck ball radius is about 10.40 Gpc and can be compared with current observations of 14 Gpc. Current cosmic sphere seems to constitute around 14 Hubble spheres and needs further study with respect to the Bayesian model average estimate of >251 Hubble spheres proposed by Vardanyan et al (2011).
- 7) With reference to ordinary matter density ratio, dark matter density ratio and the proposed relation $\left(\frac{3H_t^2 c^2}{8\pi G(aT_t^4)}\right)^{\frac{1}{4}} \cong \sqrt{\gamma_t} \cong \sqrt{1 + \ln\left(\frac{H_{pl}}{H_t}\right)} \cong ((Z_T)_t + 1)$, we are working on understanding the scope, physical significance and validity of the proposed ad-hoc coefficients, $\beta_t \equiv \left(\frac{1 + \sqrt{\gamma_t}}{2}\right) \cong \left(1 + \frac{(Z_T)_t}{2}\right)$ and $\beta_t^2 \equiv \left(\frac{1 + \sqrt{\gamma_t}}{2}\right)^2 \cong \left(1 + \frac{(Z_T)_t}{2}\right)^2$ where $\frac{(Z_T)_t}{2} \approx \ln(\gamma_t)$ and $\beta_t \approx 1 + \ln(\gamma_t)$.
- 8) Independent of galactic redshift data, we are working on finding alternative tools for understanding the cosmic expansion rate. In future, with advanced science, engineering and technology, by considering $\frac{d(T_0)}{dt}$ or $\frac{d(H_0)}{dt}$ or $\frac{d(\Omega_{OM} + \Omega_{DM})_0}{dt}$, absolute cosmic rate of expansion can be estimated.

ACKNOWLEDGEMENTS

It is great pleasure to express our deep gratitude to Dr.E.Terry Tatum for his valuable technical discussions and kind encouragement. Author Seshavatharam is indebted to professors brahmashri M. Nagaphani Sarma, Chairman, shri K.V. Krishna Murthy, founder Chairman, Institute of Scientific Research in Vedas (I-SERVE), Hyderabad, India and Shri K.V.R.S. Murthy, former scientist IICT (CSIR), Govt. of India, Director, Research and Development, I-SERVE, for their valuable guidance and great support in developing this subject.

REFERENCES

- Abhas Mitra, 2014, MNRAS **442**, 382–387
 Abhas Mitra, 2011, Journal of Modern Physics, 2, 1436-1442
 Annala A, 2012, Mon. Not. R. Astron. Soc. 423, 1973–1977
 Arbab. A.I, 2001, Spacetime & Substance, Vol. 2, No. 2 (7), pp. 55–62
 Arbab. A.I, 2004, Gen.Rel.Grav.36:2465-2479.
 Balakrishna S. Haridasu et al. 2017, A&A 600, L1
 Berman MS, 2008, Astrophys.SpaceSci.318:269-272, 2008
 Berman MS, 2009, Int J Theor Phys 48: 3257–3261
 Bojowald M, 2015, Rep. Prog. Phys. 78, 023901.
 Chechin L.M. Astron.Rep. 60 no.6, 535-541. (2016)
 Chechin L.M, Gravit. Cosmol. (2017) 23
 Christos G Tsagas, 2011, J. Phys.: Conf. Ser. 283 012040 (2011)
 D. Rubin and B. Hayden, 2016, The Astrophysical Journal Letters, 833:L30 (5pp).
 Deng Wang and Xin-He Meng, 2017, Phys. Rev. D 96, 103516
 Einstein A, 1918, Ann. Phys. 360, 241.
 Gamow G, 1948, Nature, 162, 680-682.
 Gamow G, 1948, Phys. Rev. 74, 505-506
 G'abor R'acz et al. 2017, Mon Not R Astron Soc Lett slx026.
 Godlowski W and Szydlowski Marek. Gen.Rel.Grav. 37 (2005) 907-936
 Godlowski W. International Journal of Modern Physics D 20 1643 (2011)
 Gong-Bo Zhao et al, 2017, Nature Astronomy, 2017; 1 (9): 627
 Guth. A.H, 1981, Phys. Rev.;D23:347.

- Ijjas A, Steinhardt P. J and Abraham Loeb. 2014, Physics Letters B 736, 142-146.
- Joel Smoller et al, 2017, Proceedings of the Royal Society A: Mathematical, Physical and Engineering Science
- John M.V. 2016, Mon.Not.Roy.Astron.Soc. 000: 1-12.
- Jun-Jie Wei, et al, 2015, The Astronomical Journal, 149:102 (11pp).
- Jungjai Lee and Hyun Seok Yang., 2017. Preprint (arXiv:1709.04914)
- Korotky V.A and Obukhov Yu.N, JETP 81 (6) 1031-1035 (1995)
- Lasha Berezhiani et al, 2017, Phys. Rev. D 95, 123530.
- Lawrence H. Dam et al., 2017, Mon.Not.Roy.Astron.Soc, 472 (2017) 835-851
- Li-Xin Li. General Relativity and Gravitation (1998) 30: 497
- Linde A. 1982, Physics Letters B. 108 (6): 389–393.
- Longo M, Phys Lett B 2011; 699: 224.
- Miyazaki A, 2001, arXiv:gr-qc/0101112v1
- Melia F. 2012, Astron. J. 144.
- Melia F, et al.2016,Mon.Not.Roy.Astron.Soc. 456(4): 3422-3431.
- Neilsen J.T. et al., 2016, Scientific Reports 6: 35596.
- Obukhov Y. N. ar-Xiv:astro-ph/0008106v1 7 Aug 2000.
- Padmanabhan T., 2005, Curr.Sci.88:1057.
- Patrick Grahn et al, 2018, AIP Advances 8, 035028
- Perlmutter S et al. 1999, Astrophys.J.517:565-586
- Planck Collaboration: Planck 2015 Results. XIII. Cosmological Parameters.
- Rennan Barkana, 2018. Nature 555 71-74
- Riess AG et al. 1998, Astron.J.116:1009-1038.
- Riess AG et al. 2016, Astrophys.J. 826 no.1.
- Rong-Gen Cai and Zhong-Liang Tuo, JCAP 1202 (2012) 004
- Saulo Carneiro. General Relativity and Gravitation, Vol. 34, No. 6, 793-805, 2002
- Serkan Zorba. Mod. Phys. Lett. A, 27, 1250106 (2012)
- Seshavatharam U. V. S & Lakshminarayana S, 2017, Physical Science International Journal,14(2):1-17.
- Seshavatharam U. V. S & Lakshminarayana S, 2017, Physical Science International Journal 15(4): 1-13.
- Seshavatharam U. V. S & Lakshminarayana S, 2018, International Journal of Astronomy, Astrophysics and Space Science, 5(1), 1-13
- Seshavatharam U. V. S & Lakshminarayana. S, 2018, Prespacetime journal, Vol 9, Issue 4, Page 326-342.
- Sivaram C and Kennath Arun, The Open Astronomy Journal, 2012, 5, 7-11
- Sivaram C., 2000, Current Science, Vol. 79, No. 4, 41-420, 25.
- Steinhardt. P.J., 2011, Scientific American.;304(4):18-25.
- Tatum, E.T et al. (2015) Journal of Applied Physical Science International, 4, 18-26.
- Tutusaus I et al, 2018. arXiv:1803.06197 [astro-ph.CO]
- Vardanyan M. et al., 2011. MNRAS Lett 413, 1, L91-L95.
- Weinberg DH et al, 2013, Physics Reports 530, 87–255.
- Zhengxiang Li et al, 2011, Phys.Lett.B695:1-8.