

Review

Regulation of DNA Double-Strand Break Repair by Non-Coding RNAs

Roopa Thapar^{1*}

¹ Department of Molecular and Cellular Oncology, University of Texas M.D. Anderson Cancer Center, Houston, TX 77030, USA; rthapar@mdanderson.edu

* Correspondence: rthapar@mdanderson.edu; Tel.: +1-713-201-7875

Received: date; Accepted: date; Published: date

Abstract: DNA double-strand breaks (DSBs) are deleterious lesions that are generated in response to ionizing radiation or replication fork collapse that can lead to genomic instability and cancer. Eukaryotes have evolved two major pathways, namely homologous recombination (HR) and non-homologous end joining (NHEJ) to repair DSBs. Whereas the roles of protein-DNA interactions in HR and NHEJ have been fairly well defined, the functions of small and long non-coding RNAs and RNA-DNA hybrids in the DNA damage response is just beginning to be elucidated. This review summarizes recent discoveries on the identification of non-coding RNAs and RNA-mediated regulation of DSB repair.

Keywords: DNA repair; long non-coding RNA; microRNA; DNA damage; double-strand breaks; NHEJ; HR.

1. Introduction

One of the most remarkable advances in molecular biology during the last decade is the annotation of the transcriptome in numerous organisms and the discovery that ~75% of the genome is transcribed into noncoding RNAs, but only ~2% of the transcriptome accounts for proteins[1-3]. These noncoding RNAs are involved in a diverse array of biological functions that are only beginning to be understood. A strong connection has been established between several small and long noncoding RNAs (lncRNAs) and human disease, especially cancer. MicroRNA signatures are linked to cancer metastasis and tumor progression. Several microRNAs and lncRNAs are oncogenes or tumor suppressors that regulate the DNA damage response and genomic stability. Understanding the mechanisms by which they act opens up new opportunities for intervention in cancer.

Double-strand DNA breaks (DSBs) are the most lethal form of DNA damage in cells and are predominantly repaired either by Non-Homologous DNA End Joining (NHEJ)[4-6] or Homologous Recombination (HR) pathways[7,8]. DSBs can occur endogenously as a result of unrepaired mutations that lead to stalled replication forks, due to inhibition of DNA replication culminating in breaks during mitosis, or because of the harmful effects of ionizing radiation[9-11]. Large-scale genome rearrangements such as deletions, insertions, and translocations that are highly genotoxic, are a consequence of defects in DSB repair and are positively correlated with cancer progression[12]. Germ line mutations in proteins involved in HR can cause several diseases such as Fanconi Anemia (FA)[13,14], breast and ovarian cancers[15,16], Bloom syndrome[17,18], Werner syndrome[19], and ataxia telangiectasia/Nijmegen breakage syndrome[20,21], whereas defects in NHEJ can result in chromosomal instability, promoting tumorigenesis, radiosensitive severe combined immunodeficiency (RS-SCID)[22,23], microcephaly, and growth defects in children[24-26]. A third mechanism of repair called Microhomology-mediated end joining (MMEJ) or Alt-NHEJ is a minor

pathway of DSB repair, and is triggered in the event that HR is not capable of repairing breaks[27-30]. Pathway choice for double strand break repair is complex and is dependent on the type of DSB induced, the cell cycle, and the activity of the repair components[31-33]. Understanding the molecular mechanisms involved in repair of DSBs has broad implications for a range of human diseases such as cancer, as well as aging[8,34].

In mammals, NHEJ is the primary DSB repair pathway that is active throughout the cell cycle, although it is more prominent during the G0 and G1 phases of the cell cycle[35-38]. NHEJ is used to repair double-strand DNA breaks (DSBs) that arise due to V (D) J recombination[39,40], ionizing radiation[41], and reactive oxygen species[42], without the need for a template DNA. The first step in NHEJ involves binding of Ku70–Ku80 heterodimers to both ends of a DNA break[43-45], followed by the recruitment and activation of the catalytic subunits of DNA dependent protein kinase (DNA-PKcs)[46,47]. DNA end processing involves phosphorylation of nucleases such as Artemis, and DNA polymerases to replace damaged bases; followed by ligation of blunt DNA ends by the XLF-XRCC4-DNA ligase IV complex. Recently, a triple-strand break repair model has been proposed in which ribonucleotide incorporation at the break termini is important for the ligation step of NHEJ[48]. Due to direct joining of the DNA ends that frequently involves DNA end resection, NHEJ is an error-prone DNA repair mechanism.

Unlike NHEJ, HR occurs only in rapidly dividing cells during the S and G2/M phases of the cell cycle and requires a sister chromatid as a template for repair[35,49,50]. The early events in HR can be characterized as (1) signaling to the DSB (2) relaxing chromatin to open the site of the break for repair, and (3) resection of the 5' end of the DNA end[51-53]. The kinetics of the reaction measured *in vivo*, based on fluorescence recovery after photobleaching (FRAP) experiments after UV-based laser irradiation experiments, is rapid[54]. Within ~1 sec of the damage, the DSB is recognized by poly (ADP-ribose) polymerase 1 (PARP1), followed by recruitment of the chromatin remodeler Alc1 to the PAR chains, relaxation of chromatin allowing access of the helicase-nuclease “sensor” Mre11-Rad50-Nbs1 (MRN) complex to the site of damage[55-57]. Recruitment of MRN to the DSB site has been reported to occur within ~13 sec and appears to be via direct association with PAR chains. Single-molecule localization experiments show that the temporal dynamics and spatial distribution of MRN foci are cell type-specific, and the MRN complex is recruited after γ H2AX foci have already formed[58,59]. In addition to MRN, the phosphatidylinositol 3-kinase-like protein kinase (PIKK) ATM is activated by autophosphorylation[60]. ATM phosphorylates the DNA repair factors BRCA1, CtIP, EXO1 that act downstream in the reaction mechanism to promote resection, as well as the histone variant H2AX, which is phosphorylated at Ser139 to generate γ H2AX[50]. Deposition of γ H2AX occurs at the site of the DSB and is amplified, spreading for hundreds of kilobases from the damage site. These initial factors recruited to the sites of DNA damage form distinct DNA repair foci, the nature of which remains ambiguous. The amplification of this damage is dependent on formation of the γ H2AX-MDC1 complex that recruits the ubiquitin E3 ligase RNF8[61]. RNF8-mediated ubiquitylation of histones H2A and H2AX results in decondensation of chromatin and recruitment of RAP80, ABRA1, and BRCA1[61]. RNF8 also interacts with other chromatin remodeling complexes such as CHD4, a component of the nucleosome remodeling and deacetylase complex, NuRD[62]. After repair has occurred, chromatin is remodeled to its initial state. Resection of the 5' ends of the DSB is promoted by MRN, BRCA1, CtIP, and Exo1 nuclease and helicase activities, leaving ssDNA, which is rapidly bound by replication protein A (RPA)[50,63]. RPA is then exchanged for Rad51 in the presence of BRCA2[50]. The later steps in HR involve strand invasion to the homologous template followed by DNA synthesis. This can occur by either the double-strand break repair pathway (DSBR) that results in chromosomal crossover, or the synthesis-dependent strand-annealing (SDSA) pathway in which there is strand displacement after DNA synthesis. Precise assembly of a multitude of protein factors and macromolecular machines is required for repair to occur. Regulators such as kinases, ubiquitin ligases, and acetyl-transferases that remodel chromatin are also essential to temporally and spatially control DSB repair.

HR and NHEJ DNA repair mechanisms are conserved in evolution and occur in both eukaryotes and prokaryotes. There are several excellent reviews on the detailed mechanisms and protein factors involved in DSB repair[64-69]. Recent studies indicate that non-coding RNAs (ncRNAs), that include long ncRNAs (lncRNAs) as well as small ~21 nt RNAs such as microRNAs (miRNAs), DNA damage induced RNAs (diRNAs), and Drosha- and Dicer-dependent small RNAs (DDRNs) play important roles in the DNA damage response[70-73]. This review focuses on emerging themes in the roles of ncRNAs in DSB repair that are observed in humans and have direct cancer relevance.

2. LncRNAs

Long non-coding RNAs are >200 nt RNA Pol-II transcripts that, in general, are not translated into proteins[74,75]. It is projected that >30,000 lncRNAs are expressed in humans, greater than the number of protein-coding genes (~20,000) in the genome[76-78]. These transcripts are polyadenylated, capped at the 5' end, and spliced into mature RNAs, exported into the cytoplasm and are regulated like protein-coding mRNAs, although the mechanisms of RNA processing might differ from mRNAs[79]. Based on their proximity to protein coding genes, they have been classified as sense, antisense, pseudogenes, intronic, and intergenic (also called lincRNAs)[80,81]. Although the functions of most lncRNAs are poorly understood, in several cases such as the Xist RNA[82,83] that is required for mammalian dosage compensation and X-chromosome inactivation, they are not artifacts of pervasive transcription of "junk DNA", and play important biological roles in control of the cell cycle, in development, and have been linked to cancer progression[84,85].

The mechanisms by which lncRNAs act are varied. Some lncRNAs exert epigenetic functions by acting as scaffolds for chromatin-modifying complexes, e.g. *HOTAIR* is associated with chromatin and recruits the polycomb repressive complex 2 (PRC2) and the LSD1/CoREST/REST demethylase complexes to specific loci in the genome for gene silencing[86,87]. Chromatin associated lncRNAs such as promotor RNAs (pRNAs) are tethered to specific sites in the genome by forming RNA-DNA hybrids thereby acting in *cis*, where they recruit PARP1, the ATP-remodeling complex NoRC, and the methyltransferase Dnmt3b to silence rDNA[88,89]. LncRNAs can also directly regulate transcription by acting as transcription coregulators as in the case of the steroid receptor RNA activator *SRA* that activates transcription with nuclear receptors[90,91]. An example of a corepressor is the ncRNAs transcribed from the *CCND1* gene (cyclin D1) locus that can bind the TLS protein that in turn turns-off the histone acetyl transferase activity of p300/CBP[92]. Another role of lncRNAs is in formation of nuclear bodies or foci. Paraspeckles are nuclear bodies that trap adenosine-to-inosine edited RNAs and paraspeckle formation relies on the lncRNA *NEAT1*[93,94]. Similarly, *MALAT1* (or *NEAT2*) is a component of nuclear speckles where it retains serine/arginine (SR) splicing factors and affects splicing[95,96]. Besides their roles in the nucleus, several antisense lncRNAs can hybridize with the 3' untranslated regions (3' UTRs) of mRNAs to regulate their stability in the cytoplasm and/or interact with RNA decay factors and miRNAs. Examples of antisense transcripts are lncRNAs produced from the 3' UTR of *iNOS*[97] and half-STAU1-binding site RNAs (1/2sbsRNAs)[98]. The lncRNAs *PTENP1* and *KRAS P1*[99] sequester miRNAs and act as miRNA sponges thereby affecting RNA levels of a range of mRNAs in the cell.

3. LncRNAs and the DNA damage response

Given the large number of lncRNAs and their diverse functions, it is not surprising that several lncRNAs are induced upon DNA damage and have been proposed to play a role in double-strand break repair (Table 1). Oncogenes as well as tumor suppressor lncRNAs have been documented. The first lncRNAs (ncRNA_{CCND1}) identified as being transcribed in response to DNA damage signals (ionizing radiation), were from the *CCND1* gene (cyclin D1) locus[92,100]. The ncRNA_{CCND1} (between 200 – 330 nts and greater) exist on chromatin as RNA: DNA hybrids as well as ssRNA.

Table 1. LncRNAs involved in homologous recombination (HR) or non-homologous end joining (NHEJ)

lncRNA	Mechanism	DSB repair pathway	References
1. p53 -linked lncRNAs			
<i>(lincRNA)-p21</i>	Recruits hnRNPK to repress p21 transcription	HR	[101]
<i>PINT</i>	Interacts with PRC2 to silence transcription		[102,103]
<i>PANDA</i>	Negatively regulates apoptosis by sequestering the transcription factor NF-YA from pro-apoptotic gene site	HR	[104]
<i>DINO</i>	Interacts directly with p53 to stabilize it, inducing p53 target genes	HR	[105]
<i>LINP1</i>	May interact with Ku80/70 and DNAPKcs	NHEJ	[106]
<i>WRAP53</i>	Antisense lncRNA to p53 that regulates p53 levels	HR, NHEJ	[107]
<i>APELA</i>	Binds hnRNPL to block interaction with p53	HR	[104]
<i>MEG3</i>	Increases p53 levels	HR	
<i>LincROR</i>	Inhibits p53 translation after DNA damage	HR, NHEJ	[108,109]
<i>MALAT1</i>	Directly binds PARP1 and LIG3 to promote DNA repair; may promote p53 deacetylation via SIRT1 impairing its function	HR, NHEJ, Alt-NHEJ	[110,111]
<i>TUG-1</i>	Induced by p53 and binds PRC2 to repress cell-cycle genes	HR	[112,113]
<i>loc285194</i>	P53 target, tumor suppressor, down regulates miR-211	HR	[114]
2. p53 independent lncRNAs			
<i>DDSR1</i>	Sequesters the BRCA1-Rap80 complex via direct interactions with BRCA1	HR	[115]
<i>PCAT-1</i>	Represses BRCA2 expression in prostate cancer cells	HR	[116-118]
<i>lncRNA JADE</i>	Induced by ATM activation. Increases transcription of Jade1, a component of the HBO1 histone acetylation complex. Promotes H4 acetylation at K5, K8, K12	HR	[119]
<i>ANRIL</i>	Induced by ATM-mediated E2F1 activation. Regulates cell cycle checkpoints and apoptosis	HR	[120]
<i>BARD1 9'L</i>	Increases expression of a subset of BARD1 isoforms by sequestering miRNAs that normally destabilize BARD1 mRNAs	HR	[121]
<i>TERRA</i>	Interacts with Ku70/Ku80; facilitates Exo1 mediated DNA resection; promotes interaction of Mre11 with LSD1	HR, NHEJ	[122,123]
<i>TODRA</i>	Increases Rad51 transcription	HR	[124]
<i>MDC1-AS</i>	Upregulates the expression of the chromatin adaptor MDC1	HR	[125]
<i>Evf2</i>	Directly binds BRG1 and inhibits its ATPase dependent chromatin remodeling activity, may prevent Rad51 loading onto ssDNA via BRG1	HR	[126,127]
<i>CUPID1 and CUPID2</i>	Involved in pathway choice in switching from NHEJ to HR; promotes Rad51 recruitment to DSBs	HR	[128]

RNA immunoprecipitation and ChIP experiments established TLS (translocated in liposarcoma) as the protein that was recruited to ncRNA_{ACCND1}. RNA bound TLS results in an allosteric change that allows TLS to bind and inhibit the histone acetyltransferase complex CBP/p300, resulting in transcription repression. Since then, several damage-induced lncRNAs have been identified. The mechanisms by which lncRNAs modulate DSB repair include (i) modulating the activity of p53 either at the level of transcription, translation or posttranslational modifications (ii) by acting as tethers for recruitment of chromatin remodeling complexes to the site of damage (iii) by sequestering negative regulators of DNA repair away from the damage site as decoys (iv) by acting as scaffolds that directly interact with several DNA repair proteins such as Ku70/Ku80, BRCA1, 53BP1, Mre11, PARP1, (v) by sequestering miRNAs that regulate the stability of DNA repair proteins, thereby modulating mRNA expression levels. As noted below and in Table 1, a subset of lncRNAs are part of a p53 network i.e. either under direct p53 control or they modulate the activity or expression of p53. In this section, I highlight some of the unique lncRNAs that have clear cancer relevance. A summary of the roles of DNA damage induced lncRNAs in specific disease states and development is in Table 2.

3.1 p53 linked lncRNAs

The transcription factor p53 plays key roles in normal cell proliferation and tumor suppression and is mutated in over 50% of human cancers. In response to DNA damage, it is involved in transactivation or repression of target genes to induce either cell cycle arrest or apoptosis. Not surprisingly, several lncRNAs have been discovered that are either transcription targets of p53 or regulate the p53-dependent gene expression signature[129,130]. Notably, the intergenic lncRNA that resides ~15 Kb upstream of the p53 target *CDKN1A* (or p21), called lincRNA-p21[101,104,131] was identified as a repressor of the p53 transcriptional response, decreasing gene expression of hundreds of p53 target genes and triggering apoptosis. lincRNA-p21 is a large ~3.1 Kb transcript that is a distinct gene with its own promoter and is transcribed in the opposite orientation to *CDKN1A* in response to DNA damage. LincRNA-p21 likely functions through its interaction with




Figure 1. The antisense lncRNA *WRAP53α* (shown in blue) directly interacts with the p53 mRNA (shown in red) to stabilize it, affecting p53 protein levels. The *WRAP53β* (in green) recruits and stabilizes RNF8 at double strand breaks.

heterogeneous nuclear ribonucleoprotein-K (hnRNP-K), which is a component of repressor complexes, and is recruited to promoters of p53 target genes. In addition to its roles in the nucleus, lincRNA-p21 is also reported to regulate translation in the cytoplasm[132]. A similar mechanism is employed by the DNA damage induced lncRNA called *PANDA* (for p21 associated ncRNA DNA damage activated) which is also induced by p53 and is transcribed antisense ~5 Kb upstream of *CDKN1A*[104]. *PANDA* acts as a decoy for the nuclear transcription factor Y subunit alpha (NF-YA) to repress the pro-apoptotic genes *FAS* and *BIK*. The *APELA* lncRNA is a positive regulator of p53[104,133]. It binds to heterogeneous nuclear ribonucleoprotein-L (hnRNP-L) to inhibit the hnRNP-L-p53 interaction, thereby promoting p53-induced apoptosis. A more recent study[105] identified the lncRNA *DINO* (Damage Induced Noncoding), a ~951 bp RNA that is induced by p53 and is also located upstream of *CDKN1A*. Intriguingly, *DINO* interacts directly with p53 and stabilizes it, promoting the transactivation functions of p53. *DINO* knockout mice are also deficient in p53-dependent gene expression in response to DNA damage. Several lncRNAs have been shown to bind the repressor PRC2 that has H3K27 trimethylase activity. *PINT* is a lincRNA that is transcriptionally activated by p53 and is a positive regulator of cell proliferation and cell survival in mouse cells and a negative regulator in human cells. In both mice and humans, *PINT* targets PRC2 to specific gene loci for repression, thereby affecting cell proliferation, although the outcomes appear to be different in mice vs. humans[102,103]. *TUG1* is a lincRNA transcript that is upregulated by p53 and may act on chromatin to downregulate p53 mediated transcriptional pathways[112,113]. *TUG1* interacts with PRC2 and additional corepressor complexes that include histone methyltransferases, demethylases, and chromatin modifiers[112,113].

3.1.1. WRAP53 (WD repeat containing antisense to p53)

A fascinating example of a lncRNA that is partially antisense to *p53* and induces p53 levels upon DNA damage is *WRAP53*[134,135] (Figure 1). Antisense lncRNA transcripts of *p53* that overlap with the first *p53* exon are called *WRAP53α*[135] and those that overlap with the first intron are called *WRAP53γ*. The *WRAP53α* RNA directly interacts with the p53 mRNA via RNA-RNA interactions to affect p53 protein levels[135]. *WRAP53* is an interesting example of gene that encodes both a lncRNA and is also transcribed into the protein *WRAP53β* (also known as *WDR79* or *TCAB1*)[136-138]. *WRAP53β* has multiple functions in the cell one of which is to target RNF8 to DNA double-strand breaks. Therefore both the protein, as well as the lncRNA is involved in the DNA damage response, albeit through different mechanisms.

3.1.2. LINP1

The long non-coding RNA “lncRNA in non-homologous end joining pathway” LINP1[106,139,140] forms an RNA scaffold that, at least in cell extracts, interacts with both Ku70–Ku80 and DNA-PKcs on chromatin to promote end joining (Figure 2). LINP1 is overexpressed in triple negative breast cancer (TNBC) tumors, in the TNBC cell lines MDA-MB-231 and MDA-MB-468, and in triple-negative immortalized mammary cells MCF10A. LINP1 is not detected in ER-positive MCF7 cells. NHEJ reporter assays show that knockdown of LINP1 in TNBC cells decreases NHEJ and overexpression of LINP1 in MCF7 cells increases NHEJ activity[106]. Knockdown of LINP1 also sensitizes mice to ionizing radiation due to defects in DNA repair via the NHEJ pathway. Activation of EGFR signaling via MEK and JNK kinases up regulates LINP1 whereas tumor suppressor p53 represses LINP1 via the microRNA miR-29. Recent studies show that LINP1 levels are also increased in cervical cancer[141] as well as advanced prostate cancer[139] and are correlated with poor prognosis.

3.1.3. MALAT1 or NEAT2

Metastasis-associated lung adenocarcinoma transcript 1 or *MALAT1*[110,111] is a ~7 Kb lincRNA that is evolutionarily conserved and is associated with many cancers[142]. It was originally identified in a screen for lncRNAs in early-stage non-small cell lung cancers that were metastatic. *MALAT1* performs a wide array of functions. It co-localizes with nuclear speckles[143] and is important for pre-mRNA processing and splicing[96,144,145]. The longer

258 *MALAT1* transcript can be further processed into a short 61 nt tRNA-like fragment called
259 *MALAT1*-associated small cytoplasmic RNA (mascRNA) that regulates translation in the
260 cytoplasm[146]. *MALAT1* is one of the few lncRNAs for which structural information is available
261 for at least a portion of the RNA. The 3' end of *MALAT1* forms an expression and nuclear




Figure 2. *LINP1* is a 917 nt long RNA that promotes the assembly of NHEJ factors to enhance DNA repair in Triple Negative Breast Cancers (TNBC). *LINP1* directly associates with Ku70/Ku80 as well as DNA-PKcs to increase NHEJ activity in TNBC tumors. Activation of EGFR signaling via MEK and JNK kinases up regulates *LINP1* whereas the tumor suppressor TP53 represses *LINP1* via the microRNA miR-29.

262 retention element (ENE)+A, a triple helix element for which a 3.1 Å crystal structure is available
263 (Figure 3A). The structure shows how the triple helix sequesters the 3' end of *MALAT1* in a U-A-U
264 base triple, and protects the 3' end of the lncRNA from decay[147].

265 *Like TUG1, MALAT1* is also upregulated by p53 upon DNA damage. A proteomics screen has
266 identified numerous proteins involved in transcription, RNA processing, translation, protein
267 degradation, and metabolism as *MALAT1* interacting proteins[148]. *MALAT1* indirectly regulates
268 p53 protein transactivation function by sequestering DBC1, a partner for the deacetylase SIRT1,
269 which targets p53 for deacetylation (Figure 3B). Therefore similar to several lncRNAs discussed
270 here, p53 creates a feedback loop to regulate the expression of several lncRNAs, that in turn control
271 p53 function. Besides regulating p53 activity, *MALAT1* has also been reported to exist in a
272 complex with PARP1 and Lig3 *in vivo*, and may regulate the Alt-NHEJ pathway of double-strand
273 break repair[149].

274
275
276 **3.2 p53-independent lncRNAs**

277 Several lncRNAs (Table 1) are transcribed independent of p53, and act indirectly in DDR by
278 altering signaling pathways or via epigenetic mechanisms that repress transcription of DNA repair
279 genes. The lncRNAs *ANRIL* (antisense noncoding RNA in the INK4 locus), *MDC1-AS* and
280 *lncRNA-JADE* are induced by ATM activation. *ANRIL* is encoded from the INK4B-ARF-INK4A locus
281 on chromosome 9p21 and this locus is the most frequent copy number alteration across tumors.
282 *ANRIL* recruits the PRC complex to repress transcription of tumor suppressors INK4B, ARF, INK4A at

this locus and indirectly affects HR by altering cell cycle checkpoints. The *CUPID1* and *CUPID2* lncRNAs are transcribed in hormone-positive breast cancers and regulate pathway choice *in vivo*, switching repair from the error-prone NHEJ to HR. However the exact mechanism is unclear. The *TODRA* RNA (transcribed in opposite direction of *RAD51*) increases the expression of *RAD51* in an E2F1-dependent manner and hence facilitates HR.




Figure 3. (A) Structure of the 3' end of *MALAT1* reveals a triple helix with the 3' end sequestered in a U-A-U base triple, protecting the 3' end from nucleolytic degradation (B) *MALAT1* sequesters the negative regulator DBC1, releasing the deacetylase SIRT1 to activate p53 for transactivation.

3.2.1 DNA damage-sensitive RNA1 (*DDSR1*)

The lincRNA *DDSR1* was identified in microarray screens of RNA isolated from hTert-immortalized human skin fibroblasts treated with the DNA damaging agents neocarzinostatin, camptothecin, or etoposide[115]. *DDSR1* was identified as a 1616 nt inter-genic transcript that was upregulated by ~2.5 fold in all samples treated with the DNA damaging drugs. *DDSR1* expression is induced by ATM and under control of NF- κ B, however p53 is not required for expression. Although *DDSR1* expression is not under p53 control, *DDSR1* was found to regulate the expression of p53 target genes and *DDSR1* knockdown significantly upregulated p53 targeted mRNAs, particularly those involved in cell proliferation and cell survival. Intriguingly, *DDSR1* knockdown impaired HR by ~50% and *DDSR1* knockdown cells were also more sensitive to the PARP1 inhibitor Olaparib. RNA pull-down experiments identified heterogeneous nuclear ribonucleoprotein U-like 1 (hnRNPUL1) as an RNA binding protein that specifically associated with *DDSR1*, is recruited to sites of double-strand breaks and is important for end resection during HR[150]. *In vivo* imaging experiments show that depletion of *DDSR1* or hnRNPUL1 leads to an

increase in BRCA1-RAP80 at DSBs. This leads to a model in which the *DDSR1*-hnRNPUL1 complex regulates recruitment of BRCA1-RAP80 to sites of DNA breaks.

Table 2. Relevance of lncRNAs to development, disease, and inborn errors of metabolism

lncRNA	Role in Disease and Development	References
1. p53 -linked lncRNAs		
<i>(lincRNA)-p21</i>	Type 2 diabetes, multiple cancer types	[151,152]
<i>PINT</i>	Breast cancer, pancreatic cancer, neuronal development, acute myocardial infarction	[153-155]
<i>PANDA</i>	Pancreatic cancer, Type 2 diabetes, osteosarcoma	[151,156,157]
<i>DINO</i>	multiple sclerosis	[158]
<i>LINP1</i>	Triple negative breast cancer, cervical cancer, prostate cancer	[106,139-141]
<i>WRAP53</i>	Unknown	
<i>APELA</i>	Ovarian cancer	[159]
<i>MEG3</i>	Huntington’s disease, gliomas	
<i>LincROR</i>	Non-small-cell lung cancer	[160]
<i>MALAT1</i>	multiple sclerosis	[158]
<i>TUG-1</i>	multiple sclerosis	[158]
<i>loc285194</i>	Osteosarcoma, ischemic heart failure	[161-163]
2. p53 independent lncRNAs		
<i>DDSR1</i>	Unknown	
<i>PCAT-1</i>	Multiple cancers including prostate cancer, bladder cancer, gastric cancer	[116-118,164,165]
<i>lncRNA JADE</i>	Unknown	
<i>ANRIL</i>	Coronary artery disease, COPD, multiple cancer types Type 2 diabetes, multiple sclerosis	[151,156,166,167]
<i>BARD1 9’L</i>	Multiple cancer types	[168]
<i>TERRA</i>	Alternative lengthening of telomeres via homology directed repair (ALT) cancers	[169]
<i>TODRA</i>	Epithelial ovarian cancer	[124]
<i>MDC1-AS</i>	Bladder cancer, gliomas, gastric cancer	[125,170,171]
<i>Evf2</i>	Important for embryonic neuronal development in mice	[172,173]
<i>CUPID1 and CUPID2</i>	Breast cancer	[128]

3.2.2 Prostate cancer associated transcript 1 (*PCAT-1*)

PCAT-1 is the first lincRNA identified that participates in DSB repair in prostate cancer[118], and extends the “BRCA-ness” paradigm to include lncRNAs, in addition to mutations in DNA repair genes. Increased *PCAT-1* levels downregulate *BRCA2* and impairs HR, resulting in an increase in γ -H2AX foci formation[116]. *PCAT-1* is being used as a biomarker for a predictive response to treatment with PARP1 inhibitors due to synthetic lethality. *PCAT-1* levels are inversely correlated with *RAD51* foci formation when prostate cancer cells are treated with the PARP1 inhibitors Olaparib or ABT-888. The mechanism by which *PCAT-1* down regulates *BRCA2* is unique compared to other lincRNAs that act predominantly via epigenetic pathways. *PCAT-1* is a predominantly cytoplasmic lincRNA and the first 250 nt from the 5’ end of the *PCAT-1* RNA were sufficient to downregulate the 3’ untranslated region of *BRCA2* mRNA via posttranscriptional mechanisms, likely at the level of RNA stability, although the exact mechanism is unknown. *PCAT-1*

can also regulate c-Myc levels via a microRNA sequestering mechanism[117]. It will be interesting to see whether a similar mechanism of regulation applies to BRCA2 mRNA.

3.2.3 Telomeric repeat-containing RNAs (TERRA)

Telomeres are structures assembled at the ends of chromosomes that protect the chromosome ends from being recognized as double-strand breaks and hence inhibit activation of the DNA damage response[174]. Telomeres have a unique “end problem” in that activation of DDR could either result in end degradation or trigger incorrect recombination events. Telomere ends also need to be replicated, and progressive shortening of telomeres is correlated with aging. TERRA RNAs are unique class of lncRNAs that are transcribed from the ends of chromosomes and consist of the G-rich telomeric repeats (TTAGGG)_n in mammalian cells. TERRA RNAs form stacked G-quadruplex structures[175] that can interact with a number of proteins that are part of the shelterin complex for telomere capping, as well as DNA repair factors[123,176,177]. TERRA lncRNAs regulate the activity of telomerase, form heterochromatin at chromosome ends, and are also involved in capping of chromosome ends to protect telomere genomic integrity. During telomere replication, the ssDNA is bound by RPA, which triggers the ATR checkpoint for repair. The shelterin component POT1 is believed to compete with RPA and inhibit the ATR mediated checkpoint[178]. TERRA RNAs play a role in switching RPA for POT1. In addition, upon depletion of the shelterin protein, telomeric repeat factor 2, TERRA RNA levels increase and TERRA interacts with the histone demethylase LSD1[122]. This interaction stimulates the exonuclease activity of Mre11 to trim the 3' G overhangs of uncapped telomeres and form heterochromatin at telomeric ends[122,177]. Although TERRA RNAs are a very specialized lncRNA family, structure/function studies of TERRA interactions with chromatin modifiers and DNA repair proteins provides important insights into lncRNA function in DSB repair.

4. Small ncRNAs

In mammals, two classes of small non-coding RNAs have been identified as major players in the DNA damage response. The first class consists of microRNAs (miRNAs) which are small, phylogenetically conserved ~22 nt RNAs that negatively regulate the translation and stability of mRNAs via their association with Argonaute (Ago) proteins. miRNAs associate with 3' untranslated regions (3'UTRs) of mRNAs via RNA-RNA interactions to trigger translational repression or mRNA decay. The second class consisting of ~21 nt small RNAs has recently been identified in response to DNA damage, and are produced from sequences around the DSB. These ncRNAs are called DSB-induced small RNAs or diRNAs in plants and Drosha- and Dicer-dependent small RNAs (DDRNs) in mammals. Although the same enzymes that are involved in the miRNA pathway generate DDRNs and diRNAs, their biogenesis and mechanism is distinct from miRNAs, and is discussed in greater detail below.

4.1. miRNAs involved in the DNA damage response

miRNAs are generated from precursor RNA Pol II transcripts that can either be intergenic or intragenic in origin[179,180]. Intergenic miRNAs have their own promoters and are transcribed into pri-miRNA precursors that are capped, and polyadenylated. More than 50% of miRNAs can be transcribed from intergenic regions as multicistronic transcripts[181]. Intragenic miRNAs are predominantly transcribed from introns of a host gene and released during splicing of the host gene. The pri-miRNAs have stem-loop structures that harbor the mature miRNA. The mature miRNA is generated by endonucleolytic cleavage by the nuclear Drosha/DGCR8 heterodimer to release a ~70 nt pre-miRNA hairpin with a 2 nt 3' overhang. Exportin-5 (XPO5) and Ran-GTP facilitate nuclear export of the pre-miRNA hairpin by recognizing the 3' overhang. In the cytoplasm, the pre-miRNA hairpin is cleaved by the RNase III enzyme Dicer to generate the mature ~22 nt miRNA duplex. These mature miRNA duplexes associate with Ago proteins to form the RNA-induced silencing complex (RISC). Assembly of RISC requires degradation of one of the miRNA strands (passenger strand) while retaining the guide strand that ultimately base pairs with the mRNA 3' UTR. The

loading of the Ago-miRNA complex onto the mRNA recruits the mRNA degradation machinery that then targets the mRNA for either translational repression or decay (Figure 4A).




Figure 4. (A) pathway of miRNA processing to regulate mRNA decay. Pri-miRNA precursors are transcribed and processed before assembling into a macromolecular complex containing the Drosha/DGCR8 heterodimer to release a ~70 nt pre-miRNA hairpin in the nucleus. The pre-miRNA hairpin is exported to the cytoplasm by XPO5 and Ran-GTP. In the cytoplasm, the pre-miRNA hairpin is cleaved by Dicer to generate the mature ~22 nt miRNA duplex that associates with Ago2 leading to the formation of the RNA-induced silencing complex (RISC). Activation of mRNA decay results in base-pairing of the guide strand to the mRNA 3' UTR and recruitment of mRNA decay factors such as GW182. (B) Activation of the DNA damage response by DDRNAs also involves miRNA processing factors Drosha and Dicer that are required to generate small ~20-35 nt RNAs from the sequences around the site of damage. Formation of DNA damage foci and recruitment of DNA damage factors requires DDRNAs, Drosha, and Dicer.

The miRNA expression profile or “signatures” are predictors of cancer prognosis and are cancer hallmarks[182,183]. miRNA dependent pathways can be regulated at multiple levels upon DNA damage[181,184]. Similar to lncRNAs, the transcription of several miRNA genes, such as the miR-34 family in under the control of p53 and is altered upon DNA-damage[185]. Several miRNAs have been identified that directly regulate the abundance of DSB repair proteins at the posttranscriptional level (Table 3). The up- or down-regulation of miRNAs can affect protein stability of sensors of DNA damage such as γ H2AX[186,187], decreasing γ H2AX containing damage foci. The kinase ATM, a key regulator of the DNA damage response, is also under the control of several miRNAs (Table 3). Upregulation of miR-18a decreases the DNA damage response due to decreased ATM levels[188] whereas increased levels of miR-421, miR-100 and miR-101 leads to

increased radiosensitivity[188-190]. The expression of DNA repair effectors is also under microRNA regulation (Table 3). In particular, the tumor suppressor BRCA1 and ssDNA binding protein RAD51, which are important for strand invasion in HR are tightly regulated by a number of miRNAs. MicroRNAs also control protein levels of cell cycle checkpoint factors and apoptosis regulators as has been reviewed [181,184]. Intriguingly, miRNAs can be secreted into body fluids such plasma and urine via exosomes and may be important mediators of DNA damage from radiation-targeted cells to abscopal normal cells, leading to the bystander effect[191]. Exosomes have been reported to contain as many as 764 miRNAs[192,193]. The miR-1246 is an example of a radiation-induced miRNA that is packaged into exosomes and is delivered to non-irradiated cells to decrease NHEJ efficiency by targeting the 3' UTR of LIG4[194]. These diverse roles of miRNAs in the DNA damage response and cancer have made them attractive targets for cancer therapy.

Table 3. miRNAs that directly target genes involved in homologous recombination (HR) or non-homologous end joining (NHEJ)

miRNAs	Target gene	DSB repair pathway	References
miR-27a, miR-421, miR-101, miR-100, miR-18a, miR-181	ATM	HR	[188-190,195,196]
miR-101	DNA-PKcs	NHEJ	[190]
miR-124, miR-622	Ku70	NHEJ	[197,198]
miR-623, miR-526b, miR-622	Ku80	NHEJ	[198-200]
miR-1246	LIG4	NHEJ	[194]
miR-138, miR-24	γ H2AX	HR, NHEJ	[186,187]
miR-182-5p, miR-146a, miR-146b-5p, mir-1255b, miR-148b, miR-193b, miR-99, miR-28, let-7	BRCA1	HR	[188,201-205]
miR-19a, miR-19b, miR-1255b, miR-148b, miR-193b, let-7	BRCA2	HR	[204-206]
miR-96, miR-193a-3p, miR-506, miR-155, miR-1255b, miR-148b, miR-193b, miR-222, miR-107	RAD51	HR	[204,207-212]
let-7	FANCD2	HR	[205]
miR-210	RAD52	HR	[213]
miR-335	CTIP	HR	[214]

4.2. Drosha- and Dicer-dependent small RNAs (DDRNs)

Recent studies have shown that small ~21 nt RNAs are induced from the sequences around the site of damage in response to ionizing radiation. These RNAs are called diRNAs in *Arabidopsis thaliana*[71], qiRNAs in *Neurospora crassa*[215], endo-siRNAs in *Drosophila melanogaster*[216], and DDRNs in human cells[217,218]. These RNAs are generated in a Dicer-dependent manner in all organisms and are distinct from miRNAs (Figure 4B). In mammals, down-regulation or inactivation of Drosha and Dicer impairs formation of DNA damage foci containing phosphor-ATM, MDC1, and 53BP1 in response to ionizing radiation[217,218]. Knockdown of Dicer and Drosha also results in a loss of the G1/S and G2/M checkpoints. Formation of DNA damage foci is RNA-dependent and removal of RNaseA and inhibiting transcription reduces foci formation. Using a chromosomally integrated reporter system and deep sequencing, recent studies show that foci formation requires small RNAs (20-35 nt) called DDRNs that are generated by Drosha and Dicer from precursor RNAs at the damage site. The precise mechanism by which DDRNs act is unclear, and it has been proposed that they act later in the DNA damage response and may function

to recruit factors such as MDC1 and 53BP1 during HR[217]. A more recent study[219] shows that Drosha has an early role in DSB repair and is required for both HR and NHEJ. Drosha facilitates formation of RNA-DNA hybrids before resection can occur, and may interact directly with BRCA1. However, no DDRNAs were observed in this study[219] that examined endogenous break loci. However, all studies are in agreement for the role of RNA at DNA damage breaks and the requirement for miRNA processing enzymes such as Dicer and Drosha for efficient repair.

5. Concluding Remarks

Small RNA and lncRNAs play intricate roles in the DNA damage response, although the mechanisms by which they act remain unclear. Understanding their modes of regulation in DDR provides new opportunities in cancer therapy. lncRNAs can act as guides, scaffolds, decoys in DNA repair. They may also be important for spatial regulation of DNA repair complexes by formation of DNA repair foci. Use of the latest deep sequencing technologies, imaging tools, combined with molecular and structural information on RNA and RNA-protein complexes is needed to understand non-coding RNA function in DNA repair.

Acknowledgments: I would like to acknowledge support from the M.D. Anderson Cancer Center Knowledge GAP Award. I gratefully acknowledge Dr. John A. Tainer for sponsoring our ongoing research efforts (NIH R35CA22043) to understand the roles of lncRNAs in the DNA damage response.

Conflicts of Interest: I declare no conflict of interest.

References

1. Lander, E.S. Initial impact of the sequencing of the human genome. *Nature* **2011**, *470*, 187-197, doi:10.1038/nature09792.
2. Djebali, S.; Davis, C.A.; Merkel, A.; Dobin, A.; Lassmann, T.; Mortazavi, A.; Tanzer, A.; Lagarde, J.; Lin, W.; Schlesinger, F., et al. Landscape of transcription in human cells. *Nature* **2012**, *489*, 101-108, doi:10.1038/nature11233.
3. International Human Genome Sequencing, C. Finishing the euchromatic sequence of the human genome. *Nature* **2004**, *431*, 931-945, doi:10.1038/nature03001.
4. Helleday, T.; Petermann, E.; Lundin, C.; Hodgson, B.; Sharma, R.A. DNA repair pathways as targets for cancer therapy. *Nat Rev Cancer* **2008**, *8*, 193-204, doi:10.1038/nrc2342.
5. Lieber, M.R. The mechanism of double-strand DNA break repair by the nonhomologous DNA end-joining pathway. *Annu Rev Biochem* **2010**, *79*, 181-211, doi:10.1146/annurev.biochem.052308.093131.
6. Lord, C.J.; Ashworth, A. The DNA damage response and cancer therapy. *Nature* **2012**, *481*, 287-294, doi:10.1038/nature10760.
7. Ciccia, A.; Elledge, S.J. The DNA damage response: making it safe to play with knives. *Mol Cell* **2010**, *40*, 179-204, doi:10.1016/j.molcel.2010.09.019.
8. Jackson, S.P.; Bartek, J. The DNA-damage response in human biology and disease. *Nature* **2009**, *461*, 1071-1078, doi:10.1038/nature08467.
9. Helleday, T.; Eshtad, S.; Nik-Zainal, S. Mechanisms underlying mutational signatures in human cancers. *Nat Rev Genet* **2014**, *15*, 585-598, doi:10.1038/nrg3729.

- 508 10. Chatterjee, N.; Walker, G.C. Mechanisms of DNA damage, repair, and mutagenesis. *Environ*
509 *Mol Mutagen* **2017**, *58*, 235-263, doi:10.1002/em.22087.
- 510 11. Gaillard, H.; Garcia-Muse, T.; Aguilera, A. Replication stress and cancer. *Nat Rev Cancer*
511 **2015**, *15*, 276-289, doi:10.1038/nrc3916.
- 512 12. Aparicio, T.; Baer, R.; Gautier, J. DNA double-strand break repair pathway choice and
513 cancer. *DNA Repair (Amst)* **2014**, *19*, 169-175, doi:10.1016/j.dnarep.2014.03.014.
- 514 13. Walden, H.; Deans, A.J. The Fanconi anemia DNA repair pathway: structural and functional
515 insights into a complex disorder. *Annu Rev Biophys* **2014**, *43*, 257-278,
516 doi:10.1146/annurev-biophys-051013-022737.
- 517 14. Nalepa, G.; Clapp, D.W. Fanconi anaemia and cancer: an intricate relationship. *Nat Rev*
518 *Cancer* **2018**, *18*, 168-185, doi:10.1038/nrc.2017.116.
- 519 15. Petrucelli, N.; Daly, M.B.; Feldman, G.L. Hereditary breast and ovarian cancer due to
520 mutations in BRCA1 and BRCA2. *Genet Med* **2010**, *12*, 245-259,
521 doi:10.1097/GIM.0b013e3181d38f2f.
- 522 16. Levy-Lahad, E.; Friedman, E. Cancer risks among BRCA1 and BRCA2 mutation carriers. *Br J*
523 *Cancer* **2007**, *96*, 11-15, doi:10.1038/sj.bjc.6603535.
- 524 17. Bischof, O.; Kim, S.H.; Irving, J.; Beresten, S.; Ellis, N.A.; Campisi, J. Regulation and
525 localization of the Bloom syndrome protein in response to DNA damage. *J Cell Biol* **2001**, *153*,
526 367-380.
- 527 18. Deans, A.J.; West, S.C. FANCM connects the genome instability disorders Bloom's
528 Syndrome and Fanconi Anemia. *Mol Cell* **2009**, *36*, 943-953, doi:10.1016/j.molcel.2009.12.006.
- 529 19. Bernstein, K.A.; Gangloff, S.; Rothstein, R. The RecQ DNA helicases in DNA repair. *Annu*
530 *Rev Genet* **2010**, *44*, 393-417, doi:10.1146/annurev-genet-102209-163602.
- 531 20. Moynahan, M.E.; Jasin, M. Mitotic homologous recombination maintains genomic stability
532 and suppresses tumorigenesis. *Nat Rev Mol Cell Biol* **2010**, *11*, 196-207, doi:10.1038/nrm2851.
- 533 21. Rothblum-Oviatt, C.; Wright, J.; Lefton-Greif, M.A.; McGrath-Morrow, S.A.; Crawford, T.O.;
534 Lederman, H.M. Ataxia telangiectasia: a review. *Orphanet J Rare Dis* **2016**, *11*, 159,
535 doi:10.1186/s13023-016-0543-7.
- 536 22. van der Burg, M.; Ijspeert, H.; Verkaik, N.S.; Turul, T.; Wiegant, W.W.; Morotomi-Yano, K.;
537 Mari, P.O.; Tezcan, I.; Chen, D.J.; Zdzienicka, M.Z., et al. A DNA-PKcs mutation in a
538 radiosensitive T-B- SCID patient inhibits Artemis activation and nonhomologous
539 end-joining. *J Clin Invest* **2009**, *119*, 91-98, doi:10.1172/JCI37141.
- 540 23. Moshous, D.; Callebaut, I.; de Chasseval, R.; Corneo, B.; Cavazzana-Calvo, M.; Le Deist, F.;
541 Tezcan, I.; Sanal, O.; Bertrand, Y.; Philippe, N., et al. Artemis, a novel DNA double-strand
542 break repair/V(D)J recombination protein, is mutated in human severe combined immune
543 deficiency. *Cell* **2001**, *105*, 177-186.
- 544 24. Woodbine, L.; Gennery, A.R.; Jeggo, P.A. The clinical impact of deficiency in DNA
545 non-homologous end-joining. *DNA Repair (Amst)* **2014**, *16*, 84-96,
546 doi:10.1016/j.dnarep.2014.02.011.
- 547 25. de Bruin, C.; Mericq, V.; Andrew, S.F.; van Duyvenvoorde, H.A.; Verkaik, N.S.; Losekoot,
548 M.; Porollo, A.; Garcia, H.; Kuang, Y.; Hanson, D., et al. An XRCC4 splice mutation
549 associated with severe short stature, gonadal failure, and early-onset metabolic syndrome. *J*
550 *Clin Endocrinol Metab* **2015**, *100*, E789-798, doi:10.1210/jc.2015-1098.

- 551 26. Chistiakov, D.A.; Voronova, N.V.; Chistiakov, A.P. Ligase IV syndrome. *Eur J Med Genet*
552 **2009**, *52*, 373-378, doi:10.1016/j.ejmg.2009.05.009.
- 553 27. Higgins, G.S.; Boulton, S.J. Beyond PARP-POLtheta as an anticancer target. *Science* **2018**, *359*,
554 1217-1218, doi:10.1126/science.aar5149.
- 555 28. Deriano, L.; Roth, D.B. Modernizing the nonhomologous end-joining repertoire: alternative
556 and classical NHEJ share the stage. *Annu Rev Genet* **2013**, *47*, 433-455,
557 doi:10.1146/annurev-genet-110711-155540.
- 558 29. Bennardo, N.; Cheng, A.; Huang, N.; Stark, J.M. Alternative-NHEJ is a mechanistically
559 distinct pathway of mammalian chromosome break repair. *PLoS Genet* **2008**, *4*, e1000110,
560 doi:10.1371/journal.pgen.1000110.
- 561 30. Dueva, R.; Iliakis, G. Alternative pathways of non-homologous end joining (NHEJ) in
562 genomic instability and cancer. *Transl. Cancer. Res.* **2013**, *2*, 163-177.
- 563 31. Dutta, A.; Eckelmann, B.; Adhikari, S.; Ahmed, K.M.; Sengupta, S.; Pandey, A.; Hegde, P.M.;
564 Tsai, M.S.; Tainer, J.A.; Weinfeld, M., et al. Microhomology-mediated end joining is
565 activated in irradiated human cells due to phosphorylation-dependent formation of the
566 XRCC1 repair complex. *Nucleic Acids Res* **2017**, *45*, 2585-2599, doi:10.1093/nar/gkw1262.
- 567 32. Shibata, A.; Moiani, D.; Arvai, A.S.; Perry, J.; Harding, S.M.; Genois, M.M.; Maity, R.; van
568 Rossum-Fikkert, S.; Kertokallio, A.; Romoli, F., et al. DNA double-strand break repair
569 pathway choice is directed by distinct MRE11 nuclease activities. *Mol Cell* **2014**, *53*, 7-18,
570 doi:10.1016/j.molcel.2013.11.003.
- 571 33. Schipler, A.; Iliakis, G. DNA double-strand-break complexity levels and their possible
572 contributions to the probability for error-prone processing and repair pathway choice.
573 *Nucleic Acids Res* **2013**, *41*, 7589-7605, doi:10.1093/nar/gkt556.
- 574 34. Hoeijmakers, J.H. DNA damage, aging, and cancer. *N Engl J Med* **2009**, *361*, 1475-1485,
575 doi:10.1056/NEJMr0804615.
- 576 35. Branzei, D.; Foiani, M. Regulation of DNA repair throughout the cell cycle. *Nat Rev Mol Cell*
577 *Biol* **2008**, *9*, 297-308, doi:10.1038/nrm2351.
- 578 36. Shibata, A.; Conrad, S.; Birraux, J.; Geuting, V.; Barton, O.; Ismail, A.; Kakarougkas, A.;
579 Meek, K.; Taucher-Scholz, G.; Lobrich, M., et al. Factors determining DNA double-strand
580 break repair pathway choice in G2 phase. *EMBO J* **2011**, *30*, 1079-1092,
581 doi:10.1038/emboj.2011.27.
- 582 37. Kakarougkas, A.; Jeggo, P.A. DNA DSB repair pathway choice: an orchestrated handover
583 mechanism. *Br J Radiol* **2014**, *87*, 20130685, doi:10.1259/bjr.20130685.
- 584 38. Chapman, J.R.; Taylor, M.R.; Boulton, S.J. Playing the end game: DNA double-strand break
585 repair pathway choice. *Mol Cell* **2012**, *47*, 497-510, doi:10.1016/j.molcel.2012.07.029.
- 586 39. Bassing, C.H.; Swat, W.; Alt, F.W. The mechanism and regulation of chromosomal V(D)J
587 recombination. *Cell* **2002**, *109 Suppl*, S45-55.
- 588 40. Schatz, D.G.; Swanson, P.C. V(D)J recombination: mechanisms of initiation. *Annu Rev Genet*
589 **2011**, *45*, 167-202, doi:10.1146/annurev-genet-110410-132552.
- 590 41. Wang, C.; Lees-Miller, S.P. Detection and repair of ionizing radiation-induced DNA double
591 strand breaks: new developments in nonhomologous end joining. *Int J Radiat Oncol Biol Phys*
592 **2013**, *86*, 440-449, doi:10.1016/j.ijrobp.2013.01.011.

- 593 42. Hoeijmakers, J.H. Genome maintenance mechanisms for preventing cancer. *Nature* **2001**,
594 411, 366-374, doi:10.1038/35077232.
- 595 43. Radhakrishnan, S.K.; Jette, N.; Lees-Miller, S.P. Non-homologous end joining: emerging
596 themes and unanswered questions. *DNA Repair (Amst)* **2014**, 17, 2-8,
597 doi:10.1016/j.dnarep.2014.01.009.
- 598 44. Wang, J.L.; Duboc, C.; Wu, Q.; Ochi, T.; Liang, S.; Tsutakawa, S.E.; Lees-Miller, S.P.; Nadal,
599 M.; Tainer, J.A.; Blundell, T.L., et al. Dissection of DNA double-strand-break repair using
600 novel single-molecule forceps. *Nat Struct Mol Biol* **2018**, 25, 482-487,
601 doi:10.1038/s41594-018-0065-1.
- 602 45. Hammel, M.; Yu, Y.; Radhakrishnan, S.K.; Chokshi, C.; Tsai, M.S.; Matsumoto, Y.;
603 Kuzdovich, M.; Remesh, S.G.; Fang, S.; Tomkinson, A.E., et al. An Intrinsically Disordered
604 APLF Links Ku, DNA-PKcs, and XRCC4-DNA Ligase IV in an Extended Flexible
605 Non-homologous End Joining Complex. *J Biol Chem* **2016**, 291, 26987-27006,
606 doi:10.1074/jbc.M116.751867.
- 607 46. Sibanda, B.L.; Chirgadze, D.Y.; Ascher, D.B.; Blundell, T.L. DNA-PKcs structure suggests an
608 allosteric mechanism modulating DNA double-strand break repair. *Science* **2017**, 355,
609 520-524, doi:10.1126/science.aak9654.
- 610 47. Sibanda, B.L.; Chirgadze, D.Y.; Blundell, T.L. Crystal structure of DNA-PKcs reveals a large
611 open-ring cradle comprised of HEAT repeats. *Nature* **2010**, 463, 118-121,
612 doi:10.1038/nature08648.
- 613 48. Pryor, J.M.; Conlin, M.P.; Carvajal-Garcia, J.; Luedeman, M.E.; Luthman, A.J.; Small, G.W.;
614 Ramsden, D.A. Ribonucleotide incorporation enables repair of chromosome breaks by
615 nonhomologous end joining. *Science* **2018**, 361, 1126-1129, doi:10.1126/science.aat2477.
- 616 49. Shibata, A. Regulation of repair pathway choice at two-ended DNA double-strand breaks.
617 *Mutat Res* **2017**, 803-805, 51-55, doi:10.1016/j.mrfmmm.2017.07.011.
- 618 50. San Filippo, J.; Sung, P.; Klein, H. Mechanism of eukaryotic homologous recombination.
619 *Annu Rev Biochem* **2008**, 77, 229-257, doi:10.1146/annurev.biochem.77.061306.125255.
- 620 51. Bekker-Jensen, S.; Lukas, C.; Kitagawa, R.; Melander, F.; Kastan, M.B.; Bartek, J.; Lukas, J.
621 Spatial organization of the mammalian genome surveillance machinery in response to DNA
622 strand breaks. *J Cell Biol* **2006**, 173, 195-206, doi:10.1083/jcb.200510130.
- 623 52. Kruhlak, M.J.; Celeste, A.; Deltaille, G.; Fernandez-Capetillo, O.; Muller, W.G.; McNally, J.G.;
624 Bazett-Jones, D.P.; Nussenzweig, A. Changes in chromatin structure and mobility in living
625 cells at sites of DNA double-strand breaks. *J Cell Biol* **2006**, 172, 823-834,
626 doi:10.1083/jcb.200510015.
- 627 53. Lukas, C.; Melander, F.; Stucki, M.; Falck, J.; Bekker-Jensen, S.; Goldberg, M.; Lerenthal, Y.;
628 Jackson, S.P.; Bartek, J.; Lukas, J. Mdc1 couples DNA double-strand break recognition by
629 Nbs1 with its H2AX-dependent chromatin retention. *EMBO J* **2004**, 23, 2674-2683,
630 doi:10.1038/sj.emboj.7600269.
- 631 54. Haince, J.F.; McDonald, D.; Rodrigue, A.; Dery, U.; Masson, J.Y.; Hendzel, M.J.; Poirier, G.G.
632 PARP1-dependent kinetics of recruitment of MRE11 and NBS1 proteins to multiple DNA
633 damage sites. *J Biol Chem* **2008**, 283, 1197-1208, doi:10.1074/jbc.M706734200.
- 634 55. D'Amours, D.; Jackson, S.P. The Mre11 complex: at the crossroads of dna repair and
635 checkpoint signalling. *Nat Rev Mol Cell Biol* **2002**, 3, 317-327, doi:10.1038/nrm805.

- 636 56. Williams, G.J.; Lees-Miller, S.P.; Tainer, J.A. Mre11-Rad50-Nbs1 conformations and the
637 control of sensing, signaling, and effector responses at DNA double-strand breaks. *DNA*
638 *Repair (Amst)* **2010**, *9*, 1299-1306, doi:10.1016/j.dnarep.2010.10.001.
- 639 57. Paull, T.T. 20 Years of Mre11 Biology: No End in Sight. *Mol Cell* **2018**, *71*, 419-427,
640 doi:10.1016/j.molcel.2018.06.033.
- 641 58. Nelms, B.E.; Maser, R.S.; MacKay, J.F.; Lagally, M.G.; Petrini, J.H. In situ visualization of
642 DNA double-strand break repair in human fibroblasts. *Science* **1998**, *280*, 590-592.
- 643 59. Eryilmaz, M.; Schmitt, E.; Krufczik, M.; Theda, F.; Lee, J.H.; Cremer, C.; Bestvater, F.;
644 Schaufler, W.; Hausmann, M.; Hildenbrand, G. Localization Microscopy Analyses of MRE11
645 Clusters in 3D-Conserved Cell Nuclei of Different Cell Lines. *Cancers (Basel)* **2018**, *10*,
646 doi:10.3390/cancers10010025.
- 647 60. Uziel, T.; Lerenthal, Y.; Moyal, L.; Andegeko, Y.; Mittelman, L.; Shiloh, Y. Requirement of
648 the MRN complex for ATM activation by DNA damage. *EMBO J* **2003**, *22*, 5612-5621,
649 doi:10.1093/emboj/cdg541.
- 650 61. Mailand, N.; Bekker-Jensen, S.; Fastrup, H.; Melander, F.; Bartek, J.; Lukas, C.; Lukas, J.
651 RNF8 ubiquitylates histones at DNA double-strand breaks and promotes assembly of repair
652 proteins. *Cell* **2007**, *131*, 887-900, doi:10.1016/j.cell.2007.09.040.
- 653 62. Luijsterburg, M.S.; Acs, K.; Ackermann, L.; Wiegant, W.W.; Bekker-Jensen, S.; Larsen, D.H.;
654 Khanna, K.K.; van Attikum, H.; Mailand, N.; Dantuma, N.P. A new non-catalytic role for
655 ubiquitin ligase RNF8 in unfolding higher-order chromatin structure. *EMBO J* **2012**, *31*,
656 2511-2527, doi:10.1038/emboj.2012.104.
- 657 63. Wold, M.S. Replication protein A: a heterotrimeric, single-stranded DNA-binding protein
658 required for eukaryotic DNA metabolism. *Annu Rev Biochem* **1997**, *66*, 61-92,
659 doi:10.1146/annurev.biochem.66.1.61.
- 660 64. Seeber, A.; Hauer, M.H.; Gasser, S.M. Chromosome Dynamics in Response to DNA
661 Damage. *Annu Rev Genet* **2018**, *10.1146/annurev-genet-120417-031334*,
662 doi:10.1146/annurev-genet-120417-031334.
- 663 65. Sung, P. Introduction to the Thematic Minireview Series: DNA double-strand break repair
664 and pathway choice. *J Biol Chem* **2018**, *293*, 10500-10501, doi:10.1074/jbc.TM118.003212.
- 665 66. Her, J.; Bunting, S.F. How cells ensure correct repair of DNA double-strand breaks. *J Biol*
666 *Chem* **2018**, *293*, 10502-10511, doi:10.1074/jbc.TM118.000371.
- 667 67. Pannunzio, N.R.; Watanabe, G.; Lieber, M.R. Nonhomologous DNA end-joining for repair
668 of DNA double-strand breaks. *J Biol Chem* **2018**, *293*, 10512-10523,
669 doi:10.1074/jbc.TM117.000374.
- 670 68. Wright, W.D.; Shah, S.S.; Heyer, W.D. Homologous recombination and the repair of DNA
671 double-strand breaks. *J Biol Chem* **2018**, *293*, 10524-10535, doi:10.1074/jbc.TM118.000372.
- 672 69. Sallmyr, A.; Tomkinson, A.E. Repair of DNA double-strand breaks by mammalian
673 alternative end-joining pathways. *J Biol Chem* **2018**, *293*, 10536-10546,
674 doi:10.1074/jbc.TM117.000375.
- 675 70. Sharma, V.; Misteli, T. Non-coding RNAs in DNA damage and repair. *FEBS Lett* **2013**, *587*,
676 1832-1839, doi:10.1016/j.febslet.2013.05.006.

- 677 71. Wei, W.; Ba, Z.; Gao, M.; Wu, Y.; Ma, Y.; Amiard, S.; White, C.I.; Rendtlew Danielsen, J.M.;
678 Yang, Y.G.; Qi, Y. A role for small RNAs in DNA double-strand break repair. *Cell* **2012**, *149*,
679 101-112, doi:10.1016/j.cell.2012.03.002.
- 680 72. Yang, Y.G.; Qi, Y. RNA-directed repair of DNA double-strand breaks. *DNA Repair (Amst)*
681 **2015**, *32*, 82-85, doi:10.1016/j.dnarep.2015.04.017.
- 682 73. d'Adda di Fagagna, F. A direct role for small non-coding RNAs in DNA damage response.
683 *Trends Cell Biol* **2014**, *24*, 171-178, doi:10.1016/j.tcb.2013.09.008.
- 684 74. Wilusz, J.E.; Sunwoo, H.; Spector, D.L. Long noncoding RNAs: functional surprises from the
685 RNA world. *Genes Dev* **2009**, *23*, 1494-1504, doi:10.1101/gad.1800909.
- 686 75. Mercer, T.R.; Dinger, M.E.; Mattick, J.S. Long non-coding RNAs: insights into functions. *Nat*
687 *Rev Genet* **2009**, *10*, 155-159, doi:10.1038/nrg2521.
- 688 76. Consortium, F.; the, R.P.; Clst; Forrest, A.R.; Kawaji, H.; Rehli, M.; Baillie, J.K.; de Hoon, M.J.;
689 Haberle, V.; Lassmann, T., et al. A promoter-level mammalian expression atlas. *Nature* **2014**,
690 *507*, 462-470, doi:10.1038/nature13182.
- 691 77. Consortium, E.P. An integrated encyclopedia of DNA elements in the human genome.
692 *Nature* **2012**, *489*, 57-74, doi:10.1038/nature11247.
- 693 78. Lander, E.S.; Linton, L.M.; Birren, B.; Nusbaum, C.; Zody, M.C.; Baldwin, J.; Devon, K.;
694 Dewar, K.; Doyle, M.; FitzHugh, W., et al. Initial sequencing and analysis of the human
695 genome. *Nature* **2001**, *409*, 860-921, doi:10.1038/35057062.
- 696 79. Schlackow, M.; Nojima, T.; Gomes, T.; Dhir, A.; Carmo-Fonseca, M.; Proudfoot, N.J.
697 Distinctive Patterns of Transcription and RNA Processing for Human lincRNAs. *Mol Cell*
698 **2017**, *65*, 25-38, doi:10.1016/j.molcel.2016.11.029.
- 699 80. Andersson, R.; Gebhard, C.; Miguel-Escalada, I.; Hoof, I.; Bornholdt, J.; Boyd, M.; Chen, Y.;
700 Zhao, X.; Schmidl, C.; Suzuki, T., et al. An atlas of active enhancers across human cell types
701 and tissues. *Nature* **2014**, *507*, 455-461, doi:10.1038/nature12787.
- 702 81. Kung, J.T.; Colognori, D.; Lee, J.T. Long noncoding RNAs: past, present, and future. *Genetics*
703 **2013**, *193*, 651-669, doi:10.1534/genetics.112.146704.
- 704 82. Penny, G.D.; Kay, G.F.; Sheardown, S.A.; Rastan, S.; Brockdorff, N. Requirement for Xist in X
705 chromosome inactivation. *Nature* **1996**, *379*, 131-137, doi:10.1038/379131a0.
- 706 83. Brown, C.J.; Hendrich, B.D.; Rupert, J.L.; Lafreniere, R.G.; Xing, Y.; Lawrence, J.; Willard,
707 H.F. The human XIST gene: analysis of a 17 kb inactive X-specific RNA that contains
708 conserved repeats and is highly localized within the nucleus. *Cell* **1992**, *71*, 527-542.
- 709 84. Huarte, M. The emerging role of lncRNAs in cancer. *Nat Med* **2015**, *21*, 1253-1261,
710 doi:10.1038/nm.3981.
- 711 85. Ling, H.; Vincent, K.; Pichler, M.; Fodde, R.; Berindan-Neagoe, I.; Slack, F.J.; Calin, G.A. Junk
712 DNA and the long non-coding RNA twist in cancer genetics. *Oncogene* **2015**, *34*, 5003-5011,
713 doi:10.1038/onc.2014.456.
- 714 86. Rinn, J.L.; Kertesz, M.; Wang, J.K.; Squazzo, S.L.; Xu, X.; Brugmann, S.A.; Goodnough, L.H.;
715 Helms, J.A.; Farnham, P.J.; Segal, E., et al. Functional demarcation of active and silent
716 chromatin domains in human HOX loci by noncoding RNAs. *Cell* **2007**, *129*, 1311-1323,
717 doi:10.1016/j.cell.2007.05.022.

- 718 87. Gupta, R.A.; Shah, N.; Wang, K.C.; Kim, J.; Horlings, H.M.; Wong, D.J.; Tsai, M.C.; Hung, T.;
719 Argani, P.; Rinn, J.L., et al. Long non-coding RNA HOTAIR reprograms chromatin state to
720 promote cancer metastasis. *Nature* **2010**, *464*, 1071-1076, doi:10.1038/nature08975.
- 721 88. Guetg, C.; Lienemann, P.; Sirri, V.; Grummt, I.; Hernandez-Verdun, D.; Hottiger, M.O.;
722 Fussenegger, M.; Santoro, R. The NoRC complex mediates the heterochromatin formation
723 and stability of silent rRNA genes and centromeric repeats. *EMBO J* **2010**, *29*, 2135-2146,
724 doi:10.1038/emboj.2010.17.
- 725 89. Guetg, C.; Scheifele, F.; Rosenthal, F.; Hottiger, M.O.; Santoro, R. Inheritance of silent rDNA
726 chromatin is mediated by PARP1 via noncoding RNA. *Mol Cell* **2012**, *45*, 790-800,
727 doi:10.1016/j.molcel.2012.01.024.
- 728 90. Lanz, R.B.; Razani, B.; Goldberg, A.D.; O'Malley, B.W. Distinct RNA motifs are important for
729 coactivation of steroid hormone receptors by steroid receptor RNA activator (SRA). *Proc*
730 *Natl Acad Sci U S A* **2002**, *99*, 16081-16086, doi:10.1073/pnas.192571399.
- 731 91. Lanz, R.B.; McKenna, N.J.; Onate, S.A.; Albrecht, U.; Wong, J.; Tsai, S.Y.; Tsai, M.J.; O'Malley,
732 B.W. A steroid receptor coactivator, SRA, functions as an RNA and is present in an SRC-1
733 complex. *Cell* **1999**, *97*, 17-27.
- 734 92. Wang, X.; Arai, S.; Song, X.; Reichart, D.; Du, K.; Pascual, G.; Tempst, P.; Rosenfeld, M.G.;
735 Glass, C.K.; Kurokawa, R. Induced ncRNAs allosterically modify RNA-binding proteins in
736 cis to inhibit transcription. *Nature* **2008**, *454*, 126-130, doi:10.1038/nature06992.
- 737 93. Chen, L.L.; Carmichael, G.G. Altered nuclear retention of mRNAs containing inverted
738 repeats in human embryonic stem cells: functional role of a nuclear noncoding RNA. *Mol*
739 *Cell* **2009**, *35*, 467-478, doi:10.1016/j.molcel.2009.06.027.
- 740 94. Clemson, C.M.; Hutchinson, J.N.; Sara, S.A.; Ensminger, A.W.; Fox, A.H.; Chess, A.;
741 Lawrence, J.B. An architectural role for a nuclear noncoding RNA: NEAT1 RNA is essential
742 for the structure of paraspeckles. *Mol Cell* **2009**, *33*, 717-726, doi:10.1016/j.molcel.2009.01.026.
- 743 95. Nakagawa, S.; Naganuma, T.; Shioi, G.; Hirose, T. Paraspeckles are subpopulation-specific
744 nuclear bodies that are not essential in mice. *J Cell Biol* **2011**, *193*, 31-39,
745 doi:10.1083/jcb.201011110.
- 746 96. Bernard, D.; Prasanth, K.V.; Tripathi, V.; Colasse, S.; Nakamura, T.; Xuan, Z.; Zhang, M.Q.;
747 Sedel, F.; Jourdain, L.; Couplier, F., et al. A long nuclear-retained non-coding RNA regulates
748 synaptogenesis by modulating gene expression. *EMBO J* **2010**, *29*, 3082-3093,
749 doi:10.1038/emboj.2010.199.
- 750 97. Matsui, K.; Nishizawa, M.; Ozaki, T.; Kimura, T.; Hashimoto, I.; Yamada, M.; Kaibori, M.;
751 Kamiyama, Y.; Ito, S.; Okumura, T. Natural antisense transcript stabilizes inducible nitric
752 oxide synthase messenger RNA in rat hepatocytes. *Hepatology* **2008**, *47*, 686-697,
753 doi:10.1002/hep.22036.
- 754 98. Gong, C.; Maquat, L.E. lncRNAs transactivate STAU1-mediated mRNA decay by duplexing
755 with 3' UTRs via Alu elements. *Nature* **2011**, *470*, 284-288, doi:10.1038/nature09701.
- 756 99. Poliseno, L.; Salmena, L.; Zhang, J.; Carver, B.; Haveman, W.J.; Pandolfi, P.P. A
757 coding-independent function of gene and pseudogene mRNAs regulates tumour biology.
758 *Nature* **2010**, *465*, 1033-1038, doi:10.1038/nature09144.
- 759 100. Klein, E.A.; Assoian, R.K. Transcriptional regulation of the cyclin D1 gene at a glance. *J Cell*
760 *Sci* **2008**, *121*, 3853-3857, doi:10.1242/jcs.039131.

- 761 101. Huarte, M.; Guttman, M.; Feldser, D.; Garber, M.; Koziol, M.J.; Kenzelmann-Broz, D.; Khalil,
762 A.M.; Zuk, O.; Amit, I.; Rabani, M., et al. A large intergenic noncoding RNA induced by p53
763 mediates global gene repression in the p53 response. *Cell* **2010**, *142*, 409-419,
764 doi:10.1016/j.cell.2010.06.040.
- 765 102. Marin-Bejar, O.; Mas, A.M.; Gonzalez, J.; Martinez, D.; Athie, A.; Morales, X.; Galduroz, M.;
766 Raimondi, I.; Grossi, E.; Guo, S., et al. The human lncRNA LINC-PINT inhibits tumor cell
767 invasion through a highly conserved sequence element. *Genome Biol* **2017**, *18*, 202,
768 doi:10.1186/s13059-017-1331-y.
- 769 103. Marin-Bejar, O.; Marchese, F.P.; Athie, A.; Sanchez, Y.; Gonzalez, J.; Segura, V.; Huang, L.;
770 Moreno, I.; Navarro, A.; Monzo, M., et al. Pint lincRNA connects the p53 pathway with
771 epigenetic silencing by the Polycomb repressive complex 2. *Genome Biol* **2013**, *14*, R104,
772 doi:10.1186/gb-2013-14-9-r104.
- 773 104. Hung, T.; Wang, Y.; Lin, M.F.; Koegel, A.K.; Kotake, Y.; Grant, G.D.; Horlings, H.M.; Shah,
774 N.; Umbricht, C.; Wang, P., et al. Extensive and coordinated transcription of noncoding
775 RNAs within cell-cycle promoters. *Nat Genet* **2011**, *43*, 621-629, doi:10.1038/ng.848.
- 776 105. Schmitt, A.M.; Garcia, J.T.; Hung, T.; Flynn, R.A.; Shen, Y.; Qu, K.; Payumo, A.Y.;
777 Peres-da-Silva, A.; Broz, D.K.; Baum, R., et al. An inducible long noncoding RNA amplifies
778 DNA damage signaling. *Nat Genet* **2016**, *48*, 1370-1376, doi:10.1038/ng.3673.
- 779 106. Zhang, Y.; He, Q.; Hu, Z.; Feng, Y.; Fan, L.; Tang, Z.; Yuan, J.; Shan, W.; Li, C.; Hu, X., et al.
780 Long noncoding RNA LINP1 regulates repair of DNA double-strand breaks in
781 triple-negative breast cancer. *Nat Struct Mol Biol* **2016**, *23*, 522-530, doi:10.1038/nsmb.3211.
- 782 107. Mahmoudi, S.; Henriksson, S.; Corcoran, M.; Mendez-Vidal, C.; Wiman, K.G.; Farnebo, M.
783 Wrap53, a natural p53 antisense transcript required for p53 induction upon DNA damage.
784 *Mol Cell* **2009**, *33*, 462-471, doi:10.1016/j.molcel.2009.01.028.
- 785 108. Rezaei, M.; Emadi-Baygi, M.; Hoffmann, M.J.; Schulz, W.A.; Nikpour, P. Altered expression
786 of LINC-ROR in cancer cell lines and tissues. *Tumour Biol* **2016**, *37*, 1763-1769,
787 doi:10.1007/s13277-015-3933-x.
- 788 109. Zhang, A.; Zhou, N.; Huang, J.; Liu, Q.; Fukuda, K.; Ma, D.; Lu, Z.; Bai, C.; Watabe, K.; Mo,
789 Y.Y. The human long non-coding RNA-RoR is a p53 repressor in response to DNA damage.
790 *Cell Res* **2013**, *23*, 340-350, doi:10.1038/cr.2012.164.
- 791 110. Ji, P.; Diederichs, S.; Wang, W.; Boing, S.; Metzger, R.; Schneider, P.M.; Tidow, N.; Brandt, B.;
792 Buerger, H.; Bulk, E., et al. MALAT-1, a novel noncoding RNA, and thymosin beta4 predict
793 metastasis and survival in early-stage non-small cell lung cancer. *Oncogene* **2003**, *22*,
794 8031-8041, doi:10.1038/sj.onc.1206928.
- 795 111. Lin, R.; Maeda, S.; Liu, C.; Karin, M.; Edgington, T.S. A large noncoding RNA is a marker for
796 murine hepatocellular carcinomas and a spectrum of human carcinomas. *Oncogene* **2007**, *26*,
797 851-858, doi:10.1038/sj.onc.1209846.
- 798 112. Khalil, A.M.; Guttman, M.; Huarte, M.; Garber, M.; Raj, A.; Rivea Morales, D.; Thomas, K.;
799 Presser, A.; Bernstein, B.E.; van Oudenaarden, A., et al. Many human large intergenic
800 noncoding RNAs associate with chromatin-modifying complexes and affect gene
801 expression. *Proc Natl Acad Sci U S A* **2009**, *106*, 11667-11672, doi:10.1073/pnas.0904715106.

- 802 113. Yang, L.; Lin, C.; Liu, W.; Zhang, J.; Ohgi, K.A.; Grinstein, J.D.; Dorrestein, P.C.; Rosenfeld,
803 M.G. ncRNA- and Pc2 methylation-dependent gene relocation between nuclear structures
804 mediates gene activation programs. *Cell* **2011**, *147*, 773-788, doi:10.1016/j.cell.2011.08.054.
- 805 114. Liu, Q.; Huang, J.; Zhou, N.; Zhang, Z.; Zhang, A.; Lu, Z.; Wu, F.; Mo, Y.Y. LncRNA
806 loc285194 is a p53-regulated tumor suppressor. *Nucleic Acids Res* **2013**, *41*, 4976-4987,
807 doi:10.1093/nar/gkt182.
- 808 115. Sharma, V.; Khurana, S.; Kubben, N.; Abdelmohsen, K.; Oberdoerffer, P.; Gorospe, M.;
809 Misteli, T. A BRCA1-interacting lncRNA regulates homologous recombination. *EMBO Rep*
810 **2015**, *16*, 1520-1534, doi:10.15252/embr.201540437.
- 811 116. Prensner, J.R.; Chen, W.; Iyer, M.K.; Cao, Q.; Ma, T.; Han, S.; Sahu, A.; Malik, R.;
812 Wilder-Romans, K.; Navone, N., et al. PCAT-1, a long noncoding RNA, regulates BRCA2
813 and controls homologous recombination in cancer. *Cancer Res* **2014**, *74*, 1651-1660,
814 doi:10.1158/0008-5472.CAN-13-3159.
- 815 117. Prensner, J.R.; Chen, W.; Han, S.; Iyer, M.K.; Cao, Q.; Kothari, V.; Evans, J.R.; Knudsen, K.E.;
816 Paulsen, M.T.; Ljungman, M., et al. The long non-coding RNA PCAT-1 promotes prostate
817 cancer cell proliferation through cMyc. *Neoplasia* **2014**, *16*, 900-908,
818 doi:10.1016/j.neo.2014.09.001.
- 819 118. Prensner, J.R.; Iyer, M.K.; Balbin, O.A.; Dhanasekaran, S.M.; Cao, Q.; Brenner, J.C.; Laxman,
820 B.; Asangani, I.A.; Grasso, C.S.; Kominsky, H.D., et al. Transcriptome sequencing across a
821 prostate cancer cohort identifies PCAT-1, an unannotated lincRNA implicated in disease
822 progression. *Nat Biotechnol* **2011**, *29*, 742-749, doi:10.1038/nbt.1914.
- 823 119. Wan, G.; Hu, X.; Liu, Y.; Han, C.; Sood, A.K.; Calin, G.A.; Zhang, X.; Lu, X. A novel
824 non-coding RNA lncRNA-JADE connects DNA damage signalling to histone H4
825 acetylation. *EMBO J* **2013**, *32*, 2833-2847, doi:10.1038/emboj.2013.221.
- 826 120. Wan, G.; Mathur, R.; Hu, X.; Liu, Y.; Zhang, X.; Peng, G.; Lu, X. Long non-coding RNA
827 ANRIL (CDKN2B-AS) is induced by the ATM-E2F1 signaling pathway. *Cell Signal* **2013**, *25*,
828 1086-1095, doi:10.1016/j.cellsig.2013.02.006.
- 829 121. Pilyugin, M.; Irminger-Finger, I. Long non-coding RNA and microRNAs might act in
830 regulating the expression of BARD1 mRNAs. *Int J Biochem Cell Biol* **2014**, *54*, 356-367,
831 doi:10.1016/j.biocel.2014.06.018.
- 832 122. Porro, A.; Feuerhahn, S.; Lingner, J. TERRA-reinforced association of LSD1 with MRE11
833 promotes processing of uncapped telomeres. *Cell Rep* **2014**, *6*, 765-776,
834 doi:10.1016/j.celrep.2014.01.022.
- 835 123. Maringele, L.; Lydall, D. EXO1-dependent single-stranded DNA at telomeres activates
836 subsets of DNA damage and spindle checkpoint pathways in budding yeast yku70Delta
837 mutants. *Genes Dev* **2002**, *16*, 1919-1933, doi:10.1101/gad.225102.
- 838 124. Gazy, I.; Zeevi, D.A.; Renbaum, P.; Zeligson, S.; Eini, L.; Bashari, D.; Smith, Y.; Lahad, A.;
839 Goldberg, M.; Ginsberg, D., et al. TODRA, a lncRNA at the RAD51 Locus, Is Oppositely
840 Regulated to RAD51, and Enhances RAD51-Dependent DSB (Double Strand Break) Repair.
841 *PLoS One* **2015**, *10*, e0134120, doi:10.1371/journal.pone.0134120.
- 842 125. Xue, Y.; Ma, G.; Zhang, Z.; Hua, Q.; Chu, H.; Tong, N.; Yuan, L.; Qin, C.; Yin, C.; Zhang, Z.,
843 et al. A novel antisense long noncoding RNA regulates the expression of MDC1 in bladder
844 cancer. *Oncotarget* **2015**, *6*, 484-493, doi:10.18632/oncotarget.2861.

- 845 126. Cajigas, I.; Leib, D.E.; Cochrane, J.; Luo, H.; Swyter, K.R.; Chen, S.; Clark, B.S.; Thompson, J.;
846 Yates, J.R., 3rd; Kingston, R.E., et al. Evf2 lncRNA/BRG1/DLX1 interactions reveal
847 RNA-dependent inhibition of chromatin remodeling. *Development* **2015**, *142*, 2641-2652,
848 doi:10.1242/dev.126318.
- 849 127. Kwon, S.J.; Park, J.H.; Park, E.J.; Lee, S.A.; Lee, H.S.; Kang, S.W.; Kwon, J. ATM-mediated
850 phosphorylation of the chromatin remodeling enzyme BRG1 modulates DNA
851 double-strand break repair. *Oncogene* **2015**, *34*, 303-313, doi:10.1038/onc.2013.556.
- 852 128. Betts, J.A.; Moradi Marjaneh, M.; Al-Ejeh, F.; Lim, Y.C.; Shi, W.; Sivakumaran, H.; Tropee, R.;
853 Patch, A.M.; Clark, M.B.; Bartonicek, N., et al. Long Noncoding RNAs CUPID1 and CUPID2
854 Mediate Breast Cancer Risk at 11q13 by Modulating the Response to DNA Damage. *Am J*
855 *Hum Genet* **2017**, *101*, 255-266, doi:10.1016/j.ajhg.2017.07.007.
- 856 129. Sanchez, Y.; Segura, V.; Marin-Bejar, O.; Athie, A.; Marchese, F.P.; Gonzalez, J.; Bujanda, L.;
857 Guo, S.; Matheu, A.; Huarte, M. Genome-wide analysis of the human p53 transcriptional
858 network unveils a lncRNA tumour suppressor signature. *Nat Commun* **2014**, *5*, 5812,
859 doi:10.1038/ncomms6812.
- 860 130. Leveille, N.; Melo, C.A.; Rooijers, K.; Diaz-Lagares, A.; Melo, S.A.; Korkmaz, G.; Lopes, R.;
861 Akbari Moqadam, F.; Maia, A.R.; Wijchers, P.J., et al. Genome-wide profiling of
862 p53-regulated enhancer RNAs uncovers a subset of enhancers controlled by a lncRNA. *Nat*
863 *Commun* **2015**, *6*, 6520, doi:10.1038/ncomms7520.
- 864 131. Dimitrova, N.; Zamudio, J.R.; Jong, R.M.; Soukup, D.; Resnick, R.; Sarma, K.; Ward, A.J.; Raj,
865 A.; Lee, J.T.; Sharp, P.A., et al. lincRNA-p21 activates p21 in cis to promote Polycomb target
866 gene expression and to enforce the G1/S checkpoint. *Mol Cell* **2014**, *54*, 777-790,
867 doi:10.1016/j.molcel.2014.04.025.
- 868 132. Yoon, J.H.; Abdelmohsen, K.; Srikantan, S.; Yang, X.; Martindale, J.L.; De, S.; Huarte, M.;
869 Zhan, M.; Becker, K.G.; Gorospe, M. lincRNA-p21 suppresses target mRNA translation. *Mol*
870 *Cell* **2012**, *47*, 648-655, doi:10.1016/j.molcel.2012.06.027.
- 871 133. Li, M.; Gou, H.; Tripathi, B.K.; Huang, J.; Jiang, S.; Dubois, W.; Waybright, T.; Lei, M.; Shi, J.;
872 Zhou, M., et al. An Apela RNA-Containing Negative Feedback Loop Regulates
873 p53-Mediated Apoptosis in Embryonic Stem Cells. *Cell Stem Cell* **2015**, *16*, 669-683,
874 doi:10.1016/j.stem.2015.04.002.
- 875 134. Farnebo, M. Wrap53, a novel regulator of p53. *Cell Cycle* **2009**, *8*, 2343-2346,
876 doi:10.4161/cc.8.15.9223.
- 877 135. Mahmoudi, S.; Henriksson, S.; Corcoran, M.; Mendez-Vidal, C.; Wiman, K.G.; Farnebo, M.
878 Wrap53, a Natural p53 Antisense Transcript Required for p53 Induction upon DNA
879 Damage. *Mol Cell* **2016**, *64*, 1009, doi:10.1016/j.molcel.2016.11.027.
- 880 136. Henriksson, S.; Rassoolzadeh, H.; Hedstrom, E.; Coucoravas, C.; Julner, A.; Goldstein, M.;
881 Imreh, G.; Zhivotovsky, B.; Kastan, M.B.; Helleday, T., et al. The scaffold protein
882 WRAP53beta orchestrates the ubiquitin response critical for DNA double-strand break
883 repair. *Genes Dev* **2014**, *28*, 2726-2738, doi:10.1101/gad.246546.114.
- 884 137. Tycowski, K.T.; Shu, M.D.; Kukoyi, A.; Steitz, J.A. A conserved WD40 protein binds the
885 Cajal body localization signal of scaRNP particles. *Mol Cell* **2009**, *34*, 47-57,
886 doi:10.1016/j.molcel.2009.02.020.

- 887 138. Zhong, F.; Savage, S.A.; Shkreli, M.; Giri, N.; Jessop, L.; Myers, T.; Chen, R.; Alter, B.P.;
888 Artandi, S.E. Disruption of telomerase trafficking by TCAB1 mutation causes dyskeratosis
889 congenita. *Genes Dev* **2011**, *25*, 11-16, doi:10.1101/gad.2006411.
- 890 139. Wu, H.F.; Ren, L.G.; Xiao, J.Q.; Zhang, Y.; Mao, X.W.; Zhou, L.F. Long non-coding RNA
891 LINP1 promotes the malignant progression of prostate cancer by regulating p53. *Eur Rev*
892 *Med Pharmacol Sci* **2018**, *22*, 4467-4476, doi:10.26355/eurev_201807_15498.
- 893 140. Liang, Y.; Li, Y.; Song, X.; Zhang, N.; Sang, Y.; Zhang, H.; Liu, Y.; Chen, B.; Zhao, W.; Wang,
894 L., et al. Long noncoding RNA LINP1 acts as an oncogene and promotes chemoresistance in
895 breast cancer. *Cancer Biol Ther* **2018**, *19*, 120-131, doi:10.1080/15384047.2017.1394543.
- 896 141. Wang, X.; Liu, H.; Shi, L.; Yu, X.; Gu, Y.; Sun, X. LINP1 facilitates DNA damage repair
897 through non-homologous end joining (NHEJ) pathway and subsequently decreases the
898 sensitivity of cervical cancer cells to ionizing radiation. *Cell Cycle* **2018**, *17*, 439-447,
899 doi:10.1080/15384101.2018.1442625.
- 900 142. Gutschner, T.; Hammerle, M.; Diederichs, S. MALAT1 -- a paradigm for long noncoding
901 RNA function in cancer. *J Mol Med (Berl)* **2013**, *91*, 791-801, doi:10.1007/s00109-013-1028-y.
- 902 143. Hutchinson, J.N.; Ensminger, A.W.; Clemson, C.M.; Lynch, C.R.; Lawrence, J.B.; Chess, A. A
903 screen for nuclear transcripts identifies two linked noncoding RNAs associated with SC35
904 splicing domains. *BMC Genomics* **2007**, *8*, 39, doi:10.1186/1471-2164-8-39.
- 905 144. Lamond, A.I.; Spector, D.L. Nuclear speckles: a model for nuclear organelles. *Nat Rev Mol*
906 *Cell Biol* **2003**, *4*, 605-612, doi:10.1038/nrm1172.
- 907 145. Tripathi, V.; Ellis, J.D.; Shen, Z.; Song, D.Y.; Pan, Q.; Watt, A.T.; Freier, S.M.; Bennett, C.F.;
908 Sharma, A.; Bubulya, P.A., et al. The nuclear-retained noncoding RNA MALAT1 regulates
909 alternative splicing by modulating SR splicing factor phosphorylation. *Mol Cell* **2010**, *39*,
910 925-938, doi:10.1016/j.molcel.2010.08.011.
- 911 146. Wilusz, J.E.; Freier, S.M.; Spector, D.L. 3' end processing of a long nuclear-retained
912 noncoding RNA yields a tRNA-like cytoplasmic RNA. *Cell* **2008**, *135*, 919-932,
913 doi:10.1016/j.cell.2008.10.012.
- 914 147. Brown, J.A.; Bulkley, D.; Wang, J.; Valenstein, M.L.; Yario, T.A.; Steitz, T.A.; Steitz, J.A.
915 Structural insights into the stabilization of MALAT1 noncoding RNA by a bipartite triple
916 helix. *Nat Struct Mol Biol* **2014**, *21*, 633-640, doi:10.1038/nsmb.2844.
- 917 148. Chen, R.; Liu, Y.; Zhuang, H.; Yang, B.; Hei, K.; Xiao, M.; Hou, C.; Gao, H.; Zhang, X.; Jia, C.,
918 et al. Quantitative proteomics reveals that long non-coding RNA MALAT1 interacts with
919 DBC1 to regulate p53 acetylation. *Nucleic Acids Res* **2017**, *45*, 9947-9959,
920 doi:10.1093/nar/gkx600.
- 921 149. Hu, Y.; Lin, J.; Fang, H.; Fang, J.; Li, C.; Chen, W.; Liu, S.; Ondrejka, S.; Gong, Z.; Reu, F., et al.
922 Targeting the MALAT1/PARP1/LIG3 complex induces DNA damage and apoptosis in
923 multiple myeloma. *Leukemia* **2018**, 10.1038/s41375-018-0104-2, doi:10.1038/s41375-018-0104-2.
- 924 150. Polo, S.E.; Blackford, A.N.; Chapman, J.R.; Baskcomb, L.; Gravel, S.; Rusch, A.; Thomas, A.;
925 Blundred, R.; Smith, P.; Kzhyshkowska, J., et al. Regulation of DNA-end resection by
926 hnRNP-like proteins promotes DNA double-strand break signaling and repair. *Mol Cell*
927 **2012**, *45*, 505-516, doi:10.1016/j.molcel.2011.12.035.

- 928 151. Sathishkumar, C.; Prabu, P.; Mohan, V.; Balasubramanyam, M. Linking a role of lncRNAs
929 (long non-coding RNAs) with insulin resistance, accelerated senescence, and inflammation
930 in patients with type 2 diabetes. *Human genomics* **2018**, *12*, 41, doi:10.1186/s40246-018-0173-3.
- 931 152. Chen, S.; Liang, H.; Yang, H.; Zhou, K.; Xu, L.; Liu, J.; Lai, B.; Song, L.; Luo, H.; Peng, J., et al.
932 LincRNA-p21: function and mechanism in cancer. *Medical oncology (Northwood, London,
933 England)* **2017**, *34*, 98, doi:10.1007/s12032-017-0959-5.
- 934 153. Sauvageau, M.; Goff, L.A.; Lodato, S.; Bonev, B.; Groff, A.F.; Gerhardinger, C.;
935 Sanchez-Gomez, D.B.; Hacısuleyman, E.; Li, E.; Spence, M., et al. Multiple knockout mouse
936 models reveal lincRNAs are required for life and brain development. *Elife* **2013**, *2*, e01749,
937 doi:10.7554/eLife.01749.
- 938 154. Li, L.; Zhang, G.Q.; Chen, H.; Zhao, Z.J.; Chen, H.Z.; Liu, H.; Wang, G.; Jia, Y.H.; Pan, S.H.;
939 Kong, R., et al. Plasma and tumor levels of Linc-pint are diagnostic and prognostic
940 biomarkers for pancreatic cancer. *Oncotarget* **2016**, *7*, 71773-71781,
941 doi:10.18632/oncotarget.12365.
- 942 155. Zhu, J.; Gu, H.; Lv, X.; Yuan, C.; Ni, P.; Liu, F. LINC-PINT Activates the Mitogen-Activated
943 Protein Kinase Pathway to Promote Acute Myocardial Infarction by Regulating
944 miR-208a-3p. *Circulation journal : official journal of the Japanese Circulation Society* **2018**,
945 10.1253/circj.CJ-18-0396, doi:10.1253/circj.CJ-18-0396.
- 946 156. Permuth, J.B.; Chen, D.T.; Yoder, S.J.; Li, J.; Smith, A.T.; Choi, J.W.; Kim, J.; Balagurunathan,
947 Y.; Jiang, K.; Coppola, D., et al. Linc-ing Circulating Long Non-coding RNAs to the
948 Diagnosis and Malignant Prediction of Intraductal Papillary Mucinous Neoplasms of the
949 Pancreas. *Sci Rep* **2017**, *7*, 10484, doi:10.1038/s41598-017-09754-5.
- 950 157. Kotake, Y.; Goto, T.; Naemura, M.; Inoue, Y.; Okamoto, H.; Tahara, K. Long Noncoding
951 RNA PANDA Positively Regulates Proliferation of Osteosarcoma Cells. *Anticancer research*
952 **2017**, *37*, 81-85, doi:10.21873/anticancer.11292.
- 953 158. Fenoglio, C.; Oldoni, E.; Serpente, M.; De Riz, M.A.; Arcaro, M.; D'Anca, M.; Pietroboni,
954 A.M.; Calvi, A.; Lecchi, E.; Goris, A., et al. LncRNAs expression profile in peripheral blood
955 mononuclear cells from multiple sclerosis patients. *Journal of neuroimmunology* **2018**, *324*,
956 129-135, doi:10.1016/j.jneuroim.2018.08.008.
- 957 159. Yi, Y.; Tsai, S.H.; Cheng, J.C.; Wang, E.Y.; Anglesio, M.S.; Cochrane, D.R.; Fuller, M.; Gibb,
958 E.A.; Wei, W.; Huntsman, D.G., et al. APELA promotes tumour growth and cell migration in
959 ovarian cancer in a p53-dependent manner. *Gynecologic oncology* **2017**, *147*, 663-671,
960 doi:10.1016/j.ygyno.2017.10.016.
- 961 160. Yang, Y.; Huang, J.; Xie, N.; Huang, H.; Xu, S.; Cai, J.; Qi, S. lincROR influences the stemness
962 and crizotinib resistance in EML-ALK(+) non-small-cell lung cancer cells. *Onco Targets Ther*
963 **2018**, *11*, 3649-3657, doi:10.2147/ott.s165290.
- 964 161. Greco, S.; Zaccagnini, G.; Perfetti, A.; Fuschi, P.; Valaperta, R.; Voellenkle, C.; Castelvechio,
965 S.; Gaetano, C.; Finato, N.; Beltrami, A.P., et al. Long noncoding RNA dysregulation in
966 ischemic heart failure. *Journal of translational medicine* **2016**, *14*, 183,
967 doi:10.1186/s12967-016-0926-5.
- 968 162. Li, Z.; Yu, X.; Shen, J. Long non-coding RNAs: emerging players in osteosarcoma. *Tumour*
969 *Biol* **2016**, *37*, 2811-2816, doi:10.1007/s13277-015-4749-4.

- 970 163. Li, Z.; Dou, P.; Liu, T.; He, S. Application of Long Noncoding RNAs in Osteosarcoma:
971 Biomarkers and Therapeutic Targets. *Cellular physiology and biochemistry : international journal*
972 *of experimental cellular physiology, biochemistry, and pharmacology* **2017**, *42*, 1407-1419,
973 doi:10.1159/000479205.
- 974 164. Liu, L.; Liu, Y.; Zhuang, C.; Xu, W.; Fu, X.; Lv, Z.; Wu, H.; Mou, L.; Zhao, G.; Cai, Z., et al.
975 Inducing cell growth arrest and apoptosis by silencing long non-coding RNA PCAT-1 in
976 human bladder cancer. *Tumour Biol* **2015**, *36*, 7685-7689, doi:10.1007/s13277-015-3490-3.
- 977 165. Cui, W.C.; Wu, Y.F.; Qu, H.M. Up-regulation of long non-coding RNA PCAT-1 correlates
978 with tumor progression and poor prognosis in gastric cancer. *Eur Rev Med Pharmacol Sci*
979 **2017**, *21*, 3021-3027.
- 980 166. Aguilo, F.; Di Cecilia, S.; Walsh, M.J. Long Non-coding RNA ANRIL and Polycomb in
981 Human Cancers and Cardiovascular Disease. *Current topics in microbiology and immunology*
982 **2016**, *394*, 29-39, doi:10.1007/82_2015_455.
- 983 167. Simion, V.; Haemmig, S.; Feinberg, M.W. LncRNAs in vascular biology and disease. *Vascular*
984 *pharmacology* **2018**, 10.1016/j.vph.2018.01.003, doi:10.1016/j.vph.2018.01.003.
- 985 168. Heeke, A.L.; Pishvaian, M.J.; Lynce, F.; Xiu, J.; Brody, J.R.; Chen, W.J.; Baker, T.M.; Marshall,
986 J.L.; Isaacs, C. Prevalence of Homologous Recombination-Related Gene Mutations Across
987 Multiple Cancer Types. *JCO precision oncology* **2018**, *2018*, doi:10.1200/po.17.00286.
- 988 169. Harley, C.B. Telomerase and cancer therapeutics. *Nat Rev Cancer* **2008**, *8*, 167-179,
989 doi:10.1038/nrc2275.
- 990 170. Qin, Y.; Zhuang, S.; Wen, J.; Zheng, K. Long non-coding RNA MDC1-AS inhibits human
991 gastric cancer cell proliferation and metastasis through an MDC1-dependent mechanism.
992 *Exp. Ther. Med.* **2018**, *15*, 191-197.
- 993 171. Yue, H.; Zhu, J.; Xie, S.; Li, F.; Xu, Q. MDC1-AS, an antisense long noncoding RNA, regulates
994 cell proliferation of glioma. *Biomed Pharmacother* **2016**, *81*, 203-209.
- 995 172. Cajigas, I.; Chakraborty, A.; Swyter, K.R.; Luo, H.; Bastidas, M.; Nigro, M.; Morris, E.R.;
996 Chen, S.; VanGompel, M.J.W.; Leib, D., et al. The Evf2 Ultraconserved Enhancer lncRNA
997 Functionally and Spatially Organizes Megabase Distant Genes in the Developing Forebrain.
998 *Mol Cell* **2018**, *71*, 956-972 e959, doi:10.1016/j.molcel.2018.07.024.
- 999 173. Bond, A.M.; Vangompel, M.J.; Sametsky, E.A.; Clark, M.F.; Savage, J.C.; Disterhoft, J.F.;
1000 Kohtz, J.D. Balanced gene regulation by an embryonic brain ncRNA is critical for adult
1001 hippocampal GABA circuitry. *Nat Neurosci* **2009**, *12*, 1020-1027, doi:10.1038/nn.2371.
- 1002 174. de Lange, T. Shelterin-Mediated Telomere Protection. *Annu Rev Genet* **2018**,
1003 10.1146/annurev-genet-032918-021921, doi:10.1146/annurev-genet-032918-021921.
- 1004 175. Patel, D.J.; Phan, A.T.; Kuryavyi, V. Human telomere, oncogenic promoter and 5'-UTR
1005 G-quadruplexes: diverse higher order DNA and RNA targets for cancer therapeutics.
1006 *Nucleic Acids Res* **2007**, *35*, 7429-7455, doi:10.1093/nar/gkm711.
- 1007 176. Cusanelli, E.; Chartrand, P. Telomeric repeat-containing RNA TERRA: a noncoding RNA
1008 connecting telomere biology to genome integrity. *Front Genet* **2015**, *6*, 143,
1009 doi:10.3389/fgene.2015.00143.
- 1010 177. Azzalin, C.M.; Lingner, J. Telomere functions grounding on TERRA firma. *Trends Cell Biol*
1011 **2015**, *25*, 29-36, doi:10.1016/j.tcb.2014.08.007.

- 1012 178. Flynn, R.L.; Centore, R.C.; O'Sullivan, R.J.; Rai, R.; Tse, A.; Songyang, Z.; Chang, S.;
1013 Karlseder, J.; Zou, L. TERRA and hnRNP A1 orchestrate an RPA-to-POT1 switch on
1014 telomeric single-stranded DNA. *Nature* **2011**, *471*, 532-536, doi:10.1038/nature09772.
- 1015 179. Bartel, D.P. Metazoan MicroRNAs. *Cell* **2018**, *173*, 20-51, doi:10.1016/j.cell.2018.03.006.
- 1016 180. Bartel, D.P. MicroRNAs: genomics, biogenesis, mechanism, and function. *Cell* **2004**, *116*,
1017 281-297.
- 1018 181. Di Leva, G.; Garofalo, M.; Croce, C.M. MicroRNAs in cancer. *Annu Rev Pathol* **2014**, *9*,
1019 287-314, doi:10.1146/annurev-pathol-012513-104715.
- 1020 182. Calin, G.A.; Croce, C.M. MicroRNA signatures in human cancers. *Nat Rev Cancer* **2006**, *6*,
1021 857-866, doi:10.1038/nrc1997.
- 1022 183. Lu, J.; Getz, G.; Miska, E.A.; Alvarez-Saavedra, E.; Lamb, J.; Peck, D.; Sweet-Cordero, A.;
1023 Ebert, B.L.; Mak, R.H.; Ferrando, A.A., et al. MicroRNA expression profiles classify human
1024 cancers. *Nature* **2005**, *435*, 834-838, doi:10.1038/nature03702.
- 1025 184. Bottai, G.; Pasculli, B.; Calin, G.A.; Santarpia, L. Targeting the microRNA-regulating DNA
1026 damage/repair pathways in cancer. *Expert Opin Biol Ther* **2014**, *14*, 1667-1683,
1027 doi:10.1517/14712598.2014.950650.
- 1028 185. Hermeking, H. MicroRNAs in the p53 network: micromanagement of tumour suppression.
1029 *Nat Rev Cancer* **2012**, *12*, 613-626, doi:10.1038/nrc3318.
- 1030 186. Lal, A.; Pan, Y.; Navarro, F.; Dykxhoorn, D.M.; Moreau, L.; Meire, E.; Bentwich, Z.;
1031 Lieberman, J.; Chowdhury, D. miR-24-mediated downregulation of H2AX suppresses DNA
1032 repair in terminally differentiated blood cells. *Nat Struct Mol Biol* **2009**, *16*, 492-498,
1033 doi:10.1038/nsmb.1589.
- 1034 187. Wang, Y.; Huang, J.W.; Li, M.; Cavenee, W.K.; Mitchell, P.S.; Zhou, X.; Tewari, M.; Furnari,
1035 F.B.; Taniguchi, T. MicroRNA-138 modulates DNA damage response by repressing histone
1036 H2AX expression. *Mol Cancer Res* **2011**, *9*, 1100-1111, doi:10.1158/1541-7786.MCR-11-0007.
- 1037 188. Tessitore, A.; Ciciarelli, G.; Del Vecchio, F.; Gaggiano, A.; Verzella, D.; Fischietti, M.;
1038 Vecchiotti, D.; Capece, D.; Zazzeroni, F.; Alesse, E. MicroRNAs in the DNA Damage/Repair
1039 Network and Cancer. *Int J Genomics* **2014**, *2014*, 820248, doi:10.1155/2014/820248.
- 1040 189. Hu, H.; Du, L.; Nagabayashi, G.; Seeger, R.C.; Gatti, R.A. ATM is down-regulated by
1041 N-Myc-regulated microRNA-421. *Proc Natl Acad Sci U S A* **2010**, *107*, 1506-1511,
1042 doi:10.1073/pnas.0907763107.
- 1043 190. Yan, D.; Ng, W.L.; Zhang, X.; Wang, P.; Zhang, Z.; Mo, Y.Y.; Mao, H.; Hao, C.; Olson, J.J.;
1044 Curran, W.J., et al. Targeting DNA-PKcs and ATM with miR-101 sensitizes tumors to
1045 radiation. *PLoS One* **2010**, *5*, e11397, doi:10.1371/journal.pone.0011397.
- 1046 191. Chen, X.; Liang, H.; Zhang, J.; Zen, K.; Zhang, C.Y. Secreted microRNAs: a new form of
1047 intercellular communication. *Trends Cell Biol* **2012**, *22*, 125-132, doi:10.1016/j.tcb.2011.12.001.
- 1048 192. Perez-Gonzalez, R.; Gauthier, S.A.; Kumar, A.; Saito, M.; Saito, M.; Levy, E. A Method for
1049 Isolation of Extracellular Vesicles and Characterization of Exosomes from Brain
1050 Extracellular Space. *Methods Mol Biol* **2017**, *1545*, 139-151, doi:10.1007/978-1-4939-6728-5_10.
- 1051 193. Beach, A.; Zhang, H.G.; Ratajczak, M.Z.; Kakar, S.S. Exosomes: an overview of biogenesis,
1052 composition and role in ovarian cancer. *J Ovarian Res* **2014**, *7*, 14, doi:10.1186/1757-2215-7-14.

- 1053 194. Mo, L.J.; Song, M.; Huang, Q.H.; Guan, H.; Liu, X.D.; Xie, D.F.; Huang, B.; Huang, R.X.;
1054 Zhou, P.K. Exosome-packaged miR-1246 contributes to bystander DNA damage by
1055 targeting LIG4. *Br J Cancer* **2018**, *119*, 492-502, doi:10.1038/s41416-018-0192-9.
- 1056 195. Di Francesco, A.; De Pitta, C.; Moret, F.; Barbieri, V.; Celotti, L.; Mognato, M. The
1057 DNA-damage response to gamma-radiation is affected by miR-27a in A549 cells. *Int J Mol Sci*
1058 **2013**, *14*, 17881-17896, doi:10.3390/ijms140917881.
- 1059 196. Bisso, A.; Faleschini, M.; Zampa, F.; Capaci, V.; De Santa, J.; Santarpia, L.; Piazza, S.;
1060 Cappelletti, V.; Daidone, M.; Agami, R., et al. Oncogenic miR-181a/b affect the DNA damage
1061 response in aggressive breast cancer. *Cell Cycle* **2013**, *12*, 1679-1687, doi:10.4161/cc.24757.
- 1062 197. Zhu, F.; Liu, J.L.; Li, J.P.; Xiao, F.; Zhang, Z.X.; Zhang, L. MicroRNA-124 (miR-124) regulates
1063 Ku70 expression and is correlated with neuronal death induced by ischemia/reperfusion. *J*
1064 *Mol Neurosci* **2014**, *52*, 148-155, doi:10.1007/s12031-013-0155-9.
- 1065 198. Choi, Y.E.; Meghani, K.; Brault, M.E.; Leclerc, L.; He, Y.J.; Day, T.A.; Elias, K.M.; Drapkin, R.;
1066 Weinstock, D.M.; Dao, F., et al. Platinum and PARP Inhibitor Resistance Due to
1067 Overexpression of MicroRNA-622 in BRCA1-Mutant Ovarian Cancer. *Cell Rep* **2016**, *14*,
1068 429-439, doi:10.1016/j.celrep.2015.12.046.
- 1069 199. Zhang, Z.Y.; Fu, S.L.; Xu, S.Q.; Zhou, X.; Liu, X.S.; Xu, Y.J.; Zhao, J.P.; Wei, S. By
1070 downregulating Ku80, hsa-miR-526b suppresses non-small cell lung cancer. *Oncotarget* **2015**,
1071 *6*, 1462-1477, doi:10.18632/oncotarget.2808.
- 1072 200. Wei, S.; Zhang, Z.Y.; Fu, S.L.; Xie, J.G.; Liu, X.S.; Xu, Y.J.; Zhao, J.P.; Xiong, W.N.
1073 Hsa-miR-623 suppresses tumor progression in human lung adenocarcinoma. *Cell Death Dis*
1074 **2016**, *7*, e2388, doi:10.1038/cddis.2016.260.
- 1075 201. Krishnan, K.; Steptoe, A.L.; Martin, H.C.; Wani, S.; Nones, K.; Waddell, N.; Mariasegaram,
1076 M.; Simpson, P.T.; Lakhani, S.R.; Gabrielli, B., et al. MicroRNA-182-5p targets a network of
1077 genes involved in DNA repair. *RNA* **2013**, *19*, 230-242, doi:10.1261/rna.034926.112.
- 1078 202. Moskwa, P.; Buffa, F.M.; Pan, Y.; Panchakshari, R.; Gottipati, P.; Muschel, R.J.; Beech, J.;
1079 Kulshrestha, R.; Abdelmohsen, K.; Weinstock, D.M., et al. miR-182-mediated
1080 downregulation of BRCA1 impacts DNA repair and sensitivity to PARP inhibitors. *Mol Cell*
1081 **2011**, *41*, 210-220, doi:10.1016/j.molcel.2010.12.005.
- 1082 203. Shen, J.; Ambrosone, C.B.; DiCioccio, R.A.; Odunsi, K.; Lele, S.B.; Zhao, H. A functional
1083 polymorphism in the miR-146a gene and age of familial breast/ovarian cancer diagnosis.
1084 *Carcinogenesis* **2008**, *29*, 1963-1966, doi:10.1093/carcin/bgn172.
- 1085 204. Choi, Y.E.; Pan, Y.; Park, E.; Konstantinopoulos, P.; De, S.; D'Andrea, A.; Chowdhury, D.
1086 MicroRNAs down-regulate homologous recombination in the G1 phase of cycling cells to
1087 maintain genomic stability. *Elife* **2014**, *3*, e02445, doi:10.7554/eLife.02445.
- 1088 205. Johnson, C.D.; Esquela-Kerscher, A.; Stefani, G.; Byrom, M.; Kelnar, K.; Ovcharenko, D.;
1089 Wilson, M.; Wang, X.; Shelton, J.; Shingara, J., et al. The let-7 microRNA represses cell
1090 proliferation pathways in human cells. *Cancer Res* **2007**, *67*, 7713-7722,
1091 doi:10.1158/0008-5472.CAN-07-1083.
- 1092 206. Mogilyansky, E.; Clark, P.; Quann, K.; Zhou, H.; Londin, E.; Jing, Y.; Rigoutsos, I.
1093 Post-transcriptional Regulation of BRCA2 through Interactions with miR-19a and miR-19b.
1094 *Front Genet* **2016**, *7*, 143, doi:10.3389/fgene.2016.00143.

- 1095 207. Wang, Y.; Huang, J.W.; Calses, P.; Kemp, C.J.; Taniguchi, T. MiR-96 downregulates REV1
1096 and RAD51 to promote cellular sensitivity to cisplatin and PARP inhibition. *Cancer Res* **2012**,
1097 72, 4037-4046, doi:10.1158/0008-5472.CAN-12-0103.
- 1098 208. Shen, L.; Wang, Q.; Liu, R.; Chen, Z.; Zhang, X.; Zhou, P.; Wang, Z. LncRNA lnc-RI regulates
1099 homologous recombination repair of DNA double-strand breaks by stabilizing RAD51
1100 mRNA as a competitive endogenous RNA. *Nucleic Acids Res* **2018**, 46, 717-729,
1101 doi:10.1093/nar/gkx1224.
- 1102 209. Liu, G.; Xue, F.; Zhang, W. miR-506: a regulator of chemo-sensitivity through suppression of
1103 the RAD51-homologous recombination axis. *Chin J Cancer* **2015**, 34, 485-487,
1104 doi:10.1186/s40880-015-0049-z.
- 1105 210. Gasparini, P.; Lovat, F.; Fassan, M.; Casadei, L.; Cascione, L.; Jacob, N.K.; Carasi, S.; Palmieri,
1106 D.; Costinean, S.; Shapiro, C.L., et al. Protective role of miR-155 in breast cancer through
1107 RAD51 targeting impairs homologous recombination after irradiation. *Proc Natl Acad Sci U S*
1108 *A* **2014**, 111, 4536-4541, doi:10.1073/pnas.1402604111.
- 1109 211. Neijenhuis, S.; Bajrami, I.; Miller, R.; Lord, C.J.; Ashworth, A. Identification of miRNA
1110 modulators to PARP inhibitor response. *DNA Repair (Amst)* **2013**, 12, 394-402,
1111 doi:10.1016/j.dnarep.2013.02.003.
- 1112 212. Huang, J.W.; Wang, Y.; Dhillon, K.K.; Calses, P.; Villegas, E.; Mitchell, P.S.; Tewari, M.;
1113 Kemp, C.J.; Taniguchi, T. Systematic screen identifies miRNAs that target RAD51 and
1114 RAD51D to enhance chemosensitivity. *Mol Cancer Res* **2013**, 11, 1564-1573,
1115 doi:10.1158/1541-7786.MCR-13-0292.
- 1116 213. Jung, E.J.; Santarpia, L.; Kim, J.; Esteva, F.J.; Moretti, E.; Buzdar, A.U.; Di Leo, A.; Le, X.F.;
1117 Bast, R.C., Jr.; Park, S.T., et al. Plasma microRNA 210 levels correlate with sensitivity to
1118 trastuzumab and tumor presence in breast cancer patients. *Cancer* **2012**, 118, 2603-2614,
1119 doi:10.1002/cncr.26565.
- 1120 214. Martin, N.T.; Nakamura, K.; Davies, R.; Nahas, S.A.; Brown, C.; Tunuguntla, R.; Gatti, R.A.;
1121 Hu, H. ATM-dependent MiR-335 targets CtIP and modulates the DNA damage response.
1122 *PLoS Genet* **2013**, 9, e1003505, doi:10.1371/journal.pgen.1003505.
- 1123 215. Lee, H.C.; Chang, S.S.; Choudhary, S.; Aalto, A.P.; Maiti, M.; Bamford, D.H.; Liu, Y. qiRNA
1124 is a new type of small interfering RNA induced by DNA damage. *Nature* **2009**, 459, 274-277,
1125 doi:10.1038/nature08041.
- 1126 216. Michalik, K.M.; Bottcher, R.; Forstemann, K. A small RNA response at DNA ends in
1127 *Drosophila*. *Nucleic Acids Res* **2012**, 40, 9596-9603, doi:10.1093/nar/gks711.
- 1128 217. Francia, S.; Cabrini, M.; Matti, V.; Oldani, A.; d'Adda di Fagagna, F. DICER, DROSHA and
1129 DNA damage response RNAs are necessary for the secondary recruitment of DNA damage
1130 response factors. *J Cell Sci* **2016**, 129, 1468-1476, doi:10.1242/jcs.182188.
- 1131 218. Francia, S.; Michelini, F.; Saxena, A.; Tang, D.; de Hoon, M.; Anelli, V.; Mione, M.; Carninci,
1132 P.; d'Adda di Fagagna, F. Site-specific DICER and DROSHA RNA products control the
1133 DNA-damage response. *Nature* **2012**, 488, 231-235, doi:10.1038/nature11179.
- 1134 219. Lu, W.T.; Hawley, B.R.; Skalka, G.L.; Baldock, R.A.; Smith, E.M.; Bader, A.S.; Malewicz, M.;
1135 Watts, F.Z.; Wilczynska, A.; Bushell, M. Drosha drives the formation of DNA:RNA hybrids
1136 around DNA break sites to facilitate DNA repair. *Nat Commun* **2018**, 9, 532,
1137 doi:10.1038/s41467-018-02893-x.