

Article

Untangling Cosmic Magnetic Fields: Faraday Tomography at Metre Wavelengths with LOFAR

Shane P. O'Sullivan ¹ , M. Brüggen ¹, C. L. Van Eck ² , M. J. Hardcastle ³, M. Haverkorn ⁴, T. W. Shimwell ⁵, C. Tasse ⁶, V. Vacca ⁷, C. Horellou ⁸, G. Heald ⁹

¹ Hamburger Sternwarte, Universität Hamburg, Gojenbergsweg 112, D-21029 Hamburg, Germany; shane@hs.uni-hamburg.de

² Dunlap Institute for Astronomy and Astrophysics, University of Toronto, 50 St. George Street, Toronto, ON M5S 3H4, Canada;

³ Centre for Astrophysics Research, School of Physics, Astronomy and Mathematics, University of Hertfordshire, College Lane, Hatfield AL10 9AB, UK;

⁴ Department of Astrophysics / IMAPP, Radboud University Nijmegen, PO Box 9010, 6500 GL Nijmegen, the Netherlands;

⁵ ASTRON, The Netherlands Institute for Radio Astronomy, Postbus 2, 7990 AA Dwingeloo, The Netherlands;

⁶ GEPI & USN, Observatoire de Paris, Université PSL, CNRS, 5 Place Jules Janssen, 92190 Meudon, France;

⁷ INAF – Osservatorio Astronomico di Cagliari, Via della Scienza 5, 09047 Selargius (CA), Italy;

⁸ Dept. of Space, Earth and Environment, Chalmers University of Technology, Onsala Space Observatory, 43992 Onsala, Sweden;

⁹ CSIRO Astronomy and Space Science, PO Box 1130, Bentley, WA 6102, Australia.

Abstract: The technique of Faraday tomography is a key tool for the study of magnetised plasmas in the new era of broadband radio polarisation observations. In particular, observations at metre-wavelengths provide significantly better Faraday depth accuracies compared to traditional cm-wavelength observations. However, the effect of Faraday depolarisation makes the polarised signal very challenging to detect at metre wavelengths (MHz frequencies). In this work, Faraday tomography is used to characterise the Faraday rotation properties of polarised sources found in data from the LOFAR Two-Metre Sky Survey (LoTSS). Of the 76 extragalactic polarised sources analysed here, we find that all host a radio-loud AGN. The majority of the sources ($\sim 64\%$) are large FR II radio galaxies with a median projected linear size of 710 kpc and median radio luminosity at 144 MHz of $4 \times 10^{26} \text{ W Hz}^{-1}$. In several cases, both hotspots are detected in polarisation at an angular resolution of $\sim 20''$. One such case allowed a study of intergalactic magnetic fields on scales of 3.4 Mpc. Other detected source types include an FRI radio galaxy and at least 8 blazars. Most sources display simple Faraday spectra, however, we highlight one blazar that displays a complex Faraday spectrum, with two close peaks in the Faraday dispersion function.

Keywords: magnetic fields; Faraday tomography; large-scale structure; AGN; Milky Way

1. Introduction

Advances in radio receiver technology that enable observations spanning wide, continuous frequency ranges, at high spectral resolution, has opened a new parameter space for studies of cosmic magnetism. In particular, it brings the technique of Faraday tomography to the fore, which allows detailed studies of the distribution of linearly polarised radiation as a function of Faraday depth, $F(\phi)$, which probes the physical properties of magnetised plasma along the line of sight. In this formalism, the complex linearly polarised intensity, $P(\lambda^2)$, is expressed as

$$P(\lambda^2) = \int_{-\infty}^{\infty} F(\phi) e^{2i\phi\lambda^2} d\phi, \quad (1)$$

where $F(\phi)$ is commonly known as the Faraday dispersion function (FDF), and the Faraday depth $\phi = 0.81 \int_L n_e B_{\parallel} dl \text{ rad m}^{-2}$, encodes the amount of Faraday rotation caused by a magnetoionic region of electron number density (n_e, cm^{-3}), line-of-sight magnetic field strength ($B_{\parallel}, \mu\text{G}$) and path length (l, pc).

Since the accuracy with which one can reconstruct $F(\phi)$ depends on the total wavelength-squared (λ^2) coverage, broadband polarisation observations at metre-wavelengths [1–4] can provide a typical improvement close to two orders of magnitude in Faraday depth accuracy compared to observations at centimetre-wavelengths [5–8]. However, the distribution of Faraday depths within the synthesised beam of the telescope can cause Faraday depolarisation, which makes the polarised signal fainter and more difficult to detect at long wavelengths [9].

In the context of the linearly polarised synchrotron emission from extragalactic radio-loud AGN, there are many possible contributions to the net observed Faraday rotation measure (RM) and Faraday depolarisation along the line of sight: from magnetised plasma internal to the radio emitting source [e.g. 10], from the turbulent magnetic fields associated with the hot ionised gas of the ambient group or cluster environment [e.g. 11], magnetoionic gas potentially associated with the filamentary large scale structure of the universe outside of clusters of galaxies [e.g. 12], the magnetised disks and halos of intervening galaxies [e.g. 13], the magnetised interstellar medium of the Milky Way [e.g. 14], and the time variable Faraday rotation contribution from the Earth’s ionosphere [e.g. 15]. The technique of Faraday tomography at radio wavebands, in addition to complementary observations at other wavebands, is a powerful tool to help isolate the contributions from all these regions of magnetoionic material.

The Faraday tomography results presented in this paper are based on data from the Low Frequency Array [LOFAR; 16], which is a radio interferometer capable of observing from 10 to 90 MHz with Low Band Antennas (LBA) and with High Band Antennas (HBA) from 110 to 250 MHz. The array is composed of ‘core stations’ that provide many short baselines up to $\sim 2 \text{ km}$, with more sparsely distributed ‘remote stations’ extending out to $\sim 100 \text{ km}$ from the core. There are also several international stations spread throughout Europe that can provide baselines of $\sim 1000 \text{ km}$. In particular, we make use of data from the ongoing LOFAR Two-Metre Sky Survey [LoTSS; 17], coupled with a preliminary catalog of linearly polarised sources [18]. The LoTSS survey aims to observe the whole northern sky above 0° Declination using the LOFAR High-Band Antenna (HBA) from 120 to 168 MHz.

The LoTSS observations are currently ongoing ($\sim 20\%$ complete), but there is an initial data release (LoTSS-DR1) of images and catalogs for 325,694 radio sources, detected above 5 times the noise level, covering 424 square degrees in the HETDEX Spring Field (RA: $10^{\text{h}}45^{\text{m}}$ to $15^{\text{h}}30^{\text{m}}$, Dec: 45° to 57°) [19]. The median sensitivity achieved in images of this region is $\sim 70 \mu\text{Jy beam}^{-1}$ at an angular resolution of $6''$, and diffuse, extended radio emission can be recovered with high fidelity. This provides a radio source surface density ~ 10 times higher than the best existing wide-area radio surveys and is comparable, for typical radio sources, to that expected from the planned ASKAP-EMU [20] and APERTIF [21] surveys. The LoTSS-DR1 release also comes with host galaxy identifications for 72% of the radio sources, and provides spectroscopic and photometric redshifts for 70% of these sources [22,23].

The polarisation catalog of this region found 92 polarised radio components (including 1 pulsar) after imaging at an angular resolution of $4.3'$ [18]. The focus of this paper is on the further investigation of the physical properties of the extragalactic polarised sources, making use of the calibrated data, images ($20''$ and $6''$ angular resolution) and value-added data products (radio source flux densities, host identifications, redshifts and source sizes) provided by the LoTSS-DR1 team.

2. Methods

The radio visibility data were calibrated using the PREFACTOR pipeline <https://github.com/lofar-astron/prefactor>, [15,17], which includes the ionospheric RM correction using RMEXTRACT <https://github.com/lofar-astron/RMextract> [53]. The estimated residual ionospheric RM correction

errors are ~ 0.1 to 0.3 rad m^{-2} [18,54]. To create Stokes Q and U cubes for each of the known polarised sources, the calibrated data were phase-shifted to the source position and averaged in time to reduce the data size using NDPPP [55] <https://support.astron.nl/LOFARImagingCookbook/>. The imaging software WSCLEAN [56] <https://sourceforge.net/projects/wsclean> was used to create channel images at 97.6 kHz resolution, with a minimum uv-range of 150λ and a maximum uv-range of $18k\lambda$, producing channel images with an angular resolution of $\sim 20''$. RM synthesis and RMCLEAN [57,58] were applied using PYRMSYNTH <https://github.com/mrbell/pyrmsynth> to create RM cubes with a range of $\pm 150 \text{ rad m}^{-2}$, sampled at 0.15 rad m^{-2} . The frequency range of 120 to 168 MHz, with a channel resolution of 97.6 kHz, provides a theoretical RM resolution of $\sim 1.1 \text{ rad m}^{-2}$, with a maximum scale of $\sim 1 \text{ rad m}^{-2}$, and a maximum detectable $|RM|$ of $\sim 450 \text{ rad m}^{-2}$. See [18], [28] and [39] for full details of the polarisation and Faraday rotation imaging and analysis methods, and [19, and references therein] for the data calibration and total intensity imaging procedures.

3. Results

3.1. Polarisation and Faraday rotation properties

Of the 91 extragalactic polarised sources found by [18] in a region of 570 deg^2 , 76 reside within the LoTSS-DR1 release area of 424 deg^2 [19]. It is the properties of these 76 sources that we present in this paper. This is only $\sim 0.4\%$ of the 19,233 radio sources with total flux densities greater than 10 mJy in the LoTSS-DR1 area. This emphasises the scarcity of polarised sources at 144 MHz, with a polarised source sky density of ~ 0.18 per square degree¹. The faintest polarised source was detected at $\sim 0.8 \text{ mJy beam}^{-1}$, and the brightest at $\sim 98 \text{ mJy beam}^{-1}$, with the majority of sources detected around $3.5 \text{ mJy beam}^{-1}$ (Figure 1, left panel).

A comparison with the RM of the same sources at 1.4 GHz [24], derived from the NRAO VLA Sky Survey [NVSS; 25], found that the LOFAR RM distribution was significantly narrower, likely due to the smaller errors, but also because sources near Faraday depths of 0 rad m^{-2} are missing due to contamination from uncorrected instrumental polarisation, making the LOFAR RM catalog incomplete [18]. The scatter in the difference between the RM values derived at 1.4 GHz and 144 MHz was also found to be significantly larger than expected from the derived errors, suggesting that regions of large RM variance seen at 1.4 GHz were completely depolarised at 144 MHz [18].

The integrated degree of polarisation at 144 MHz was also found to be lower than at 1.4 GHz in all cases, as expected due to the effect of Faraday depolarisation. However, even though the median degree of polarisation is ~ 8 times lower at 144 MHz than at 1.4 GHz [18], there are still 33 polarised sources found at 144 MHz that are not in the NVSS RM catalog [24]. This is due to a combination of small amounts of Faraday depolarisation for some sources, coupled with the LoTSS survey being much deeper than the NVSS. The noise level in Stokes Q and U in the LoTSS data at $4.3'$ is $\sim 0.15 \text{ mJy beam}^{-1}$ [18], which is ~ 10 times more sensitive than the NVSS for steep spectrum polarised radio sources (assuming a spectral index of -0.7). In addition, the NVSS RM catalog uses an $8\sigma_{QU}$ cutoff compared to the $\sim 5\sigma_{QU}$ threshold used at 144 MHz.

¹ Note that sources with an RM near zero ($\pm 2.5 \text{ rad m}^{-2}$) have been excluded due to contamination from instrumental polarisation in this range, and sources with an RM magnitude greater than $\sim 450 \text{ rad m}^{-2}$ are strongly affected by bandwidth depolarisation due to the channel spacing of 97.6 kHz. Furthermore, higher polarised source densities are expected at higher angular resolution, as demonstrated in a pointed observation of one of the fields by [39], which found a source density of ~ 0.3 per deg^2 at $\sim 18''$.




Figure 1. Left panel: histogram of the peak polarised intensity of the 76 polarised sources in LoTSS-DR1. **Right panel:** histograms of spectroscopic (solid line) and photometric (dashed line) redshifts of the polarised sources.

3.2. Radio luminosity, morphology and projected linear size

Based on the photometric [23] and spectroscopic [22] redshift catalogs of LoTSS-DR1, there are redshifts for $\sim 80\%$ of the 76 polarised sources (31 spectroscopic redshifts and 30 photometric redshifts), while 8 of the sources without a redshift also lack an optical identification of the host galaxy. All identified sources are consistent with being radio-loud AGN. The spectroscopic and photometric redshift distributions of the polarised sources are shown in Figure 1, right panel. They range from $z \sim 0.1$ to $z \sim 1.5$, with a median redshift of 0.5. This is similar to the redshift distribution of all radio-loud AGN in the LoTSS-DR1 area [26]. The radio luminosity distribution at 144 MHz ($L_{144\text{ MHz}}$) confirms the nature of these sources as powerful radio-loud AGN. It ranges from $3.6 \times 10^{24} \text{ W Hz}^{-1}$ to $1 \times 10^{28} \text{ W Hz}^{-1}$ with a median luminosity of $4 \times 10^{26} \text{ W Hz}^{-1}$ (Figure 2, left panel).

The majority of the polarised sources are resolved in the $6''$ total intensity images (62/76), with a median angular size for all polarised sources of $73''$. From inspection of the total intensity images, the majority of the polarised sources can be identified as FRII morphology radio galaxies (49/76), with the same luminosity range and median as above. One FRI radio galaxy and at least 8 blazars are identified, while the remaining compact or morphologically ambiguous sources require higher angular resolution observations to determine their source type. The projected linear size can be calculated for 49 of the resolved sources. The linear size distribution is shown in Figure 2 (right), with upper limits included for the unresolved sources. The median linear size of all sources is 415 kpc with a range from $< 50 \text{ kpc}$ up to 3.4 Mpc. As a large fraction of sources (10/76) have a linear size $> 1 \text{ Mpc}$, polarisation observations at low frequencies can be useful in selecting for ‘giant’ radio galaxies. In addition to the main population of FRII sources (with the median linear size for the 39 of those FRIIs with redshifts being 720 kpc), there is also a smaller population of compact sources, the majority of which can be identified as blazars [27].




Figure 2. Left panel: histograms of the radio luminosity at 144 MHz ($L_{144\text{MHz}}$) for the polarised sources with spectroscopic (solid line) and photometric (dashed line) redshifts. The radio luminosities are consistent with all sources being radio-loud AGN. Right panel: histograms of the projected linear size of resolved sources with spectroscopic (solid line) and photometric (blue dashed line) redshifts. The red dashed line shows the histogram of upper limits on the projected linear size of unresolved sources with redshift measurements.

3.3. In depth study of the Faraday rotation from the largest FR II radio galaxy in the sample

The source with the largest projected linear size of 3.4 Mpc (and angular size of $11'$), ILT J123459.82+531851.0, at a redshift of $z \sim 0.34$ has been investigated in detail [see 28, for a full description]. Here we present a brief summary of the results from that study. The key result was based on the polarisation and RM distribution of the lobes at an angular resolution of $20''$. A mean RM difference of $\sim 2.5 \text{ rad m}^{-2}$ was found between the regions of polarised emission in the opposite lobes of this FR II radio galaxy. This RM difference was investigated in the context of the potential contribution of intergalactic magnetic fields (IGMF) in foreground large-scale-structure (LSS) filaments.

Due to the large linear size of the source, the polarised emission from the north-west and south-east lobes probes significantly different cosmic lines of sight, such that an excess of up to three LSS filaments were estimated to be probed by the polarised emission from North-West lobe. The LSS filaments were identified from a catalog constructed using spectroscopic observations of galaxies in the Sloan Digital Sky Survey (SDSS) [29,30]. Associating the entire RM difference to these LSS filaments implies a gas density-weighted magnetic field strength of $\sim 0.3 \mu\text{G}$. However, from comparisons with cosmological MHD simulations [31] of the expected RM signal from LSS filaments, it was found that an RM difference as large as 2.5 rad m^{-2} , on 3.4 Mpc scales, had a low probability ($\sim 5\%$) of occurring, for IGMF strengths of tens of nG. The magnetic field in the simulation was amplified from a initial magnetic field of 1 nG seeded at early cosmological epochs, which is close to the upper limit for such fields provided by the Planck satellite [32].

Furthermore, variations in the Faraday rotation of the Milky Way on scales of $11'$ likely contribute significantly to the observed RM difference between the lobes. The best available reconstructions of the Galactic RM [33] have a resolution of only ~ 1 degree, although RM structure function analyses suggest RM variations of several rad m^{-2} are possible on arcminute scales [14,34,35]. This highlights the need for the denser RM grids that are expected from the VLASS [36] and ASKAP-POSSUM [37], in the near future, to provide a better understanding of the Galactic RM variations on scales less than 1 degree [38].

3.4. High resolution Faraday rotation imaging with LOFAR

In the preliminary catalog, polarised sources from the HETDEX region of the LoTSS survey data were imaged at an angular resolution of $4.3''$ [18]. One of the reasons for imaging at low angular resolution is due to the challenging computational requirements for processing and storing polarisation data products at high angular resolution across the entire LoTSS area. However, smaller datasets can be made for a list of individual sources to be imaged at a higher angular resolution [39]. Thus, we have imaged the polarisation and RM distribution of the LoTSS polarised sources at $20''$ resolution. Figure 3 shows some examples of these high resolution images, where the grey total intensity contours are at the same resolution as the RM image ($20''$), while the black total intensity contours are at $6''$.




Figure 3. Total intensity and Faraday rotation images of four polarised sources in LoTSS-DR1. The grey contours and colourscale Faraday rotation measure (RM) images are at $20''$ resolution (indicated by the circle in the bottom left corner of the images), while the black contours are at $6''$ resolution. Contours increase by factors of two, with the low contour for each source quoted below. The RM images are clipped at 8 times the noise level in Stokes Q and U . **Top left:** The FR II radio galaxy ILT J105715.33+484108.6, which has a projected linear size of ~ 1.9 Mpc, and both hotspots are polarised. Low contour: $20''$: 5 mJy beam^{-1} , $6''$: $0.35 \text{ mJy beam}^{-1}$. **Top right:** The FR II radio galaxy ILT J134102.90+491609.1, ~ 1 Mpc in size, which is the most common type of polarised source found in the LoTSS-DR1 data. Low contour: $20''$: 5 mJy beam^{-1} , $6''$: $0.35 \text{ mJy beam}^{-1}$. **Bottom left:** The only polarised source identified as an FRI radio galaxy in the current sample, with only the inner jet region detected in polarisation. Low contour: $20''$: 5 mJy beam^{-1} , $6''$: $0.35 \text{ mJy beam}^{-1}$. **Bottom right:** The blazar ILTJ141945.53+542314.4, with both the core and the extension to the west being polarised. Low contour: $20''$: 10 mJy beam^{-1} , $6''$: $3.5 \text{ mJy beam}^{-1}$.

Figure 3 (top left) shows a large angular ($\sim 500''$) and linear size (~ 1.9 Mpc) FR II radio galaxy ($L_{144\text{MHz}} \sim 7 \times 10^{25} \text{ W Hz}^{-1}$), with the polarised hotspots displaying a difference in RM of $\sim 2.1 \text{ rad m}^{-2}$. This demonstrates the impressive capability of the LoTSS survey to image both compact and extended regions of large radio galaxies. Another advantage of imaging at higher angular resolution is that only the south-west hotspot was previously detected in polarisation [18]. Figure 3 (top right) shows another example of an FR II radio galaxy ($L_{144\text{MHz}} \sim 5 \times 10^{26} \text{ W Hz}^{-1}$), which is the most common morphology of the polarised sources currently found at 144 MHz [18], with 64% of polarised sources being identified with an FR II. Generally the polarised hotspots have simple Faraday spectra with only a single peak in the FDF (e.g. Figure 4).




Figure 4. Examples of the absolute value of the Faraday dispersion function (FDF) for the polarised hotspots of two FR II radio galaxies. **Left panel:** FDF for the south-west hotspot of ILT J105715.33+484108.6 (Figure 3, top left), with the red cross indicating the real polarised peak of $\sim 19.3 \text{ mJy beam}^{-1}$ at a Faraday depth of $\sim 16.85 \text{ rad m}^{-2}$. The other peak in the FDF near 0 rad m^{-2} is due to instrumental polarisation at a level of $\sim 2\%$ of Stokes I . **Right panel:** FDF of the south hotspot of ILT J134102.90+491609.1 (Figure 3, top right), with a peak of $\sim 5.9 \text{ mJy beam}^{-1}$ at a Faraday depth of $\sim 9.9 \text{ rad m}^{-2}$.

However, other types of radio-loud AGN are also present, for example, an FRI radio galaxy ($L_{144\text{MHz}} \sim 8 \times 10^{24} \text{ W Hz}^{-1}$) with polarised emission detected from the inner jet region only (Figure 3, bottom left). Also, several of the compact polarised sources are associated with blazars. Figure 3 (bottom right) shows a BL Lac object [40] with both the core and an extension $\sim 200 \text{ kpc}$ to the west detected in polarisation. Indeed the kpc-scale extension exhibits some evidence for Faraday complexity with multiple peaks in the Faraday dispersion function at that location (Figure 5).




Figure 5. Example FDFs from the blazar ILT J141945.53+542314.4. **Left panel:** Simple FDF of the blazar core with a polarised intensity of ~ 3 mJy beam $^{-1}$ at a Faraday depth of 17.5 rad m $^{-2}$. **Right panel:** Complex FDF of the western extension of the blazar, with a peak of ~ 1.0 mJy beam $^{-1}$ at 17.7 rad m $^{-2}$ and a second peak of ~ 0.88 mJy beam $^{-1}$ at 16.2 rad m $^{-2}$.

4. Discussion

4.1. The nature of extragalactic polarised sources at low frequencies

The most striking feature of the polarised sources is that the majority are large FR II radio galaxies, with a median linear size of 710 kpc. Furthermore, the polarised emission is mainly associated with the hotspots of these FR IIs. This means that the regions of detected polarisation extend well beyond their host galaxy environment, and likely well into the outskirts of the galaxy group or cluster in which the host galaxy resides. As the ionised gas density and magnetic field strength are known to decrease with radius in galaxy groups and clusters [41,42], this means that the effect of Faraday depolarisation is also much lower than near the centre [43]. As even small RM variations of ~ 1 rad m $^{-2}$ across the emission region can be sufficient to depolarise a source below the detection limits at LOFAR frequencies, it is not surprising that the LoTSS polarised sources are so physically large. In addition, FR IIs are also known to typically inhabit less dense environments than FR Is [44–46].

Large linear size is likely not the only consideration here, as there are also large FRI radio galaxies [47]. However, in these sources the brightness decreases with increasing distance from the inner jet out into the diffuse lobes [48], compared to FR II radio galaxies where the brightest regions are near the outer edges of the source. Furthermore, even when imaged at $20''$, the compact nature of FR II hotspots means that the emission probes a smaller Faraday depth volume than, for example, the extended lobes of an FRI radio galaxy. The large angular size of the polarised sources helps mitigate against wavelength-independent depolarisation, where variations of the intrinsic magnetic field and total intensity structure make the polarisation more difficult to detect in integrated measurements. Of course it also helps to resolve more of the variations in Faraday rotation across the emission region. Figure 3 shows three such examples, where an additional polarised component is detected in the $20''$ images compared to the $4.3'$ images. More sensitive and higher resolution observations may help detect the polarised emission from extended regions of FRI radio galaxies, unless internal Faraday rotation is strongly depolarising the emission in these sources.

4.2. Future prospects

The ability to study the physical properties of extragalactic polarised sources and the intervening intergalactic medium is significantly enhanced by the unique capability of LOFAR to image at high angular resolution ($6''$) at low frequencies, which is also crucial for the identification of the host galaxy. Soon, the WEAVE-LOFAR survey [49] will begin measuring redshifts for over a million of the LoTSS

detected radio sources. While only a small fraction of these sources will be polarised (possibly of order 10,000), it will still enable major advances in the field of cosmic magnetism, as at present, it would be the largest RM catalog with associated redshifts. Such large samples, coupled with the unrivalled Faraday depth accuracy, will enable the application of advanced statistical analyses to isolate the contributions of the different magnetoionic media along the line of sight to the net Faraday rotation and depolarisation [33,50–52].

LOFAR [16] is the Low Frequency Array designed and constructed by ASTRON. It has observing, data processing, and data storage facilities in several countries, that are owned by various parties (each with their own funding sources), and that are collectively operated by the ILT foundation under a joint scientific policy. The ILT resources have benefitted from the following recent major funding sources: CNRS-INSU, Observatoire de Paris and Université d'Orléans, France; BMBF, MIWF-NRW, MPG, Germany; Science Foundation Ireland (SFI), Department of Business, Enterprise and Innovation (DBEI), Ireland; NWO, The Netherlands; The Science and Technology Facilities Council, UK; Ministry of Science and Higher Education, Poland. SPO and MB acknowledge financial support from the Deutsche Forschungsgemeinschaft (DFG) under grant BR2026/23. Part of this work was carried out on the Dutch national e-infrastructure with the support of the SURF Cooperative through grant e-infra 160022 & 160152. The LOFAR software and dedicated reduction packages on https://github.com/apmechev/GRID_LRT were deployed on the e-infrastructure by the LOFAR e-infragroup, consisting of J. B. R. Oonk (ASTRON & Leiden Observatory), A. P. Mechev (Leiden Observatory) and T. Shimwell (ASTRON) with support from N. Danezi (SURFsara) and C. Schrijvers (SURFsara). This research has made use of data analysed using the University of Hertfordshire high-performance computing facility (<http://uhhpc.herts.ac.uk/>) and the LOFAR-UK computing facility located at the University of Hertfordshire and supported by STFC [ST/P000096/1].

Acknowledgments: This paper is based (in part) on data obtained with the International LOFAR Telescope (ILT) under project codes LC2_038 and LC3_008. This research made use of Astropy, a community-developed core Python package for astronomy [59] hosted at <http://www.astropy.org/>, of Matplotlib [60], of APLpy [61], an open-source astronomical plotting package for Python hosted at <http://aplpy.github.com/>, and of TOPCAT, an interactive graphical viewer and editor for tabular data [62].

Conflicts of Interest: The authors declare no conflict of interest. The founding sponsors had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript, or in the decision to publish the results.

Abbreviations

The following abbreviations are used in this manuscript:

RM	Rotation measure
IGMF	Intergalactic magnetic field
GRM	Galactic rotation measure
FDF	Faraday dispersion function
FRI	Fanaroff Riley I
FRII	Fanaroff Riley II
AGN	Active Galactic Nuclei
BL Lac	BL Lacertae
LSS	Large scale structure
LOFAR	Low Frequency Array
HBA	High Band Antenna
LBA	Low Band Antenna
ILT	International LOFAR Telescope
LoTSS	LOFAR Two-metre Sky Survey
DR1	Data Release 1
HETDEX	Hobby-Eberly Telescope Dark Energy Experiment
WEAVE	William Herschel Telescope Enhanced Area Velocity Explorer
APERTIF	Aperture Tile in Focus
ASKAP	Australian Square Kilometre Array Pathfinder
EMU	Evolutionary Map of the Universe
POSSUM	Polarisation Survey of the Universe's Magnetism
NRAO	National Radio Astronomy Observatory
VLA	Very Large Array
NVSS	NRAO VLA Sky Survey
VLASS	VLA Sky Survey
SDSS	Sloan Digital Sky Survey

References

1. Jelić, V.; de Bruyn, A.G.; Pandey, V.N.; Mevius, M.; Haverkorn, M.; Brentjens, M.A.; Koopmans, L.V.E.; Zaroubi, S.; Abdalla, F.B.; Asad, K.M.B.; Bus, S.; Chapman, E.; Ciardi, B.; Fernandez, E.R.; Ghosh, A.; Harker, G.; Iliev, I.T.; Jensen, H.; Kazemi, S.; Mellema, G.; Offringa, A.R.; Patil, A.H.; Vedantham, H.K.; Yatawatta, S. Linear polarization structures in LOFAR observations of the interstellar medium in the 3C 196 field. *A&A* **2015**, *583*, A137, [1508.06650]. doi:10.1051/0004-6361/201526638.
2. Lenc, E.; Gaensler, B.M.; Sun, X.H.; Sadler, E.M.; Willis, A.G.; Barry, N.; Beardsley, A.P.; Bell, M.E.; Bernardi, G.; Bowman, J.D.; Briggs, F.; Callingham, J.R.; Cappallo, R.J.; Carroll, P.; Corey, B.E.; de Oliveira-Costa, A.; Deshpande, A.A.; Dillon, J.S.; Dwarkanath, K.S.; Emrich, D.; Ewall-Wice, A.; Feng, L.; For, B.Q.; Goeke, R.; Greenhill, L.J.; Hancock, P.; Hazelton, B.J.; Hewitt, J.N.; Hindson, L.; Hurley-Walker, N.; Johnston-Hollitt, M.; Jacobs, D.C.; Kapińska, A.D.; Kaplan, D.L.; Kasper, J.C.; Kim, H.S.; Kratzenberg, E.; Line, J.; Loeb, A.; Lonsdale, C.J.; Lynch, M.J.; McKinley, B.; McWhirter, S.R.; Mitchell, D.A.; Morales, M.F.; Morgan, E.; Morgan, J.; Murphy, T.; Neben, A.R.; Oberoi, D.; Offringa, A.R.; Ord, S.M.; Paul, S.; Pindor, B.; Pober, J.C.; Prabu, T.; Procopio, P.; Riding, J.; Rogers, A.E.E.; Roshi, A.; Udaya Shankar, N.; Sethi, S.K.; Srivani, K.S.; Staveley-Smith, L.; Subrahmanyan, R.; Sullivan, I.S.; Tegmark, M.; Thyagarajan, N.; Tingay, S.J.; Trott, C.; Waterson, M.; Wayth, R.B.; Webster, R.L.; Whitney, A.R.; Williams, A.; Williams, C.L.; Wu, C.; Wyithe, J.S.B.; Zheng, Q. Low-frequency Observations of Linearly Polarized Structures in the Interstellar Medium near the South Galactic Pole. *ApJ* **2016**, *830*, 38, [1607.05779]. doi:10.3847/0004-637X/830/1/38.
3. O'Sullivan, S.P.; Lenc, E.; Anderson, C.S.; Gaensler, B.M.; Murphy, T. Faraday rotation at low frequencies: magnetoionic material of the large FRII radio galaxy PKS J0636-2036. *MNRAS* **2018**, *475*, 4263–4277, [1801.02452]. doi:10.1093/mnras/sty171.
4. Riseley, C.J.; Lenc, E.; Van Eck, C.L.; Heald, G.; Gaensler, B.M.; Anderson, C.S.; Hancock, P.J.; Hurley-Walker, N.; Sridhar, S.S.; White, S.V. The Polarised GLEAM Survey (POGS) I: First Results from a Low-Frequency Radio Linear Polarisation Survey of the Southern Sky. *ArXiv e-prints* **2018**, [1809.09327].
5. O'Sullivan, S.P.; Brown, S.; Robishaw, T.; Schnitzeler, D.H.F.M.; McClure-Griffiths, N.M.; Feain, I.J.; Taylor, A.R.; Gaensler, B.M.; Landecker, T.L.; Harvey-Smith, L.; Carretti, E. Complex Faraday depth

- structure of Active Galactic Nuclei as revealed by broadband radio polarimetry. *MNRAS* **2012**, *421*, 3300, [arXiv:astro-ph.CO/1201.3161].
6. Anderson, C.S.; Gaensler, B.M.; Feain, I.J. A Study of Broadband Faraday Rotation and Polarization Behavior over 1.3–10 GHz in 36 Discrete Radio Sources. *ApJ* **2016**, *825*, 59, [arXiv:astro-ph.HE/1604.01403]. doi:10.3847/0004-637X/825/1/59.
 7. O'Sullivan, S.P.; Purcell, C.R.; Anderson, C.S.; Farnes, J.S.; Sun, X.H.; Gaensler, B.M. Broad-band, radio spectro-polarimetric study of 100 radiative-mode and jet-mode AGN. *MNRAS* **2017**, *469*, 4034–4062, [1705.00102]. doi:10.1093/mnras/stx1133.
 8. Pasetto, A.; Carrasco-González, C.; O'Sullivan, S.; Basu, A.; Bruni, G.; Kraus, A.; Curiel, S.; Mack, K.H. Broadband radio spectro-polarimetric observations of high-Faraday-rotation-measure AGN. *A&A* **2018**, *613*, A74, [1801.09731]. doi:10.1051/0004-6361/201731804.
 9. Farnsworth, D.; Rudnick, L.; Brown, S. Integrated Polarization of Sources at $\lambda \sim 1$ m and New Rotation Measure Ambiguities. *AJ* **2011**, *141*, 191, [arXiv:astro-ph.CO/1103.4149]. doi:10.1088/0004-6256/141/6/191.
 10. Anderson, C.S.; Gaensler, B.M.; Heald, G.H.; O'Sullivan, S.P.; Kaczmarek, J.F.; Feain, I.J. Broadband Radio Polarimetry of Fornax A. I. Depolarized Patches Generated by Advected Thermal Material from NGC 1316. *ApJ* **2018**, *855*, 41, [arXiv:astro-ph.HE/1802.04812]. doi:10.3847/1538-4357/aaaec0.
 11. Guidetti, D.; Laing, R.A.; Croston, J.H.; Bridle, A.H.; Parma, P. The magnetized medium around the radio galaxy B2 0755+37: an interaction with the intragroup gas. *MNRAS* **2012**, *423*, 1335–1350, [arXiv:astro-ph.CO/1203.4582]. doi:10.1111/j.1365-2966.2012.20961.x.
 12. Akahori, T.; Ryu, D. Faraday Rotation Measure due to the Intergalactic Magnetic Field. II. The Cosmological Contribution. *ApJ* **2011**, *738*, 134, [1107.0142]. doi:10.1088/0004-637X/738/2/134.
 13. Farnes, J.S.; O'Sullivan, S.P.; Corrigan, M.E.; Gaensler, B.M. Faraday Rotation from Magnesium II Absorbers toward Polarized Background Radio Sources. *ApJ* **2014**, *795*, 63, [1406.2526]. doi:10.1088/0004-637X/795/1/63.
 14. Haverkorn, M.; Gaensler, B.M.; McClure-Griffiths, N.M.; Dickey, J.M.; Green, A.J. Magnetic Fields and Ionized Gas in the Inner Galaxy: An Outer Scale for Turbulence and the Possible Role of H II Regions. *ApJ* **2004**, *609*, 776–784, [astro-ph/0403655]. doi:10.1086/421341.
 15. de Gasperin, F.; Mevius, M.; Rafferty, D.A.; Intema, H.T.; Fallows, R.A. The effect of the ionosphere on ultra-low-frequency radio-interferometric observations. *A&A* **2018**, *615*, A179, [arXiv:astro-ph.IM/1804.07947]. doi:10.1051/0004-6361/201833012.
 16. van Haarlem, M.P.; Wise, M.W.; Gunst, A.W.; Heald, G.; McKean, J.P.; Hessels, J.W.T.; de Bruyn, A.G.; Nijboer, R.; Swinbank, J.; Fallows, R.; Brentjens, M.; Nelles, A.; Beck, R.; Falcke, H.; Fender, R.; Hörandel, J.; Koopmans, L.V.E.; Mann, G.; Miley, G.; Röttgering, H.; Stappers, B.W.; Wijers, R.A.M.J.; Zaroubi, S.; van den Akker, M.; Alexov, A.; Anderson, J.; Anderson, K.; van Ardenne, A.; Arts, M.; Asgekar, A.; Avruch, I.M.; Batejat, F.; Bähren, L.; Bell, M.E.; Bell, M.R.; van Bemmelen, I.; Bennema, P.; Bentum, M.J.; Bernardi, G.; Best, P.; Birzan, L.; Bonafede, A.; Boonstra, A.J.; Braun, R.; Bregman, J.; Breitling, F.; van de Brink, R.H.; Broderick, J.; Broekema, P.C.; Brouw, W.N.; Brüggen, M.; Butcher, H.R.; van Cappellen, W.; Ciardi, B.; Coenen, T.; Conway, J.; Coolen, A.; Corstanje, A.; Damstra, S.; Davies, O.; Deller, A.T.; Dettmar, R.J.; van Diepen, G.; Dijkstra, K.; Donker, P.; Doorduyn, A.; Dromer, J.; Drost, M.; van Duin, A.; Eislöffel, J.; van Enst, J.; Ferrari, C.; Frieswijk, W.; Gankema, H.; Garrett, M.A.; de Gasperin, F.; Gerbers, M.; de Geus, E.; Grießmeier, J.M.; Grit, T.; Gruppen, P.; Hamaker, J.P.; Hassall, T.; Hoeft, M.; Holties, H.A.; Horneffer, A.; van der Horst, A.; van Houwelingen, A.; Huijgen, A.; Iacobelli, M.; Intema, H.; Jackson, N.; Jelic, V.; de Jong, A.; Juetten, E.; Kant, D.; Karastergiou, A.; Koers, A.; Kollen, H.; Kondratiev, V.I.; Kooistra, E.; Koopman, Y.; Koster, A.; Kuniyoshi, M.; Kramer, M.; Kuper, G.; Lambropoulos, P.; Law, C.; van Leeuwen, J.; Lemaître, J.; Loose, M.; Maat, P.; Macario, G.; Markoff, S.; Masters, J.; McFadden, R.A.; McKay-Bukowski, D.; Meijering, H.; Meulman, H.; Mevius, M.; Middelberg, E.; Millenaar, R.; Miller-Jones, J.C.A.; Mohan, R.N.; Mol, J.D.; Morawietz, J.; Morganti, R.; Mulcahy, D.D.; Mulder, E.; Munk, H.; Nieuwenhuis, L.; van Nieuwpoort, R.; Noordam, J.E.; Norden, M.; Noutsos, A.; Offringa, A.R.; Olofsson, H.; Omar, A.; Orrú, E.; Overeem, R.; Paas, H.; Pandey-Pommier, M.; Pandey, V.N.; Pizzo, R.; Polatidis, A.; Rafferty, D.; Rawlings, S.; Reich, W.; de Reijer, J.P.; Reitsma, J.; Renting, G.A.; Riemers, P.; Rol, E.; Romein, J.W.; Roosjen, J.; Ruiter, M.; Scaife, A.; van der Schaaf, K.; Scheers, B.; Schellart, P.; Schoenmakers, A.; Schoonderbeek, G.; Serylak, M.; Shulevski, A.; Sluman, J.; Smirnov, O.; Sobey, C.; Spreeuw, H.; Steinmetz, M.; Sterks,

- C.G.M.; Stiepel, H.J.; Stuurwold, K.; Tagger, M.; Tang, Y.; Tasse, C.; Thomas, I.; Thoudam, S.; Toribio, M.C.; van der Tol, B.; Usov, O.; van Veelen, M.; van der Veen, A.J.; ter Veen, S.; Verbiest, J.P.W.; Vermeulen, R.; Vermaas, N.; Vocks, C.; Vogt, C.; de Vos, M.; van der Wal, E.; van Weeren, R.; Weggemans, H.; Weltevrede, P.; White, S.; Wijnholds, S.J.; Wilhelmsson, T.; Wucknitz, O.; Yatawatta, S.; Zarka, P.; Zensus, A.; van Zwieten, J. LOFAR: The LOw-Frequency ARray. *A&A* **2013**, *556*, A2, [arXiv:astro-ph.IM/1305.3550]. doi:10.1051/0004-6361/201220873.
17. Shimwell, T.W.; Röttgering, H.J.A.; Best, P.N.; Williams, W.L.; Dijkema, T.J.; de Gasperin, F.; Hardcastle, M.J.; Heald, G.H.; Hoang, D.N.; Horneffer, A.; Intema, H.; Mahony, E.K.; Mandal, S.; Mechev, A.P.; Morabito, L.; Oonk, J.B.R.; Rafferty, D.; Retana-Montenegro, E.; Sabater, J.; Tasse, C.; van Weeren, R.J.; Brügger, M.; Brunetti, G.; Chyży, K.T.; Conway, J.E.; Haverkorn, M.; Jackson, N.; Jarvis, M.J.; McKean, J.P.; Miley, G.K.; Morganti, R.; White, G.J.; Wise, M.W.; van Bemmell, I.M.; Beck, R.; Brienza, M.; Bonafede, A.; Calistro Rivera, G.; Cassano, R.; Clarke, A.O.; Cseh, D.; Deller, A.; Drabent, A.; van Driel, W.; Engels, D.; Falcke, H.; Ferrari, C.; Fröhlich, S.; Garrett, M.A.; Harwood, J.J.; Heesen, V.; Hoeft, M.; Horellou, C.; Israel, F.P.; Kapińska, A.D.; Kunert-Bajraszewska, M.; McKay, D.J.; Mohan, N.R.; Orrú, E.; Pizzo, R.F.; Prandoni, I.; Schwarz, D.J.; Shulevski, A.; Sipior, M.; Smith, D.J.B.; Sridhar, S.S.; Steinmetz, M.; Stroe, A.; Varenus, E.; van der Werf, P.P.; Zensus, J.A.; Zwart, J.T.L. The LOFAR Two-metre Sky Survey. I. Survey description and preliminary data release. *A&A* **2017**, *598*, A104, [arXiv:astro-ph.IM/1611.02700]. doi:10.1051/0004-6361/201629313.
 18. Van Eck, C.L.; Haverkorn, M.; Alves, M.I.R.; Beck, R.; Best, P.; Carretti, E.; Chyży, K.T.; Farnes, J.S.; Ferrière, K.; Hardcastle, M.J.; Heald, G.; Horellou, C.; Iacobelli, M.; Jelić, V.; Mulcahy, D.D.; O'Sullivan, S.P.; Polderman, I.M.; Reich, W.; Riseley, C.J.; Röttgering, H.; Schnitzeler, D.H.F.M.; Shimwell, T.W.; Vacca, V.; Vink, J.; White, G.J. Polarized point sources in the LOFAR Two-meter Sky Survey: A preliminary catalog. *A&A* **2018**, *613*, A58, [1801.04467]. doi:10.1051/0004-6361/201732228.
 19. Shimwell, T.W.; Röttgering, H.J.A.; Best, P.N.; Williams, W.L.; Dijkema, T.J.; de Gasperin, F.; Hardcastle, M.J.; Heald, G.H.; Hoang, D.N.; Horneffer, A.; Intema, H.; Mahony, E.K.; Mandal, S.; Mechev, A.P.; Morabito, L.; Oonk, J.B.R.; Rafferty, D.; Retana-Montenegro, E.; Sabater, J.; Tasse, C.; van Weeren, R.J.; Brügger, M.; Brunetti, G.; Chyży, K.T.; Conway, J.E.; Haverkorn, M.; Jackson, N.; Jarvis, M.J.; McKean, J.P.; Miley, G.K.; Morganti, R.; White, G.J.; Wise, M.W.; van Bemmell, I.M.; Beck, R.; Brienza, M.; Bonafede, A.; Calistro Rivera, G.; Cassano, R.; Clarke, A.O.; Cseh, D.; Deller, A.; Drabent, A.; van Driel, W.; Engels, D.; Falcke, H.; Ferrari, C.; Fröhlich, S.; Garrett, M.A.; Harwood, J.J.; Heesen, V.; Hoeft, M.; Horellou, C.; Israel, F.P.; Kapińska, A.D.; Kunert-Bajraszewska, M.; McKay, D.J.; Mohan, N.R.; Orrú, E.; Pizzo, R.F.; Prandoni, I.; Schwarz, D.J.; Shulevski, A.; Sipior, M.; Smith, D.J.B.; Sridhar, S.S.; Steinmetz, M.; Stroe, A.; Varenus, E.; van der Werf, P.P.; Zensus, J.A.; Zwart, J.T.L. The LOFAR Two-metre Sky Survey – II. First Data Release. *A&A* **2019**, *LOFAR Special Issue*, in press.
 20. Norris, R. Survey Science with ASKAP: EMU - Evolutionary Map of the Universe. American Astronomical Society Meeting Abstracts #215, 2010, Vol. 36, *Bulletin of the American Astronomical Society*, p. 1127.
 21. Verheijen, M.A.W.; Oosterloo, T.A.; van Cappellen, W.A.; Bakker, L.; Ivashina, M.V.; van der Hulst, J.M. Apertif, a focal plane array for the WSRT. The Evolution of Galaxies Through the Neutral Hydrogen Window; Minchin, R.; Momjian, E., Eds., 2008, Vol. 1035, *American Institute of Physics Conference Series*, pp. 265–271, [0806.0234]. doi:10.1063/1.2973599.
 22. Williams, W.L.; Duncan, K.J.; Hardcastle, M.J.; Shimwell, T.W. spec zs. *A&A* **2019**, *LOFAR Special Issue*, in press.
 23. Duncan, K.J.; Williams, W.L.; Hardcastle, M.J.; Shimwell, T.W. photo zs. *A&A* **2019**, *LOFAR Special Issue*, in press.
 24. Taylor, A.R.; Stil, J.M.; Sunstrum, C. A Rotation Measure Image of the Sky. *ApJ* **2009**, *702*, 1230–1236. doi:10.1088/0004-637X/702/2/1230.
 25. Condon, J.J.; Cotton, W.D.; Greisen, E.W.; Yin, Q.F.; Perley, R.A.; Taylor, G.B.; Broderick, J.J. The NRAO VLA Sky Survey. *AJ* **1998**, *115*, 1693–1716. doi:10.1086/300337.
 26. Hardcastle, M.J.; Williams, W.L.; Best, P.N.; Croston, J.H. LoTSS AGN. *A&A* **2019**, *LOFAR Special Issue*, in press.
 27. Mooney. LoTSS blazars. *A&A* **2019**, *LOFAR Special Issue*, in press.
 28. O'Sullivan, S.P. IGMF. *A&A* **2019**, *LOFAR Special Issue*, in press.

29. Chen, Y.C.; Ho, S.; Freeman, P.E.; Genovese, C.R.; Wasserman, L. Cosmic web reconstruction through density ridges: method and algorithm. *MNRAS* **2015**, *454*, 1140–1156, [[1501.05303](#)]. doi:10.1093/mnras/stv1996.
30. Chen, Y.C.; Ho, S.; Brinkmann, J.; Freeman, P.E.; Genovese, C.R.; Schneider, D.P.; Wasserman, L. Cosmic web reconstruction through density ridges: catalogue. *MNRAS* **2016**, *461*, 3896–3909, [[1509.06443](#)]. doi:10.1093/mnras/stw1554.
31. Vazza, F.; Brüggén, M.; Gheller, C.; Wang, P. On the amplification of magnetic fields in cosmic filaments and galaxy clusters. *MNRAS* **2014**, *445*, 3706–3722, [[1409.2640](#)]. doi:10.1093/mnras/stu1896.
32. Planck Collaboration.; Ade, P.A.R.; Aghanim, N.; Arnaud, M.; Arroja, F.; Ashdown, M.; Aumont, J.; Baccigalupi, C.; Ballardini, M.; Banday, A.J.; et al.. Planck 2015 results. XIX. Constraints on primordial magnetic fields. *A&A* **2016**, *594*, A19, [[1502.01594](#)]. doi:10.1051/0004-6361/201525821.
33. Oppermann, N.; Junklewitz, H.; Greiner, M.; Enßlin, T.A.; Akahori, T.; Carretti, E.; Gaensler, B.M.; Goobar, A.; Harvey-Smith, L.; Johnston-Hollitt, M.; Pratley, L.; Schnitzeler, D.H.F.M.; Stil, J.M.; Vacca, V. Estimating extragalactic Faraday rotation. *A&A* **2015**, *575*, A118, [[arXiv:astro-ph.IM/1404.3701](#)]. doi:10.1051/0004-6361/201423995.
34. Mao, S.A.; Gaensler, B.M.; Haverkorn, M.; Zweibel, E.G.; Madsen, G.J.; McClure-Griffiths, N.M.; Shukurov, A.; Kronberg, P.P. A Survey of Extragalactic Faraday Rotation at High Galactic Latitude: The Vertical Magnetic Field of the Milky Way Toward the Galactic Poles. *ApJ* **2010**, *714*, 1170–1186, [[arXiv:astro-ph.GA/1003.4519](#)]. doi:10.1088/0004-637X/714/2/1170.
35. Stil, J.M.; Taylor, A.R.; Sunstrum, C. Structure in the Rotation Measure Sky. *ApJ* **2011**, *726*, 4, [[arXiv:astro-ph.GA/1010.5299](#)]. doi:10.1088/0004-637X/726/1/4.
36. Myers, S.T.; Law, C.J.; Baum, S.A.; Chandler, C.J.; Chatterjee, S.; Lacy, M.; Murphy, E.J.; VLASS Survey Science Group. The VLA Sky Survey (VLASS): Technical Implementation Plans and Progress. AAS #227, 2016, Vol. 227, *American Astronomical Society Meeting Abstracts*, p. 324.08.
37. Gaensler, B.M.; Landecker, T.L.; Taylor, A.R.; POSSUM Collaboration. BAAS, 2010, Vol. 42, p. 470.13.
38. Boulanger, F.; Enßlin, T.; Fletcher, A.; Girichides, P.; Hackstein, S.; Haverkorn, M.; Hörandel, J.R.; Jaffe, T.; Jasche, J.; Kachelrieß, M.; Kotera, K.; Pfrommer, C.; Rachen, J.P.; Rodrigues, L.F.S.; Ruiz-Granados, B.; Seta, A.; Shukurov, A.; Sigl, G.; Steininger, T.; Vacca, V.; van der Velden, E.; van Vliet, A.; Wang, J. IMAGINE: a comprehensive view of the interstellar medium, Galactic magnetic fields and cosmic rays. *JCAP* **2018**, *8*, 049, [[1805.02496](#)]. doi:10.1088/1475-7516/2018/08/049.
39. Neld, A.; Horellou, C.; Mulcahy, D.D.; Beck, R.; Bourke, S.; Carozzi, T.D.; Chyży, K.T.; Conway, J.E.; Farnes, J.S.; Fletcher, A.; Haverkorn, M.; Heald, G.; Horneffer, A.; Nikiel-Wroczyński, B.; Paladino, R.; Sridhar, S.S.; Van Eck, C.L. Reliable detection and characterization of low-frequency polarized sources in the LOFAR M51 field. *A&A* **2018**, *617*, 136.
40. Plotkin, R.M.; Anderson, S.F.; Hall, P.B.; Margon, B.; Voges, W.; Schneider, D.P.; Stinson, G.; York, D.G. a Large Sample of BL Lac Objects from the SDSS and First. *AJ* **2008**, *135*, 2453–2469, [[arXiv:astro-ph/0803.4219](#)]. doi:10.1088/0004-6256/135/6/2453.
41. Laing, R.A.; Bridle, A.H.; Parma, P.; Murgia, M. Structures of the magnetoionic media around the Fanaroff-Riley Class I radio galaxies 3C31 and Hydra A. *MNRAS* **2008**, *391*, 521–549, [[0809.2411](#)]. doi:10.1111/j.1365-2966.2008.13895.x.
42. Bonafede, A.; Feretti, L.; Murgia, M.; Govoni, F.; Giovannini, G.; Dallacasa, D.; Dolag, K.; Taylor, G.B. The Coma cluster magnetic field from Faraday rotation measures. *A&A* **2010**, *513*, A30, [[arXiv:astro-ph.CO/1002.0594](#)]. doi:10.1051/0004-6361/200913696.
43. Strom, R.G.; Jaegers, W.J. Extensive gaseous haloes surrounding giant elliptical galaxies - Evidence from depolarization in radio galaxies. *A&A* **1988**, *194*, 79–89.
44. Best, P.N. Radio source populations: Results from SDSS. *Astronomische Nachrichten* **2009**, *330*, 184–189. doi:10.1002/asna.200811152.
45. Gendre, M.A.; Best, P.N.; Wall, J.V.; Ker, L.M. The relation between morphology, accretion modes and environmental factors in local radio AGN. *MNRAS* **2013**, *430*, 3086–3101, [[arXiv:astro-ph.CO/1301.1526](#)]. doi:10.1093/mnras/stt116.
46. Croston, J.H. Environments. *A&A* **2019**, *LOFAR Special Issue*, in press.
47. Heesen, V.; Croston, J.H.; Morganti, R.; Hardcastle, M.J.; Stewart, A.J.; Best, P.N.; Broderick, J.W.; Brüggén, M.; Brunetti, G.; Chyży, K.T.; Harwood, J.J.; Haverkorn, M.; Hess, K.M.; Intema, H.T.; Jamrozy, M.;

- Kunert-Bajraszewska, M.; McKean, J.P.; Orrú, E.; Röttgering, H.J.A.; Shimwell, T.W.; Shulevski, A.; White, G.J.; Wilcots, E.M.; Williams, W.L. LOFAR reveals the giant: a low-frequency radio continuum study of the outflow in the nearby FR I radio galaxy 3C 31. *MNRAS* **2018**, *474*, 5049–5067, [1710.09746]. doi:10.1093/mnras/stx2869.
48. Laing, R.A.; Bridle, A.H. Systematic properties of decelerating relativistic jets in low-luminosity radio galaxies. *MNRAS* **2014**, *437*, 3405–3441, [1311.1015]. doi:10.1093/mnras/stt2138.
 49. Smith, D.J.B.; Best, P.N.; Duncan, K.J.; Hatch, N.A.; Jarvis, M.J.; Röttgering, H.J.A.; Simpson, C.J.; Stott, J.P.; Cochrane, R.K.; Coppin, K.E.; Dannerbauer, H.; Davis, T.A.; Geach, J.E.; Hale, C.L.; Hardcastle, M.J.; Hatfield, P.W.; Houghton, R.C.W.; Maddox, N.; McGee, S.L.; Morabito, L.; Nisbet, D.; Pandey-Pommier, M.; Prandoni, I.; Saxena, A.; Shimwell, T.W.; Tarr, M.; van Bemmell, I.; Verma, A.; White, G.J.; Williams, W.L. The WEAVE-LOFAR Survey. SF2A-2016: Proceedings of the Annual meeting of the French Society of Astronomy and Astrophysics; Reylé, C.; Richard, J.; Cambrésy, L.; Deleuil, M.; Pécontal, E.; Tresse, L.; Vauglin, I., Eds., 2016, pp. 271–280, [1611.02706].
 50. Stasyszyn, F.; Nuza, S.E.; Dolag, K.; Beck, R.; Donnert, J. Measuring cosmic magnetic fields by rotation measure-galaxy cross-correlations in cosmological simulations. *MNRAS* **2010**, *408*, 684–694, [1003.5085]. doi:10.1111/j.1365-2966.2010.17166.x.
 51. Akahori, T.; Gaensler, B.M.; Ryu, D. Statistical Techniques for Detecting the Intergalactic Magnetic Field from Large Samples of Extragalactic Faraday Rotation Data. *ApJ* **2014**, *790*, 123, [1406.3871]. doi:10.1088/0004-637X/790/2/123.
 52. Vacca, V.; Oppermann, N.; Enßlin, T.; Jasche, J.; Selig, M.; Greiner, M.; Junklewitz, H.; Reinecke, M.; Brüggen, M.; Carretti, E.; Feretti, L.; Ferrari, C.; Hales, C.A.; Horellou, C.; Ideguchi, S.; Johnston-Hollitt, M.; Pizzo, R.F.; Röttgering, H.; Shimwell, T.W.; Takahashi, K. Using rotation measure grids to detect cosmological magnetic fields: A Bayesian approach. *A&A* **2016**, *591*, A13, [1509.00747]. doi:10.1051/0004-6361/201527291.
 53. Mevius, M. RMextract: Ionospheric Faraday Rotation calculator. *Astrophysics Source Code Library* **2018**, [1806.024].
 54. Sotomayor-Beltran, C.; Sobey, C.; Hessels, J.W.T.; de Bruyn, G.; Noutsos, A.; Alexov, A.; Anderson, J.; Asekar, A.; Avruch, I.M.; Beck, R.; Bell, M.E.; Bell, M.R.; Bentum, M.J.; Bernardi, G.; Best, P.; Birzan, L.; Bonafede, A.; Breitling, F.; Broderick, J.; Brouw, W.N.; Brüggen, M.; Ciardi, B.; de Gasperin, F.; Dettmar, R.J.; van Duin, A.; Duscha, S.; Eislöffel, J.; Falcke, H.; Fallows, R.A.; Fender, R.; Ferrari, C.; Frieswijk, W.; Garrett, M.A.; Grießmeier, J.; Grit, T.; Gunst, A.W.; Hassall, T.E.; Heald, G.; Hoeft, M.; Horneffer, A.; Iacobelli, M.; Juette, E.; Karastergiou, A.; Keane, E.; Kohler, J.; Kramer, M.; Kondratiev, V.I.; Koopmans, L.V.E.; Kuniyoshi, M.; Kuper, G.; van Leeuwen, J.; Maat, P.; Macario, G.; Markoff, S.; McKean, J.P.; Mulcahy, D.D.; Munk, H.; Orru, E.; Paas, H.; Pandey-Pommier, M.; Pilia, M.; Pizzo, R.; Polatidis, A.G.; Reich, W.; Röttgering, H.; Serylak, M.; Sluman, J.; Stappers, B.W.; Tagger, M.; Tang, Y.; Tasse, C.; ter Veen, S.; Vermeulen, R.; van Weeren, R.J.; Wijers, R.A.M.J.; Wijnholds, S.J.; Wise, M.W.; Wucknitz, O.; Yatawatta, S.; Zarka, P. Calibrating high-precision Faraday rotation measurements for LOFAR and the next generation of low-frequency radio telescopes. *A&A* **2013**, *552*, A58, [arXiv:astro-ph.IM/1303.6230]. doi:10.1051/0004-6361/201220728.
 55. van Diepen, Ger; Dijkema, T.J. DPPP: Default Pre-Processing Pipeline. *Astrophysics Source Code Library* **2011**, [1804.003].
 56. Offringa, A.R.; McKinley, B.; Hurley-Walker, others. WSClean: an implementation of a fast, generic wide-field imager for radio astronomy. *MNRAS* **2014**, *444*, 606–619. doi:10.1093/mnras/stu1368.
 57. Brentjens, M.A.; de Bruyn, A.G. Faraday rotation measure synthesis. *A&A* **2005**, *441*, 1217–1228, [arXiv:astro-ph/0507349]. doi:10.1051/0004-6361:20052990.
 58. Heald, G.; Braun, R.; Edmonds, R. The Westerbork SINGS survey. II Polarization, Faraday rotation, and magnetic fields. *A&A* **2009**, *503*, 409–435, [arXiv:astro-ph.GA/0905.3995]. doi:10.1051/0004-6361/200912240.
 59. Astropy Collaboration.; Robitaille, T.P.; Tollerud, E.J.; Greenfield, P.; Droettboom, M.; Bray, E.; Aldcroft, T.; Davis, M.; Ginsburg, A.; Price-Whelan, A.M.; Kerzendorf, W.E.; Conley, A.; Crighton, N.; Barbary, K.; Muna, D.; Ferguson, H.; Grollier, F.; Parikh, M.M.; Nair, P.H.; Unther, H.M.; Deil, C.; Woillez, J.; Conseil, S.; Kramer, R.; Turner, J.E.H.; Singer, L.; Fox, R.; Weaver, B.A.; Zabalza, V.; Edwards, Z.I.; Azalee Bostroem, K.; Burke, D.J.; Casey, A.R.; Crawford, S.M.; Dencheva, N.; Ely, J.; Jenness, T.; Labrie, K.; Lim, P.L.; Pierfederici,

- F.; Pontzen, A.; Ptak, A.; Refsdal, B.; Servillat, M.; Streicher, O. Astropy: A community Python package for astronomy. *A&A* **2013**, *558*, A33, [[arXiv:astro-ph.IM/1307.6212](https://arxiv.org/abs/1307.6212)]. doi:10.1051/0004-6361/201322068.
60. Hunter, J.D. Matplotlib: A 2D graphics environment. *Computing In Science & Engineering* **2007**, *9*, 90–95. doi:10.1109/MCSE.2007.55.
61. Robitaille, T.; Bressert, E. APLpy: Astronomical Plotting Library in Python. *Astrophysics Source Code Library*, 2012, [[1208.017](https://arxiv.org/abs/1208.017)].
62. Taylor, M.B. TOPCAT & STIL: Starlink Table/VOTable Processing Software. *Astronomical Data Analysis Software and Systems XIV*; Shopbell, P.; Britton, M.; Ebert, R., Eds., 2005, Vol. 347, *Astronomical Society of the Pacific Conference Series*, p. 29.