

# Fish protein and lipid interactions on the digestibility and bioavailability of starch and protein from durum wheat pasta

Ajay S. Desai <sup>1,2,3</sup>, Margaret A. Brennan <sup>1,2</sup> Xinbo Guo <sup>1,4</sup>, Xin-An Zeng <sup>1,4</sup> & Charles S. Brennan <sup>1,2,3,4\*</sup>

<sup>1</sup> School of Food Science and Engineering, South China University of Technology, Guangzhou 510640, China; [ajay.desai@lincoln.ac.nz](mailto:ajay.desai@lincoln.ac.nz) (A.S.D.); [charles.brennan@lincoln.ac.nz](mailto:charles.brennan@lincoln.ac.nz) (C.S.B.); [guoxinbo@scut.edu.cn](mailto:guoxinbo@scut.edu.cn) (X.G.); [xazeng@scut.edu.cn](mailto:xazeng@scut.edu.cn) (X.-A.Z.)

<sup>2</sup> Department of Wine, Food and Molecular Biosciences, Lincoln University, Christchurch 7647, New Zealand; [margaret.brennan@lincoln.ac.nz](mailto:margaret.brennan@lincoln.ac.nz) (M.A.B.)

<sup>3</sup> Riddet Research Institute, Palmerston North 4442, New Zealand. [Charles.brennan@lincoln.ac.nz](mailto:Charles.brennan@lincoln.ac.nz) (C.S.B.)

<sup>4</sup> Overseas Expertise Introduction Center for Discipline Innovation of Food Nutrition and Human Health (111 Center), Guangzhou 510640, China; [charles.brennan@lincoln.ac.nz](mailto:charles.brennan@lincoln.ac.nz) (C.S.B.); [guoxinbo@scut.edu.cn](mailto:guoxinbo@scut.edu.cn) (X.G.); [xazeng@scut.edu.cn](mailto:xazeng@scut.edu.cn) (X.-A.Z.)

\*Correspondence: [Charles.brennan@lincoln.ac.nz](mailto:Charles.brennan@lincoln.ac.nz); Tel.: +6434230637

## Abstract

This research focussed on utilisation of salmon protein and lipid to manipulate pasta glycaemic index and protein digestibility. Salmon fish (*Oncorhynchus tshawytscha*) powder (SFP) supplemented pasta flour at the from 5% to 20% (w/w). Inclusion of SFP lead to a significant reduction in starch digestibility and hence the potential glycaemic values of pasta (experimental pasta being up to 143% lower than control values). SFP addition to pasta increased the release of phenolic compounds from pasta during both a gastric digestion (179%) and pancreatic digestion (133%) in comparison to the control sample. At the same time, the antioxidant activity of the digested pasta was increased by up to 263% (gastric) and 190% (pancreatic) in comparison to durum wheat pasta alone. Interestingly, although protein levels increased with incorporation of SFP, the digestibility values of the protein decreased from 86.41% for the control pasta to 81.95% for 20% SFP pasta. This may indicate that there are interactions between phenolic and protein in the pasta samples which affect overall protein digestibility levels.

**Keywords:** pasta; salmon powder; glycaemic index; protein digestibility; polyphenols; antioxidant activity; bioaccessibility

## 1. Introduction

Numerous researchers have studied the omega-3 polyunsaturated fatty acids (LCn-3PUFAs) namely eicosapentaenoic acid (EPA) and docosahexaenoic acid (DHA) compositions of salmon (*Oncorhynchus tshawytscha*) in relation to their utilisation in human nutrition possibly related to the high antioxidant levels found associated with the astaxanthin and other carotenoids in the flesh [1]. Such research has indicated that diets which are rich in LCn-3PUFAs have reduced incidences of some chronic diseases including cardiovascular diseases, diabetes, cancer, and obesity [2]. In addition, fish have been shown to be rich both in vitamins (A, D, B6 and B12) as well as containing high levels of micronutrients such as iron, potassium and selenium [3]. The recommended weekly intake of fish as directed by the American heart association (AHA) is at least two serves, which relate to an estimated intake of approximately 200mg day<sup>-1</sup> of long chain n-3 polyunsaturated fatty acid (PUFA). Despite this suggestion many countries have dietary intakes far lower than the recommendation. It is possible that waste from the fish processing industry could be utilised to supplement existing dietary levels by fortifying food products often consumed by individuals [4]. The antioxidants found in foods has been shown to manipulate cellular oxidative stress [5], and protein fractions from fish extracts have been used by previous researchers to reduce glycaemic responses of individuals and hence regulate obesity and potentially diabetes [6, 7]. Indeed, researchers have studied the fortification of cereal foods with a range of protein sources from milk, animals and also vegetables to achieve similar regulation of disease biomarkers [8,9].

One of the most commonly used cereal food products used in the manipulation of dietary influences of food related illnesses is pasta, mainly as it is already a relatively low glycaemic index food product and hence regarded as a healthy carbohydrate rich food product [10]. When considering the factors which effect the glycaemic response of an individual, the total starch content of the food is of great importance as this is the material converted into reducing sugar components and in turn affects blood glucose levels. Hence the consumption of starchy foods (especially those which have a high level of starch which is considered readily digestible) has been related to diseases such as obesity and diabetes [11]. Generally, the recommendation is to

consumer foods which exhibit low glycaemic responses in order to avoid the risks associated with diabetes, cardiovascular and even neurodegenerative disease [12]. Two of the ways to manipulate the rate and extent of starch digestion is by altering the protein and oil content of foods as these tend to lead to a reduction in reducing sugar release post ingestion [11, 12, 13]. This may be linked to the possibility of forming amylose-lipid complexes when starch and lipids are combined [14]. Previously researchers have studied the effects of different food lipids, including butter, coconut oil, grapeseed oil and olive oil of different degree of saturation and chain lengths, on the glycaemic response of bread [15]. Lipids significantly decreased the starch hydrolysis rate, and the formation amylose-lipid complexes and protein-lipid complexes may be responsible for this observation [11]. The presence of protein in the food matrix may influence starch digestion by the encapsulation of starch granules into the protein matrix of the food [7]. The effect of meat protein interactions on the digestibility of pasta has been studied [9]. The researchers observed that starch-protein interactions increased with increasing levels of meat additions and accounted for decreasing glycaemic responses. Also, interaction between starch-protein- phenolic compound in the food product affect protein structure through precipitation and decrease the starch and protein digestibility [16]. The supplementation of pasta with other functional ingredients has received much attention. For instance pasta has been fortified with protein rich ingredients such as faba bean flour [17], meat [9], shrimp powder [18], green mussel powder (*Perna canaliculus*) [19], barely flour [20], amaranth seed flour [21], *Eruca vesicaria* leaves [22], artichoke canning by-products [23], almond flour [24] and *Nannochloropsis* sp. [25]. However, the nutritional properties of pasta enriched with partial replacement of semolina wheat flour by salmon (*Oncorhynchus tshawytscha*) powder (SFP) is still unknown. Therefore, the present investigation aimed to evaluate the effects of salmon powder as ingredients for pasta production and their contribution to *in vitro* starch, protein digestibility and antioxidant activity.

## 2. Results and Discussion

### 2.1 In Vitro Predictive Glycaemic Response

As mentioned before, starch and protein digestion (and the regulation of these chemical compositions) are of great importance to the nutritional benefit of foods such as pasta. Many studies have illustrated that the interactions between protein and fibre, or starch and fibre, even starch – protein – fibre, on the overall food structure confer effects on the rate and extent of carbohydrate and protein digestion [26]. For this reason, the research focused on using protein and oil from SFP, and incorporating SFP into pasta, to determine effects on protein, carbohydrate and antioxidant activities following a standardised 120 min *in vitro* digestion.

As can be seen from figure1 the level of reducing sugars released over the 120 min *in vitro* digestion of the pasta samples varied between samples, however incorporation of SFP significantly reduced starch digestion and sugar release during the digestion of experimental pasta samples compared to durum wheat pasta samples (the control). Similarly, SFP fortified pasta samples exhibited lower levels of rapidly digestible starch (RDS) and slowly digestible starch (SDS) compared with the control samples (Figure 2). Such results may be related to the fact that lipids have been shown to form complexes with amylose and protein and in so doing they have been shown to disrupt the enzyme adsorption sites on the surface of starch granule [11], and follows previous research on fortifying spaghetti with protein from bean flour [27], possibly due to incomplete gelatinisation of starch granules. In the present study, cooked pasta samples enriched with 5% SFP, 10% SFP, 15% SFP and 20% SFP had 0.25%, 1.25%, 2.59% and 3.69% lipid content, respectively. The reduction in digestibility may be attributed to the formation of amylose-lipid complexes (ALC) as has been demonstrated by other researchers investigating the effects of lipids on the enzymatic resistance of starch, and manipulation of starch granule swelling characteristics associated with starch gelatinisation [15, 26]. These studies, and others, have revealed that amylose-lipid interactions results in the formation of single helical structure with a conformational hindrance that restricts enzymes to hydrolyse the starch granule [14, 26, 28]. Additionally, it has been illustrated that the accessibility of starch degrading enzymes to the substrate can be hampered by the incorporation of proteins [14, 26, 28, 29]. The results of this

study confirm that protein (in the form of SFP) can be utilised to affect the starch - protein network and possibly regulate starch digestion by restricting the activity of  $\alpha$ -amylase. It is possible that the addition of SFP encapsulates starch hence reducing the accessibility of starch degrading enzymes as mentioned by previous researchers [9, 18, 30]. For instance, the addition of yam flour (*Dioscorea schimperiana*) into pasta [31] has been shown to disrupt the protein – starch matrix and restrict the access of starch degrading enzymes to starch granules. As can be observed in Figure 3 fortifying pasta with SFP reduced the standardised glycaemic AUC values when compared with the control sample. Similarly, when pasta flour was replaced with soya bean flour reductions were observed in the glycaemic values of the pasta [32]. Our work confirms that the incorporation of protein (SFP) into pasta can reduce the digestibility of starch and may be a novel option to the lowering of the glycaemic index of the carbohydrate rich food products.

## 2.2 Protein Content and In Vitro Protein Digestibility

Researchers have used the *in vitro* protein digestibility value (IVPD) to determine the rate and extent of protein digestion in food materials [10]. The protein digestibility values of SFP enriched pasta samples are shown in Table 1 and show that the addition of SFP to pasta samples increased the overall content of protein in the pasta samples. What was of interest was that the values for the uncooked and cooked pasta samples were similar and this indicated that the protein fraction in the pasta did not leach out during the cooking of pasta. However, despite the overall protein content being increased with SFP addition, the *in vitro* protein digestibility values of enriched pasta samples was lower when compared with the control pasta (reduced from 84.60 to 80.80%).

The reduction in digestibility could be due to fish protein structure, other components such as formation of protein-starch complex, cross links between proteins [33] and presence of phenolic compounds [34]. Oxidized phenolic compounds have been proposed to react with proteins and form insoluble complexes, inhibiting the activity of proteolytic enzymes and interfering with utilization of proteins [35]. Our results are supported by those previously reported [36], which

found a reduction of protein digestibility of shrimp meat and broad bean enriched pasta. Figure 4 illustrates the pH drop curves obtained from proteolytic enzymatic hydrolysis of enriched pasta, and are a result of the release of amino acids and peptides during the digestion of protein, and the release of carboxyl ( $\text{-COO}^-$ ) compounds as well as amino ( $\text{-NH}_3^+$ ) units which in turn result in a reduction in the value of the pH of the solution [37].

### *2.3 The Composition of Amino Acids Released into Intestine After In Vitro Digestion*

The quality of proteins may also be assessed by evaluating the composition of the breakdown compounds observed in the small intestine. Hence, combining both the amino acid composition and the protein digestion levels of foods can result in a clearer idea of the nutritional quality of pasta samples. The amino acid content of the SFP enriched pasta post in vitro digestion can be seen in Table 2. Of note, the content of phenylalanine, tyrosine, isoleucine and leucine from the enriched pasta decreased significantly ( $p < 0.05$ ) compared to the control, and the enrichment of SFP into pasta resulted in a decrease in non-essential amino acids in the digesta (excepting arginine, alanine and asparagine) compared to control pasta. Such results could be due to the protein - lipid interactions as discussed previously resulting in the oxidation of amino acids, as has been noted by researchers previously [38, 39, 40].

### *2.4 In vitro Bioaccessibility of Phenolic content and Antioxidant activity*

Recently there has been a lot of research attention given to the fate of phenolic compounds post digestion [38, 39, 40]. Figures 5 & 6 illustrate the effects of digestion stages on the phenolic content of digesta and illustrate that the SFP pasta samples showed significantly increased bioaccessible phenolic compounds after gastric and pancreatic digestion compared to the control pasta samples. Analysis of the antioxidant activity of the phenolic compounds was conducted using the oxygen radical absorbance capacity (ORAC) mechanism. Results illustrate an increase in antioxidant activity by approximately 20% when SFP was incorporated into pasta (from 5.20 to 13.69  $\mu\text{mol Trolox g of pasta}^{-1}$  (as observed during the gastric digestion stage) and 40.36 to 76.75  $\mu\text{mol Trolox g of pasta}^{-1}$  (as observed during the pancreatic digestion stage). The total phenolic

content of the control sample before and after digestion (1.49 mg of gallic acid/g of pasta and 0.82 mg of gallic acid g of pasta<sup>-1</sup>) and antioxidant activity (5.20  $\mu$ mol Trolox g of pasta<sup>-1</sup> and 40.36  $\mu$ mol Trolox g of pasta<sup>-1</sup>) was lower than the SFP fortified pasta. This observation may be due to possible leaching of phenolic compounds during the cooking of pasta. Indeed, previous researchers has shown a similar result in cooked faba bean flour fortified pasta [41, 42] with the phenolic compounds leaching into cooking medium and degraded due to thermal treatment. Other researchers have noted similar correlations between the phenolic level of food products and the antioxidant activity of fortified pasta [19, 43]. For instance incorporation of parley into pasta increased the total phenolic content by 126-167% and antiradical activity against ABTS by 161-246% [43]. Researchers have noted that protein and phenol may interact with each other through covalent or non-covalent interaction [35]. These interactions might lead to precipitation of protein from food matrix and an alteration in the secondary and tertiary structure of protein [44, 45].

### 3 Materials and Methods

#### 3.1 Materials

Pasta semolina flour was obtained from Sun Valley Foods (Christchurch, New Zealand) and fresh salmon was obtained locally from Akaroa Salmon Ltd (Christchurch, New Zealand).

#### 3.2 Fish Powder Preparation

The fish was prepared as described previously [1]. The dried powder was stored at -20 °C temperature until required.

#### 3.3 Pasta Preparation

Pasta was prepared by blending 500g dry ingredients and 32.5 g/100g water (tap water, 41 °C) for 20 min in a commercial pasta machine (Model: MPF15N235M; Firmer, Ravenna, Italy). Pasta samples were divided into 20 g portions and stored in polythene bags at -18 °C until required.

Prior to analysis, the pasta was defrosted for 10 min at room temperature. Five pasta formulations were prepared in the ratios (semolina/ SFP) of 100:0; 95:5:90:10; 85:15 and 80:20.

### 3.4 *In Vitro* Starch Digestibility and Glycaemic Response

An *in vitro* digestion system as described previously [46] was used to evaluate the starch digestion properties of the pasta samples. The process used stomach digestion procedures utilising 0.8 mL 1M HCL and 1 mL of 10% pepsin solution (Sigma Aldrich, Sydney, Australia) incubated at 37 °C for 30 min under constant stirring. The process also mimicked the digestion of the small intestine by the addition of 5 mL of 2.5% Pancreatin solution (Sigma Aldrich, Sydney, Australia) in 0.1 M sodium maleate buffer pH 6 at 37 °C for 120 min. Samples were analysed for reducing sugar content using the 3,5-dinitrosalicylic acid (DNS). Glucose release was calculated in mg glucose g/ sample and plotted against time and area under the curve (AUC) was calculated by dividing the graph into trapezoids.

### 3.5 *In Vitro* Protein Digestibility of pasta

The multi-enzyme technique was used for the determination of *in vitro* protein digestibility of cooked pasta samples [47]. A 50 mL of protein suspension was prepared in distilled water (6.25 mg of protein/ mL), adjusted to pH 8 with a solution of 0.1 N HCL and /or 0.1 N NaOH, and placed on magnetic heating stirring block at 37 °C. The multi-enzyme solution (1.6 mg/ mL Trypsin, 3.1 mg/ mL chymotrypsin and 1.3 mg/ mL peptidase) was maintained in an ice bath and adjusted to pH 8.0 with 0.1 N HCL and /or 0.1 N NaOH. 5 mL of the multi-enzyme solution was then added to the protein suspension, which was maintained at 37 °C. The decrease in pH was measured after the addition of an enzymatic solution at every minute for period of 10 min using a digital pH meter (S20 Seven Easy<sup>TM</sup>, Mettler Toledo, USA). The percent protein digestibility (Y) was calculated by using Eq. (1):

$$Y = 210.46 - 18.10 x, \quad (1)$$

where x is the change in pH after 10 min.



Protein availability refers to the quantity of protein digested in the pasta. It was calculated over the protein content in cooked pasta and the protein digestibility as:

$$\text{Protein availability (PA)} = (\text{Protein digestibility} \times \text{Protein content in cooked pasta}) / 100$$

### 3.6 Amino Acid Profile

The amino acid profile of the digested pasta material were evaluated using an Agilent 1100 series (Agilent Technologies, Walbronn, Germany) high-performance liquid chromatography machine as described previously [48]. The machine used a 150 X 4.6mm, C18, 3u ACE-111-1546, column and the amino acids were applied at a flow rate of 0.7 mL/min. Detection was at an excitation of 335 nm and emission of 440 nm for primary amino acids. At 22 min, the detector was switched to excitation 260 nm, emission 315 nm to detect secondary amino acid such as proline. The amino acid results are expressed in milligram amino acids/ g protein of the pasta sample [48].

### 3.7 In Vitro Gastro-Intestinal Digestion

During pepsin and pancreatic digestion, aliquots (1 mL) withdrawn after 30 and 120 min to which 1 mL ethanol was added (1:1) to arrest any further enzymatic reaction. Thereafter, samples were centrifuged at 1000 rpm for 5 min. Supernatants (gastrointestinal digested extracts) and pellets were separated and kept at -20 °C for further analysis.

### 3.8 Total Phenolic Content and Antioxidant Activity of pasta

The total phenolic content of supernatant obtained from the *in vitro* gastro-intestinal digestion was measured using the Folin-Ciocalteu method as described by [49]. Freshly prepared 2.5 mL of 0.2 N Folin Ciocalteu reagent and 7.5% Na<sub>2</sub>CO<sub>3</sub> was added to the digesta aliquots (0.5 mL) and incubated for 2 h in the dark. The absorbance of the reaction mixture was measured at 760 nm using the V-1200 model (Schimadzu, Maryland, USA). Gallic acid was used as a standard to determine total phenolic content of the samples as mg of Gallic acid equivalents (GAE)/ g sample

### 3.8 Oxygen Radical Absorbance Capacity (ORAC) Assay

### 3.8.1 Chemical Reagents and Standard Solutions

AAPH (2, 2' azobis (2-amidino-propane) dihydrochloride) (0.645 g) was completely dissolved in 10 mL of 75 mM phosphate buffer (pH 7.4) and was kept in ice bath. Fluorescein stock solution (1mM) was prepared with 0.016 g dissolved in 50 mL of 75 mM phosphate buffer (pH 7.4) was kept at 4 °C in dark condition. Trolox (6-hydroxy-2, 5, 7, 8 -tetra methylchroman-2-carboxylic acid) standard was prepared and the stock solution of Trolox was diluted with phosphate buffer to 100 µM, 50 µM, 25 µM, 12.5 µM, 6.25 µM, 3.125 µM, 1.5625 µM and 0 µM working solution.

### 3.8.2 Oxygen Radical Absorbance Capacity (ORAC) Assay

ORAC (oxygen radical absorbance capacity) was used according to the method described previously [49]. A 96 well microplate reader (FLUOstar Omega, BMG LABTECH, Germany) was used for all measurements. Trolox was used as a standard and the antioxidant capacity of the samples was expressed as mmol Trolox equivalent (TE) / g sample.

### 3.9 Statistical Analysis

All experiments were performed in triplicate. Data was subjected to a one way analysis of variance (ANOVA) and significance differences were evaluated by Tukey's comparison test ( $p < 0.05$ ). Statistical software version 16 (Minitab, Australia) was used to perform the statistical analysis of the data.

## 3. Conclusions

This study illustrates that the fortification of durum wheat pasta with SFP can improve the nutritional quality of pasta. For instance, SFP addition led to an increase in the antioxidant levels and protein content of pasta whilst reducing the predicted glycaemic index of the food product. The study also illustrated that the antioxidant activity from supplemented pasta was bioaccessible *in vitro* and significantly increased with the supplementation of SFP.

**Author Contributions:** M.A.B., C.S.B. and A.S.D. conceived and designed the experiments; A.S.D. performed the experiments; A.S.D. analysed the data; C.S.B., M.A.B. and A.S.D. were responsible for writing the manuscript.

**Funding:** The research was supported by Lincoln University, New Zealand postgraduate funding.

**Conflict of interest:** The authors declare no conflict of interest.

## References

- Ortiz, J.; Lemus-Mondaca, R.; Vega-Galvez, A.; Ah-Hen, K.; Puente-Diaz, L.; Zura-Bravo, L.; Aubourg, S. Influence of air-drying temperature on drying kinetics, colour, firmness and biochemical characteristics of Atlantic salmon (*Salmo salar* L.) fillets. *Food Chem.* **2013**, *139*, 162–169, doi:10.1016/j.foodchem.2013.01.037.
- Cascant, M. M.; Breil, C.; Fabiano-Tixier, A. S.; Chemat, F.; Garrigues, S.; de la Guardia, M. Determination of fatty acids and lipid classes in salmon oil by near infrared spectroscopy. *Food Chem.* **2018**, *239*, 865–871, doi:10.1016/j.foodchem.2017.06.158.
- Matos, J.; Lourenço, H. M.; Brito, P.; Maulvault, A. L.; Martins, L. L.; Afonso, C. Influence of bioaccessibility of total mercury, methyl-mercury and selenium on the risk/benefit associated to the consumption of raw and cooked blue shark (*Prionace glauca*). *Environ. Res.* **2015**, *143*, 123–129, doi:10.1016/j.envres.2015.09.015.
- Oliveira, I. S. de; Lourenco, L. de F. H.; Sousa, C. L.; Peixoto Joele, M. R. S.; Ribeiro, S. da C. do A. Composition of MSM from Brazilian catfish and technological properties of fish flour. *Food Control* **2015**, *50*, 38–44, doi:10.1016/j.foodcont.2014.08.018.
- Pisoschi, A. M.; Pop, A. The role of antioxidants in the chemistry of oxidative stress: A review. *Eur. J. Med. Chem.* **2015**, *97*, 55–74, doi:10.1016/j.ejmech.2015.04.040.
- Kadam, S. U.; Prabhasankar, P. Marine foods as functional ingredients in bakery and pasta products. *Food Res. Int.* **2010**, *43*, 1975–1980, doi:10.1016/j.foodres.2010.06.007.

- 289 7. Brennan, M. A.; Derbyshire, E.; Tiwari, B. K.; Brennan, C. S. Ready-to-eat snack products:  
290 the role of extrusion technology in developing consumer acceptable and nutritious snacks.  
291 *Int. J. Food Sci. Technol.* **2013**, *48*, 893–902, doi:10.1111/ijfs.12055.
- 292 8. Torres, A.; Frias, J.; Granito, M.; Vidal-Valverde, C. Germinated *Cajanus cajan* seeds as  
293 ingredients in pasta products: Chemical, biological and sensory evaluation. *Food Chem.*  
294 **2006**, *101*, 202–211, doi:10.1016/j.foodchem.2006.01.018.
- 295 9. Liu, T.; Hamid, N.; Kantono, K.; Pereira, L.; Farouk, M. M.; Knowles, S. O. Effects of  
296 meat addition on pasta structure, nutrition and *in vitro* digestibility. *Food Chem.* **2016**, *213*,  
297 108–114, doi:10.1016/j.foodchem.2016.06.058.
- 298 10. Tazart, K.; Lamacchia, C.; Zaidi, F.; Haros, M. Nutrient composition and *in vitro*  
299 digestibility of fresh pasta enriched with *Vicia faba*. *J. Food Compos. Anal.* **2016**, *47*, 8–  
300 15, doi:10.1016/j.jfca.2015.12.007.
- 301 11. Chen, X.; He, X. W.; Zhang, B.; Fu, X.; Jane, J. lin; Huang, Q. Effects of adding corn oil  
302 and soy protein to corn starch on the physicochemical and digestive properties of the starch.  
303 *Int. J. Biol. Macromol.* **2017**, *104*, 481–486, doi:10.1016/j.ijbiomac.2017.06.024.
- 304 12. Augustin, L. S. A.; Kendall, C. W. C.; Jenkins, D. J. A.; Willett, W. C.; Astrup, A.; Barclay,  
305 A. W.; Björck, I.; Brand-Miller, J. C.; Brighenti, F.; Buyken, A. E.; Ceriello, A.; La  
306 Vecchia, C.; Livesey, G.; Liu, S.; Riccardi, G.; Rizkalla, S. W.; Sievenpiper, J. L.;  
307 Trichopoulou, A.; Wolever, T. M. S.; Baer-Sinnott, S.; Poli, A. Glycemic index, glycemic  
308 load and glycemic response: An international scientific consensus summit from the  
309 international carbohydrate quality consortium (ICQC). *Nutr. Metab. Cardiovasc. Dis.* **2015**,  
310 *25*, 795–815, doi:10.1016/j.numecd.2015.05.005.
- 311 13. Foschia, M.; Peressini, D.; Sensidoni, A.; Brennan, M. A.; Brennan, C. S. Synergistic effect  
312 of different dietary fibres in pasta on *in vitro* starch digestion? *Food Chem* **2015**, *172*, 245–  
313 250, doi:10.1016/j.foodchem.2014.09.062.
- 314 14. Oyeyinka, S. A.; Singh, S.; Venter, S. L.; Amonsou, E. O. Effect of lipid types on

- 315 complexation and some physicochemical properties of bambara groundnut starch.  
 316 *Starch/Staerke* **2017**, 69, 1–10, doi:10.1002/star.201600158.
- 317 15. Lau, E.; Zhou, W.; Henry, C. J. Effect of fat type in baked bread on amylose–lipid complex  
 318 formation and glycaemic response. *Br. J. Nutr.* **2016**, 1–8, doi:10.1017 /  
 319 S0007114516001458.
- 320 16. Czubinski, J.; Dwiecki, K. A review of methods used for investigation of protein–phenolic  
 321 compound interactions. *Int. J. Food Sci. Technol.* **2017**, 52, 573–585,  
 322 doi:10.1111/ijfs.13339.
- 323 17. Coda, R.; Varis, J.; Verni, M.; Rizzello, C. G.; Katina, K. Improvement of the protein  
 324 quality of wheat bread through faba bean sourdough addition. *LWT - Food Sci. Technol.*  
 325 **2017**, 82, 296–302, doi:10.1016/j.lwt.2017.04.062.
- 326 18. Ramya N. S.; Prabhasankar, P. . Gowda, L. R.; Modi, V. K.; . Bhaskar, N. Influence of  
 327 freeze-dried shrimp meat in pasta processing qualities of Indian *T. durum* Wheat. *J. Aquat.*  
 328 *Food Prod. Technol.* **2014**, 8850, doi:10.1080/10498850.2013.796581.
- 329 19. Vijaykrishnaraj, M.; Bharath Kumar, S.; Prabhasankar, P. Green mussel (*Perna*  
 330 *canaliculus*) as a marine ingredient to enrich gluten free pasta: product quality,  
 331 microstructure and biofunctional evaluation. *J. Food Meas. Charact.* **2014**, 9, 76–85,  
 332 doi:10.1007/s11694-014-9212-5.
- 333 20. Montalbano, A.; Tesoriere, L.; Diana, P.; Barraja, P.; Carbone, A.; Spanò, V.; Parrino, B.;  
 334 Attanzio, A.; Livrea, M. A.; Cascioferro, S.; Cirrincione, G. Quality characteristics and *in*  
 335 *vitro* digestibility study of barley flour enriched ditalini pasta. *LWT - Food Sci. Technol.*  
 336 **2016**, 72, 223–228, doi:10.1016/j.lwt.2016.04.042.
- 337 21. Cardenas-Hernandez, A.; Beta, T.; Loarca-Pica, G.; Castao-Tostado, E.; Nieto-Barrera, J.  
 338 O.; Mendoza, S. Improved functional properties of pasta: Enrichment with amaranth seed  
 339 flour and dried amaranth leaves. *J. Cereal Sci.* **2016**, 72, 84–90, doi:10.1016 /  
 340 j.jcs.2016.09.014.

- 341 22. Bouacida, S.; Ben Amira, A.; Ben Haj Koubaier, H.; Blecker, C.; Bouzouita, N. Chemical  
342 composition, cooking quality, texture and consumer acceptance of pasta with *Eruca*  
343 *vesicaria* leaves. *Int. J. Food Sci. Technol.* **2017**, *52*, 2248–2255, doi:10.1111/ijfs.13504.
- 344 23. Pasqualone, A.; Punzi, R.; Trani, A.; Summo, C.; Paradiso, V. M.; Caponio, F.;  
345 Gambacorta, G. Enrichment of fresh pasta with antioxidant extracts obtained from artichoke  
346 canning by-products by ultrasound-assisted technology and quality characterisation of the  
347 end product. *Int. J. Food Sci. Technol.* **2017**, 1–10, doi:10.1111/ijfs.13486.
- 348 24. Martínez, M. L.; Marín, M. A.; Gili, R. D.; Penci, M. C.; Ribotta, P. D. Effect of defatted  
349 almond flour on cooking, chemical and sensorial properties of gluten-free fresh pasta. *Int.*  
350 *J. Food Sci. Technol.* **2017**, *52*, 2148–2155, doi:10.1111/ijfs.13493.
- 351 25. Rodríguez De Marco, E.; Steffolani, M. E.; Martínez, M.; León, A. E. The use of  
352 *Nannochloropsis* sp. as a source of omega-3 fatty acids in dry pasta: chemical, technological  
353 and sensory evaluation. *Int. J. Food Sci. Technol.* **2018**, *53*, 499–507,  
354 doi:10.1111/ijfs.13609.
- 355 26. Ren, X.; Chen, J.; Molla, M. M.; Wang, C.; Diao, X.; Shen, Q. *In vitro* starch digestibility  
356 and *in vivo* glycemic response of foxtail millet and its products. *Food Funct.* **2016**, *1*, 372–  
357 379, doi:10.1039/c5fo01074h.
- 358 27. Giuberti, G.; Gallo, A.; Cerioli, C.; Fortunati, P.; Masoero, F. Cooking quality and starch  
359 digestibility of gluten free pasta using new bean flour. *Food Chem.* **2015**, *175*, 43–49,  
360 doi:10.1016/j.foodchem.2014.11.127.
- 361 28. Annor, G. A.; Marcone, M.; Corredig, M.; Bertoft, E.; Seetharaman, K. Effects of the  
362 amount and type of fatty acids present in millets on their *in vitro* starch digestibility and  
363 expected glycemic index (eGI). *J. Cereal Sci.* **2015**, *64*, 76–81, doi:10.1016 /  
364 j.jcs.2015.05.004.
- 365 29. Singh, J.; Dartois, A.; Kaur, L. Starch digestibility in food matrix: a review. *Trends Food*  
366 *Sci. Technol.* **2010**, *21*, 168–180, doi:10.1016/j.tifs.2009.12.001.

- 367 30. Rosa-Sibakov, N.; Heiniö, R. L.; Cassan, D.; Holopainen-Mantila, U.; Micard, V.; Lantto,  
 368 R.; Sozer, N. Effect of bioprocessing and fractionation on the structural, textural and  
 369 sensory properties of gluten-free faba bean pasta. *LWT - Food Sci. Technol.* **2016**, *67*, 27–  
 370 36, doi:10.1016/j.lwt.2015.11.032.
- 371 31. Djeukeu, W. A.; Gouado, I.; Leng, M. S.; Vijaykrishnaraj, M.; Prabhasankar, P. Effect of  
 372 dried yam flour (*Dioscorea schimperiana*) on cooking quality, digestibility profile and  
 373 antioxidant potential of wheat based pasta. *J. Food Meas. Charact.* **2017**, *11*, 1421–1429,  
 374 doi:10.1007/s11694-017-9521-6.
- 375 32. Chillo, S.; Monro, J. A.; Mishra, S.; Henry, C. J. Effect of incorporating legume flour into  
 376 semolina spaghetti on its cooking quality and glycaemic impact measured *in vitro*. *Int J*  
 377 *Food Sci Nutr* **2010**, *61*, 149–160, doi:10.3109/09637480903476423.
- 378 33. Gimenez, M. A.; Drago, S. R.; Bassett, M. N.; Lobo, M. O.; Samman, N. C. Nutritional  
 379 improvement of corn pasta-like product with broad bean (*Vicia faba*) and quinoa  
 380 (*Chenopodium quinoa*). *Food Chem.* **2016**, *199*, 150–156, doi:10.1016 /  
 381 j.foodchem.2015.11.065.
- 382 34. Swieca, M.; Seczyk, L.; Gawlik-Dziki, U.; Dziki, D. Bread enriched with quinoa leaves -  
 383 The influence of protein-phenolics interactions on the nutritional and antioxidant quality.  
 384 *Food Chem.* **2014**, *162*, 54–62, doi:10.1016/j.foodchem.2014.04.044.
- 385 35. Ozdal, T.; Capanoglu, E.; Altay, F. A review on protein-phenolic interactions and  
 386 associated changes. *Food Res. Int.* **2013**, *51*, 954–970, doi:10.1016/j.foodres.2013.02.009.
- 387 36. Kadam, S. U.; Prabhasankar, P. Evaluation of cooking, microstructure, texture and sensory  
 388 quality characteristics of shrimp meat-based pasta. *J. Texture Stud.* **2012**, *43*, 268–274,  
 389 doi:10.1111/j.1745-4603.2011.00336.x.
- 390 37. Prodpran, T.; Benjakul, S.; Phatcharat, S. Effect of phenolic compounds on protein cross-  
 391 linking and properties of film from fish myofibrillar protein. *Int. J. Biol. Macromol.* **2012**,  
 392 *51*, 774–782, doi:10.1016/j.ijbiomac.2012.07.010.



38. Pastor-Cavada, E.; Drago, S. R.; González, R. J.; Juan, R.; Pastor, J. E.; Alaiz, M.; Vioque, J. Effects of the addition of wild legumes (*Lathyrus annuus* and *Lathyrus clymenum*) on the physical and nutritional properties of extruded products based on whole corn and brown rice. *Food Chem* **2011**, *128*, 961–967, doi:10.1016/j.foodchem.2011.03.126.
39. Ma, Y.; Zhou, M.; Huang, H. Changes of heat-treated soymilks in bioactive compounds and their antioxidant activities under *in vitro* gastrointestinal digestion. *Eur. Food Res. Technol.* **2014**, *239*, 637–652, doi:10.1007/s00217-014-2260-6.
40. Rustad, T.; Storror, I.; Slizyte, R. Possibilities for the utilisation of marine by-products. *Int. J. Food Sci. Technol.* **2011**, *46*, 2001–2014, doi:10.1111/j.1365-2621.2011.02736.x.
41. Turco, I.; Bacchetti, T.; Bender, C.; Zimmermann, B.; Oboh, G.; Ferretti, G. Polyphenol content and glycemic load of pasta enriched with Faba bean flour. *Funct. Foods Heal. Dis.* **2016**, *6*, 291–305.
42. Swieca, M.; Gawlik-Dziki, U.; Dziki, D.; Baraniak, B.; Czyz, J. The influence of protein-flavonoid interactions on protein digestibility *in vitro* and the antioxidant quality of breads enriched with onion skin. *Food Chem.* **2013**, *141*, 451–458, doi:10.1016 / j.foodchem.2013.03.048.
43. Seczyk, L.; Swieca, M.; Gawlik-Dziki, U.; Luty, M.; Czyz, J. Effect of fortification with parsley (*Petroselinum crispum* Mill.) leaves on the nutraceutical and nutritional quality of wheat pasta. *Food Chem.* **2016**, *190*, 419–428, doi:10.1016/j.foodchem.2015.05.110.
44. Rawel, H. M.; Meidtner, K.; Kroll, J. Binding of selected phenolic compounds to proteins. *J. Agric. Food Chem.* **2005**, *53*, 4228–4235, doi:10.1021/jf0480290.
45. Sun-Waterhouse, D.; Jin, D.; Waterhouse, G. I. N. Effect of adding elderberry juice concentrate on the quality attributes, polyphenol contents and antioxidant activity of three fibre-enriched pastas. *Food Res. Int.* **2013**, *54*, 781–789, doi:10.1016 / j.foodres.2013.08.035.
46. Gao, J.; Brennan, M. A.; Mason, S. L.; Brennan, C. S. Effect of sugar replacement with



stevianna and inulin on the texture and predictive glycaemic response of muffins. *Int. J. Food Sci. Technol.* **2016**, *51*, 1979–1987, doi:10.1111/ijfs.13143.

47. Hsu, H.; Vavak, D. A multienzyme technique for estimating protein digestibility. *J. Food Sci.* **1977**, *45*, 1269–1273, doi:10.1111/j.1365-2621.1977.tb14476.x.

48. Heems, D.; Luck, G.; Fraudeau, C.; Verette, E. Fully automated precolumn derivatization, on-line dialysis and high- performance liquid chromatographic analysis of amino acids in food, beverages and feedstuff. *J. Chromatogr. A* **1998**, *798*, 9–17, doi:10.1016/S0021-9673(97)01007-8.

49. Li, W.; Pickard, M. D.; Beta, T. Evaluation of antioxidant activity and electronic taste and aroma properties of antho-beers from purple wheat grain. *J. Agric. Food Chem.* **2007**, *55*, 8958–8966, doi:10.1021/jf071715p.

**Table 1** Protein content, *In vitro* protein digestibility and protein availability of pasta fortified with salmon fish powder (SFP).

Samples	PC in raw pasta (g/100g dry pasta)	PC in cooked pasta (g/100g dry pasta)	PD (%)	PA (g/100g dry pasta)
CP	12.60 ± 0.05 <sup>a</sup>	12.88 ± 0.06 <sup>a</sup>	86.41 ± 0.37 <sup>a</sup>	11.13± 0.07 <sup>a</sup>
SFP 5	14.34 ± 0.03 <sup>b</sup>	15.41 ± 0.17 <sup>b</sup>	84.60 ± 0.20 <sup>b</sup>	13.03 ± 0.14 <sup>b</sup>
SFP 10	17.67 ± 0.04 <sup>c</sup>	18.10 ± 0.11 <sup>c</sup>	82.97 ± 0.10 <sup>c</sup>	15.02 ± 0.09 <sup>c</sup>
SFP 15	20.73 ± 0.10 <sup>d</sup>	20.77 ± 0.09 <sup>d</sup>	81.16 ± 0.27 <sup>d</sup>	16.85 ± 0.12 <sup>d</sup>
SFP 20	22.7 ± 0.30 <sup>e</sup>	23.40 ± 0.13 <sup>e</sup>	81.95 ± 0.18 <sup>e</sup>	19.18 ± 0.07 <sup>e</sup>


PC- protein content, PD- *In vitro* protein digestibility, PA-protein availability. SFP5, SFP10, SFP15, and SFP20: pasta prepared with 5, 10, 15, and 20 g of salmon fish powder /100 g of semolina flour. CP: control pasta. Results are presented as the mean value ± standard deviation, *n* = 3; Values within a column followed by different small letters are significantly different (*p* < 0.05).

**Table 2.** Amino acid (AAs) composition (mg/g protein) from digestibility studies in the intestinal stage at 120 min of pasta enriched with different salmon fish powder (SFP) levels and control.


Amino acid	CP	SFP5	SFP10	SFP15	SFP20
Phenylalanine	18.07 ± 0.17 <sup>a</sup>	14.63 ± 1.13 <sup>bc</sup>	14.86 ± 1.09 <sup>b</sup>	13.40 ± 0.46 <sup>bc</sup>	12.68 ± 0.31 <sup>c</sup>
Tyrosine	14.03 ± 0.37 <sup>a</sup>	12.07 ± 0.96 <sup>b</sup>	12.61 ± 1.07 <sup>ab</sup>	11.40 ± 0.30 <sup>b</sup>	10.90 ± 0.39 <sup>b</sup>
Isoleucine	14.94 ± 0.10 <sup>a</sup>	12.55 ± 1.05 <sup>a</sup>	13.48 ± 1.09 <sup>ab</sup>	12.24 ± 0.31 <sup>b</sup>	11.98 ± 0.23 <sup>b</sup>
Leucine	26.82 ± 0.21 <sup>a</sup>	22.72 ± 1.78 <sup>b</sup>	23.92 ± 1.86 <sup>ab</sup>	21.87 ± 0.60 <sup>b</sup>	21.68 ± 0.29 <sup>b</sup>
Lysine	16.15 ± 0.33 <sup>b</sup>	16.72 ± 1.30 <sup>b</sup>	21.58 ± 1.58 <sup>a</sup>	21.41 ± 0.58 <sup>a</sup>	22.27 ± 1.54 <sup>a</sup>
Methionine	0.57 ± 0.35 <sup>a</sup>	0.81 ± 0.61 <sup>a</sup>	0.51 ± 0.17 <sup>a</sup>	0.47 ± 0.15 <sup>a</sup>	0.46 ± 0.08 <sup>a</sup>
Threonine	13.14 ± 0.18 <sup>a</sup>	11.67 ± 0.90 <sup>a</sup>	13.18 ± 0.98 <sup>a</sup>	12.08 ± 0.38 <sup>a</sup>	11.96 ± 0.42 <sup>a</sup>
Tryptophan	6.85 ± 0.84 <sup>a</sup>	5.56 ± 0.96 <sup>a</sup>	6.05 ± 0.24 <sup>a</sup>	5.44 ± 0.44 <sup>a</sup>	5.21 ± 0.07 <sup>a</sup>
Valine	16.42 ± 0.55 <sup>a</sup>	14.09 ± 1.44 <sup>a</sup>	15.72 ± 1.28 <sup>a</sup>	14.25 ± 0.31 <sup>a</sup>	13.92 ± 0.58 <sup>a</sup>
ΣEAAs	126.99	110.82	121.91	112.56	111.06
NEAAs					
Argine	21.64 ± 0.22 <sup>ab</sup>	18.13 ± 1.58 <sup>b</sup>	23.60 ± 1.38 <sup>a</sup>	21.04 ± 1.37 <sup>ab</sup>	20.44 ± 2.07 <sup>ab</sup>
Alanine	14.32 ± 0.09 <sup>a</sup>	13.05 ± 0.97 <sup>a</sup>	14.63 ± 1.10 <sup>a</sup>	13.71 ± 0.34 <sup>a</sup>	13.75 ± 0.22 <sup>a</sup>
Glutamic acid	85.31 ± 13.45 <sup>a</sup>	73.76 ± 5.11 <sup>ab</sup>	77.68 ± 5.86 <sup>ab</sup>	65.44 ± 0.92 <sup>ab</sup>	55.85 ± 4.59 <sup>b</sup>
Glycine	17.00 ± 0.33 <sup>a</sup>	14.61 ± 1.01 <sup>b</sup>	15.02 ± 1.22 <sup>ab</sup>	13.98 ± 0.58 <sup>b</sup>	13.53 ± 0.54 <sup>b</sup>
Proline	37.87 ± 2.08 <sup>a</sup>	23.43 ± 4.20 <sup>b</sup>	25.67 ± 2.20 <sup>b</sup>	21.64 ± 0.72 <sup>b</sup>	19.32 ± 1.90 <sup>b</sup>
Serine	16.15 ± 1.32 <sup>a</sup>	14.08 ± 1.21 <sup>ab</sup>	15.00 ± 1.22 <sup>ab</sup>	13.33 ± 0.33 <sup>ab</sup>	12.43 ± 1.00 <sup>b</sup>
Asparagine	20.58 ± 0.41 <sup>ab</sup>	19.50 ± 1.50 <sup>b</sup>	23.03 ± 1.78 <sup>a</sup>	21.29 ± 0.70 <sup>ab</sup>	21.19 ± 0.63 <sup>ab</sup>
ΣNEAAs	212.87	176.56	194.63	170.43	156.51

Histidine, Aspartic acid, Cysteine, Glutamine amino acid: not detected

SFP5, SFP10, SFP15, and SFP20: pasta prepared with 5, 10, 15, and 20 g of salmon fish powder /100 g of semolina flour. CP: control pasta. Results are presented as the mean value  $\pm$  standard deviation,  $n = 3$ ; Values within a column followed by the same superscript letter are not significantly different from each other ( $p > 0.05$ ), according to Tukey's test.




**Figure 1.** Amount of reducing sugar released during *in vitro* digestion for control (C), and pasta containing 5 % salmon fish powder (SFP), 10% SFP, 15 % SFP and 20% SFP respectively.




**Figure 2.** Starch content hydrolysed within 20 min readily digestible starch (RDS) left and within 120 min slowly digestible starch (SDS) right of pasta enriched with 5%, 10 %, 15% and 20% SFP.


The values are expressed as mean  $\pm$  SD (n=3). Different letters showed the significant difference ( $p < 0.05$ ).



**Figure 3.** Values for area under the curve (AUC) comparing control and enriched salmon fish powder (SFP) pasta samples.




**Figure 4.** The pH Vs time curves obtained by pasta made with different concentration of salmon fish powder (SFP) incubated with multi-enzymes (Trypsin, Chymotrypsin and protease).





**Figure 5.** Total phenolic content of pasta enriched with different concentration of (SFP), before digestion and at gastric and pancreatic digestion. Bar represent mean  $\pm$  SD (n=3), followed by different small (before digestion), capital (gastric) and small underlined (pancreatic digestion) letters indicate significant difference among the values at  $p < 0.05$ .



**Figure 6.** Antioxidant activity of pasta enriched with SFP determined with ORAC assay during *in vitro* gastric and pancreatic phase of digestion and before digestion. Results are expressed as Trolox ( $\mu\text{mol g}^{-1}$ ). Data are mean  $\pm$  SD (n=3), followed by small (before digestion), capital (gastric) and small underlined (pancreatic digestion) letters indicate significant difference among the values at  $p < 0.05$ .