

Article

# Improving the Conscious Consumption of Fish Through an Educational Campaign on the Detroit River (Michigan, USA)

Corey A. Krabbenhoft<sup>1</sup>, Susan Manente<sup>2</sup> and Donna R. Kashian<sup>1,\*</sup>

<sup>1</sup> Wayne State University; ckrab@wayne.edu

<sup>2</sup> Michigan Department of Health and Human Services; ManenteS@michigan.gov

\* Correspondence: dkashian@wayne.edu; Tel.: +1-313-577-8052

**Abstract:** Consumption guidelines are a common way to improve conscious consumption behaviors in areas where game fish are known to contain contaminants. However, guideline information can be difficult to distribute, and effectiveness difficult to measure. To increase the distribution and effectiveness of guideline information for the Detroit River, an educational campaign was launched in 2010, which included distribution of pamphlets with consumption information, posting of permanent signs at popular fishing locations, and hiring River Walkers to personally communicate with anglers. In 2013 and 2015, we conducted in-person surveys of active shoreline anglers to determine the effectiveness of education and outreach efforts. Results from the survey indicated that 55% of anglers were aware of the guidelines in 2013, and by 2015 36% had communicated the information to family or friends. However, anglers were often unwilling to reduce consumption of popular game species, despite high contaminant levels. Encouragingly, black anglers were most likely to supplement their diet with species lower in contaminants. Our results suggest that utilizing multiple educational strategies including reaching out directly to individual anglers may improve conscious consumption behavior among the targeted population, providing a template for educational campaigns to successfully target vulnerable populations.

**Keywords:** fish consumption; subsistence fishing; Detroit River; surveys; contamination; consumption advisories; environmental justice; mercury; PCBs; dioxins

## 1. Introduction

While fish can be an important source of daily protein, omega-3 fatty acids, and other essential nutrients, fish may also contain contaminants which are detrimental to human health [1], particularly in self-caught fish from urban areas with a history of chemical contamination. Considerable effort has been extended to inform the public of the benefits and risks of fish consumption at local, state and international (U.S. and Canada) levels. Fish consumption advisories or guidelines are designed to provide citizens with information on fish from local waters that are lower in chemical contamination and are therefore safe meal choices relative to those that are high in contamination. However, advisories typically contain complicated information which is difficult to convey without indiscriminately discouraging fish consumption. Many populations can benefit from fish as a low-cost, readily available addition to their diet; avoidance of fish altogether reduces nutritional opportunities from fish low in contaminants, a problematic potential outcome [2].

Research suggests that to achieve a change in behavior, educational approaches need to be tailored to the particular population and problem and involve direct contact with anglers to be effective [3-5]. The efficacy of advisory programs has been variable and difficult to measure [6]. Criticisms of advisory campaigns suggest that information does not often reach diverse groups of people, or even those groups who may be the most impacted by fish contamination. In particular, previous research has shown that advisory information typically does not effectively reach


minorities, women, people with low levels of educational attainment [7], or immigrant communities [8]. Further, while progress has been made on assessing the impact of advisories, many of these studies primarily include anglers who are white and have moderate income and educational backgrounds [9]. This problem is especially significant in distressed urban environments where anglers are more likely to engage in subsistence fishing due to high poverty rates. Studies have also shown that a higher proportion of those who catch species high in contaminants, keep them, and share them with family and friends are people of color [10,11]. Frequent consumption of fish with high levels of contaminants can contribute to adverse health conditions especially for fetuses, children, and adults with existing chronic health issues such as heart, thyroid or immune diseases [12].

Residents of Detroit, Michigan commonly supplement their food supply with locally caught fish, which are available at low to no cost [13]. These subsistence anglers are primarily low-income, minority individuals [10] who regularly fish for white bass (*Morone chrysops*) and walleye (*Sander vitreus*), which are two of the more contaminated species in this area. As of the 2010 U.S. Census [14], a third of Detroit residents live in poverty (more than twice the state average) and the median household income (\$29,447) was 60% below the state median. Additionally, Detroit had a 20% unemployment rate. Unfortunately, this suggests that the populations with the highest reliance on fish from the Detroit River as a food source are also those groups most difficult to target for educational and outreach efforts.



**Figure 1.** The Eat Safe Fish advisory sign posted initially in 2010 at 28 popular fishing locations along the Detroit River. This updated version was posted in 2015.

Due to possible environmental justice concerns associated with fish contaminant levels on the Detroit River, an intensive educational campaign was launched in 2010 to provide fish consumption guideline information in a targeted way to those individuals who were most susceptible, and thus most likely to benefit from the information. As part of this program, multiple outreach methods were utilized in an attempt to have the largest impact on the shoreline angler population.



**Figure 2.** Map of study location. Blue points indicate locations where 45 signs were posted (some locations had multiple signs). Orange points in inset map indicate home zip codes for anglers surveyed.

The first method was to have a permanent educational presence at fishing locations, through the installation of signs containing consumption guideline information (Figure 1) along the Detroit River in 2010 (and updated in 2015) at 28 locations known to be popular shore-fishing access points (Figure 2). Signs were designed in collaboration with the Michigan Department of Health and Human Services, community focus groups, and communication experts from Wayne State University, to provide information on contaminant levels in the various fish species in the Detroit River and provide guidance on the safest consumption practices. Local community focus groups were involved in evaluating both the signs and pamphlets during the design phase of these materials.

A second outreach method was designed to confront deep cultural preferences and people’s own interpretation of risk which often hinder behavioral change. Social norms and community practices may intercede between effective interventions and adequate uptake into daily life [15]. To overcome this challenge in information distribution, two to three Detroit residents were hired as River Walkers, beginning in 2012 and continuing through 2016 who assisted outreach efforts. By visiting shore-fishing sites and directly communicating with active anglers, River Walkers provided information on eating locally caught fish and offered hard copies of educational materials (a third outreach method). River Walkers were able to distribute materials including 1) an “Eat Safe Fish in the Detroit Area” pamphlet, 2) Eat Safe Fish Guidelines educational pamphlet 3) a “Hooked on Fish from the Great Lakes” cookbook, 4) a fishing crossword puzzle and word search for children, and 5) temporary tattoos. This personal interaction with an informed, local individual provided an easy way for anglers

to express concerns or get answers to their questions. It also provided a face to the consumption guideline campaign.

The overall objective of this study was to assess the progress, strengths, and weaknesses of an educational outreach program designed to improve conscious consumption practices of fish from the Detroit River. Specifically, this study evaluated the overall awareness of fish consumption guidelines among anglers, their knowledge of what the guidelines contained, which methods were most effective in implementing changes in behavior, and potential environmental justice issues associated with either contamination exposure or the consumption guideline campaign itself.

## 2. Materials and Methods

### 2.1 Study Location

The Detroit River is a narrow, 45 km long strait connecting Lake St. Clair and Lake Erie and forms part of the border between Canada and the United States. The cities of Detroit, Michigan and Windsor, Ontario lie along the flow path of the river. The river is an important shipping route in the Great Lakes and has long been utilized as a fishing location, particularly for subsistence anglers. Shore fishing on the Detroit River accounted for nearly a million angler-hours when it was measured in the 1980s [16]. In a 2002 creel census Thomas and Towns [17] estimated that 40% of all Great Lakes fishing occurs in only 1% of Michigan's Great Lakes waters, specifically the St. Clair System which includes the Detroit River, Lake St. Clair and the St. Clair River. The Detroit River was the primary location for consumption guideline awareness efforts and angler surveys in this study (Figure 2).

Contamination of fish in the Detroit River has a long history [18]. A large concern for local anglers are the levels of polychlorinated biphenyls (PCBs), mercury, and dioxins [19]. Some species of fish from the Detroit River contain high levels of contaminants and should not be eaten in any amount (catfish, carp), and others have consumption restrictions but can be eaten occasionally (rock bass, perch, etc.). Depending on location, fish can sometimes be safely consumed without concern for contaminants, but no species from the Detroit River fall into this category.

### 2.2 Surveys

Face-to-face angler surveys were conducted from May to September 2013 and May to August 2015, four and six years after the initial educational programs were implemented. Of the 28 areas where fish consumption signs had been posted, surveys were conducted at 12 locations in 2013 and 17 locations in 2015 (though most sites were visited multiple times). People who were actively fishing were approached and asked to participate in a survey about their knowledge of fish consumption guidelines and their fishing activity. Participants were informed of the purpose of the survey and provided the opportunity to ask questions prior to consenting to participate; questions were welcome through the duration of the survey. No compensation was given for participation, but supplemental guideline materials were provided for those participants who were interested in further information. The survey was conducted verbally and took approximately ten minutes to complete.

Survey questions were designed to evaluate the state of knowledge of the guidelines among anglers and the effectiveness of the 1) signs produced by the State of Michigan, 2) River Walkers, and 3) the Eat Safe Fish educational pamphlet. Incorporation of multiple educational methods was an important part of this study because of the diversity of anglers in the study area. Fishing behaviors of each angler and demographic questions were asked to allow for an evaluation of potential environmental justice issues surrounding fish consumption advisories on the Detroit River [10]. Specifically, anglers were asked about their awareness of river contaminants and the Eat Safe Fish Guidelines, how s/he learned of that information, which material/s s/he found to be most helpful, or whether the information had led to any behavioral change on her/his part (e.g., the fish species consumed, the cooking methods, etc.). No names or otherwise identifying information were collected from the angler.




2.3 Data Analysis

Surveys were analyzed for general information and for between year differences. Comparisons of answers from 2013 to 2015 were analyzed using Chi-square ( $X^2$ ) goodness of fit tests. Contingency tables of survey responses and demographic information were assessed for correlation using Fisher's Exact Tests for categorical data that included low numbers for some responses (e.g., income range) and Yates' chi-squared tests for those that had more equal distributions among response variables (e.g., gender). For continuous data (e.g., number of meals consumed per week), Kruskal-Wallis tests were used to determine difference between years and simple logistic regressions were used to determine correlation with demographic variables. Diversity of fish species consumed was calculated using Shannon's Diversity Index [20]. Difference in the timing of surveys between years was assessed using the Mann-Whitney-Wilcoxon test for non-normal distributions with unequal variances on the Julian dates for surveys from each year.

All data analysis was done using the statistical software R [21]. The add-on package MASS [22] was used to complete chi-squared tests.

3. Results



**Figure 3.** Demographics of survey participants in 2013 and 2015. (A) Angler age ranges (by decade); (B) angler educational attainment; (C) angler racial identity; (D) angler gender identity; and (E) angler household income range (in increments of \$25,000). U.S. census data combines


161 ‘White/Caucasian’ and ‘Arab/Middle Eastern’ into a single category; our data was similarly combined  
162 for comparison. ‘Asian’ includes the Far East, southeastern Asia, and India. ‘American Native’  
163 includes first peoples of North America including Alaska Natives and native Pacific Islanders.

164 A total of 431 shoreline anglers were surveyed (200 in 2013 and 231 in 2015). The anglers  
165 represented 87 of 1160 Michigan zip codes and one Ohio zip code, most of which are located within  
166 25 km of the Detroit River (Figure 2). For each year, the majority of the survey participants self-  
167 identified as African American/Black (68% in 2013; 62% in 2015), with smaller proportions identifying  
168 as White/Caucasian, Asian/Pacific Islander, Arab/Middle Eastern, Hispanic/Latinx, American Indian  
169 or Alaskan native, or other (Figure 3C). Most of the anglers surveyed were male (82% in 2013; 70% in  
170 2015), though an increase in female anglers from 18% in 2013 to 30% in 2015 was observed (Figure  
171 3D). Anglers ranged in age from 18 to 85, with the largest proportion of anglers being in their 50s in  
172 2013, and their 30s in 2015 (Figure 3A). In 2013, 34% of the surveyed anglers reported an annual  
173 household income less than \$25,000 with 34% reporting a household income in the range of \$25,000-  
174 49,999 and 32% reporting a household income higher than \$50,000 (Figure 3E). Income ranges were  
175 similar in 2015 with 39% reporting an income less than \$25,000, 42% reporting a household income  
176 between \$25,000-\$49,000, and 20% reporting an income greater than \$50,000 (Figure 3E). In 2013, 10%  
177 of anglers had less than a high school diploma, while 43% of anglers had received a high school  
178 diploma or GED, and 48% had at least some college or post-high school education (Figure 3B).  
179 Similarly, in 2015, 19% had less than a high school diploma, 37% had a high school diploma or GED,  
180 and 44% of anglers had at least some post-high school education (Figure 3B).

181 *3.1 Fish Consumption*


182 Most anglers surveyed (77% in 2013 and 68% in 2015) reported that fish are at least somewhat  
183 important in their diets. In both survey years, anglers reported consuming the most fish in the  
184 summer and spring seasons, which coincide with typical spawning runs of most species. During this  
185 time, anglers were observed in the greatest numbers along the shoreline. The number of meals per  
186 week consumed was much higher in 2015 than 2013 ( $P < 0.001$ ;  $X^2 = 146.78$ ). However, this difference  
187 may be due to the addition of serving size estimates included in the 2015 survey. Because a ‘serving  
188 size’ may be different among individuals, the 2015 survey included a definition of ‘serving size’ as  
189 being approximately the size of one’s hand. This would increase the total number of servings  
190 reported if those servings were generally multiple hand-sized portions, which were reported as a  
191 single serving in the 2013 survey. Our data shows that the mean number of servings, as defined by a  
192 ‘hand-sized’ serving of fish, consumed per week by the anglers in 2015 was  $7.5 \pm 0.25$  in spring and  
193  $6.2 \pm 0.25$  in summer, or roughly one serving per day. Consumption fell to  $0.99 \pm 0.25$  servings per  
194 week in the fall and  $0.11 \pm 0.26$  in the winter, or less than one serving per week.  
195

Anglers reported frequently providing fish to family and friends with 76% in 2013 and 69% in 2015 reporting giving fish away (Figure 4A). Of that which is given away, 52-59% of anglers reported providing fish to children, 78-83% to women, and 61-92% to men (depending on survey year- Figure 4B). Encouragingly, almost all anglers (94-98%) reported removing the head, skin, fat, and/or organs prior to cooking. In 2013, typical cooking methods were also addressed. Ninety percent of anglers regularly fried their fish (a cooking method not recommended due to its inability to eliminate fats containing PCBs and dioxins). Baking and grilling (more preferred methods) were less common with only 34% and 21% of anglers, respectively, cooking their fish this way. Finally, fewer than 5% of anglers reported boiling, broiling, smoking, and/or steaming their catch.



**Figure 4.** (A) Percent of surveyed anglers who reported giving away some of their catch in 2013 and 2015. (B) Of those anglers who gave fish away, the percent who gave their fish to Children, Women, and/or Men (total number of anglers who answered questions differed between years).

White bass (*Morone chrysops*; also locally referred to as silver bass or occasionally stripe bass) was the most common species taken home by anglers. Walleye (*Sander vitreus*) was the next most common species followed by yellow perch (*Perca flavescens*), smallmouth bass (*Micropterus dolomieu*; locally brown bass) and catfish (typically channel catfish, *Ictalurus punctatus*). Some differences were identified in the species anglers commonly took home between 2013 and 2015 (Figure 5). Specifically, white bass, walleye, and smallmouth bass were all more commonly kept in 2015 than 2013 ( $P = 0.003$ ;  $X^2 = 39.28$ ). Additionally, a larger diversity of species was kept in 2013 than 2015 ( $H' = 3.00$  in 2013,  $H' = 2.33$  in 2015). Species with higher restrictions (as indicated by the Eat Safe Fish program) were more commonly kept in 2015 than 2013 ( $P = 0.021$ ;  $X^2 = 9.70$ ; Figure 7). However, this trend was driven by an increase in walleye (46% to 72%) and white bass (75% to 90%), while consumption of catfish decreased (35% to 32%). Additionally, the mean timing of surveys was later in 2015 than in 2013 ( $P < 0.001$ ) by a mean of 6.21 Julian days (median difference of 16 Julian days), which may account for skewing of data toward a particular subset of species (as spawning runs for specific species are typically distinct events and attract a large number of anglers, particularly for white bass).



**Figure 5.** Fish species taken home by anglers in 2013 (A) and 2015 (B) with their relative suggested serving frequencies indicated.

### 3.2 Consumption Guideline Awareness

During the two-year period between surveys, general awareness of the fish consumption guidelines did not change (55% in 2013, 57% in 2015;  $P = 0.766$ ,  $X^2 = 0.088$ ). A larger proportion of anglers saw the signs and/or encountered the River Walkers in 2015 than 2013 ( $P = 0.005$ ,  $X^2 = 7.76$ ;  $P < 0.001$ ,  $X^2 = 54.26$ ). However, fewer people who saw the sign in 2015 actually read it ( $P = 0.011$ ,  $X^2 = 6.45$ ), perhaps suggesting some decline in interest over time as people become accustomed to the signs the longer they are present. Demographic factors had little impact on anglers' awareness of the guidelines, with race, income, and gender having insignificant correlation with angler awareness of the guidelines ( $P = 0.814$ ,  $0.198$ ,  $0.149$ , respectively). However, education was correlated with awareness ( $P = 0.007$ ); individuals with higher educational achievement (some college, trade school, or more) were more commonly aware of the guidelines (69%) than individuals with lower educational achievement (high school degree, GED, or less; 46%). Encouragingly, anglers who were aware of the guidelines were 5% more likely to supplement their diet with species lower in contaminants. Additionally, 36% of anglers shared information about the guidelines with friends or family.


The different methods utilized in the study were differentially effective between years. The River Walkers, signs, and educational pamphlets were all reportedly more effective in 2013 than in 2015 (Table 1). In 2013, all methods were reported as helpful to the anglers, with no significant differences for any pairwise comparison. Pamphlets had an influence on 51% of anglers in 2013, but this number fell to 13% in 2015. In 2015, the signs were reportedly more helpful to anglers than were the River Walkers. However, a higher percentage of anglers reported that the sign was confusing in 2015 than in 2013 (up from 15% to 23%). No correlation with race, education, or income was found with a positive response to any particular outreach method ( $P = 0.119 - 0.955$ ).

**Table 1.** Statistical summary of outreach methods. Data was derived from "Yes" or "No" answers regarding whether anglers found each method to be helpful to them. Percent of angler



responses indicating a method was helpful are indicated and comparisons between methods and years are included. *P* indicates p-value from chi-squared or Fisher’s Exact Test comparisons. Significant comparisons appear in bold.

Method	% anglers who reported method as helpful		<i>P</i>
	2013	2015	2013 vs. 2015
River Walkers	94	47	<b>&lt;0.001</b>
Signs	93	65	<b>&lt;0.001</b>
Pamphlets	89	52	<b>&lt;0.001</b>
<hr/>			
Pairwise Comparisons	<i>P</i>	<i>P</i>	<i>P</i>
	2013	2015	2013 vs. 2015
RW vs. Signs	0.999	<b>0.006</b>	-
RW vs. Pamphlets	0.531	0.460	-
Signs vs. Pamphlets	0.462	0.052	-
RW 2013 vs. 2015	-	-	<b>&lt;0.001</b>
Signs 2013 vs. 2015	-	-	<b>&lt;0.001</b>
Pamphlets 2013 vs. 2015	-	-	<b>&lt;0.001</b>



**Figure 6.** The percentage of anglers who reported various outreach methods as helpful (A), and the behavioral changes implemented by anglers in 2015 (B).

Anglers did report having implemented some behavioral changes due to consumption suggestions. The largest reported change was in the species consumed by anglers (29%). Cooking method (21%) and fishing method or location (9%) were also impacted by the consumption campaign (Figure 6). Walleye and white bass were the most popular species consumed by all anglers, but those who reported having made changes to the species they consume were 40% more likely to supplement their diet with species with lower contamination levels in 2015. Education, income, and gender had no significant impact on changes in behavior. However, race was significantly correlated with reports of changes in species anglers chose to consume ( $P = 0.002$ ). Specifically, participants who self-identified as White/Caucasian were less likely to have changed the species they consumed than those who were Hispanic/Latinx ( $P = 0.008$ ), though the small sample sizes for all races other than African American/Black may have influenced this outcome. This trend was thus driven by African American/Black participants who reported changing the species they consume at a higher rate than other participants (34% as opposed to a combined 16% for all other participants).

**4. Discussion**

The history of contamination in Detroit and the surrounding area make understanding the relative benefits and risks of fish consumption difficult to grasp [18]. Despite challenges in designing effective educational campaigns, methods tailored to the specific population of anglers as described in this study have demonstrated gains in general knowledge among Detroit anglers. Following a multifaceted educational campaign that began in 2010, and included one-on-one interactions with anglers beginning in 2012, we were able to document awareness of fish consumption advisories by 2013, a period just three years after the start of the program. Anglers reported significant behavioral changes in the fish species they consumed by 2015. Despite awareness of the consumption guidelines not having changed over our two-year survey period, there have been significant improvements in the conscious consumption of fish among Detroit anglers.

Contrary to previous findings [23], African American/black anglers were more likely to supplement their diet with lower-risk species than anglers of other races. This finding is encouraging given that minorities are disproportionately affected by contaminants through fish consumption [23,24,9]. The survey results also indicated areas where educational efforts may be improved; for example, all outreach methods were reported as highly effective in the 2013 survey but were less helpful in 2015. Specifically, the educational pamphlet influenced anglers at the highest rate initially, but significantly decreased in influence by 2015. Further, fewer anglers in 2015 read the signs, and we suspect they may have become part of the ‘background’ of the landscape. This may indicate some level of saturation of knowledge following initial efforts, which corresponds to a drop in new interest in subsequent years. Overall, these trends demonstrate that outreach efforts need to vary over time to reach a broad audience and be maximally effective.

In terms of angler behavior, fish species of greater concern were still some of the most consumed in the later year of the surveys. Overall, this may indicate relative willingness to adjust behaviors with respect to specific species, particularly those like walleye which have deep cultural importance for recreational fishing in this region [25,26]. Indeed, walleye and white bass were still the most commonly consumed species among anglers who reported having made changes to the species they consume; however, anglers who were aware of the guidelines or reported making behavioral changes were also willing to supplement their diets with species reported to have lower contaminant levels. This suggests that fish species which are not specifically sought after (due to local importance or lower abundance) may provide greater opportunity for angler behavior change. For example, anglers may be unwilling to remove walleye from their diet but might consume yellow perch instead of catfish. Species like catfish, which have relatively high contaminant loads but are not favorites among anglers may thus provide the greatest opportunity for overall improvement in consumption trends, as resistance to decreasing consumption of those species will be lower.

Race and education were both correlated with overall consumption guideline awareness and the implementation of behavioral change. This may indicate some cultural implications in the perception of messages [15]. In some cases, despite being aware of guideline suggestions, anglers were not

308 amenable to the overall message, potentially indicating a mistrust of the information [27]. The  
309 suggestions provided may confront generational or cultural tradition, which can make receipt of the  
310 information difficult and can hinder implementation of behavioral suggestions [28]. Incorporation of  
311 outreach methods which address familial and cultural concerns over the guideline suggestions need  
312 to be considered in designing outreach efforts in this and other systems. Importantly, anglers did  
313 report relaying consumption guidelines to friends and family members. Encouraging dissemination  
314 of information to friends and family may be a way to improve overall awareness as it allows  
315 information to flow through inherently trustworthy sources. Further, utilizing relationships with  
316 stakeholder groups in the area could also improve translation of research to active anglers as well as  
317 provide educational campaigns with appropriate techniques to effectively engage the public [29].

318 In this study, several techniques were assessed for their value in informing local anglers. Signs  
319 were posted, River Walkers were hired to engage anglers, and pamphlets were provided with  
320 information on consumption guidelines. The assessment using surveys allowed direct feedback on  
321 guideline efforts from the target population. Of the outreach efforts utilized, the highest percentage  
322 of anglers reported that the signs were most helpful in the later survey year. This suggests that  
323 location-specific visual aids which anglers can engage on their own time are important in reaching  
324 anglers and that efforts including such resources may be more effective [30]. However, the percent of  
325 anglers who found each method to be helpful changed between years which may indicate anglers are  
326 not always amenable to the guideline message or perhaps a decline in the number of anglers who  
327 were amenable to the message were encountered due to survey efforts that took place later in the  
328 fishing season. Over the course of this study, we identified an increase in the proportion of female  
329 anglers as well as a decrease in the mean age of the angler population. This is consistent with general  
330 trends observed in angling communities [31] and will be important in the design of future advisories,  
331 particularly as women of child-bearing age are increasingly engaging in sport fishing, especially in  
332 the Great Lakes [32]. As seen in this study, outreach methods are not equally effective, and better  
333 understanding the changing demographics of the audience will aid in designing more effective  
334 educational programs.

335 Importantly, this study surveyed the same geographic population of anglers (though  
336 individuals varied) over multiple years to assess changes in behavior associated with educational  
337 outreach efforts. Although consumption studies are relatively common, and those which survey  
338 anglers have produced similar data [e.g., 33], few occur over multiple years to assess longer term  
339 changes and retention of guideline information within a population [34]. This type of repeated  
340 sampling is necessary to ensure ongoing impact of consumption guidelines, particularly as  
341 information is updated and the angler demographics change. In the case of the Detroit River,  
342 consumption guidelines are updated annually so it is imperative that anglers are made aware of  
343 recommendations on a continual basis.

344 This study adopted a unique approach to evaluating the progress of fish consumption guidelines  
345 awareness. The face-to-face interaction with individuals who were actively fishing ensured that the  
346 target group was reached [35]. This strategy differs from a majority of previous efforts which relied  
347 on phone surveys [36,7], online questionnaires, or face-to-face surveys occurring in general public  
348 areas not specific to fishing activities [37]. Further, the anglers who participated in the surveys  
349 occupied demographic groups (low income and education, racial minorities) traditionally missed in  
350 these types of studies despite being at high risk. This unique design is particularly important for areas  
351 with high rates of poverty, such as Detroit, where literacy and access to communication services may  
352 be low. Further, this study focused on a specific body of water rather than obtaining data on a larger,  
353 regional scale [e.g., 36]. Data collected from surveys at a regional scale may be difficult to extrapolate  
354 to local water bodies and fishing activities [e.g., 7]. Providing consumption materials and conducting  
355 surveys in person at locations where anglers are most likely to be affected by contamination ensures  
356 that those individuals who are most likely to be impacted are receiving the information, in addition  
357 to providing the greatest possible accuracy in measuring the impact of consumption guideline efforts.  
358 This practice also allows for the ability to regularly adapt education and outreach strategies based on  
359 feedback directly provided by anglers.

Data from anglers fishing the Detroit River is invaluable in setting priorities in fish consumption education and outreach that are meaningful and appropriate for this population and in designing future activities. Specifically, identifying the most successful methods of outreach for particular groups and over time is an important contribution of this study. Educating anglers and their families about the benefits of eating safer fish and the risks of eating the most contaminated fish can affect behavior change and will help protect the health of urban anglers who depend upon local fish as a food source. Educating anglers about choosing the safest fish to eat and the appropriate cooking and cleaning methods has the potential to improve conscious consumption behaviors among Detroit residents and create a more informed public over time. This study can be used as a template going forward to continue the efforts to improve educational campaigns in Detroit, as well as a guideline for evaluating fish consumption advisories elsewhere.

**Author Contributions:** Conceptualization, S.M. and D.K.; methodology, S.M. and D.K.; formal analysis, C.K.; investigation, D.K. and C.K.; resources, D.K.; data curation, C.K. and D.K.; writing—original draft preparation, C.K.; writing—review and editing, C.K., S.M. and D.K.; visualization, C.K.; supervision, D.K.; project administration, D.K.; funding acquisition, D.K.

**Funding:** Funding for this study was provided by the Erb Family Foundation 12-0433, and the Michigan Department of Health and Human Services 20142506-00.

**Acknowledgments:** Prior to this study the ground work for this project was initiated with a grant to D. Kashian funded by Michigan Sea Grant. Thanks to our surveyors Karim Alame, Krystal Bakkila, and Ali Shakoor and our River Walkers, Robert Woods, Shameka Walker, and James Bridgforth. This project was guided by the efforts of the Detroit River Fish Advisory Community Advisory Board: Robert Burns, Friends of the Detroit River; Mary Bohling, Michigan Sea Grant; Sherri Briller, Andrea Sankar, and Mark Luborsky, Wayne State University; Kory Groetsch, L Gossiaux, and Jennifer Gray, Michigan Department of Health and Human Services; Paul Max and Paul Harbin, City of Detroit Department of Health and Wellness Promotion; Charles Stokes, Detroiters Working for Environmental Justice; Liz Hay-Chmielewski, Michigan Department of Natural Resources; Mike Reed, Detroit Zoo/Belle Isle Nature Center; Robert Woods, Shameka Walker, and Jim Bridgforth, River Walkers; Bob Reider and Matt Shalkeford, DTE Energy; and Joe Bohr, Michigan Department of Environmental Quality. This study was conducted under IRB # 058012B3E. Data from this study will be submitted to Dryad digital repository upon notification of acceptance for publication.

**Conflicts of Interest:** The authors declare no conflict of interest.

References

1. U.S. Department of Health and Human Services and U.S. Department of Agriculture (USDHHS and USDA). 2015-2020 Dietary Guidelines for Americans. 8th ed; 2015. Available online: <https://health.gov/dietaryguidelines/2015/guidelines/> (accessed 25 October 2018).
2. Oken, E.; Choi, A.L.; Karagas, M.R.; Mariën, K.; Rheinberger, C.M.; Schoeny, R.; Sunderland, R.; Korrick, S. Which fish should I eat? Perspectives influencing fish consumption choices. *Environ. Health Perspect.* **2012**, *120*, 790-798.
3. Glanz, K.; Rimer, B.K.; Lewis, F.M. *Health Behavior and Health Education: Theory, Research and Practice*, 3rd ed.; Wiley & Sons: San Francisco, USA, 2002, B01K0S0ZBU.
4. Kelley, M.A.; Baldyga, W.; Barajas, F.; Rodriguez-Sanchez, M. Capturing change in a community-university partnership: the ¡Sí Se Puede! project. *Prev. Chronic Dis.* **2005**, *2*, A22.
5. McLeroy, K.; Norton, B.L.; Kegler, M.C.; Burdine, J.N.; Sumaya, C.V. Community based interventions. *Am. J. Public Health* **2003**, *93*, 529-533.
6. Reinert, R.E.; Knuth, B.A.; Kamrin, M.A.; Stober, Q.J. Risk assessment, risk management, and fish consumption advisories in the United States. *Fisheries* **1991**, *19*, 5-12.
7. Tilden, J.; Hanrahan, L.P.; Anderson, H.; Palit, C.; Olson, J.; Kenzie, W.M.; and the Great Lakes Sport Fish Consortium. Health advisories for consumers of Great Lakes sport fish: is the message being received? *Environ. Health Perspect.* **1997**, *105*, 1360-1365.
8. Lauber, T.B.; Connelly, N.A.; Niederdeppe, J.; Knuth, B.A. Urban anglers in the Great Lakes region: fish consumption patterns, influences, and responses to advisory messages. *Sci. Total Environ.* **2017**, *590*, 495-501.



9. Lauber, T.B.; Connelly, N.A.; Niederdeppe, J.; Knuth, B.A. Urban anglers' adherence to fish consumption advisories in the Great Lakes region. *J. Great Lakes Res.* **2017**, *43*, 180-186.
10. Kalkirtz, V.; Martinez, M.; Teague, A. Environmental Justice and Fish Consumption Advisories on the Detroit River Area of Concern. M.S. Thesis, University of Michigan, Ann Arbor, MI, U.S., May 2008.
11. Michigan Department of Community Health. Fish consumption survey of people fishing and harvesting fish from the Saginaw Bay Watershed; Report for the Saginaw Bay Watershed Initiative Network: Bay City, MI, USA, 2007. Available online: [www.michigan.gov/documents/mdch/FCS\\_Final\\_rpt\\_061407\\_199288\\_7.pdf](http://www.michigan.gov/documents/mdch/FCS_Final_rpt_061407_199288_7.pdf) (accessed on 26 October 2018).
12. Food and Nutrition Board, Institute of Medicine (IOM). Seafood Safety; Ahmed, F.E., Ed.; National Academy Press: Washington, D.C., USA, 1991, 0-309-04387-5.
13. West, P.; Fly, M.; Larkin, F.; Marans, R.W. Minority anglers and toxic fish consumption: evidence from a statewide survey of Michigan. In *Race and the incidence of environmental hazards: a time for discourse*; Bryant, B., Mohai, P., Eds. Westview Press: Boulder, CO, USA, 1992; 081338513X.
14. U.S. Census Bureau, American Fact Finder. Available online: [https://factfinder.census.gov/faces/nav/jsf/pages/community\\_facts.xhtml?src=bkmk](https://factfinder.census.gov/faces/nav/jsf/pages/community_facts.xhtml?src=bkmk) (accessed 25 October 2018).
15. Dellinger, J.A. Exposure assessment and initial intervention regarding fish consumption of tribal members of the Upper Great Lakes Region in the United States. *Environ. Res.* **2004**, *95*, 325-340.
16. Haas, R.; Bryant, W.; Smith, K.; Nuhfer, A. Movement and harvest of fish in the Lake St. Clair, St. Clair River, and Detroit River. Michigan Department of Natural Resources, Fisheries Division. Winter Navigation Study – Final Report. 1985; Available online: [www.dtic.mil/cgi-bin/GetTRDoc?AD=ADA204616](http://www.dtic.mil/cgi-bin/GetTRDoc?AD=ADA204616) (accessed on 25 October 2018).
17. Thomas, M.; Towns, G. Summary of creel results for the St. Clair River, Lake St. Clair, and the Detroit River, 2002-2005. State of Michigan Department of Natural Resources; 2011. Available online: [http://www.michigan.gov/documents/dnr/2011-1tr\\_362659\\_7.pdf](http://www.michigan.gov/documents/dnr/2011-1tr_362659_7.pdf) (accessed online 25 October 2018).
18. Fallon, M.E.; Horvath, F.J. Preliminary assessment of contaminants in soft sediments of the Detroit River. *J. Great Lakes Res.* **1985**, *11*, 373-378.
19. Gandhi, N.; Drouillard, K.G.; Arhonditsis, G.B.; Gewurtz, S.B.; Bhavsar, S.P. Are fish consumption advisories for the Great Lakes adequately protective against chemical mixtures. *Environ. Health Perspect.* **2016**, *125*, 586-593.
20. Shannon, C.E.; Weaver, W. *The Mathematical Theory of Communication*; University of Illinois Press: Urbana, IL, USA; 1963. 9780252725487.
21. R Core Team. R: a language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. Available online: <https://www.R-project.org/> (accessed 25 October 2018).
22. Venables, W.N.; Ripley, B.D. *Modern Applied Statistics*, 4th ed.; Springer: New York, USA, 2002; 0387954570.
23. Beehler, G.P.; McGuinness, B.M.; Vena, J.E. Polluted fish, sources of knowledge, and the perception of risk: contextualizing African American anglers' sport fishing practices. *Hum. Organ.* **2001**, *60*, 288-297.
24. Chess, C.; Burger, J.; McDermott, M.H. Speaking like a state: environmental justice and fish consumption advisories. *Soc. Nat. Resour.* **2005**, *18*, 267-278.
25. Hopper, M.; Power, G. The fisheries of an Ojibwa community in northern Ontario. *Arctic* **1991**, 267-274.
26. Hudson, J.C.; Ziegler, S.S. Environment, culture, and the Great Lakes fisheries. *Geogr. Rev.* **2014**, *104*, 391-413.
27. Hornbarger, K.; MacFarlene, C.; Pompa, C.R. Target audience analysis: recommendations for effectively communicating toxic fish consumption advisories to anglers on the Detroit River. In Natural Resources Sociology Lab Technical Report #11; Natural Resource Sociology Research Lab, University of Michigan: Ann Arbor, MI, USA, 1994.
28. Habron, G.; Barbier, M.; Kinnunen, R. Local understanding of fish consumption advisory risks in Michigan's Upper Peninsula: the role of structure, culture, and agency. *Rural Sociol.* **2008**, *73*, 275-299.
29. Kashian, D.R.; Krause, A.E.; Drouillard, K.G.; Sano, L.L.; Nowell, B. Capacity building in stakeholders around Detroit River fish consumption advisory issues. *Freshw. Sci.* **2014**, *33*, 674-678.



464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484

30. Shubat, P.J.; Raatz, K.A.; Olson, R.A. Fish consumption advisories and outreach programs for southeast Asian immigrants. *Toxicol. Ind. Health* **1996**, *12*, 427-434.

31. Winkler, R.L.; Burkett, E.M. Upper Great Lakes States' Angler Estimates and Map Book Documentation, 1999-2015. Michigan Technological University: Houghton, MI, USA, 2018. Available online: <https://www.mtu.edu/greatlakes/fishery/pdfs/angler-database-documentation-2018.pdf> (accessed 25 October 2018).

32. Connelly, N.A.; Lauber, T.B.; Niederdeppe, J.; Knuth, B.A. Fish consumption among women anglers of child bearing age in the Great Lakes region. *Environ. Res.* **2016**, *150*, 213-218.

33. Burger J.; Gochfeld, M. Knowledge about fish consumption advisories: a risk communication failure within a university population. *Sci. Total Environ.* **2008**, *390*, 346-354.

34. Burger J. Fishing, fish consumption, and awareness about warnings in a university community in central New Jersey in 2007, and comparisons with 2004. *Environ. Res.* **2008**, *108*, 107-116.

35. Burger J.; Campbell, K.R. Fishing and consumption patterns of anglers adjacent to the Oak Ridge Reservation, Tennessee: higher income anglers ate more fish and are more at risk. *J. Risk Res.* **2008**, *11*, 335-350.

36. Imm P.; Knobeloch, L.; Anderson, H.A.; the Great Lakes Sport Fish Consortium. Fish consumption and advisory awareness in the Great Lakes Basin. *Environ. Health Perspect.* **2005**, *113*, 1325-1329.

37. Burger, J. Fishing, fish consumption, and knowledge about advisories in college students and others in central New Jersey. *Environ. Res.* **2005**, *98*, 268-275.