

Article

Physicochemical and sensory characteristics of a Chagalapoli fruit (*Ardisia compressa*) beverage fermented using *Saccharomyces cerevisiae*

Ana Flores-García¹, Rubén Márquez-Meléndez¹, Erika Salas-Muñoz¹, Guillermo Ayala-Soto¹, Iván Salmerón-Ochoa¹, León Hernández-Ochoa^{1*}.

¹ Universidad Autónoma de Chihuahua, Facultad de Ciencias Químicas, Departamento de Posgrado. Circuito Universitario, Campus Universitario #2, Chihuahua, México. C.P. 31125.

* Correspondence: lhernandez@uach.mx, Tel/Fax: +52(614)2366000 ext.4254.

Abstract: Chagalapoli fruit (*Ardisia compressa*) is similar to *Vaccinium myrtillus* (berries) with high-polyphenol content. The objective of this study was to evaluate the physicochemical properties of Chagalapoli fruit and to determine the conditions for the preparation of a fermented beverage using *Saccharomyces cerevisiae* yeast, evaluating the impact on sensory properties. The fermentation process lasted 4 days at 27 °C, with absence of light and a fixed pH of 3.8. The phenolic contents obtained in samples were 1.27 epicatechin mg/mL in filtered juice, 1.59 epicatechin mg/mL in filtered fermented beverage, 1.91 epicatechin mg/mL in partially filtered juice and 3.19 epicatechin mg/mL in partially filtered fermented beverage. An affective test was carried out to determine the sensory acceptability of the final product, evaluating the flavor, color and aroma parameters. The fermented beverage with the greatest preference on color and flavor attributes was the partially filtered fermented beverage.

Keywords: Chagalapoli (*Ardisia compressa*); fermented beverage; phenolic compounds

1. Introduction

Chagalapoli (*Ardisia compressa* subsp. *Myrsinaceae*) belongs to the *Myrsinaceae* family, an extensive family of trees with approximately 500 species [1]. The chagalapoli fruit has a round shape of approximately 12 mm in diameter, with a purple red color that turns black once it reaches maturity; its shell is smooth and delicate, its seed represents up to 50% of its total weight, and the ripe fruit has a bittersweet, mildly astringent taste [2]. In Mexico it is distributed in the states of Chihuahua, Chiapas, San Luis Potosí, Tlaxcala, Hidalgo, Aguascalientes, Puebla, Tamaulipas and Veracruz [3]. Chagalapoli fruit has a similar composition to common berries with a high phenolic content; that is the reason for the high interest in this fruit from a health point of view [4]. The fruit is used to make food products such as juices, jams, and liqueurs, considered to possess antioxidant and antimicrobial properties. In the state of Veracruz the fruit is used against digestive diseases [5]. Chandra & Mejia [6] quantified the total polyphenols of the *Ardisia compressa* leaf extract, obtaining values of 0.58 mg of gallic acid equivalent per mL mgGAE/mL DL. They identified the presence of gallic acid, catechin, epicatechin gallate, ardisin and kaempferol. Heredia-Vasquez [3] determined the total polyphenol content of chagalapoli fruit and obtained 1.74 mgGAE/mL in *Ardisia compressa kunth*; Joaquín-Cruz, et al., [4] obtained 1.051±0.43 mgEAG/g and identified the polyphenols of flavonols, flavan-3-ols (catechin and proanthocyanidin dimers) and hydroxycinnamoyl derivatives. Jácome-Hernandez [2] obtained 1,638.12±74.98 mgGAE/kg in dry weight (DW). Fermented beverages, such as grape wines, have been associated with health benefits due to the presence of large amounts of phenolic compounds, which are claimed to possess antioxidant properties and have an important role in the prevention of deleterious processes such as ageing, diabetes, cancer, neurological disorders, atherosclerosis and cardiovascular diseases [7]. Moreover, several studies

have shown that these beverages induce relaxation in isolated vessels, which makes them important allies for cardiovascular protection [8]. This means that it is necessary to carry out research studies focused on the physicochemical characterization of the fruit and the process of making a fermented beverage that will promote the potential of the chagalapoli fruit. For this reason, the aim of this work was to evaluate the physicochemical properties of the chagalapoli (*Ardisia compressa*) fruit and determine the conditions for the preparation of a fermented beverage using *Saccharomyces cerevisiae*, evaluating the impact on the physicochemical and sensory properties of the final product.

2. Material and methods

2.1. Materials

Acetonitrile and formic acid were high-performance liquid chromatography grade. Glucose, Epicatechin, Kaempferol, Quercetin, Catechin, Chlorogenic and Folin-Ciocalteu reagents were provided by Sigma-Aldrich. Milli-Q water was produced using an Elix Millipore water purification system.

2.2. Raw material

Chagalapoli (*Ardisia compressa*) fruit was collected in Aguascalientes, Mexico. 20 kg were manually selected and washed with chlorine (0.05 mL/L). Samples were stored at 4 °C for further analysis.

2.3. Characterization of raw material

Proximal analysis of chagalapoli (*Ardisia compressa*) was determined according to the AOAC methods; protein (976.05) – determined by Micro Kjeldahl (Nx6.25) –, moisture (934.06), ash (942.05), fat (920.39), crude fiber (962.09) and carbohydrates.

2.4. Must preparation

Chagalapoli fruit was extracted using a destoner machine (Bertuzzi, Brugherio, Milan, Italy) with a 0.2 sieve separating the seed. The experiment was divided into two batches: Filtered juice (all crushed fruit was filtered), and partially filtered juice (75% filtered crushed fruit and 25% unfiltered fruit). The filtration consisted in separating residues from the juice by centrifugation. Must was prepared according to the methods of Dias et al. (2007) [9], with minor modifications. Tartaric acid was added to adjust pH to 3.8 in order to inhibit bacterial growth, and sulfur dioxide was added in the form of potassium metabisulfite at concentrations of 100 mg/L.

2.5. Yeast strains

Commercial *Saccharomyces cerevisiae* yeast was used and the inoculum was done following the recommendations of the manufacturer (Red Star brand, provided by Maltas e Insumos Cerveceros S.A. de C.V.). 0.173 g of dried yeast was added to 173 mL of warm water with 1.73 g of sucrose at 35-38°C for 20 min, it was left to cool for 10 min. Initial cell density of yeast was $1.9 \times 10^6 \pm 0.2$ cells/mL.

2.6. Fermentation conditions

Once the must preparation was done according to 2.4 and the yeast inoculum was done according to 2.5, they were put in a 1L bioreactor (BioBundle, Applikon Biotechnology, Netherlands) at 27 °C. The fermentation process was evaluated every 24 h and considered complete when the specific gravity was stable. At the end of the fermentation, the beverage was transferred to a glass bottle and stored at 10 °C. After 24 h, the beverage was transferred to a new bottle and after 10 days the beverage was filtered and stored at 5 °C in 350 mL glass bottles filled to the top to avoid oxygen entrance [11].

2.7. Analysis of juice and fermented beverage

Titratable acidity (TAC), specific gravity, pH, cell counts (Neubauer chamber, Celeromics France) total soluble solids (TSS) were determined according to García and Xirau (2000) [12]. Sulfur dioxide, volatile acidity (VA) and alcoholic strength by volume were determined after fermentation and storage using methods of the International Organisation of Vine and Wine [13]. Colorimetric properties were measured in terms of CIELAB parameters using a colorimeter model CR400 (Konica Minolta sensing NJ USA) and expressed in terms of rectangular (L^* , a^* , b^*) color coordinates. L^* indicates lightness ranging from black (0) to white (100). a^* value ranges from red to green and b^* ranges from yellow to blue.

2.8. Determination of reducing sugar

The reducing sugar was obtained using the method of Miller [14]. Briefly, 0.5 mL of sample was added with 0.5 mL of 3,5-dinitrosalicylic acid (DNS) reagent, kept in a boiling water bath for 5 minutes, then the reaction was stopped by placing the test tubes in a cold water bath, 5 mL of distilled water was added and it was left to rest for 15 minutes. Absorbance was determined at 540 nm in an Absorbance microplate reader model ELx808 (Biotek VT USA). All the experiments were performed in triplicate. The reducing sugar content was calculated based on a standard glucose curve.

2.9. Determination of total sugar

The total sugar content was determined using the method of Dubois [15] and Chow & Landhauser [16]. 2 mL of sample was diluted in 2 mL of 5% phenol solution; it was boiled in a water bath for 5 min and then cooled with ice. Then 5 mL of H_2SO_4 was added and it was stirred, then it was left to rest during 30 min. Absorbance was determined at 490 nm in an Absorbance microplate reader. All experiments were performed in triplicate.

2.10. Total phenolic content

The total phenolic content was determined using the method of Folin-Ciocalteu [17]. 30 μ L of sample was diluted in 3 mL of water and 200 μ L of Folin-Ciocalteu reagent and 600 μ L of a sodium carbonate solution was added. Afterwards this sample was warmed up at 40 °C for 20 min, then it was cooled to room temperature and after 15 min the absorbance was measured at 760 nm. Total phenolic content was calculated as gallic acid equivalent based on a standard gallic acid curve. All the experiments were performed in triplicate.

2.11. HPLC-DAD analysis

Samples were filtered through a 0.45 μ m membrane filter (Millipore Corporation). Samples were analyzed on an Agilent 1100 HPLC (Agilent, Technologies CA USA) equipped with a diode detector, with wavelengths at 280 nm, 320 nm and 360 nm. A C18 column (Phenomenex) was used for the stationary phase. The mobile phase consisted of two solvents: (A) water/formic acid (99:1; v/v), and (B) acetonitrile (100%). The gradient employed was: isocratic 0% B for 8 min, 10% B for 2 min, 20% B for 13 min, 30% B for 7 min, 40% B for 15 min, 80% B for 5 min, 100% B for 5 min. Flow rate was set at 1 mL/min and a temperature of 25 °C. Phenolic identities were assigned based on their retention characteristics and UV-visible spectra. For the identification and quantification of total phenolics compounds external standard calibration curves of the reported compounds were used (Epicatechin, Kaempferol, Quercetin, Catechin, Chlorogenic).

2.12. Sensory analysis

The final beverages were evaluated by a panel of 32 adult men and women. 20 mL samples at 10 °C were given in transparent glasses marked with three-digit random numbers. Fermented beverages were evaluated by acceptability for appearance (color), aroma and flavor according to a

5-point hedonic scale where 0 corresponds to least appreciated (extremely dislike) and 5 corresponds to most appreciated (extremely like).

2.13 Statistical analysis

Data for all of the measurements were obtained in triplicate and expressed as mean \pm standard deviation. Statistical analyses during fermentation (pH, relative density, total soluble solids (TSS) and total titratable acidity (TAC)) were performed with a one-way analysis of variance and means comparison (Tukey). Fermented beverages (filtered and partially filtered beverages) were compared with the filtered and partially filtered juices by means of a t-test to determine the significant difference between samples. A significance level $p < 0.05$ was adopted. Different letters were used to label significantly different values. This statistical treatment was carried out using Minitab 17 Statistical Software.

3. Results and discussion

3.1. Physicochemical characteristics of *Ardissia compressa*

Results obtained in the proximal composition are shown in Table 1. The water content of the fruit was 80.5%, similar to values reported by Joaquin-Cruz, et al., [4] who reported 86.8% in *Ardissia compressa* from Veracruz. The Chagalapoli fruit used in this research had a pH level of 4.2 ± 0.1 . This result was higher than the values reported by Jacome-Hernandez [2] (pH 2.91) and Joaquin-Cruz, et al., [4] (pH 2.73). Likewise, total soluble solids were different from those reported by Jacome-Hernandez [2] (10.5%), while our results were $17.7 \pm 0.05\%$. This could be attributed to the fact that each fruit comes from different locations, weather conditions, harvest season, storage, etc. that affect the characteristics of each fruit. Low pH could contribute to a decrease of general sensory quality of the beverage to an unacceptable level [18].

Table 1. Proximal composition and physicochemical characteristics of chagalapoli (*Ardissia compressa*) fruit

Components ^a	Value
Water (%)	80.52 ± 0.8
Ether extract (%)	0.55 ± 0.05
Ash (%)	0.66 ± 0.01
Protein (%)	8.58 ± 0.4
Crude Fiber (%)	3.58 ± 0.005
Carbohydrates (%)	11.9
Water activity	0.97 ± 0.001
TSS ^b (%)	17.7 ± 0.05
pH	4.2 ± 0.1

Results are expressed as mean \pm SE (triplicate).^a Proximal components are expressed in fresh weight (FW) ^bTotal soluble solids ($^{\circ}$ Brix).

3.2. Analysis during fermentation

Table 2 shows the results obtained of pH, specific gravity, total soluble solids (TSS) and total titratable acidity (TAC). It was observed that pH values of the beverages did not have a significant difference during the fermentation process, in fermented filtered chagalapoli the change was from 3.84 ± 0.04 to 3.62 ± 0.09 and in fermented partially filtered it was from 3.87 ± 0.2 to 3.54 ± 0.1 . The TAC did not exhibit significant difference in either of the batches (8.9 g/L to 8.6 g/L for the filtered sample and 8.9 g/L to 9.2 g/L tartaric acid for the partially filtered sample); in fermented fruit the limit values are 5.5 g/L to 9 g/L (Norma Oficial Mexicana PROY-NOM-199-SCFI-2015) which indicates that TAC obtained in the filtered sample is within the permitted limits. Specific gravity showed significant difference between samples, which showed a tendency to decrease with time;

batches had the same value at the end of fermentation. During alcoholic fermentation of both batches of *Ardisia compressa*, initial value of yeast was $1.9 \times 10^6 \pm 0.2$ cells/mL. Number of microorganisms increases due to environmental conditions and sugar content decreases rapidly. Maximum value of yeasts was obtained on the second day with a density of $4.1 \times 10^7 \pm 0.1$ cells/mL; this is relevant because it suggests that at this point the yeast was in optimal conditions that favor the production of alcohol. Cell density decrease was observed after the third day and fermentation concluded without a significant difference on the fourth day with a value of $4.4 \times 10^6 \pm 0.2$ cells/mL.

Table 2. Physicochemical properties of chagalapoli (*Ardisia compressa*) fruit filtered and partially filtered beverage during fermentation

Days	Filtered fermented beverage					Partially filtered fermented beverage				
	0	1	2	3	4	0	1	2	3	4
pH	3.84±0.04 ^a	3.75±0.06 ^{ab}	3.65±0.07 ^{ab}	3.61±0.09 ^b	3.62±0.09 ^b	3.87±0.2 ^a	3.74±0.07 ^a	3.5±0.1 ^a	3.51±0.1 ^a	3.54±0.1 ^a
SP. Gr ¹	1.072±0.005 ^a	1.048±0.007 ^b	1.028±0.001 ^c	1.009±0.001 ^d	1.004±0 ^d	1.074±0 ^a	1.044±0.002 ^b	1.030±0.003 ^c	1.006±0.001 ^d	1.004±0.001 ^d
TSS ² (%)	17.5±0.1 ^a	11.16±0.2 ^b	7.09±0.3 ^c	2.34±0.2 ^d	1.04±0 ^d	17.9±0 ^a	11.03±0.4 ^b	7.74±0.8 ^c	1.7±0.2 ^d	1.2±0.2 ^d
TAC ³ (g/L tartaric acid)	8.9±0.1 ^a	8.6±0.2 ^a	8.6±0.08 ^a	8.6±0.1 ^a	8.6±0.08 ^a	8.9±0.1 ^a	9.1±0.2 ^a	9.3±0.05 ^a	9.2±0.1 ^a	9.2±0.2 ^a

Results are expressed as mean ± SE (triplicate). Values with same letter are not statistically different (p<0.05). ¹Specific Gravity. ²Total soluble solids. ³Titrateable acidity.

3.3. Color parameters

Table 3 shows CIELAB color parameters obtained. L* value was 27.5 to 30.34 in the filtered sample and 30.75 to 31.09 in the partially filtered sample. It was observed that values increased during fermentation, which indicates the samples are darker. a* obtained in filtered samples was 8.87 to 8.42 and in partially filtered samples was 7.15 to 6.92. It was observed that values decreased during fermentation which indicates a red tone. b* values were 14.50 to 17.50 in filtered samples and 16.59 to 16.28 in partially filtered samples, which indicates that b* values increased in the filtered samples and decreased in partially filtered samples, obtaining a yellow color. Angle of hue and chromaticity defined the sample within the red color, presenting significant differences in all samples. Heredia-Vasquez [3] reported different values in L*, a* and b*, obtained low luminosity and more tendency towards red color. Differences can be justified by the type of fruit used as well as by climatic condition, region, harvest time, etc.

Table 3. Chagalapoli (*Ardisia compressa*) juice and fermented beverage color analysis

	L*	a*	b*	Chroma	Angle of hue (h°)
Filtered juice	27.50 ± 0.005 ^d	8.87 ± 0.02 ^a	5.63 ± 0.02 ^c	14.50 ± 0.01 ^d	0.56 ± 0.002 ^d
Filtered fermented beverage	30.34 ± 0.01 ^c	8.42 ± 0.02 ^b	9.07 ± 0.02 ^b	17.50 ± 0.02 ^a	0.79 ± 0.002 ^c
Partially filtered juice	30.75 ± 0.04 ^b	7.15 ± 0.03 ^c	9.44 ± 0.02 ^a	16.59 ± 0.03 ^b	0.87 ± 0.002 ^b
Partially filtered fermented beverage	31.09 ± 0.09 ^a	6.92 ± 0.06 ^d	9.36 ± 0.09 ^a	16.28 ± 0.03 ^c	0.87 ± 0.006 ^a

Results are expressed as mean ± SE (n=5). Values with same letter are not statistically different (p<0.05).

3.4. Analysis of fermented beverage

Fermented filtered and partially filtered beverage of chagalapoli (*Ardisia compressa*) fruit obtained a similar result in alcoholic degree method, 6% and 6.4% respectively. Official Mexican Standard NOM-199-SCFI-2017 specifies the limit values for fermented fruit beverages from 6% to 12%; the values obtained from chagalapoli (*Ardisia compressa*) fermented beverage are within the established range. Chowdhury & Ray [19] reported a 6% of alcohol content in a fermented jamun berry (*Syzygium cumini* L.) beverage.

In relation to volatile acidity 0.32g/L was the value obtained for filtered fermented beverage and 0.36g/L for partially filtered beverage; Mena et al., [10] determined the volatile acidity of fermented beverages of 3 pomegranate varieties, for which they obtained values of 0.33, 0.36 and 0.26 g/L. These results suggest that sugar consumption produces different acids (citric, malic, acetic and tartaric acids).

The Official Mexican Standard NOM-199-SCFI-2017 allows a maximum 50 mg/L of free sulfur dioxide and 350 mg/L for total sulfur dioxide. To determine the sulfur dioxide in fermented beverages the free sulfur dioxide was calculated first; 17.06±0.73 mg/L was determined for free sulfur dioxide and 18.77±4.8 mg/L for total sulfur dioxide. For the total sulfur dioxide there was no significant difference between the samples, 57.6±3.3 and 58.88±3.8 were obtained, under the standard the samples are within the permitted limits. Oliveira, et al. (2011) [11] reported similar results in cagaita wines (*Eugenia dysenterica* DC), they obtained from 22mg/mL to 12 mg/L of total sulfur dioxide. Kelebek, et al. (2009) [21] reported 8.2 ± 0.08mg/L of free sulfur dioxide and 73.3 ± 0.12mg/mL of total sulfur dioxide in orange wine.

3.5. Reducing sugar

Results are shown in Table 4. The values for filtered juice and partially filtered juice were 132.08 ± 1 and 96.63 ± 2 mg/mL, respectively. For filtered and partially filtered fermented beverage, the values decreased to 3.42 ± 0.3 mg/mL and 3.58 ± 0.3 mg/mL, respectively. Similar results were reported by Oliveira et al., [11] where values of 1.2 mg/mL and 2.4 mg/mL of reducing sugars were obtained in Cagaita fruit wine.

Table 4. Reducing sugar, total sugar and total polyphenol contents in filtered and partially filtered Chagalapoli (*Ardisia compressa*) fruit juice and filtered and partially filtered fermented beverages

	Reducing sugar mg (glucose)/mL		Total sugar mg (glucose)/mL		Total polyphenols			
	Juice	Fermented	Juice	Fermented	Folin-Ciocalteu mg (GAE)/mL		HPLC-DAD mg (EPI)/mL	
					Juice	Fermented	Juice	Fermented
Filtered Chagalapoli	132.08 ± 1^a	3.42 ± 0.3^b	137.82 ± 0.9^a	3.74 ± 0.1^b	1.8 ± 0.03^a	1.11 ± 0.02^b	1.07 ± 0.04	1.47 ± 0.09
Partially filtered Chagalapoli	96.63 ± 2^a	3.58 ± 0.3^b	83.04 ± 0.6^a	3.27 ± 0.03^b	1.75 ± 0.1^a	1.59 ± 0.19^b	1.43 ± 0.1	2.86 ± 0.01

Results are expressed as mean \pm SE (triplicate). Values with same letter are not statistically different ($p < 0.05$).

3.6. Total sugar contents

Total sugar contents results are shown in Table 4. Filtered juice had more total sugars than partially filtered juice with 137.83 ± 0.9 mg/mL and 83.04 ± 0.6 mg/mL, respectively. These results may be compared to the report by Kosseva, et al., [20] where blueberry presented values of 155.2 g/kg to 164.7 g/kg, while in red grapes it was around 200 g/kg or more at maturity. Comparing these results with reported values of blueberry, chagalapoli (*Ardisia compressa*) sugars are lower. There was a reduction during the fermentation, and at the end of samples 3.74 ± 0.1 mg/mL and 3.27 ± 0.03 mg/mL of total sugars were obtained in final products of fermented beverages. Kelebek et al., [21] reported 120.19 mg/mL in orange juice and 48 mg/mL in fermented orange juice. These results could differ since the orange is considered a citrus fruit and because the sugars present in each fruit are different. Since the methods for reducing and total sugars are different, results are not considered complementary.

The results obtained for total sugar content and reducing sugar content in the partially filtered samples showed the limitation of sugar desimentation, this may explain the variation in the results obtained.

3.7. Analysis of phenolic compounds

3.7.1. Folin-Ciocalteu method

The total polyphenol content by the Folin-Ciocalteu method in chagalapoli (*Ardisia compressa*) juices was 1.8 ± 0.03 mgGAE/mL for filtered juice and 1.75 ± 0.1 mgGAE/mL for partially filtered juice. Similar values were previously reported by Heredia-Vásquez [3], who obtained values of 1.74 mgGAE/mL in *Ardisia compressa kunth*; Joaquín-Cruz, et al., [4] obtained 1.051 ± 43.5 mgGAE/g in chagalapoli fruit. In filtered and partially filtered fermented beverage the values decreased to 1.11 ± 0.02 mgGAE/g and 1.59 ± 0.19 mgGAE/g, respectively. Martins de Sá, et al., [7] reported 1.105 ± 57 mgGAE/mL in a fermented jaboticaba (*Myrciaria jaboticaba*) beverage. Johnson, et al., [22] reported a total polyphenol content of 0.3754 mgGAE/mL to 0.6571 mgGAE/mL in blackberry (*Vaccinium sps*) fermented beverage.

3.7.2. HPLC-DAD analysis

Total polyphenols are shown in Table 4. An increase in the total polyphenols was observed in both samples of Chagalapoli (*Ardisia compressa*) fermented beverages. At the beginning of the

fermentation, the value was 1.07 ± 0.04 epicatechin mg/mL, with increases towards the end of fermentation with a final value of 1.47 ± 0.09 epicatechin mg/mL. The partially filtered chagalapoli sample initially presented 1.43 ± 0.1 epicatechin mg/mL and at the end of fermentation had increased to 2.86 ± 0.01 epicatechin mg/mL. Recent studies demonstrated the same behavior in fermentations in different grape varieties. In the fermentation process, several enzymes are excreted during the metabolism of the yeast, and these enzymes could act on the conjugated phenolic compounds to release free phenolic compounds, and thus change their composition [23]. Figure 1 show the compounds identified in *Ardisia compressa* samples at 280 nm wavelength. Peaks 1 and 2 correspond to catechin and epicatechin; catechin was found at 7.12 min up to 7.45 min, epicatechin was found at 8.2 min up to 8.54 min in the first chromatograms, while in the chromatogram of the partially filtered fermented beverage it was observed at 11.24 min. Figure 2 shows the juices and fermented beverage chromatograms at 320 nm wavelength. In partially filtered fermented beverage, a major peak is shown, which was identified as chlorogenic at 12.48 min, 14.49 min and 13.73 min for filtered beverages and at 14.52 min in partially filtered juice. Figure 3 shows compounds identified in *Ardisia compressa* beverages at 360 nm wavelength. Different derivatives of quercetin were identified between 24.65 min and 24.81 min for filtered beverages (juice and fermented beverages) and for partially filtered beverages they were identified at 23.23 min up to 23.92 min. Kaempferol was identified at 26.6 min in the filtered samples and in partially filtered samples it was found at 25.72 min and 24.48 min. Joaquín-Cruz, et al. [4] identified 6 derivatives of quercetin with retention times of 18.61 min, 19.49 min, 19.92 min, 20.21 min, 22.98 min and 23.72 min. For the identification of kaempferol, retention times of 16.94 min and 17.24 min were obtained.




Figure 1 Chagalapoli (*Ardisia compressa*) HPLC Chromatograms at 280 nm: (A) filtered juice, (B) partially filtered juice, (C) filtered fermented beverage, (D) partially filtered fermented beverage




Figure 2 Chagalapoli (*Ardisia compressa*) HPLC chromatograms at 320 nm: (A) filtered juice, (B) partially filtered juice, (C) filtered fermented beverage, (D) partially filtered fermented beverage




Figure 3 Chagalapoli (*Ardisia compressa*) HPLC chromatograms at 360 nm: (A) filtered juice, (B) partially filtered juice, (C) filtered fermented beverage, (D) partially filtered fermented beverage

3.8. Sensory evaluation

Fermented beverages from both batches were subjected to a sensory analysis to evaluate the degree of satisfaction of the final product, carried out with a panel of 32 untrained tasters older than 18 years. Table 5 shows the results of acceptance of each sample, it was observed that the partially filtered beverage obtained a higher acceptance value in color and flavor parameters with a

difference of 0.87 and 1.41, respectively. Aroma did not present a significant difference between samples; this means that there was not a preference for any of the two fermented beverages.

Table 5. Sensory analysis of chagalapoli (*Ardisia compressa*) fruit filtered and partially filtered fermented beverages

	Filtered fermented beverage	Partially filtered fermented beverage
Color	3.47 ± 0.8 ^b	4.34 ± 0.6 ^a
Aroma	3.69 ± 0.9 ^a	3.72 ± 1.2 ^a
Flavour	2.9 ± 0.1 ^b	4.31 ± 0.8 ^a

Results are expressed as mean ± SE (n=32). Values with same letter are not statistically different (p<0.05).

4. Conclusions

The results of the present study demonstrated that chagalapoli fruit has the potential to be used to produce fermented beverages. It was also revealed that chagalapoli fermented beverages represent a rich source of phenolic compounds. This beverage can be considered as a dry spirit since the content of reducing sugars was lower than 4 mg/mL. Based on the characteristics of the produced chagalapoli fermented beverages, it was concluded that there should be more extensive research with chagalapoli fruit due to its potential for development into a marketable beverage.

Funding: This research received no external funding.

Conflicts of Interest: The authors declare no conflict of interest.

Reference

1. Kobayashi, H. and De Mejía, E. The genus *Ardisia*: a novel source of health-promoting compounds and phytopharmaceuticals. *J. Ethnopharmacol* **2005**, 96, 347-354.
2. Jácome-Hernández, S.P. Propiedades antioxidantes y estabilidad de una bebida elaborada a partir del fruto de capulin (*Ardisia compressa*). Universidad Veracruzana. (Master's Thesis) Xalapa, Veracruz. **2012**.
3. Heredia-Vásquez, J. Propiedades fisicoquímicas y antioxidantes de polvos nanoestructurados de *Ardisia compressa* kunth. (Master's Thesis) Universidad Veracruzana. Veracruz, Mexico. **2013**.
4. Joaquín-Cruz, E., Dueñas, M., García-Cruz, L., Salinas-Moreno, Y., Santos-Buelga, C., García-Salinas, C. Anthocyanin and phenolic characterization, chemical composition and antioxidant activity of Chagalapoli (*Ardisia compressa* K.) fruit: A tropical source of natural pigments. *Food Res Int* **2015**, 70, 151-157.
5. Colorado-Vargas, R. Caracterización de Chagalapoli (*Ardisia compressa* Subsp. Myrsinaceae), para su posible uso en la elaboración de un alimento funcional (Master's Thesis) Universidad Autónoma de Chihuahua. Chihuahua, México. **2013**.
6. Chandra, S. and De Mejia-Gonzalez, E. Polyphenolic Compounds, Antioxidant Capacity, and Quinone Reductase Activity of an Aqueous Extract of *Ardisia compressa* in Comparison to Mate (*Ilex paraguariensis*) and Green (camellia sinensis) Teas. *J Agric Food Chem* **2004**, 52, 3583-3589.
7. Martins de Sá, L.Z.C., Castro P.F.S., Lino, F.M.A., Bernardes M.J.C., Viegas, J.C.J., Dinis, T. C.P.D., Santana, M. J., Romao, W., Vaz, B. G., Liao, L. M., Ghedini, P. C., Rocha, M. L., Gil, E. S. Antioxidant potential and vasodilatory activity of fermented beverages of jaboticaba berry (*Myrciaria jaboticaba*). *J Funct. Foods* **2014**, 8, 169-179.
8. Li, H., and Förstermann, U. Red wine and cardiovascular health. *Circ. Res* **2012**, 111, 959–961.

9. Dias, D. R., Schwan, R. F., Freire, E. S., & Serôdio, R. S. Elaboration of a fruit wine from cocoa (*Theobroma cacao* L.). *International Journal of Food Science and Technology* **2007**, *42*, 319-329.
10. Mena, P., Gironès-Vilaplana, A., Marti, N., Garcia-Viguera, C., Pomegranate varietal wines: Phytochemical composition and quality parameters. *Food Chemistry* **2012**, *133*, 108-115.
11. Oliveira, M.E.S., Pantoja, L., Duarte, W.F., Collela, C.F., Valarelli, L.T., Schwan, R.F., and Dias, D.R. Fruit wine produce from cagaita (*Eugenia dysenterica* DC) by both free and immobilised yeast cell fermentation. *Food Res Int* **2011**, *44*, 2391-2400.
12. García, J. Cazorla, Xirau, M. Vayreda. *Técnicas usuales de análisis en enología* **2000**. Barcelona: Panreac Química, S. A.
13. Office Internationale de la Vigne et du Vin (OIV). *Recueil des Méthodes Internationales D'analyse des Vins* **1990**. Paris, France: OIV.
14. Miller, G. Use of dinitrosalicylic acid reagent for determination of reducing sugar. *Anal. Chem.* **1959**, *31*, 426-428.
15. Dubois M., Gilles K.A, Hamilton J.K, Rebers P.A, & Smith F. Colorimetric method for determination of sugars and related substances, *Anal. Chem.* **1956**, *28*, 350-356.
16. Chow P. S. and Landhausser S.M. A method for routine measurements of total sugar and starch content in woody plant tissues. *Tree Physiol* **2004**, *24*, 1129-1136.
17. Singleton, V.L., & Rossi, J.A. Colorimetry of total phenolics with phosphomolybdic-phosphotungstic acid reagents. *Am J Enol and Vitic* **1965**, *16*, 144-158.
18. Neffe-Skocinska, K., Sionek, B., Scibisz, I. & Kolozyn-Krajewska, D. Acid contents and the effect of fermentation condition of Kombucha tea beverages on physicochemical, microbiological and sensory properties. *J food* **2017**, *15*, 601-607.
19. Chowdhury, P. and Ray, R. C. Fermentation of jamun (*Syzygium cumini* L.) fruits to form red wine. *ASEAN Food Journal* **2007**, *14*, 15-23.
20. Kosseva, M., Joshi, V.K., Panesar, P.S. Science and Technology of Fruit Wine Production. Edit. Academic press. London United Kingdom. **2016**.
21. Kelebek, H., Selli, S., Canbas, A., Cabaroglu, T. HPLC determination of organic acids, sugars, phenolic compositions and antioxidant capacity of orange juice and orange wine made from a Turkish cv. Kozan. *Microchem J* **2009**, *91*, 187-192.
22. Johnson M. H., Lucius A., Meyer T., De Mejia, E.G. Cultivar evaluation and effect of fermentation on antioxidant capacity and in vitro inhibition of α -amylase and α -glucosidase by highbush blueberry (*Vaccinium corombosum*). *J Agric Food Chem.* **2011**, *59*, 8923-8930.
23. Zhang, Y., Chang S.K.C., Stringer S.J., Zhang, Y. Characterization of titratable acids, phenolic compounds, and antioxidant activities of wine made from eight Mississippi-grown muscadine varieties during fermentation. *LWT - Food Sc and Technol* **2017**, *86*, 302-311.