

Article

SALMONELLA SURVEILLANCE SYSTEMS in SWINE and HUMANS in SPAIN: a REVIEW

Marta Martínez-Avilés ^{1,*}, Macarena Garrido-Esteba ¹, Julio Álvarez ^{2,3} and Ana de la Torre ¹

¹ Animal Health Research Center (INIA-CISA), Ctra Algete a El Casar s/n, 28130, Valdeolmos, Madrid, Spain

² VISAVET Animal Health Surveillance Center, Complutense University of Madrid, Madrid, 28040, Spain

³ Animal Health Department, Veterinary School, Complutense University of Madrid, Madrid, 28040, Spain

* Correspondence: martinez.marta@inia.es; Tel.: +34-91-620-2300

Abstract: Non-typhoid salmonellosis is a common and problematic foodborne zoonotic disease in which pork and pork products can be an important potential source of infection. In order to prevent this disease important efforts to monitor the situation in the main source, livestock, are conducted in most developed countries. In the European Union EFSA and ECDC compile information at the member state level, even though important differences in production systems and surveillance systems exist. Here, *Salmonella* surveillance systems in one of the main sources of foodborne salmonellosis, swine, and humans in Spain were reviewed to identify potential gaps and discuss potential ways of integration under a One Health approach. Despite the extensive information generated through the surveillance activities source attribution can be only routinely performed through ad-hoc outbreak investigations, and national reports on human outbreaks do not provide sufficiently detailed information to gain a better understanding of the epidemiology of the pathogen. Human and animal monitoring of *Salmonella* would benefit from a better exchange of information and collaboration. Analysis of spatio-temporal trends in livestock and humans could help to identify likely sources of infection and to target surveillance efforts in areas with higher prevalence or where specific strains are found.

Keywords: Zoonoses; food-borne; disease control; public health; domestic livestock; pig; One health

1. Introduction

Salmonella is a ubiquitous genus of bacteria commonly found in the intestines of healthy birds, reptiles and mammals that can cause one of the most common food-borne illness in humans [1]. According to the European Food Safety Agency (EFSA), the main species in the genus, *Salmonella enterica*, is one of the top agents involved in foodborne outbreaks in Europe including Spain, even though disease burden is likely severely underestimated because infection can be asymptomatic or not sufficiently severe to prompt testing [2]. Additionally, non-severe cases can be treated without further investigation of the subtype.

Poultry *Salmonella* is under an official control programme in the European Union (EU) since 2004. Consequently, there was a significant reduction of *Salmonella* in humans and poultry during the period 2008-2016 [3], particularly due to *S. Enteritidis*. In contrast, there was an apparent increase of notifications of *S. Typhimurium* cases, which are less likely associated with the consumption of eggs and egg products and more predominantly found in pork and pork products [4].

Spain is one of the main swine producing countries, currently ranking first in number of swine in the EU (with 28.3 million animals in 2015) [5]. The production of pork in 2015 reached 3.8 million

tonnes, with more than 45 million animals being slaughtered. Worldwide, Spain is the fourth largest pork producer after China, United States and Germany. Mainly an exporting country, Spain has also become the EU's third largest exporter of swine after Germany and Denmark. The swine industry accounts for 14% of the final agricultural production in Spain, and it is the most important livestock species in economic terms, representing 37% of the final livestock production.

Pork is the main source of human salmonellosis after poultry in the EU [6], given that it is the third most frequently contaminated meat after fresh chicken and turkey [7], and it is widely consumed. Because of this, monitoring and surveillance activities have been implemented along the food chain in order to assess the risk posed by pork and pork products as a source of *Salmonella* for the general public and to prevent outbreaks. In addition, the threat posed by the increasing occurrence of infections caused by antimicrobial resistant *Salmonella* strains in humans is another reason to perform "One health" surveillance efforts to control *Salmonella* at its source [8].

Swine can acquire *Salmonella* infection from a contaminated environment or feed, or through direct contact with infected animals. Infected pigs can remain carriers of *Salmonella* and shed the bacteria via the faeces intermittently for many months [9]. There is a risk of cross-contamination of carcasses with faeces of infected or carrier animals at slaughter. Prevention of *Salmonella* infection in pigs at the farm is performed through the regular monitoring of pig feed and implementation of basic biosecurity measures and in certain cases vaccination. However, environmental persistence, high turnover of young stock and incoming replacement stock pose significant barriers to eliminate *Salmonella* at the farm. The *Salmonella* status of a pig can be monitored through serological tests performed on meat-juice or serum samples [10] or, more frequently, through the bacteriological analysis of faeces collected at either the farm or the slaughterhouse, where mesenteric lymph nodes can be also collected [7]. There are also microbiological tests conducted on meat and carcasses to verify hygiene practices through the food chain, but in the case of positive results it is often not possible to establish whether the tissue contamination originated from an infected pig farm or occurred as a result of a contaminated environment [11] or insufficient hygiene practices during meat processing [12].

Spain is divided into 19 regions. Health competencies are transferred to the regions, who report to the national authorities. Animal health surveillance is the responsibility of the Regional Departments and the Ministry of Agriculture. However, meat inspection falls under the responsibility of the Regional Departments and Ministry of Health, together with human disease surveillance. Since 2004, EFSA analyses comparable data on zoonotic foodborne diseases from all EU Member States and harmonised prevalence targets are set. Member States inform EFSA and the European Centre for Disease Control (ECDC), in agreement with the Directive 2003/99/EC, of the results of the monitoring systems in place (compulsory or voluntary monitoring programmes, surveys, other procedures of sampling and lab reports), which is published in an Annual EU Summary report.

The EU-funded research project "One health"- European Joint Programme number 773830 "NOVA" [13] seeks to develop new surveillance tools and methods and aims to harmonise and optimise the use of existing surveillance data on zoonotic foodborne diseases. Under this context, we review and describe the *Salmonella* surveillance systems in Spain "from farm to fork", that is, from its animal source (here swine) up to the identification of *Salmonella* of animal origin in human outbreaks. Our aim is to identify potential gaps and to assess the feasibility of a more integrated approach in a One Health framework. We argue the differences between systems and potential ways to integrate surveillance across sectors if needed for the benefit of all parties.

2. Search and review strategy

We performed a non-systematic review on the ongoing monitoring programs for *Salmonella* in the animal reservoir (swine) and in humans across health institutions in Spain. Since the scope of the

paper is limited to Spain, a list of the institutions' and systems' acronyms used (mostly of Spanish origin) is provided in Table 1.

Table 1. List of acronyms of institutions and systems used in this paper.

| Acronym | Full name |
|---------|---|
| AECOSAN | Spanish Agency of Consumption, Food Security and Nutrition |
| ISCIII | Carlos III Institute of Health, which hosts the National Centre of Epidemiology and the National Centre of Microbiology. In this paper, ISCIII refers to the former, unless stated otherwise. |
| MAPA | Ministry of Agriculture |
| MSSSI | Ministry of Health |
| RASFF | Rapid Alert System for Food and Feed (EU) |
| RENAVE | National Network of Epidemiological Surveillance |
| SCIRI | Coordinated System for the Exchange of Information (National) |
| SIM | Microbiological Information System |
| VISAVET | Veterinary Health Surveillance Centre |

Surveillance of *Salmonella* in swine is coordinated by MAPA, and laboratory testing is conducted in the national and regional official laboratories and MAPA-approved animal health laboratories, including VISAVET. Official sources of information on monitoring in swine are listed in the MAPA website dedicated to antimicrobial resistance and zoonosis surveillance (Table 2), which includes legislation and links to MAPA reports (2012-2016) and EFSA's Spain reports (2004-2016) on zoonosis and antimicrobial resistance annual surveillance results. EFSA's annual country reports include information on the reporting and monitoring system on zoonoses sent by a Reporting Officer from that country. MAPA (Animal Health Section) is the Reporting Officer from Spain, and gathers information from other National Reporters, namely AECOSAN, ISCIII, other sections within MAPA (Farm and Traceability Registries; Animal Feed), VISAVET, and the Regional Animal Health Departments. Here, we selected those reports that provided detailed information on the monitoring design and results of *Salmonella* in pigs to assess the potential uses of that information. In consequence, only EFSA annual country reports from 2007 to 2012 were selected. The information on human surveillance contained in the EFSA annual country reports is very brief. Therefore, we tracked the websites of other national institutions mentioned in EFSA's Spain annual reports (ISCIII, AECOSAN) to retrieve further information on *Salmonella* surveillance in humans and in meat and meat products, respectively. Both ISCIII and AECOSAN belong to the Ministry of human health (MSSSI). Also, we completed our search by reading the pertinent legislation on *Salmonella* surveillance and by searching in PubMed and in the scientific bibliography archive of the Complutense University of Madrid scientific articles and doctoral thesis on *Salmonella* in pigs, particularly in Spain (combination of search terms "*Salmonella*", "pigs", "Spain", "epidemiology"). This way, general information on *Salmonella* epidemiology was also retrieved.

The Spanish National Epidemiological Surveillance Network (RENAVE) is responsible of the surveillance and control of infectious diseases in Spain. This network is composed by the 19 Spanish autonomous regions and it is coordinated by the MSSSI with the scientific and technical support of ISCIII. The Notifiable Diseases regional data is merged at ISCIII where the National Database is maintained. At ISCIII, the information is compiled, analyzed and disseminated. One of the authors of this manuscript (MGE) worked for more than ten years at ISCIII and participated in the definition of the surveillance protocols for Notifiable Diseases in Spain. Protocols for each Notifiable Disease on use by the RENAVE are publicly available and were approved by the Inter-territorial Council of the National Health System (Table 2). Each protocol includes a general description of the epidemiology of the disease, standard definitions of cases classification, outbreak, and notification system and actions after a case or an outbreak have been confirmed. ISCIII holds three different databases: Notifiable Diseases (incidence on human salmonellosis that is part of this list only since 2015); Food-

Borne Outbreak Investigation Database (human salmonellosis outbreaks information since 2006); and an additional database created with the national microbiological laboratories data: SIM (identification and distribution of *Salmonella* serovars that are circulating in Spain reported voluntarily by participant laboratories since 1995). The results of the three databases are published in two annual reports (SIM reports; and RENAVE reports, that combine Notifiable Diseases + Outbreak Investigation) available at the ISCIII website.

Tracking the legislation and the scientific literature [14, 15], we reached the information gathered by the hospitals and primary health centres, which are the first locations where a suspect case is recorded. The information gathered by primary health centres and hospitals is available through the Statistics Portal of the MSSSI. Regional authorities can access these data and recapture any pending cases to notify them through RENAVE.

AECOSAN compiles and publishes the analysis results of *Salmonella* alerts in food through SCIRI, and then disseminates the results with an annual report available online. For the purpose of this manuscript, we downloaded the reports corresponding to 2014 and 2015, because these were the years in which *Salmonella* cases started to be reported by the Regional authorities to the RENAVE.

Based on the authors' experience on epidemiological surveillance, we searched, across the existing monitoring systems in Spain on swine-related *Salmonella*, for information on sampling strategy, frequency of sampling and testing, information included in the dissemination and, most particularly, whether there was evidence of communication or actions relative to the results obtained with other stakeholders involved in *Salmonella* monitoring and results of any potential joint epidemiological analysis. For the purpose of the "One health-EJP" project we also searched for evidence on spatial dissemination of results.

3. Findings

We first present the sources in which the information on *Salmonella* surveillance of swine origin in Spain can be found, followed by an analysis of the information contained in them at the animal, meat and human source.

3.1. Sources of information

Table 2 shows the websites visited for the non-systematic review of surveillance sources on *Salmonella* of swine origin of interest for Spain.

Table 2. Institutional websites with information on swine-related *Salmonella* relevant for Spain

| Institutions | Websites URL (last visited: 22.11.2018) | Information retrieved |
|--------------|--|--|
| EFSA | https://efsa.onlinelibrary.wiley.com https://www.efsa.europa.eu/en/topics/topic/salmonella | EU Summary reports on trends and sources of zoonoses, zoonotic agents and food-borne outbreaks (EFSA&ECDC) (2007-2012) |
| EFSA | https://www.efsa.europa.eu/en/efsajournal/pub/1377 https://www.efsa.europa.eu/en/efsajournal/pub/2329 | Analysis of the baseline survey on the prevalence of <i>Salmonella</i> holdings with breeding pigs in the EU, 2008- Part A: prevalence estimates- Part B: factors associated with pen positivity |
| EFSA | https://www.efsa.europa.eu/en/efsajournal/pub/rn-135 https://www.efsa.europa.eu/en/efsajournal/pub/rn-206 | Analysis of the baseline survey on the prevalence of <i>Salmonella</i> in slaughter pigs in the EU, 2006-2007- Part A: prevalence estimates –Part B: factors associated with lymphnode positivity, surface contamination of carcasses and serovar distribution |

| | | |
|---------|---|---|
| MAPA | https://www.mapa.gob.es/es/ganaderia/temas/sanidad-animal-higiene-ganadera/sanidad-animal/zoonosis-resistencias-antimicrobianas/zoonosis.aspx | Spain Reports by EFSA on zoonoses (2007-2012) |
| MSSSI | http://pestadistico.inteligenciadegestion.msssi.es/publicoSNS/comun/DefaultPublico.aspx | Hospital and Primary Health Centres Databases |
| ISCIII | http://www.isciii.es/isciii/es/contenidos/fd-servicios-cientifico-tecnicos/fd-vigilancias-alertas/fd-sistema-informacion-microbiologica/informes-generales.shtml | Annual SIM Reports (2014-15) |
| ISCIII | https://publicaciones.isciii.es/unit.jsp?unitId=cne | Annual RENAVE reports (2014-15) |
| AECOSAN | http://www.aecosan.msssi.gob.es/AECOSAN/web/seguridad_alimentaria/subseccion/SCIRI.htm | Annual SCIRI reports (2014-15) |

The legislation consulted is summarized in Table 3: a) European (OJ= Official Journal) and b) Spanish (BOE= State Official Bulletin, acronym in Spanish).

Table 3. Relevant legislation for swine-related *Salmonella* surveillance:

a) In the EU

| EU Legislation | Subject | Reference |
|---------------------------------|--|-------------------------|
| Directive 2003/99/EC | Surveillance of <i>Salmonella</i> and other zoonotic agents | OJ, L325/31, 12.12.2003 |
| Regulation 2160/2003 | Control of <i>Salmonella</i> and other food-borne diseases | OJ, L325/1, 12.12.2003 |
| Regulation 2073/2005 | Microbiological criteria of food products | OJ, L338/1, 22.12.2005 |
| Regulation 1441/2007 | Microbiological criteria of food products (modification) | OJ, L322/12, 07.12.2007 |
| Commission Decision 2006/668/EC | Design of <i>Salmonella</i> baseline survey in slaughter pigs | OJ, L275/51, 06.10.2006 |
| Commission Decision 2008/55/EC | Design of <i>Salmonella</i> prevalence survey in breeding pigs | OJ, L14/10, 17.01.2008 |
| Regulation 16/2011 | Rapid alert system for food and feed (RASFF) | OJ, L6/7, 11.01.2011 |
| Regulation 217/2014 | Analysis of microbiological samples in carcasses | OJ, L69/93, 08.03.2014 |
| Regulation 218/2014 | Checks on the operator's microbiological tests to control <i>Salmonella</i> through the food chain | OJ, L69/95, 08.03.2014 |

b) In Spain

| Spanish Legislation | Subject | Reference |
|---------------------|---------|-----------|
|---------------------|---------|-----------|

| | | |
|------------------------|---|--------------------------------|
| Royal Decree 2210/1995 | National network of epidemiological surveillance | BOE 21, 2153-58, 24.01.1996 |
| Royal Decree 1943/2004 | Transposition of Directive 2003/99/EC | BOE 237, 32772-77, 01.10.2004 |
| Order SCO/3270/2006 | National network of epidemiological surveillance for food-borne salmonellosis | BOE 255, 37238-9, 25.10.2006 |
| Law 17/2011 | Food security and national coordinated system for the rapid exchange of information (SCIRI) | BOE 160, 71283-319, 06.07.2011 |
| Royal Decree 69/2015 | Registry of the activity of primary health centres | BOE 35, 10789-809, 10.02.2015 |
| Order SSI/445/2015 | Salmonellosis as a human notifiable disease | BOE 65, 24012-15, 17.03.2015 |


3.2. Surveillance at the pig source

Routine surveillance of *Salmonella* in pigs in Spain is carried out at slaughter through the annual sampling of faeces or lymph nodes of carcasses, in agreement with EC Regulation 2160/2003. The overall aim of the EC Regulation 2160/2003 is to reduce the incidence of *Salmonella* across the food chain through harmonised sampling schemes to obtain comparable prevalence estimates.

The slaughterhouses selected for sampling every year process at least 50-60% of the annual slaughter pigs, and slaughterhouses from at least 50% of regions in Spain are included (Figure 1). Sampling is then stratified by slaughterhouse based on its annual throughput, and one or more samples (typically faecal samples) are collected from a variable number of farms (ranging between 160 and 400 farms sampled annually in 2002-2015) and processed individually or as pools.

In addition, in the last decades larger surveys have been conducted at the request of the EU. A large survey was conducted in slaughter pigs (Commission Decision 2006/668/EC) to establish the baseline prevalence of infection in the different member states of the EU. For this purpose 2,619 lymph node samples from pigs originating from different farms were collected in Spain from slaughterhouses accounting for > 80% of the of all slaughtered fattening pigs.

Both data sources (annual monitoring programme, EFSA baseline survey) revealed a prevalence of infection around 30% in slaughter pigs, though a wider range is observed when looking at the results from the annual programme (generated through a smaller sample size) (Figure 1).



Number of slaughterhouses (SH) sampled for *Salmonella* in pigs per province

Figure 1. Number and distribution of slaughterhouses sampled and results of the *Salmonella* annual monitoring programme in slaughter pigs in Spain. In black, number of total holdings and slaughterhouses (SH) sampled (time period of sampling). In red, positive (+) results to *Salmonella*. (Self-creation from the information retrieved from the reports in MAPA website).

The serotypes with a higher average proportion of isolates from 2008 to 2013 in slaughter pigs were S.1,4,[5],12:i:- (9.08%), followed by *S. Rissen* (7.99%), *S. Typhimurium* (7.36%), *S. 4,5:i:-* (6.06%) and *S. Derby* (5.09%). However, when disaggregating the slaughterhouse results by year of sampling (Figure 2), *S. Typhimurium* was the only serotype isolated every year, reaching the highest proportion of isolates in 2009 (11.66%). *S. Rissen* and *S. Derby* are the other two serotypes most frequently isolated in slaughterhouses, only missing in 2010. The highest proportion of *S. Rissen* (9.89%) also happened in 2009. S.1,4,[5],12:i:- was only isolated in 2012 and 2013 (8.59% in 2012 and 9.57% in 2013), and S.4,5:i:- only in 2011. Curiously, *S. Anatum* was isolated from slaughterhouses only in 2008 and in 2011, and with a low proportion (<2%).




Figure 2. Distribution by serotype of the proportion of positive holdings to *Salmonella* (of total holdings tested per year) in slaughter pigs per year. This information is based on the annual Spain country reports (2008-2012).

A single national-level survey was conducted in 2008 to estimate the baseline prevalence of *Salmonella* infection in breeding farms as part of a larger effort at a European level (DC 2008/55/EC; EFSA, 2009). In Spain 359 holdings (150 breeding and 209 production holdings, sample size set to estimate a 50% expected prevalence with a 7.5% accuracy at a 95% confidence level) and 3,590 pens were sampled. A farm-level prevalence of 64.0 and 53.1% in breeding and production holdings was found, respectively, placing Spain as the country with the highest and second highest prevalence of *Salmonella* infection in each subpopulation [1]. The serotypes with a higher proportion of isolates in breeding farms were *S. Rissen* (15.88%), *S. Typhimurium* (13.09%), *S. Anatum* (9.47%) and *S. Derby* (8.08%).

3.3. *Salmonella* in meat and meat products

Salmonella status in meat is monitored at the slaughterhouse through the analysis of tissue samples in swine carcasses (20 cm² in 4 tissue samples using an abrasive sponge) and checks of the operator's microbiological tests to control *Salmonella* through the food chain, in accordance with EC Regulations 217/2014, 218/2014, 1441/2007, and 2073/2005. If positive results are obtained, AECOSAN communicates it to MAPA so that further action can be pursued at the farm of origin, in particular regarding biosecurity practices and *Salmonella* testing.

In Spain, a minimum of 50 carcasses are randomly sampled per year per slaughterhouse by the official veterinarian at the slaughterhouse, each one from a different farm (unless it is a small slaughterhouse, in which case the number of samples depends on the result of a risk assessment). If 3 or more samples are positive (out of the 50), corrective action is taken. The official veterinarian informs its region of the results of this sampling, together with the information about the number of positive samples obtained from the additional carcass sampling performed by slaughterhouse operators. The corrective action plan can include an investigation of the origin of the animals and the farm biosecurity measures, which implies a close collaboration with MAPA which is the Competent Authority on animal health. Meat product testing is the responsibility of the business operator of slaughterhouses or establishments producing minced meat, meat preparations or mechanically separated meat, and is regulated by law (EC Regulation 20173/2005). This regulation allows the business operator to decide the sampling frequency according to its HACCP plan, but specifies a minimum requirement to guarantee a comparable level of control with other EU countries. This regulation also specifies the diagnostic test that should follow ISO 6579 for *Salmonella* or any other validated analytic method.

The information on pork meat monitoring contained in the annual country reports, provided to EFSA by the Regional Health Services, includes the number of units tested (each unit corresponds to 25 g of tissue sample) and the number that resulted positive to *Salmonella* in fresh meat (at the processing plant, at retail and at the slaughterhouse), in meat products raw but intended to be eaten cooked (at the processing plant and at retail) and occasionally, in raw but ready to be eaten raw meat products. The reports also specify the serotype when available. There is a low percentage of units positive to *Salmonella* (<5% of total units tested). Similar to slaughter pigs, *S. Typhimurium* has been detected every year (of the same period analysed for pig status: 2008-2012), with the overall average prevalence of *S. Typhimurium* during this period being less than 2%. However one cannot deduct from these data how many constituted a positive sample according to the regulation that led to corrective action.

Surveillance at the pig and pig meat sources is summarized in Figure 3.




Figure 3. *Salmonella* surveillance “from-farm-to-fork” (from pig source). AECOSAN notifies MAPA of the detection of meat contaminated with *Salmonella*, so that corrective action can be taken (indicated with a dotted line in Figure 1). In addition, the two surveys carried out by EFSA request are indicated, the results of which are kept by MAPA.

Table 4 shows the most relevant results on food monitoring provided by SCIRI from 2014 to 2016. During this period 7.75% of the overall alerts were related to *Salmonella* spp. presence in animal products; 0.39% of the information notifications were associated to *Salmonella* spp. biological risk in Spanish animal products; and 0.22% of the border rejections were related to *Salmonella* spp. presence in animal products.

Table 4. Overall biological risks and *Salmonella* spp. related notifications reported by SCIRI

| Notification type | Notification definition | Year | | | |
|----------------------------------|--|------|------|------|-------|
| | | 2014 | 2015 | 2016 | Total |
| Alert | Overall food & drinks alerts | 194 | 184 | 203 | 581 |
| | Biological risk alerts related to animal products | 44 | 42 | 53 | 139 |
| | Biological risk alerts related to animal products with <i>Salmonella</i> spp. isolation | 12 | 17 | 16 | 45 |
| Information | Overall information notifications | 1321 | 1333 | 1478 | 4132 |
| | Biological risk information notifications | 430 | 451 | 504 | 1385 |
| | Biological risk information notifications related to animal Spanish products (Spanish involvement) | 38 | 19 | 53 | 110 |
| | Biological risk information notifications related to <i>Salmonella</i> spp. biological risk in animal Spanish products (Spanish involvement) | 8 | 0 | 8 | 16 |
| Border products rejection | Overall frontier products rejection | 1296 | 1310 | 1078 | 3684 |
| | Biological risk frontier products rejection related to animal products | 242 | 242 | 154 | 638 |
| | Frontier products rejection related to <i>Salmonella</i> spp. detection in animal products with <i>Salmonella</i> spp. isolation | 6 | 1 | 1 | 8 |

3.4. Cases and outbreak investigation in humans

Figure 4 summarizes the surveillance of *Salmonella* in humans, which starts with the onset of clinical signs in a suspect patient.




Figure 4. *Salmonella* surveillance in suspect patients in Spain (from ingestion of contaminated product or other exposure route). It summarizes the surveillance of *Salmonella* in humans, which starts with the onset of clinical signs in a suspect patient.

Patients exhibiting diarrhoea, fever, abdominal pain or vomits who seek primary or specialist care at the local public outpatient health centre or are admitted/referred to a hospital, are registered as “cases” if, in addition, *Salmonella enterica* (other than Typhi or Paratyphi) is confirmed in the laboratory by isolation in faeces, clinical specimens (infected wound, etc.) or any sterile tissue / body fluid (blood, urine, etc.). An outbreak is defined as two or more *Salmonella* cases with a common source as background exposure. Besides the hospital and regional reference laboratories, the National Centre of Microbiology, also at the ISCIII, is the Reference Laboratory for *Salmonella* and *Shigella* in Spain, and can provide deeper microbiological analyses than the other laboratories mentioned. ISCIII holds clinical information (symptoms onset and outcome), epidemiological data (outbreak association and outbreak identifier code) and microbiological variables (sample type and microbiological identification to serovar level). Information on hospitalised cases can be found in the hospitalisations Minimum Basic Data Set (MBDS) maintained since 1987 by the MSSSI. When *Salmonella* outbreaks are microbiologically confirmed to be of food origin, AECOSAN notifies it, when applicable, nationally through SCIRI and internationally through RASFF, ensuring the exchange of verified information and follow-up actions within a network of EU Member States.

During the year 2014, 13 of the 19 Spanish regions (55.5% of the overall population in the country) reported salmonellosis cases (7,295 in total). The number of regions reporting increased in 2015, with 15 regions (70.4% of the overall population in the country) notifying 9,069 cases to ISCIII (Table 1).

Table 1. Summary of results from the different sources, years 2014 and 2015.

| Source | Institution | Cases definition | Launch date of the dataset | Coverage | Year | |
|--------------------------|-------------|---|------------------------------------|-------------------------|---|---|
| | | | | | 2014 | 2015 |
| Notifiable Diseases (ND) | ISCIH | Non-typhoid, non-paratyphoid salmonellosis | 2015 (2014 volunteer notification) | National | 7,295 | 9,069 |
| Outbreaks Database | ISCIH | <i>Salmonella</i> confirmed outbreaks | 2006 | National | 241 outbreaks (11.7% related to meat and derivatives ¹) | 281 outbreaks (10.2% related to meat and derivatives ¹) |
| SIM | ISCIH | Any non-typhoid, non-paratyphoid <i>Salmonella</i> isolates | 1995 | 72 labs from 11 regions | 5,001 | 5,215 |
| Hospital registry (MBDS) | MSSSI | Hospitalisations in patients with gastroenteritis caused by <i>Salmonella</i> spp. (non-typhoid, non-paratyphoid) | 1987 | National | 3,526 | 3,776 |

¹ Percentage refers to those cases with information of the source of infection

Table 2 shows the *Salmonella* spp. serovars distribution reported by RENAVE.

Table 2. *Salmonella* spp. related notifications reported by RENAVE (% from the overall cases with information)

| Year | Number of cases with available information | S.Typhimurium | S.Enteritidis | S. Typhimurium Monophasic | S.Newport | <i>Salmonella</i> , other serotypes |
|------|--|---------------|---------------|---------------------------|------------|-------------------------------------|
| 2014 | 3,877 (50.5%) | 1,640 (44.5%) | 1,220 (33.1%) | 83 (2.3%) | 17 (0.46%) | 69 (1.87%) |
| 2015 | 5,299 (58.4%) | 2,152 (61.1%) | 1,066 (30.3%) | 153 (4.3%) | 28 (0.79%) | 122 (3.5%) |

The MBDS compiles information from more than 95% of the public/private hospitals and of 99% of the discharges in Spain, and in some regions 100%. Clinical information about severe cases of salmonellosis requiring hospitalisation can be obtained from it. Hospitals confirm the cases in their own laboratory or by sending the sample to the national or the regional reference laboratories, but no detailed information of serotypes is provided in the dataset. The MBDS was launched in 1987, with constant improvement in the data collection. The last improvement was made in 2016 in which the dataset started to include ambulatory surgeries and procedures and switched to ICD-10 coding (Royal Decree 69/2015). Non-typhi, non-Paratyphi *Salmonella* diagnosis are coded under the ICD-10 code A02. Overall, there were 3526 recorded hospitalisations related to non-typhi, non-Paratyphi *Salmonella* in 2014 and in 2922 (82.9%) hospitalisations *Salmonella* infection was the main diagnosis.

In 2015, there were 3776 hospitalisations related to non-typhi, non-Paratyphi *Salmonella* and in 3185 (84.3%) hospitalisations *Salmonella* infection was the main diagnosis (Table 1).

The outbreaks dataset maintained by ISCIII contains further information about the outbreak including food item/s investigated. Outbreak information can also be linked to the individual patient's information collected in the ND database through the outbreak identifier code provided in both, outbreaks and individual notifications. In 2014, 241 *Salmonella* associated outbreaks involving 1,681 individuals were notified by the regions to ISCIII (Table 1). Among them, 249 (14.8%) were hospitalised and 7 (0.4%) died. Half of the outbreaks were caused by *S. Enteritidis* (53.9%), followed by *S. Typhimurium* (17%). In total, 73% (176/241) of the outbreaks were foodborne, one of the 176 transmitted by water. Some suspect food was identified in 72.7% of the outbreaks. Among suspect foods, the most commonly implicated food was the egg and its derivatives (68.8% of the outbreaks with food identification), followed at a great distance by meat and meat products (11.7%). In 2015, 281 *Salmonella* associated outbreaks involving 1,920 individuals were notified (Table 1), 35 (1.8%) of them were hospitalized and 2 (0.1%) died. In 116 (77.3%) outbreaks the agent was *S. Enteritidis*, followed by *S. Typhimurium* with 30 (20%) outbreaks. Some suspect food was identified in 59.4% of the outbreaks and, among them, the most commonly implicated food was the egg and its derivatives (77.2% of the outbreaks with food identification). In 10.2% of the outbreaks the suspect food was meat and meat products.

SIM reports contain information on circulating *Salmonella* serovars. For all pathogens, SIM coverage of the Spanish population was 34% in 2014 and 30% in 2015. During 2014 and 2015, 72 microbiology laboratories of 11 Spanish regions participated in the SIM providing non-typhi, non-Paratyphi *Salmonella* isolates information. In 2014 and 2015 respectively, 5,001 and 5215 non-typhi, non-Paratyphi *Salmonella* isolates were reported.

3.4.1. Datasets linkage

Table 3 shows the summary of the results from the different human *Salmonella* surveillance data sources by region.

Regions notifying cases at national level to the ND system present a very similar number of reported cases to the positive isolates reported across SIM, showing that both systems are providing similar information. Some differences in numbers, especially when there are more positive isolates reported in the SIM, could be related to more than one isolate recovered from the same patient. The table also shows that the proportion of hospitalisations compared with the number of cases reported to the ND system was 30.5% (4999 hospitalisation vs. 16364 cases reported). This implies that most of the regions are reporting mild *Salmonella* cases attending primary care across both systems, ND and SIM. It is expected that the four pending regions will join soon and report data from 2015 onwards to the ND System.

Table 3. Summary of results from the different sources by region.

| Region | 2014 | | | 2015 | | |
|-----------|------------|----------------|-------------------------|------------|----------------|-------------------------|
| | Cases (ND) | Isolates (SIM) | Hospitalisations (MBDS) | Cases (ND) | Isolates (SIM) | Hospitalisations (MBDS) |
| Andalusia | | | 523 | | | 503 |
| Aragon | 442 | 461 | 194 | 555 | 557 | 190 |

| | | | | | | |
|---------------------|-------------|-------------|-------------|-------------|-------------|-------------|
| Asturias | 437 | 442 | 132 | 376 | 380 | 113 |
| Balearic Islands | | | 69 | | | 89 |
| Canary Islands | 568 | 575 | 130 | 370 | 378 | 83 |
| Cantabria | | | 44 | 83 | | 38 |
| Castile-La Mancha | 149 | 150 | 178 | 177 | 177 | 210 |
| Castile and Leon | 693 | 457 | 339 | 811 | 436 | 367 |
| Catalonia | 1778 | 1806 | 489 | 1968 | 2090 | 522 |
| Valencian Community | 2217 | | 385 | 2539 | | 422 |
| Extremadura | 79 | 81 | 81 | 287 | 66 | 97 |
| Galicia | | | 151 | | | 197 |
| Madrid | | | 431 | 778 | | 547 |
| Murcia | | | 143 | | | 153 |
| Navarra | 280 | 282 | 52 | 337 | 327 | 49 |
| Basque Country | 432 | 533 | 147 | 625 | 616 | 135 |
| La Rioja | 185 | 189 | 24 | 148 | 149 | 33 |
| Ceuta | 24 | 25 | 9 | 3 | 39 | 17 |
| Melilla | 11 | | 5 | 12 | | 11 |
| Total | 7295 | 5001 | 3526 | 9069 | 5215 | 3776 |

4. Discussion

This study reviews the information across the different reports and institutions participating in the surveillance of *Salmonella* in Spain. We have selected the swine reservoir given the importance of this livestock species in Spain and the lack of official control programme for *Salmonella*, in an attempt to connect the main stakeholders of surveillance in animals and humans in Spain, which are ultimately MAPA and MSSSI. The EFSA and the ECDC request EU Member States to report the design and results of food-borne zoonotic diseases surveillance by gathering such information directly from each of the stakeholders involved. The most extensive and detailed information in swine origin reports corresponds to the status of *Salmonella* at the animal source, where there seems to be a higher degree of collaboration between the MAPA and MSSSI than in the event of *Salmonella* detection in food items in human cases or outbreaks. The surveillance of *Salmonella* in pigs, meat and humans are not comparable *per se*, since they are designed for different purposes. In pigs and in meat, the objective is to routinely monitor the burden of *Salmonella* infection to reduce the risk of human illness through foodborne contamination or from contact with infected pigs [16]. The risk reduction measures in pigs and meat differ, since meat contamination could be also a result of poor hygiene practices at slaughter or processing. In humans, the objective is to identify *Salmonella* in clinical suspects as soon as possible to be able to prevent new cases or outbreaks by the enforcement of hygienic practices and by withdrawing or treating the suspect source of infection [17].

Structurally, all surveillance systems on swine-related *Salmonella* include similar information on sampling strategy, types of samples and, frequency of sampling, laboratory testing and results and dissemination. Under the prism of a collaborative One Health surveillance system, there is, however, ample room for improvement. Bordier et al [18] propose 6 degrees of possible collaboration in a multi-sectoral surveillance system in the planning, data collection (sampling and lab testing), data sharing,

results sharing, data analysis/interpretation, dissemination to decision makers and communication to surveillance actors and end users. After the analyses of the information in the different datasets and websites, we found very limited evidence of a high degree of collaboration in the steps proposed for a cooperative surveillance process. More likely, all steps are being taken separately for each sector, but the results are being shared periodically. The only link we found between sectors was in the procedures regarding a positive result in meat, but we could not find any report in which such collaboration was reflected.

In addition, we found at least 7 different databases (4 from the human sector: SIM, ND, Outbreak investigation, and MBDS; 1 from the meat sector: SCIRI-AECOSAN; and 2 from the veterinary sector: national monitoring, EFSA surveys) with information on *Salmonella* surveillance in Spain. Better knowledge of these databases to support an integrated "One Health" surveillance approach could help to better focus the efforts in *Salmonella* control, for example by performing a joint analysis of the distribution and trends of *Salmonella* serovars in time and space and to determine more precisely the burden of each *Salmonella* serovar. For example, in Canada, weekly counts of *Salmonella* from farm animals, meat and humans allowed to create baseline models and identify significant clusters across the different sectors [19]. While identifying "hot-spots" of *Salmonella* specific serovar distribution would be desirable, the substantial differences in disease dynamics among *Salmonella* serotypes must be taken into account for a correct epidemiological interpretation of results. Arnedo-Pena et al [4] highlight some differential characteristics regarding the incubation period, outbreak duration, attack rate, hospitalization rate, probability of reaching a microbiological diagnosis, and human behaviour that influence the variation in disease dynamics of foodborne outbreaks of *S. Typhimurium* and *S. Enteritidis* in Spain. It would be desirable to characterize also other serovars identified in animal health surveillance results (i.e. in pigs: *S. Rissen* and *S. Derby*) to target interventions. Human behaviour is so predominant in the exposure to food-borne diseases that the inclusion of social sciences in the multidisciplinary approach to investigate *Salmonella* changes over time have already been suggested [8].

Linking outbreak investigation with farm status can be very time-consuming and an association with the sufficient level of confidence hard to obtain. So far, *Salmonella* in pigs is not under an official control programme in Spain. Spain is classified as a "high prevalence" country according to EFSA. The monitoring of *Salmonella* in slaughter pigs is representative of the production system in Spain, in which more than 80% of the total census are fattening pigs [20]. The detailed census of pig holdings available in Spain at the coordinate level would in principle facilitate spatial analysis of the surveillance results, at least at the descriptive level. A deeper epidemiological analysis of the surveillance data is underway to assess their usefulness for identification of spatial and temporal trends in detection of *Salmonella*/specific strains, and for the early detection of emerging strains.

Salmonella surveillance in meat and meat products is the responsibility of the business operator, and the MSSSI vet officially checks and oversees the surveillance. Again, we have not found any evidence of a joint epidemiological analysis comparing prevalence trends between MAPA with AECOSAN surveillance results that could lead to a change in the surveillance strategy to detect and correct any potential problem. In other countries, like in Denmark, the joint analysis of the results from carcass swabs, serological surveillance of meat juices, ceacal samples and antibiotic treatment are performed by the Agriculture & Food Council to investigate any alerts deriving from any of the surveillance systems mentioned [21]. Even if the time of infection might cause mismatches among the results of these surveillance systems, since some detect antibodies and others are directed at isolating the bacteria, meaningful interpretations can be obtained over time.

Early detection of human cases before they become an epidemic is crucial and the MSSSI, in an effort to combat *Salmonella* to decrease underreporting, has declared human salmonellosis notifiable since 2015. This means that now each human case that is detected (clinical signs + microbiological confirmation) must be notified to ISCIII, whether or not it leads to an outbreak. This way, there is now a complete database with information about occurrence of cases including location, symptomatology and serovar involved. Prior to 2015, the only source of clinical information of the

disease was the National Registry for Hospitalisations and data sources for microbiological information were scattered among different databases (regional surveillance, laboratory and outbreaks databases). Surveillance systems implementation, however, requires a start-up period and a continuous evaluation of the quality of the system. Hospitalisation registries have been used as a tool not only to evaluate the epidemiology of the most severe cases of *Salmonella* infections, but to support the evaluation of the quality and performance of the surveillance systems [22]. Regions now have the appropriate mechanisms to compile the cases' microbiological and clinical information sent to the ND system. Consequently, this will increase the quality of the information and simplify the information in a unique dataset covering all aspects of the disease. Also, the inclusion in the ND dataset of primary care cases' clinical and demographic information will allow estimating a more realistic impact of the disease. This means that information on *Salmonella* human cases at National level should improve notably in the next years. Still, there is not an active surveillance system for human *Salmonellosis* implemented in Spain as there is in other countries like the USA (FoodNet), where laboratory, hospital and population surveys are conducted to estimate the burden and the attribution of foodborne illness [23].

Of the range of methods described in the literature to attribute the source to human salmonellosis [24], only outbreak investigation is carried out in Spain. This information is published in aggregated form. To be able to identify relevant risk factors for human infections, a closer collaboration between animal and human health stakeholders is required to access and analyse these data. Similarly, the information on pig status that is publicly available allows a basic analysis to detect temporal variations in the serovars present, but there is no further investigation to identify also a spatial variation, potential exposure risks for humans or quantitative microbial risk assessments to investigate the impact of different control scenarios and help decide the best strategy (e.g. reducing carcass load versus farm prevalence).

It is very likely that both the human and the animal monitoring of *Salmonella* would benefit from a better exchange of information and collaboration. For MSSSI, such exchanges could help to map the areas at higher risk of exposure, being able to alert the population to take measures to reduce that risk (e.g. enhance hygiene, cook thoroughly, etc.). For MAPA, it could improve the overall prevalence picture and help economizing routine surveillance efforts and expenses by identifying and targeting those areas from which more cases associated with pork meat arise.

Author Contributions: All authors participated in the creation of the argument line of this text and approved the final version. Conceptualization, Marta Martínez-Avilés and Ana de la Torre; Funding acquisition, Ana de la Torre; Investigation, Marta Martínez-Avilés, Macarena Garrido-Esteba and Julio Álvarez; Methodology, Marta Martínez-Avilés; Project administration, Ana de la Torre; Resources, Marta Martínez-Avilés, Macarena Garrido-Esteba, Julio Álvarez and Ana de la Torre; Supervision, Ana de la Torre; Visualization, Macarena Garrido-Esteba and Ana de la Torre; Writing – original draft, Marta Martínez-Avilés; Writing – review & editing, Macarena Garrido-Esteba and Julio Álvarez.

Acknowledgments: We acknowledge the European Union's Horizon 2020 Research and Innovation programme under Grant Agreement No 773830 (One Health European Joint Programme) and the Community of Madrid project S2013/ABI-2747 for financial support.

Conflicts of Interest: The authors declare no conflict of interest

References

1. Heredia, N.; García, S. Animals as sources of food-borne pathogens: A review. *Animal Nutrition* **2018**, *4* (3), 250-255.
2. De Jong, B.; Ekdahl, K. The comparative burden of Salmonellosis in the European Union member states associated and candidate countries. *BMC Public Health* **2006**, *6*, 4-4.
3. EFSA & ECDC. The European Union summary report on trends and sources of zoonoses, zoonotic agents and food-borne outbreaks in 2016. *EFSA Journal* **2017**, *15* (12), e05077.

4. Arnedo-Pena, A.; Bellido-Blasco, B.B.; Romeu-Garcia, M.A.; Meseguer-Ferrer, N. Detection of foodborne *Salmonella* Typhimurium outbreaks. *Enfermedades Infecciosas y Microbiología Clínica* **2017**, *35*(7), 469-473.
5. Ministry of Agriculture (MAPA). Pork sector figures: Main economic indicators [El sector de la carne en cifras: Principales indicadores económicos]. Livestock Products Subdirectorate. **2017**, 108 pp.
6. Pires, S.M.; de Kneegt, L.; Hald, T. Estimation of the relative contribution of different food and animal sources to human *Salmonella* infections in the European Union. Question No. EFSA-Q.2010-00685. **2011**. 80 pp.
7. EFSA. Technical specifications on the harmonised monitoring and reporting of antimicrobial resistance in *Salmonella*, *Campylobacter* and indicator *Escherichia coli* and *Enterococcus spp.* bacteria transmitted through food. *EFSA Journal* **2012**, *10*, 2742
8. Newell, D.G.; Koopmans, M.; Verhoef, L.; Duizer, E.; Aidara-Kane, A.; Sprong, H.; Opsteegh, M.; Langelaar, M.; Threlfall, J.; Sheutz, F.; van der Giessen, J.; Kruse, H. Food-borne diseases – The challenges of 20 years ago still persist while new ones continue to emerge. *International Journal of Food Microbiology* **2010**, *139*, S3-S15.
9. Pires, A.F.A.; Funk, J.A.; Bolin, C.A. Longitudinal study of *Salmonella* shedding in naturally infected finishing pigs. *Epidemiology and Infection* **2013**, *141*, 1928-1936
10. Alban, L.; Barfod, K.; Petersen, J.V.; Dahl, J.; Ajufo, J.C.; Sando, G.; Krog, H.H.; Aabo, S. Description of extended pre-harvest pig *Salmonella* surveillance-and-control programme and its estimated effect on food safety related to pork. *Zoonoses and Public Health* **2010**, *57* Suppl 1, 6-15
11. Casanova-Higes, A.; Andres-Barranco, S.; Mainar-Jaime, R.C. Influence of on-farm pig salmonella status on salmonella shedding at slaughter. *Zoonoses and Public Health* **2017**, *64*, 328-336
12. Schroeder, S.; Harries, M.; Prager, R.; Höfig, A.; Ahrens, B.; Hoffmann, L.; Rabsch, W.; Mertens, E.; Rimek, D. A prolonged outbreak of *Salmonella* Infantis associated with pork products in central Germany, April–October 2013. *Epidemiology and Infection* **2016**, *144*, 1429-1439
13. European Research Programme. <https://onehealth.ejp.eu/structure/jrp6-nova/> (05.12.2019)
14. Gil Prieto, R.; Alejandre, C.G.; Meca, A.Á.; Barrera, V.H.; de Miguel, Á.G. Epidemiology of hospital-treated salmonella infection; data from a national cohort over a ten-year period. *Journal of Infection* **2009**, *58*, 175-181
15. Garrido-Esteba, M.; Latasa, P.; Ordóñez-León, G.Y.; Martínez-Avilés, M.; de la Torre, A.; García-Comas, L. Non-typhi, non-Paratyphi *Salmonella*-related hospitalisations in Spain: trends, clinical aspects, risk factors for worse prognosis and hospital costs. *European Journal of Clinical Microbiology & Infectious Diseases* **2018**. <https://doi.org/10.1007/s10096-018-3433-1>
16. OIE. Prevention and control of *Salmonella* in commercial pig production systems. *Terrestrial Animal Health Code* **2018**, Chapter 6.14
17. ISCIII. Protocols of the National Epidemiological Surveillance Network. RENAVE **2015**. 680 pp. Agencia Estatal Boletín Oficial del Estado, Madrid. ISBN: 725-14-00-19.
18. Bordier, M.; Delavenne, C.; Uea-Anuwong, T.; Nguyen, D.T.; Goutard, F.; Hendriks, P. Characterisation of One Health surveillance based on a systematic literature review and an expert-opinion elicitation. The 15th *International Symposium of Veterinary Epidemiology and Economics* **2018** November 12, Chiang Mai, Thailand.
19. Vrbova, L.; Patrick, D.M.; Stephen, C.; Robertson, C.; Koehoorn, M.; Parnley, E.J.; DE With N.I.; Galanis, E. Utility of algorithms for the analysis of Surveillance data. *Epidemiology and Infection* **2016**, *144* (10), 2165-2175.
20. Martinez, M.; Iglesias, I.; Bosch, J.; Jurado, C.; Vicente, J.; Sánchez-Vizcaíno, J.M.; de la Torre, A. Protecting extensive pig systems in Spain [Proteger al sector porcino extensivo en España]. *Euroganadería* **2018**
21. Mogelmoose, V.; Lindegaard, L.L.; Dahl, J. A sudden increase in *Salmonella*- how the surveillance system reacted: a Danish experience. 12th *International Symposium on the Epidemiology and Control of Biological, Chemical and Physical Hazards in Pigs and Pork* **2017** August 21-24, Foz do Iguaçu, Brazil.
22. Mor, S.M.; DeMaria Jr, A.; Naumova, E.N. Hospitalization records as a tool for evaluating performance of food- and water-borne disease surveillance systems: A Massachusetts case study. *PLoS One* **2014**, *9*, e93744
23. <https://www.cdc.gov/foodborneburden/>
24. Pires, S.M.; Vieira, A.R.; Hald, T.; Cole, D. Source attribution of human salmonellosis: An overview of methods and estimates. *Foodborne Pathogens and Diseases* **2014**, *11*, 667-676