

Title: The Energy Homeostasis Principle: Neuronal Energy Regulation Drives Local Network Dynamics Generating Behavior.

Authors: Rodrigo C. Vergara¹, Sebastián Jaramillo-Riveri², Alejandro Luarte³, Cristóbal Moënnelocoz⁴, Rómulo Fuentes⁴, Andrés Couve³, Pedro E. Maldonado^{1*}.

Running head: The Energy Homeostasis Principle

¹Neurosystems Laboratory, Biomedical Neuroscience Institute, Faculty of Medicine, Universidad de Chile, Santiago, Chile.

²Institute of Cell Biology, School of Biological Sciences, University of Edinburgh, Edinburgh, UK.

³Cellular and Molecular Neurobiology laboratory, Biomedical Neuroscience Institute, Faculty of Medicine, Universidad de Chile, Santiago, Chile.

⁴Motor Control Laboratory, Biomedical Neuroscience Institute, Faculty of Medicine, Universidad de Chile, Santiago, Chile.

***Correspondence:**

Pedro E. Maldonado. Ph.D.
Departamento de Neurociencia
Facultad de Medicina
Universidad de Chile
Independencia 1027
Santiago, Chile
pedro@uchile.cl

Keywords: homeostasis, energy, neuronal networks, behavior, emergent properties.

Abstract

A major goal of neuroscience is understanding how neurons arrange themselves into neural networks that result in behavior. Most theoretical and experimental efforts have focused on a top-down approach which seeks to identify neuronal correlates of behaviors. This has been accomplished by effectively mapping specific behaviors to distinct neural patterns, or by creating computational models that produce a desired behavioral outcome. Nonetheless, these approaches have only implicitly considered the fact that neural tissue, like any other physical system, is subjected to several restrictions and boundaries of operations.

Here, we propose a new, bottom-up conceptual paradigm: The Energy Homeostasis Principle, where the balance between energy income, expenditure, and availability are the key parameters in determining the dynamics of the found neuronal phenomena from molecular to behavioral levels. Neurons display high energy consumption relative to other cells, with metabolic consumption of the brain representing 20% of the whole-body oxygen uptake, contrasting with this organ representing only 2% of the body weight. Also, neurons have specialized surrounding tissue providing the necessary energy which, in the case of the brain, is provided by astrocytes. Moreover, and unlike other cell types with high energy demands such as muscle cells, neurons have strict aerobic metabolism. These facts indicate that neurons are highly sensitive to energy limitations, with Gibb's free energy dictating the direction of all cellular metabolic processes. From this activity, the largest energy, by far, is expended by action potentials and post-synaptic potentials; therefore, plasticity can be reinterpreted in terms of their energy context. Consequently, neurons, through their synapses, impose energy demands over post-synaptic neurons in a close loop-manner, modulating the dynamics of local circuits. Subsequently, the energy dynamics end up impacting the homeostatic mechanisms of neuronal networks. Furthermore, local energy

management also emerges as a neural population property, where most of the energy expenses are triggered by sensory or other modulatory inputs. Local energy management in neurons may be sufficient to explain the emergence of behavior, enabling the assessment of which properties arise in neural circuits and how. Essentially, the proposal of the Energy Homeostasis Principle is also readily testable for simple neuronal networks.

Introduction

Throughout evolution, the development of the nervous system has enabled animals with the capacity to manifest ever-growing complex behavior, which has helped them survive in a changing environment. Understanding how neurons arrange themselves into neural networks that work at producing different behaviors has always been a major goal of neuroscience. Various conceptual frameworks have aimed to explain how behavior emerges from neuronal activity. Arguably, the most relevant is the Neuron Doctrine, proposed by Santiago Ramón y Cajal and further developed by Heinrich Waldeyer-Hartz and Horace Barlow (Bock 2013; Barlow 1972). Since then, the same logic has spread into coding paradigms (Fairhall 2014; Lettvin et al. 1959; Yuste 2015), especially in information processing frameworks (Friston 2002; Lorenz, Jeng, and Deem 2011; Robbins 2010; Fodor 1983), and has been scaled from neurons up to neural networks (Yuste 2015). A common and key element of these conceptual approaches has been to find neuronal correlates of behaviors, effectively associating specific behaviors with distinct neural patterns. This top-down approach (using behavior as a reference to be mapped into neuronal circuits) has been very successful in providing single-unit or network models that can implement the observed behaviors, yet simultaneously, may make difficult the capture of the emergence of behavior, which is by-large a bottom-up phenomenon. This methodological approach also limits our capacity of predicting the boundaries of the capabilities or the spectrum of behaviors of a given system, because we map or associate only those behaviors that have been well characterized. More importantly, all theoretical approaches, to our knowledge, have only implicitly addressed the fact that neural tissue, like any other physical system, is subjected to several restrictions and boundaries of operations.

Cells use energy to stay alive and at the same time, maintain some reserves to respond and adapt to dynamic situations, maintaining their *homeostasis*. For neurons, energy availability would be further important, as their energy expenses are high, as compared to other somatic cells (Attwell and Laughlin 2001; Shulman et al. 2004). Indeed, the metabolic consumption of the brain, which represents 20% of whole-body oxygen consumption, contrasts with the neural tissue representing only 2% of whole body weight (Shulman et al. 2004). Interestingly, the total brain energy consumption increases proportionally with the number of neurons among different species, including humans (Herculano-Houzel 2011), and the total energy expenditure associated to a neuron during the signaling and resting states is constant in different mammalian species (Hyder, Rothman, and Bennett 2013). Thus, neurons seem to present a highly specialized system for managing their energy demands.

Several evidences demonstrate that it is reasonable to assume a constant value for energy availability for neurons over the long term (energetic homeostasis). For instance, cultured neurons exhibit a steady value for free adenosine triphosphate (ATP) in basal conditions, which transiently decrease during the induction of glutamatergic synaptic activity through various energy challenges (Marcaida et al. 1997; Marcaida et al. 1995; Lange et al. 2015; Rangaraju, Calloway, and Ryan 2014). This tight energy management suggests a relevant role for neuronal energy homeostasis on neuronal and network functional properties.

Here, we propose a new bottom-up conceptual paradigm for neuronal networks: The Energy Homeostasis Principle. Under this principle, the condition of maintaining neuronal homeostasis triggers synaptic changes in the individual but connected neurons, resulting in the local energy balance scaling up to a network property. This conceptual framework supposes that

energy management might be critical in determining plasticity, network functional connectivity, and ultimately behavior.

Cellular Homeostasis and Gibbs Free Energy

In this article, we propose that behavior may arise as an emergent property rooted in energy requirement of neurons, thus, we would like to start from the level of biochemistry and metabolism. As such, we will begin with the fact that cells are dynamic molecular machines that require the nutrient intake to stay alive. Many biological processes are thermodynamically unfavorable, and through metabolism, cells draw energy from nutrients and generate metabolic resources necessary to drive their cellular activities (Hofmeyr and Cornish-Bowden 2000) (for a schematic, see Figure 1A). Cellular homeostasis can be defined as a state where the production and consumption of metabolic resources balance each-other, and thus their concentration is constant in time. For our specific context, balancing the intake and consumption of metabolic resources will unavoidably have a global impact on the cellular processes. The network of metabolic processes is large and complex, limiting, to some extent, our capacity to predict cellular behavior using basic principles. Nonetheless, biochemical reactions must be consistent with the laws of thermodynamics.

Thermodynamics can help us understand how a system evolves in time through the comparison of the thermodynamic potential between an initial and final state. For processes at a constant temperature and pressure, the thermodynamic potential is given by Gibbs Free Energy (G). This thermodynamic potential will dictate a directional bias of chemical reactions. The Gibbs Free Energy—supporting cellular processes—is provided with a finite amount of metabolic resources. Thus, there is a trade-off between the potential for metabolic work and metabolic expenses, which, we propose, may explain some well-established phenomenology of how cells respond to external perturbations. Additionally, the change in the Gibbs Free Energy (ΔG) and the

rate associated with chemical transformations are related (Crooks 1999). To illustrate the relation between thermodynamics and kinetics, for a reversible reaction $X \rightleftharpoons Y$, the following relation constrains kinetic rates:

$$\frac{\text{rate}(X \rightarrow Y)}{\text{rate}(X \leftarrow Y)} = e^{G(X) - G(Y)} \text{ (eq. 1)}$$

In simple terms, the difference $\Delta G\{X \Rightarrow Y\}$ can be thought of as a “directional bias,” indicating how favorable one direction is over the other. In more detail, the Gibbs Free Energy is divided into two components, Enthalpy (H) and Entropy (S):

$$G(X) = H(X) - TS(X) \text{ (eq. 2)}$$

where, T is the absolute temperature (Silbey, Alberty, and Mounji 2004). In the context of chemical transformations, Enthalpy is a measure of the energy required to form a given substance, disregarding interactions with other molecules; whereas Entropy can be interpreted as a correction accounting for all possible combinations by which molecules can react (Danos and Oury 2013). Given the combinatorial nature of entropy, it can also be interpreted as a measure of disorder or information, but certain care must be taken for this interpretation to have physical meaning (Jaynes 1965). We wish to recognize that the direct application of thermodynamics to biology has many challenges, particularly in describing macro-molecular processes (Cannon 2014; Cannon and Baker 2017; Ouldrige 2018), combining large systems of reactions (e.g. kinetic parameters may be required), and accounting for fluctuations from average behavior (Marsland and England 2018).

A long-standing observation in biology, rooted in thermodynamic laws is that for cells to function, they must couple unfavorable reactions ($\Delta G > 0$) with more favorable ones ($\Delta G < 0$). Common examples of unfavorable processes are the synthesis of macromolecules, and the maintenance of membrane potential; which are coupled with the hydrolysis of ATP, and GTP

providing more favorable ΔG (Nicholls 2013). In turn, ATP, GTP, and monomers for macromolecules are synthesized from nutrients through metabolism (Nicholls 2013). For instance, the maximum free energy provided by ATP hydrolysis is related to the concentration of ATP, ADP and phosphate.

$$\Delta G(ATP \rightarrow ADP + Pi) = \Delta G^\circ + RT(\log[ADP] + \log[Pi] - \log[ATP]) \text{ (eq. 3)}$$

where, ΔG° is the standard free energy, and \log , the natural logarithm. For generality, we will call hereafter “energy resources” the set of reactants that allow cells to maintain unfavorable reactions in the direction conducive to cellular functioning and survival. We wish to emphasize that balancing the internal production and consumption of metabolic resources by different reactions is critical, given that metabolic resources are finite and shared by many cellular processes. Thus, cells must manage their internal production and consumption of metabolic resources to stay alive and remain functional, which may be of special consequence to cellular activities with high energy demands, such as synaptic activity in neurons. Given that neurons are active most of the time, it is reasonable to expect that current and future disposal of energy resources is privileged, which may be reflected in the regulatory mechanisms responsible for synaptic, plastic changes. In the following section, we will explain how current evidence regarding neuron plasticity appears to support a relatively simple rule: Maintain the levels of energy disposal constant, by reducing the consumption of energy resources (e.g., reducing discharge rate, postsynaptic potential), or by increasing high-energy molecule production (e.g. mitochondria, and interactions with glia).

Energy management of brain neurons

Neurons are the paramount example of energy expenditure for their function and survival. This situation is reflected in their large metabolic rates and by the comparatively higher sensibility of brain tissues to oxygen and glucose deprivation (Ames 2000). Reactions controlling the

conversion of nutrients into available cytosolic levels of ATP are important to generate the potential metabolic work that is available to a neuron at any given time. During normal conditions, the primary energy substrate in the brain for neurons is blood-derived glucose; however, when at elevated levels in the blood, ketone bodies and lactate can be used as energy sources as well (Pierre J. Magistretti and Allaman 2018). The glycolytic pathway is the first step to glucose processing, where two pyruvates and two ATPs are generated from one molecule of glucose. In addition, the pyruvate could either be reduced to lactate or enter the Krebs cycle to produce metabolic reducing intermediates that will generate nearly 29 additional ATP molecules per glucose (through oxidative phosphorylation in the mitochondria). Although neurons and astrocytes are capable of glucose uptake and performing both glycolysis and the Krebs cycle, accumulated evidence supports the hypothesis that neurons may “outsource” glycolytic activity to astrocytes under activity conditions (Weber and Barros 2015). In addition, the CNS is provided with small glycogen reserves, which are predominantly present in astrocytes (A. M. Brown and Ransom 2007), but also found in neurons (Saez et al. 2014). In any case, the lactate derived from glycogen break-down may also provide ATP to the neurons under ischemic or sustained electric activity conditions (A. M. Brown and Ransom 2007).

ATP sources change dynamically with neuronal activity and several mechanisms account for this fine-tuning response. First, neuronal mitochondria are capable of raising ATP synthesis in response to increased synaptic stimuli (Jekabsons and Nicholls 2004; Connolly et al. 2014; Toloe et al. 2014; Lange et al. 2015; Rangaraju, Calloway, and Ryan 2014). Although the molecular mediators for this activation are not completely elucidated, the increase of the respiratory rate of an isolated mitochondria correlates well with the ADP concentration (G. C. Brown 1992), and neuronal mitochondrial function has been satisfactorily modeled considering the changes in ATP

and ADP levels (Berndt, Kann, and Holzhütter 2015). As an alternative mechanism, it has been reported that operating on milder stimulation conditions, the activity of Na-pump rapidly induces ATP synthesis of the mitochondria, in response to neuronal activity independent from changes in the adenosine nucleotides (Baeza-Lehnert et al. 2018). Second, neuronal activity is known to elicit local increases in blood flow (neurovascular coupling), glucose uptake, and oxygen consumption (Sokoloff 2008). Coherently, glucose uptake and glycolytic rate of astrocytes are further increased in response to the activity of excitatory neurons, potentially as a consequence of the local rise of glutamate, ammonium (NH₄), nitric oxide (NO), and importantly, K⁺ (Pierre J. Magistretti and Allaman 2018). As such, an increased glycolytic rate on astrocytes leads to lactate accumulation that is shuttled into neurons which generate ATP through oxidative phosphorylation. Thus, in CNS neurons, different neuronal and non-neuronal ATP sources work “on demand”, depending on the local levels of synaptic activity.

What is ATP used for in neurons?

Neurons are perhaps the largest eukaryotic cell in nature, their surface may be up to 10,000 times larger than an average cell (Horton and Ehlers 2003). The large size of neurons supposes that structural processes, such as protein and lipid synthesis or the traffic of subcellular organelles, should be sustained by high levels of ATP synthesis. In addition to this fact, energy consumption during signaling is far more important. Indeed, it has been estimated that nearly 75% of the gray-matter energy budget is used during signaling; a number that is coherent with the decrease of energy consumption, observed under anesthesia, and is estimated to be around 20% of the total energy budget (Harris, Jolivet, and Attwell 2012; Attwell and Laughlin 2001).

Most of the neuron's energy budget during signaling is used to restore ion gradients across the plasma membrane, mediated by the action of different ATP-dependent pumps. For example,

assuming an average firing rate of 4 Hz, a presynaptic neuron's ATP is mostly used for restoring the Na⁺ gradient due to action potentials, and to sustain the resting potential (22% and 20% of energy consumption, respectively). Meanwhile, at the postsynaptic neuron, ATP is primarily used to extrude ions participating in postsynaptic currents—about 50% of the energy consumption (Harris, Jolivet, and Attwell 2012). More detailed descriptions of the neuron energy budget is provided in Figure 1B.

Neuron's ATP availability is tightly regulated.

All cellular organizations require a minimum amount of ATP for survival. It is well known that when ATP levels decrease below a certain threshold for different eukaryotic cells, apoptosis or necrosis is induced (Eguchi, Shimizu, and Tsujimoto 1997). Nevertheless, determining the maximum and minimum thresholds of a cell's ATP requirement for not only to survive but to realize a specialized function, is less apparent. In any case, this feature must be necessarily shaped by evolutionary adaptations of cells to their specific tissue environment. It is not completely clear how a neuron's ATP levels, during rest and upon activity, may impact its structure and function. Interestingly, by computational and mathematical modeling, it has been proposed that a compromise among energy consumption and information processing capacity has shaped the fundamental features of neuronal structure and physiology, including neuronal body size, ion channel density, and the size and frequency of synaptic inputs (Sengupta et al. 2013). For example, a larger neuronal body has a better capacity to discriminate and respond to different synaptic inputs (coding capacity), but at the cost of higher energy consumption. On the other hand, with a fixed size for the soma, the ion channel density required to obtain maximum energy efficiency is at a lower value than the density needed to maximize the coding capacity. Similarly, although small synaptic inputs at low frequencies are energetically more efficient, better coding capacity arises

with larger inputs and rates. These energy constraints may have introduced important consequences during cellular evolution, such that neurons with similar shape and function may harbor similar metabolic features, even across different species.

Remarkably, it has been found that energy consumption of neurons, across the brains of varying species, is constant (Herculano-Houzel 2011). This result supposes a critical restriction for the function of neuronal networks and their coding properties. For example, sparse coding i.e., brain computations that emerge from the increased firing rate of a few neurons during a task, has been proposed as a mechanistic solution to the limited energy availability for brain neurons (Laughlin 2001; Lennie 2003; Attwell and Laughlin 2001; Weber and Barros 2015). Thus, it is also possible that variables such as the ATP cytosolic concentration may have been finely tuned during evolution to allow for the emergence of fundamental properties, including some forms of synaptic plasticity.

Accumulating evidence supports that neurons, in time, harbor a narrow window of ATP cytosolic concentration availability ($A(t)$). Despite not having dynamic measurements with absolute values of $A(t)$, different experimental approaches on cultured neurons show that this variable tends to remain constant at resting conditions and after momentary synaptic challenges. Accordingly, 60 minutes of different sorts of glutamatergic stimulation leads to a nearly five-fold decrease of $A(t)$ (Marcaida et al. 1997; Marcaida et al. 1995), but when a brief glutamatergic or electric stimulation is applied, only a transient and reversible decrease on ATP levels occurs and the $A(t)$ is subsequently restored to basal levels (Lange et al. 2015; Rangaraju, Calloway, and Ryan 2014).

Tight management of $A(t)$ also operates on axonal compartments with important functional consequences. For instance, isolated axons from the optical nerve, under low glucose conditions,

demonstrate a pronounced decay of ATP levels during high-frequency stimulation (50–100 Hz) (Trevisiol et al. 2017). Interestingly, compound action potentials (CAPs), generated by those stimulated axons, are reduced to the same extent and in high coincidence as the $A(t)$, suggesting that electric activity depends on $A(t)$ (Trevisiol et al. 2017). In addition, isolated axons exhibit a constant value for $A(t)$, which immediately and steeply decays after the inhibition of glycolysis and oxidative phosphorylation, in concomitance with CAPs. However, when inhibitors are washed out, both $A(t)$ and CAPs return to basal levels, further supporting that the system tends to reach a constant value for $A(t)$. The tendency of the system to set a constant value for $A(t)$ is also manifest in conditions where expenditures are highly reduced. For example, $A(t)$ remains constant on pre-synaptic terminals of cultured hippocampal neurons, despite the inhibition of action potential firing due to incubation with the Na^+ channel blocker Tetrodotoxin (TTX) (Rangaraju, Calloway, and Ryan 2014). Conversely, the same study showed that electrical stimulation of 10 Hz by one minute, concomitantly evokes ATP synthesis on pre-synaptic terminals, restoring $A(t)$ to basal levels (Rangaraju, Calloway, and Ryan 2014). From now on, we will call the basal value of $A(t)$ as the homeostatic availability of ATP (A_H).

Mechanisms accounting for the intrinsic control of A_H in neurons are less explored than in other cells. In the short term, there is a direct and fast effect of ATP molecules and their hydrolysis products, such as AMP/ADP, over the activity of different metabolic enzymes and ion channels. Indeed, neurons are largely known for being extremely, even disproportionately, sensitive to decreases in ATP sources, leading to a fast and significant inhibition of electrical activity (Ames 2000). For example, ATP-sensitive K^+ channels open during decreased ATP levels, hyperpolarizing the neuron to reduce endocytosis and the opening of voltage-sensitive Na^+ channels, thus preventing the ATP expenditure associated to both processes (Ben-Ari, Krnjević,

and Crépel 1990). On the other hand, it has been elegantly shown that action potential firing on presynaptic terminals' gate activity-driven ATP production is also required to allow proper synaptic transmission (Rangaraju, Calloway, and Ryan 2014). This close dependency of ATP levels to synaptic functioning has suggested that the affinity constant for ATP (e.g., K_m) of different presynaptic enzymes, might be close to certain resting ATP levels (Rangaraju, Calloway, and Ryan 2014). It is tempting to speculate that the fine-tuning of the affinity constant from key enzymes might be a broader phenomenon in neurons. In addition, it is known that calcium entry, which is transiently modified by electrical activity, is capable of orchestrating changes in ATP production. For example, synaptic stimulation with brief NMDA pulses, not only lead to pronounced increases of cytosolic calcium levels, but also of the mitochondrial matrix, whose ATP producing enzymes are known to be stimulated by calcium increases (Tarasov, Griffiths, and Rutter 2012; Lange et al. 2015). Indeed, transient increases of calcium levels are thought to be a sort of metabolic alarm which prepares cells to confront high energy demands by increasing ATP production by the mitochondria (Bhosale et al. 2015).

As a complementary mechanism, changes in the ATP and AMP ratio gate the activity of other metabolic sensors which, in turn, induce a specific signaling cascade for short and long-term adaptations of neuronal functions. For example, all known eukaryotic cells, including neurons, harbor energy sensors, such as AMP-activated protein kinase (AMPK), which tend to restore ATP concentration by decreasing anabolic and/or energy consuming processes, while increasing energy production through catabolism post energy challenges (Hardie 2011; Hardie, Ross, and Hawley 2012; Potter et al. 2010). AMPK is a highly evolutionary-conserved serine/threonine kinase enzyme that is activated either by diminished cellular energy (high AMP/ATP ratio) and/or through increased calcium (Hardie, Ross, and Hawley 2012). Recent evidence shows that in dorsal

root ganglion neurons—which express the transient receptor potential ankyrin 1 (TRPA1) channel for thermal and pain transduction—the AMPK activation results in a fast, down-regulation of membrane-associated TRPA1 and its channel activity within minutes, which is consistent with lowering energy expenditure by diminishing post-synaptic currents (S. Wang et al. 2018). Furthermore, it has been demonstrated that calcium overload, induced by an excitotoxic NMDA stimulus on cultured cortical neurons, can be reduced by the activation of AMPK, which would save the energy involved in the reversal of a Ca^{++} potential (Anilkumar et al. 2013). Interestingly, the actions of the catalytic subunit of neuronal AMPK also includes the inhibition of axon outgrowth and dendritic arborization during neuronal development, for adapting to metabolic stress mediated by the suppression of Akt and mTOR signaling pathways (Ramamurthy et al. 2014). This result suggests that AMPK may also operate in mediating structural synaptic changes during the activity of mature neurons, contributing to control energy expenditures in the long-term. Furthermore, it has been shown that the maintenance of long-term potentiation (LTP), which is energetically demanding, is dampened when AMPK activity is pharmacologically activated (mimicking a low ATP/AMP ratio), or conversely, LTP could be rescued when an ATP mimetic, ara-A, was added during an energy challenge. Thus, under low energy conditions, neuronal AMPK tends to inhibit changes on ionic gradients and reduce changes on cytoarchitecture, which can upregulate the value of $A(t)$, impacting plastic capacity as well.

Summarizing, each neuron has a certain amount of ATP available to them, which is constantly consumed by their different functions which can mostly be explained using ion gradient changes on axons and dendrites. At the same time, ATP production will compensate the ATP expenditure reaching an A_H that should remain constant until another specific synaptic challenge

arrives (Figure 2). In the next section, we will discuss the potential functional consequences of these adaptations in special cases of neuronal plasticity.

Revisiting neuronal plasticity under the perspective of energy constraints

A narrow window of ATP cytosolic concentration across time supports a bottom-up view of neuronal energy constraints, which may explain some well-described plastic adaptations from the literature. Measurements of glucose and oxygen consumption (reflecting energy consumption) have not distinguished between the contribution from glial and neuronal metabolism and the total energy expenditure attributed to one neuron (Hyder, Rothman, and Bennett 2013). Nonetheless, neurons would keep energy availability during the increment of energy demands, which include action potentials, potential propagation or dendritic depolarization, by dynamically sharing expenses with astrocytes glial cells (Hyder et al. 2006; Barros 2013). It is worth mentioning that energy management is partly performed by these latter cells (P. J. Magistretti and Allaman 2015; P. J. Magistretti 2006). Indeed, we must consider that ATP neural production is provided by the local pyruvate and glial lactate. Where a theoretical model aimed to explain brain energy availability from rat and human brains, it indirectly suggested that glial and neuron lactate sources may dynamically vary across different species and activity levels, with the condition of maintaining a rather constant energy production (Hyder, Rothman, and Bennett 2013).

We will follow a very simplified view of ATP metabolism characterized by two collections of processes: Those that produce ATP (e.g. from local pyruvate and glial lactate), and those that consume ATP (e.g. recovery of ion gradients, structural and functional synapse maintenance). We can formalize the effect of these processes on ATP concentration (A) by a simply differential equation:

$$\frac{\partial A}{\partial t} = P(t, A, \dots) - C(t, A, \dots) \text{(eq. 4)}$$

where, $P(t, A, \dots)$ is a function that represents the sum of all reaction rates that produce ATP (e.g. anaerobic and aerobic metabolism), whereas $C(t, A, \dots)$ is the sum of all reaction rates that consume ATP (e.g. membrane repolarization, structural and functional synapse maintenance). Both production (P) and consumption (C) rates are dynamic (they depend on time), but, more importantly, they depend on the levels of ATP available (A). Homeostasis will be achieved when production and consumption rates are equal, and the concentration of ATP is constant in time. We will represent the homeostatic concentration of ATP by A_H .

We can interpret the observations of relatively constant ATP concentrations in neurons, as reflecting the action of feedbacks that adjust ATP production (P) and consumption (C) rates, compensating deviations of ATP (A), such that neurons return to homeostatic ATP levels (A_H). We can expect that in case cells have an excess of ATP, they would respond by decreasing production or/and increasing consumption; and analogously, in case ATP levels are reduced, cells would respond by increasing production or/and decreasing consumption. We will call this regulation the neuron “energy management”, and summarize it mathematically using these equations:

$$\begin{cases} A > A_H \Rightarrow \frac{\partial P}{\partial t} \leq 0, \frac{\partial C}{\partial t} \geq 0 \\ A = A_H \Rightarrow \frac{\partial P}{\partial t} = 0, \frac{\partial C}{\partial t} = 0 \text{ (Eq. 5)} \\ A < A_H \Rightarrow \frac{\partial P}{\partial t} \geq 0, \frac{\partial C}{\partial t} \leq 0 \end{cases}$$

meaning that the differences between A with A_H determines whether ATP production (P) and consumption (C) processes increase, decrease, or maintain their rates over time. Note that we also consider the possibility that neurons may respond to energy challenges by adjusting production and consumption, but it must be at least one of those variables.

It is critical to notice that this formalization makes some important simplifications. First, we understand that in addition to ATP, the concentrations of ADP, AMP, and other energy resources do determine homeostasis and influence neuronal changes. ATP is a reasonable departure point, given its prevalence in metabolism and the evidence supporting its role in synaptic plasticity, and therefore, will be the main example of energy resource exploit in our argument. An additional reasonable assumption is that the magnitude of the change in reaction rates should correlate with the magnitude of the distance to homeostasis, which we have omitted from the equations but becomes relevant later in our argument for proposing experiments. We expect to expand towards a more detailed formalism in future work. Despite its simplicity, we think that our model can help to understand several previous studies and propose some experiments aimed at empirically evaluating the relation between energy resource availability and neural plasticity. We expect that simple phenomenological models, such as ours, will encourage both theoretical and experimental efforts, provided they can be readily falsified empirically, and be compared to theoretical derivations from biochemical first principles.

The tendency to set ATP at A_H might be compatible with homeostatic plastic changes that return a neuronal network to a basal firing rate, after prolonged periods of increased or decreased synaptic activity (homeostatic synaptic plasticity). Accordingly, it has been theoretically proposed that the excitability threshold of neurons might be a direct function of ATP (Huang et al. 2007). For example, the K_{ATP} channel-opener diazoxide decreases bursting and regular firing activity of the immature entorhinal cortex neurons (Lemak et al. 2014), which is coherent with a tight association of firing rates with contingent ATP concentration. Also, theoretically, neuronal circuits governed by purely Hebbian-plasticity rules are predicted to converge on instability, or to the opposite—total inactivity (Miller and MacKay 1994). One possible solution to enable neuronal

circuits to remain responsive is to limit the amount of synaptic strength per neuron. At least on excitatory synapses, this problem has shown to be solved by another form of synaptic plasticity termed “homeostatic synaptic plasticity,” and more specifically, “synaptic scaling” (Turrigiano et al. 1998; Turrigiano 2012). Synaptic scaling emerges to counteract the effects of long periods of increased or decreased synaptic activity in a multiplicative manner, thus allowing neurons to continuously reset the weight of their synaptic inputs to remain responsive to new environmental and cellular contexts. In the long term, the consequence of this regulation is that the firing rate of cortical cells in culture is sustained to an average set point (Turrigiano 2012). As far as we know, no attempt has been made to relate or prove the influence of neuronal energy load or $A(t)$ on this phenomenon.

Simple experiments on synaptic scaling could be performed to examine whether the tendency to reach A_H has a predictive value on the synaptic activity of neuronal networks. As shown in the seminal experiments of Turrigiano’s group, when a GABAergic inhibitor bicuculline (Bic) is acutely added to cultured neurons, it produces a significant increase in average firing rate. However, during 48 hours of stimulation, firing rates return to control values. On the other hand, neuronal firing rates can be completely abolished soon after adding either tetrodotoxin (TTX) or 6-cyano-7-nitroquinoxaline-2,3-dione (CNQX). Nevertheless, during the 48 hours of incubation, activity levels also return to a basal value (Turrigiano et al. 1998). The observed adaptive changes that operate in the long-term makes this experiment ideal for manipulating energy parameters.

Similar to the experiment of synaptic scaling performed by Turrigiano’s group, in our thought experiment, cultured neurons would be submitted to 48 hours of synaptic activity stimulation with Bic. During the stimulus, $A(t)$ will transiently decrease, inducing plastic changes on the network that will return ATP concentration to A_H , in a given time period (t_1) (Figure 3A).

In all conditions, $P(t)$ and $C(t)$ change accordingly with $A(t)$, following equations 4 and 5. However, if, during the stimulus, the neurons were pharmacologically modified to partially decrease ATP production (e.g., by blocking oxidative phosphorylation with sodium azide), expenditures ($C(t)$) are expected to be rapidly lowered and the time window required to return to the A_H value will be shortened (Figure 3B). Conversely, one could “enlarge” the theoretical value of A_H on cultured neurons by adding an ATP mimetic, such as ara-A. Here we assume that ara-A would cause inhibition of AMPK signaling, and that concentrations employed are low enough not to disturb the ATP synthesis. Thus, we propose that the neurons will take more time to return to A_H (Figure 3C). Under these three conditions, the firing rate of neurons should also be adapted to the same level as in the initial state, before stimulation, as well as ATP concentrations should return to the homeostatic value A_H .

From Molecules to Behavioral Homeostasis

In the previous sections, we have discussed how the energy homeostasis can affect synaptic plasticity in one neuron. Subsequently, this plasticity can impact other neurons that will trigger the same control systems to keep their A_H . Since energy demands are transferred through synapses, and synapses appear or disappear according to energy demands, a network homeostasis comes into play. In this section, we argue that energy constraints scale up a level of organization and how homeostasis in one level is affected by homeostasis in the others.

From neurons to a neural network.

The first level is single neuron homeostasis, which is the balance between $C(t)$ and $P(t)$ in single neurons. Importantly, as far as an action potential producing a post synaptic potential goes, it necessarily imposes an increment in $C(t)$ for the post synaptic neuron. As such, neurons manage their energy needs which also present an external demand from presynaptic neurons, and also

imposes an energy demand over the post synaptic neurons. The fact that a local increase in the $C(t)$ can produce a change in post synaptic neuron's $C(t)$ supports that energy management is also a neural population property, which we will name network homeostasis. The single neuron homeostasis is closely related to the network homeostasis through a two-way directional interaction, where the network structure imposes constraints on the range of possible homeostatic states that a neuron can achieve, which will, in turn, pose stress on the network through interactions with neighboring neurons. In the same way, these neurons will respond by modifying their synaptic weights (also known as network connectivity), the number of their synapses, or the location of their synapses, thus changing the functionality of the neural network structure (maybe even micro-anatomically). In any condition that causes an imbalance between $C(t)$ and $P(t)$, the neurons will tend to change. Since neurons activate each other through synapses, this means that the activity of the presynaptic neurons will induce metabolic work in the postsynaptic ones. In turn, a postsynaptic neuron will modulate its synaptic weight to couple the input from the presynaptic neuron to its own metabolic needs. This process will continue recursively until the neurons balance their $C(t)$ and $P(t)$, in which case the network would have reached homeostasis. Essentially, network homeostasis is driven by the needs of each neuron, as each of them will change in an attempt to reach their own A_H . Note that it is not necessary that every neuron should reach its own A_H , as the connectivity of activity within the network may not allow them to improve further. However, every single neuron must have enough $P(t)$ to devote towards maintenance processes required to stay alive. As such, network homeostasis becomes a neural population property.

Network homeostasis is tightly related to single neuron homeostasis; therefore, neural network homeostasis will be only achieved when several of the neurons that compose it individually can maintain themselves within homeostatic ranges (e.g. achieving A_H). It is known

that synaptic and dendrite pruning are a part of healthy development (Riccomagno and Kolodkin 2015; Huttenlocher 1990), which we could interpret as adjustments required to couple with the trade-off between maintaining the structure versus the energy spent in action and postsynaptic potentials. In worse cases where suboptimal conditions are imposed on a single neuron by the neural network homeostasis, we expect to find neuron death. This phenomenon is documented as a part of normal brain development in some species (Huttenlocher 1990), and also in pathological conditions (Pino et al. 2014; Kostrzewa and Segura-Aguilar 2003; Perry et al. 1991).

From neural networks to behavior.

Behavior can be broadly described as the set of actions performed by an organism, or anything that an organism does that involves movement and response to stimulation. These actions are adaptive when they increase the survival and reproduction probability. In a top-down interpretation of behavior, these actions are the result of the activation of the neuronal circuit that developed evolutionary to fulfill a need. Nonetheless, according to the Energy Homeostasis Principle, at the neural circuitry level, the actions performed by an organism are out of spatial and temporal context, since all the cells experiences are perturbations of the network activity. For a given neuron, the activity dynamics is dependent on the cumulative synaptic currents, regardless of the type of presynaptic cells that evoked them, or in the case of sensory receptors, the type of energy that is transduced. Similarly, it makes no difference for a given neuron to have neuron-neuron or neuro-muscular/endocrine synapses. Conversely, we can reinterpret behavior as the observed consequence of the homeostatic activity of an extended neural network (brain) which interacts with the environment. Sensory input and motor outputs can thus be viewed as “environmental synapses”, in which neurons receive synaptic activity from sensors, and send

activity through motor neurons and muscular tissue back to the environment. As such, we may argue that for the brain, environment can be seen as another neural network.

However, the interaction with the environment has an important difference that will impact the energy balance in the neuronal network. We can operationalize behavior in a neural system as a set of inputs and outputs that occur in a closed-loop manner. For instance, when we move our eyes, the brain is generating output activity, which in turn modifies the subsequent input to the retina. These dynamics occur for all sensory systems, where motor acts modify sensory inputs (Ahissar and Assa 2016). In this process, for each brain action, we should expect changes to occur in some sensory inputs. In other words, behavior can be seen as one of the ways in which the brain stimulates itself.

In principle, this closed-loop scheme would enable the brain to completely predict the sensory consequence of the motor actions. This processes of active inference is in line with previous proposals such as Friston's free energy principle and predictive coding (Schroeder et al. 2010; Friston 2010). It is crucial to note that Friston's Free Energy Principle used an informational approach where aspects such as temperature do not refer to the absolute temperature measured in an experimental setting. As such, the Energy Homeostasis Principle does not conceptually overlap with Friston's proposal; they can be considered as complementary. From a bottom-up view, Friston's proposal answers the epiphenomena, which can be related to information processing, rather omitting the underlying physiological constraints. However, in any of these proposals, there is an agreement that the brain is capable of predicting sensory input, and that it seems to reduce uncertainty as far as possible. In the case of Friston's proposal, it refers to the reduction of informational uncertainty, while in the Energy Homeostasis Principle, it refers to the reduction of energy sensorial input uncertainty.

Parsimoniously, the brain cannot fully predict the sensory inputs that occur after every motor act, because changes that are independent of the action of the organism also occur in the environment, and these changes may be critical to its survival. According to the Energy Homeostasis Principle, we should expect that neural networks will operate in a way that will favor the behavioral input activities within homeostatic energy ranges. If a given input is energetically too demanding, we should expect a change in behavior. If a given set of motor activity consistently produces an energy stressor input, it will cause synaptic changes in the brain, as the energy balance processes are spread over the neuronal network.

Sensory input represents the major energy challenge in the brain, while the motor output is the only way a neural network can modify this input. This way, the neuron in the network has the chance to regulate its $C(t)$, given the pressure representing sensory input. Neural networks will restrict the palette of behaviors that can be observed, while behavior will impose energy demands that the neural network will couple with by modifying behavior. For these reasons, behavior can also be considered as a phenomenon which affects the energy homeostasis in a two-way direction. Thus, an at least three-level nested system can be depicted, where each level will have a two-way interaction with each other (see Figure 4).

According to the Energy Homeostasis Principle, a key aspect to explaining adaptive behavior must reside within the brain's macro-structure and evolutionary mechanisms. Given a certain palette of sensory specializations, set of effectors, and brain structures, it will impose the range of possible energy disposal attractors that can emerge. For instance, the number of sensory cells, and their sensitivity to stimuli will determine the energy input imposed on neural tissue. The effectors given will determine the space in which a neural network must control that input. The macrostructure of the brain will then impose general restrictions on the neural network

homeostasis. For instance, the visual cortex mainly receives visual input, and the communication through brain regions is mainly achieved using major tracts. As such, the series of $C(t)$, imposed by one neuron on another, must follow the paths allowed by the macrostructure. This means, that neural network homeostasis will not only be a function of energy input provided by the sensory cells, and the chances to control it using effectors, but also of macro anatomical restrictions produced by genetic mechanisms (Gilbert. 2000a; Gilbert. 2000b).

Evolutionary pressures must act over all traits—genetic, physiological, and behavioral—of the organism (Darwin 1859). As such, evolutionary pressures have selected a sensory and effectors pallet, as well as a brain macrostructure. We conjecture that from the set of behaviors that satisfy the energy constrictions, behaviors that statistically improve the chances of surviving will be selected. We propose that the macro-anatomical structures impose a certain system of dynamic energy management among neural networks, which force the emergence of a certain set of energy attractors producing, in turn, a specific set of behaviors. It is important to consider that in animals that display a large set of behaviors, probably what is selected is the ability to learn. This concretely would mean that the selected is not a specific behavior, rather the flexibility with which an organism must adapt behaviorally during its own life.

Given this bottom-up view, we conclude the existence of behaviors strictly required for survival, and others which might present adaptive advantages given the specific context of the organism. In human primates, for example, there is a vast set of behaviors that are not strictly for the survival of the single individual in any context, yet they exist, such as leisure activities, those related to the production of art in its multiple forms, and even pathological behaviors which might directly impact the individual's health or survival. As far as these non-strictly adaptive behaviors do not impact the organism's life, they might be highly adaptive in certain contexts.

In any case, evolutionary mechanisms will shape the nervous system's macrostructure and behavior so that both are aligned in a way where solving the energy constraints of a single neuron and the neural network will lead to survival. If not, that macrostructure is expected to be lost, as those organisms will die. In fact, there is no need that all these levels work in alignment per se, rather they must only be aligned to survive. Evolutionary pressures will remove the organisms where the three level systems present goals that don't benefit each other. Since we can only see the animals that present all three level goals aligned, we have historically thought that neurons, neural networks, and organisms share the same goal. We proposed here that evolution shaped organisms, so when a neuron solves their needs, the behavior emerges as an epiphenomenon, which enables the organism to solve its needs, hence surviving.

Perspectives of Reinterpretation

In this section, we aim to contrast our proposal with evidence and highlight the corollary aspects which can open new avenues of research. Concretely, we will evaluate if we can reassess evidence, considering the Energy Homeostasis Principle. We think that this proposal is parsimonious as in spirit the rule is simple, what makes it complex is the wide range of interactions and properties that can emerge from the neural interaction constraint imposed by this rule. We believe that our proposal captures the essence of the concept of Braitenberg's vehicles (Braitenberg 1986)¹ and provides a plausible solution to the dynamics elaborated there. More importantly, this proposal serves new empiric avenues to study the working of the brain. For instance, plasticity has

¹ Braitenberg vehicle is a concept proposed by Valentino Braitenberg (Braitenberg, 1986), referring to a simple vehicle that has two sensors, each connected to an effector, i.e. a wheel, that provides the vehicle the ability to navigate in a given environment. The activation of the sensors can increase or decrease the speed of the respective wheel. In addition to the sensor-effector relationship, the physical configuration of the sensors and the wheels will determine the navigation "behavior" of the vehicle when the stimuli are present in the environment. One major conclusion of this concept is that complex behaviors arise from relatively simple properties (sensors, circuitry, and effectors) of the system in interaction with the environment.

always been thought to be the changes required to fix a given behavior. However, according to our proposal, plasticity is a process that takes place not only during learning but continuously, as a core component of the constant deployment of behavioral changes. As such, plasticity might not only be a determinant of behavior acquisition but a key aspect of ongoing behavior. In the following subsections, we will briefly discuss an example of the evidence interpreted considering the Energy Homeostasis Principle, and then discuss other theoretical and empirical avenues which can be reinterpreted based on this paradigm.

Hybrots: An analysis using the Energy Homeostasis Principle.

Let us discuss the energy principle proposed here in the context of a simple, in vivo network model. Empirically, one critical aspect of relating energy management with behavior is the major challenge of controlling sensory inputs. Most of experimental, in vivo animal models are not only sensible to acoustic, visual, physical, and chemical stimuli of the environment, but also to proprioceptive inputs, such as muscle contraction, tendon tension, blood acidification, hormone levels, among others. Strictly speaking, there is no way to properly control the sensory input of an animal in vivo, and the behavioral in vitro protocol seems to be unreal. Nonetheless, there are some protocols that can be considered as initial efforts of trying to build in vitro behavioral protocols. Specifically, some reports demonstrate that if we connect a neuronal culture of dissociated cortical or hippocampal neurons to an external device, coherent behavior can be obtained (Novellino et al. 2007; Tessadori et al. 2013).

Concretely, a system decodes the firing rate of the neurons in the culture and generates an output which is used to control two wheels of a vehicle. The vehicle has distance sensors. The sensor activity is coded to electrical pulses delivered back to the culture. The stimulation frequency is a function of the distance to an obstacle in front of the sensors. If the vehicle gets closer to an

obstacle, then the stimulation frequency increases. If the vehicle crashes into an obstacle, a stimulation (20 Hz for 2s) is delivered, which is previously known to trigger plasticity (le Feber, Stegenga, and Rutten 2010; Chiappalone, Massobrio, and Martinoia 2008; Madhavan, Chao, and Potter 2007; Tateno and Jimbo 1999; Jimbo, Tateno, and Robinson 1999). Leaving the vehicle in a circular maze with several obstacles under the operation of this protocol will cause it to “learn” to navigate, while avoiding impacts with obstacles (Tessadori et al. 2012). This model constitutes a protocol that enables studying the molecular, electrophysiological, and behavioral properties of neural processing simultaneously; above all, it allows the full control of the sensory input that this network will have.

Is this learning-like phenomenon compatible with the Energy Homeostasis Principle? When a single neuron is submitted to constant stimulation, we expect to have a 1-1 stimulation-action potential response. However, at a frequency stimulation as low as 10 Hz, the neurons will decay over time until they are unresponsive, or their response is importantly delayed (Gal et al. 2010). If interpreted through the Energy Homeostasis Principle we can hypothesize the following mechanism. First, we can postulate that at a frequency of 10 Hz or higher, stimulations become energetically stressful. As a response, neurons will respond with modifications in their synaptic weights in the short term, and with changes on their cytoarchitecture in the long term. Both processes will result in changes to the network structure. Each time the vehicle crashes, a stressful 20 Hz pulse will be delivered inducing plasticity. Functional restructuring is expected at each impact; leading to a random walk through different neural functional configurations, where each neuron will jump from state to state to minimize energy stress (see Figure 5). It is expected that those network configurations that decrease the effects of the sensory input will reduce energy stress due to impacting obstacles. But the best network configuration to the energy stress is indeed to

avoid it. Eventually, a network configuration will arise which will prevent the vehicle from crashing. Since no energy stress will be delivered as a sensory input with this configuration, this structure will seemingly stabilize on a configuration of homeostatic energy expenditure (Figure 5). We are aware that the above interpretation may oversimplify the actual mechanisms followed by the neurons. Neuronal changes are most likely not completely random and more complex regulations may be taking place. However, we want to point out that they can be sufficient to explain the phenomenology of the observations. As such, energy management, as a local rule, will impact the neural network structure as an emergent property, where, in turn, it will impact behavior. Critically, in this example, we have focused on sensorial input as an increment of neural activity. This might not always be the case (such as under sensorial isolation). Despite that, under this specific scenario, we propose that networks will minimize energy consumption; the goal is to arrive to A_H , not to the minimum possible energy expenditure. Therefore, if the sensorial input would move $A(t)$ below A_H , we would expect network modifications to increase expenses. In any case, the obtained behaviors must be at least compatible with the dynamic constraints imposed by $C(t)$, despite it being too high or low. In this example, behavior emerged to satisfy the energy needs of the neuron by means of $C(t)$. Finally, from all the vehicle movements, only a few, like avoiding the obstacle, might be interpreted as purposeful from an observer's point of view, the remaining ones may be considered a random trajectory. Importantly, this attribute is provided by the observer, as the neurons would only be managing their energetic demands. More research is required to evaluate what is happening with *behavior*, when the obstacles are out of the sensor's range along with the *learning* curve of the vehicle. Nonetheless, the Energy Homeostasis Principle allowed us to propose this hypothesis (Figure 5C), and it can be empirically addressed.

The Neuron Doctrine and the Energy Homeostasis Principle

Historically, the primary efforts to connect neuron activity with neural network dynamics and behavior was first proposed in 1888 (Barlow 1972; Bock 2013), which is referred to as “The Neuron Doctrine”, maintained and developed to this day (Sheppard 2016; Dehaene 2003; Moser and Moser 2015). In general terms, this theoretical proposal tries, in dual form, to solve the information coding and processing problem and has been supported by intracranial recordings, where abundant examples can be found (Moser and Moser 2015; Lettvin et al. 1959; Fairhall 2014). Specifically, neurons are expected to code for specific properties of the environment, where its activity is associated with the detection of specific stimuli. For instance, neurons in the primary visual cortex of mammals are selectively sensible to oriented bars (Taylor 1978; Hubel and Wiesel 1962), while in the Lateral Geniculate there is evidence supporting the existence of circular receptive fields representing portions of visual space (Reid and Shapley 1992). In these cases, neurons have receptive fields, which can be interpreted as a specific topologic relation of V1 with a certain retinal region; and therefore, with the image. Receptive fields with the same selectivity feature can also be found in the tactile, and auditory primary cortex, evidence of which is often interpreted as environmental stimuli being coded as a map in the brain (Ruth Clemo and Stein 1982; Penfield and Boldrey 1937; Penfield 1965). This classic evidence is theoretically in line with the recent hippocampus *where neurons*, which they represent a particular spot of a trained maze (Moser and Moser 2015).

Critically, most of the evidence supporting the neuron doctrine is associated to the neuron discharge rate. Since this discharge rate is a part of the $C(t)$, it necessarily means that most of the evidence supporting the neuron doctrine supports the Energy Homeostasis Principle as well. For this reason, it is plausible to consider that most of the neuron doctrine evidence is also evidence indicating how energy expenses of one neuron can be directly associated with behavior.

Furthermore, high discharge rates, as mentioned above, are expected to trigger plasticity mechanisms. Also, only a low percentage of neurons present high discharge rates (Olshausen and Field 2005), which should be expected under the Energy Homeostasis Principle scope. Moreover, due to the fact that high discharge rates might trigger changes in functional connectivity (synaptic weights), it should not be surprising that when presenting more complex visual scenes, classic receptive fields are no longer detectable (Fairhall 2014). We may consider that classic stimulation visual protocols impose an energy input, reflected in the high discharge rate, which needs to be managed. Importantly, in classic experiment, shadows are projected into a white screen, which from an Energy Homeostasis Principle perspective, the stimulus is everything by the shadow. Even more, plasticity might have a functional role in ongoing behavior rather than only stabilizing learned behaviors.

The neural doctrine paradigm has been closely related to information coding paradigms. The coding paradigms follow the same logic as the genetic code; the idea that information is universally coded using the same dictionary or codebook. In the case of genetics, what we call the genetic code, is an arrangement of how the sequence of nucleic acids informs specific amino acid sequences when assembling proteins. In the case of a neural code, the assumption is that environmental stimuli are translated into brain activity, which is then translated into motor output. More specifically, it is possible to map specific neuron activities to specific properties of the environment. For instance, the intensity of the stimulation can be mapped to the discharge rate of the sensory neurons (Gardner and Johnson 2013). The transduction of the stimuli is usually non-linear and sensitive to differences with previous stimulation rather than the raw value of the stimuli—Weber's law (Gardner and Johnson 2013). This adaptation law has a direct interpretation in the context of energy expenditure by neurons, as neurons coding raw stimuli would demand a

greater energy supply. Weber's law has also been extended to complex cognitive functions, such as quantity estimations (Dehaene 2003)—where discharge rates are used as the code of quantities for specific neurons—suggesting that energy saving may be a strategy widely used by neurons.

Of course, the discharge rate is far from being the only neural code proposed. Coupled with the complexity of sensory activity, temporal coding was proposed, where the exact temporal relationship between each neuron spike would be the key to understanding how environmental information is translated into brain activity (Friston 1997; Connor and Johnson 1992). Temporal coding is implicitly related to energy demands, as the time between action potentials trigger plasticity mechanisms, associated with one of the most expensive items of the neuron physiology—postsynaptic potential and plastic mechanisms (Attwell and Laughlin 2001). Another strategy is population coding (Georgopoulos, Schwartz, and Kettner 1986; Nicolelis 2003; Moxon and Foffani 2015). Population coding uses the activity of a large number of neurons, extracting properties such as discharge rate, timing, or other combined properties. As more neurons are included, more information is obtained, and we should expect that we will better predict, for instance, an arm movement. This approximation is good when the aim is to predict behavior but is not useful to understand how behavior emerges from neural activity. If reassessed using Energy Homeostasis Principle, we interpret that population coding works as it is a good assessment of neural network homeostasis, implicitly providing information about plastic changes and neural energy management. Up to some extent, all approaches have to do with when, how much, and which neurons are discharging, which in turn can be interpreted as when and how much energy is expended by individual neurons and the network.

When evaluating evidence related to a whole-brain approach, the neuron doctrine is mostly applied by associating the bold signals of brain regions to specific behaviors. Critically, the fMRI

signal is derived, to some extent, by the changes triggered through the glia to couple with the energy demands (Otsu et al. 2015). Therefore, we can interpret that energy management associated to glial function, is already associated directly with behavior. Moreover, it suggests that energy management can be mapped into networks associated to specific behaviors.

In general, the fMRI approach strongly resembles the serial symbolic programming paradigms, where a module can be homologized to a programming function, and the network would be the general architecture of the software. The loss of a programming function leads to the loss of a specific functionality of the software. This metaphor was addressed in classic literature (Hovland 1960; Searle 1980), suggesting that the brain processes information using a symbolic serial paradigm. As such, most of the neural correlates within the neurocognitive domain are interpreted as information processing, ranging from a strictly symbolic to a correlative information approach. However, using a bottom-up approach and the Energy Homeostasis Principle, those attributions are an observer's bias, as the one described in Braitenberg's vehicles (Braitenberg 1986). Behavioral functions of a neuron or the neural network would be the epiphenomena of neurons regulating their own homeostasis. In fact, as explained in the previous section, we can describe how the vehicle learns to avoid obstacles without using any informational, symbolic, or teleological explanations. Using this bottom-up approach, it is expected that an informational approach will be useful, as far as the neurons' and the neural network's needs are aligned with the organism's. However, it should be interpreted as an epiphenomenon of neural networks solving their own needs.

Reinterpreting Evidence Towards New Research Avenues.

As we have discussed above, energy management, although only implicitly considered in the literature, is a key feature of the nervous system. This necessarily means that most of our

current evidence can be reinterpreted in the light of the Energy Homeostasis Principle. We expect that this reinterpretation will trigger new ideas and strategies to understand the neural phenomena. As an example, we may try to explain the neuronal changes associated with learning processes, based on iconic paradigms such as the long-term potentiation (LTP) and depression (LTD) (Jia and Collingridge 2017; Nabavi et al. 2014). Both phenomena involve a large amount of energy expense where the ATP could be followed to understand the phenomena of plasticity as one of energy management. This is key, considering that even the Hebbian rules (Kempster, Gerstner, and Van Hemmen 1999), operates differently, depending on the neuron type (Abbott and Nelson 2000), highlighting the difficulties in predicting plasticity based on neural activity. At the same time, the calcium ion plays a critical signaling role within neural physiology, where we should ask if it might be a signal of energy expenditure. It is known that metabolic processes sense the ATP-AMP ratio (Ames 2000), however, they have not been studied in association to the plasticity phenomena.

Consequentially, we can assess energy management not solely from a molecular or electrophysiological perspective. For instance, can we consider inhibitory neurons as an adaptive feature to control brain energy expenditure? This is most intriguing if we consider that inhibitory neurons are key to increasing the neural circuits' controlling properties (e.g. negative feedback structures).

Simultaneously, the central nervous system is the only structure of the body which is actively isolated from the vascular system. It has its own system to maintain stable the neuron proximal environment. Moreover, astrocytes coordinate themselves through calcium waves, producing local changes in blood flow and hyperemia (increase on blood irrigation) (Otsu et al. 2015). The brain-blood barrier is not only a filter, but it works functionally to support the energy

demands of the neural networks. In fact, synapses are currently suggested to be tripartite structures (neuron-neuron and astrocyte) (D. D. Wang and Bordey 2008), where the glutamate-releasing excitatory synapses are proposed to control neurovascular coupling, and thus, brain energy during conditioning and behavior (Robinson and Jackson 2016). This would be a clear example of a neural activity involving external support for energy management.

Moreover, nerve cells exhibit a vast number of morphologies. It is currently unknown why some neurons display large, dendritic arborizations and short axons, while others present long axons and rather small dendritic arborizations. Similarly, there are varying basal discharge rates of activity. We think it is worth exploring whether the likelihood of particular morphologies and rate of activities are associated with energy constraints. For instance, can a neuron manage to maintain a long axon and at the same time a huge dendritic arborization where it must maintain a large number of dendritic spines? If we explore the evidence we already have, we are confident that new insights into neuron morphology will appear. Even more, if an unlikely neuron shape or size which is energetically more expensive presents itself; we should expect that those neurons would be more sensible to energy demands and may be more susceptible to neural death. In fact, Paul Bolam proposed that the reason behind Parkinson's is due to the dying out of dopaminergic neurons because of their huge size, which is very expensive in energy terms (Bolam and Pissadaki 2012; Pissadaki and Bolam 2013). It is most likely that many of these traits are genetically determined, however, energy constraints might limit the possible morphological variety. Furthermore, genetic determinants of neuron specializations may be triggered in response to the $C(t)$.

Finally, the Energy Homeostasis Principle paradigm, combined with a bottom-up view, allows us to reinterpret behavior in a much more flexible way. Animals display many behaviors

that are not intrinsically adaptive. Leisure activities are an evident example. Why the dog likes to go for the ball or follow a car? Why would we like to learn how to play the piano or to paint? Using a top-down approach would force us to interpret that evolution endorses us with a leisure activity brain module and that all behaviors are somehow beneficial. It seems more parsimonious to think that evolution restricted the system through macrostructure, so that survival-related brain functions will be selected and inherited. Above all, a wide set of diverse, seemingly useless behaviors can appear, without compromising organism survival or neural needs. Therefore, the only constraint for behavior is that the organisms must stay alive and that the sensory input can be successfully managed, in terms of its energy demand, by the neural networks and the neurons within them. As we already explained before, we think that in the cases of the vehicles controlled by neural cultures, the rules of the stimulation given is critical in understanding how they learn to avoid obstacles. From all the works that reported learning-like properties of *in vitro* dissociated cultures of neurons (Mulas and Massobrio 2010; Novellino et al. 2007; Tessadori et al. 2012), two main conclusions can be obtained: (1) learning-like properties are not dependent on a priori, highly intricate and sophisticated neural structures, and (2) there is at least one property which does not require a brain evolution argument to explain the emergence of behavior (but probably requires a neural tissue evolution argument). This would be particularly important in relation to behaviors that are not directly tied to survival.

Because of the space limitations, many of these latter considerations are laid out in a basic form. Nonetheless we stress that some of these speculations can be assessed by reviewing the current literature under the Energy Homeostasis Principle rationale. However, the proposal may encourage the development of falsifiable hypotheses, allowing for the testing of these intuitions through empiric work. Therefore, we propose the principle as a novel paradigm from which we

can reinterpret neuroscience experimental data, as well inspire the design of experiments which may connect biochemical knowledge to cognitive neuroscience.

Conflict of Interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Author Contributions

R.V., S.J., and A.L. developed the initial general argument. S.J. was the main contributor to the metabolic and biochemistry sections. A.L. was the main contributor to the single neuron's physiological sections. R.V. was the main contributor to the neural networks and behavior sections. P.M. contributed by articulating the different sections to build the manuscript. R.V. wrote the first draft. C.M. designed and built all the figures, based on all the authors' recommendations. C.M., A.C, and R.F. contributed with key observations that refined the general argument. All authors revised and edited the manuscript. R.V. and P.M. edited the final version.

Funding

This work was supported by the postdoctoral FONDECYT grant 3160403, given to R.V, the PhD scholarship by The Darwin Trust of Edinburgh, given to S.J., the regular FONDECYT grant 1151478, given to R.F., the BNI Puente postdoctoral fellowship, given to C.M., and the BNI Millennium Institute No. P09-015-F, given to A.C.

Acknowledgments

We would like to thank Dr. Felipe Barros for his valuable comments in an earlier version of the manuscript.

References

- Abbott, L. F., and Sacha B. Nelson. 2000. "Synaptic Plasticity: Taming the Beast." *Nature Neuroscience* 3 (11s): 1178–83. doi:10.1038/81453.
- Ahissar, Ehud, and Eldad Assa. 2016. "Perception as a Closed-Loop Convergence Process." *ELife* 5 (MAY2016): 1–26. doi:10.7554/eLife.12830.
- Ames, Adelbert. 2000. "CNS Energy Metabolism as Related to Function." *Brain Research. Brain Research Reviews* 34 (1–2): 42–68. doi:10.1016/S0165-0173(00)00038-2.
- Anilkumar, Ujval, Petronela Weisová, Heiko Düssmann, Caoimhín G. Concannon, Hans-Georg König, and Jochen H M Prehn. 2013. "AMP-Activated Protein Kinase (AMPK)-Induced Preconditioning in Primary Cortical Neurons Involves Activation of MCL-1." *Journal of Neurochemistry* 124 (5): 721–34. doi:10.1111/jnc.12108.
- Attwell, D, and S B Laughlin. 2001. "An Energy Budget for Signalling in the Grey Matter of the Brain." *Journal of Cerebral Blood Flow & Metabolism* 21: 1133–45.
- Baeza-Lehnert, Felipe, Aiman S Saab, Robin Gutiérrez, Valeria Larenas, Esteban Díaz, Melanie Horn, Miriam Vargas, Ladina Hösli, Jillian Stobart, and Johannes Hirrlinger. 2018. "Non-Canonical Control of Neuronal Energy Status by the Na⁺ Pump." *Cell Metabolism*. Elsevier.
- Barlow, H. B. 1972. "Single Units and Sensation: A Neuron Doctrine for Perceptual Psychology?" *Perception* 1 (4): 371–94.
- Barros, L. Felipe. 2013. "Metabolic Signaling by Lactate in the Brain." *Trends in Neurosciences*. doi:10.1016/j.tins.2013.04.002.
- Ben-Ari, Y., K. Krnjević, and V. Crépel. 1990. "Activators of ATP-Sensitive K⁺ Channels Reduce Anoxic Depolarization in CA3 Hippocampal Neurons." *Neuroscience*. doi:10.1016/0306-4522(90)90191-6.
- Berndt, Nikolaus, Oliver Kann, and Hermann Georg Holzhütter. 2015. "Physiology-Based Kinetic Modeling of Neuronal Energy Metabolism Unravels the Molecular Basis of NAD(P)H Fluorescence Transients." *Journal of Cerebral Blood Flow and Metabolism*. doi:10.1038/jcbfm.2015.70.
- Bhosale, Gauri, Jenny A. Sharpe, Stephanie Y. Sundier, and Michael R. Duchen. 2015. "Calcium Signaling as a Mediator of Cell Energy Demand and a Trigger to Cell Death." *Annals of the New York Academy of Sciences* 1350 (1): 107–16. doi:10.1111/nyas.12885.

- Bock, Ortwin. 2013. "Cajal, Golgi, Nansen, Schäfer and the Neuron Doctrine." *Endeavour* 37 (4): 228–34. doi:10.1016/j.endeavour.2013.06.006.
- Bolam, J. Paul, and Eleftheria K. Pissadaki. 2012. "Living on the Edge with Too Many Mouths to Feed: Why Dopamine Neurons Die." *Movement Disorders* 27 (12): 1478–83. doi:10.1002/mds.25135.
- Braitenberg, Valentino. 1986. *Vehicles: Experiments in Synthetic Psychology*. Edited by Valentino Braitenberg. *Bildung Und Erziehung*. First. Vol. 52. Cambridge: MIT Press. http://pale.org/taros2013.pdf%5Cnhttp://www.ecila.org/ecila_files/content/papers/ACEhom.pdf.
- Brown, Angus M., and Bruce R. Ransom. 2007. "Astrocyte Glycogen and Brain Energy Metabolism." *GLIA*. doi:10.1002/glia.20557.
- Brown, G C. 1992. "Control of Respiration and ATP Synthesis in Mammalian Mitochondria and Cells." *Biochemical Journal*. doi:10.1042/bj2840001.
- Cannon, William R. 2014. "Concepts, Challenges, and Successes in Modeling Thermodynamics of Metabolism." *Frontiers in Bioengineering and Biotechnology* 2 (November). doi:10.3389/fbioe.2014.00053.
- Cannon, William R, and Scott E Baker. 2017. "Non-Steady State Mass Action Dynamics without Rate Constants: Dynamics of Coupled Reactions Using Chemical Potentials." *Physical Biology* 14 (5): 055003. doi:10.1088/1478-3975/aa7d80.
- Chiappalone, Michela, Paolo Massobrio, and Sergio Martinoia. 2008. "Network Plasticity in Cortical Assemblies." *European Journal of Neuroscience* 28 (1): 221–37. doi:10.1111/j.1460-9568.2008.06259.x.
- Connolly, N. M. C., H. Dussmann, U. Anilkumar, H. J. Huber, and J. H. M. Prehn. 2014. "Single-Cell Imaging of Bioenergetic Responses to Neuronal Excitotoxicity and Oxygen and Glucose Deprivation." *Journal of Neuroscience*. doi:10.1523/JNEUROSCI.3127-13.2014.
- Connor, C E, and K O Johnson. 1992. "Neural Coding of Tactile Texture: Comparison of Spatial and Temporal Mechanisms for Roughness Perception." *The Journal of Neuroscience : The Official Journal of the Society for Neuroscience* 12 (9): 3414–26. doi:1527586.
- Crooks, Gavin E. 1999. "Entropy Production Fluctuation Theorem and the Nonequilibrium Work Relation for Free Energy Differences." *Physical Review E* 60 (3): 2721–26.

doi:10.1103/PhysRevE.60.2721.

Danos, Vincent, and Nicolas Oury. 2013. "Equilibrium and Termination II: The Case of Petri Nets." *Mathematical Structures in Computer Science* 23 (02): 290–307.

doi:10.1017/S0960129512000126.

Darwin, Charles. 1859. *On the Origin of Species*. Charles Darwin. Vol. 146. John Murray.

Dehaene, Stanislas. 2003. "The Neural Basis of the Weber-Fechner Law: A Logarithmic Mental Number Line." *Trends in Cognitive Sciences* 7 (4): 145–47. doi:10.1016/S1364-6613(03)00055-X.

Eguchi, Yutaka, Shigeomi Shimizu, and Yoshihide Tsujimoto. 1997. "Intracellular ATP Levels Determine Cell Death Fate by Apoptosis or Necrosis." *Cancer Research*. doi:10.1158/0008-5472.can-04-3300.

Fairhall, Adrienne. 2014. "The Receptive Field Is Dead. Long Live the Receptive Field?" *Current Opinion in Neurobiology* 25. Elsevier Ltd: 9–12. doi:10.1016/j.conb.2014.02.001.

Feber, Joost le, Jan Stegenga, and Wim L. C. Rutten. 2010. "The Effect of Slow Electrical Stimuli to Achieve Learning in Cultured Networks of Rat Cortical Neurons." Edited by Matthieu Louis. *PLoS ONE* 5 (1): e8871. doi:10.1371/journal.pone.0008871.

Fodor, J.A. 1983. *The Modularity of Mind*. 1st ed. MIT Press Cambridge, MA.

Friston, Karl. 1997. "Another Neural Code?" *NeuroImage* 5 (3): 213–20.

doi:10.1006/nimg.1997.0260.

———. 2002. "Beyond Phrenology: What Can Neuroimaging Tell Us about Distributed Circuitry?" *Annual Review of Neuroscience* 25 (January): 221–50.

doi:10.1146/annurev.neuro.25.112701.142846.

———. 2010. "The Free-Energy Principle: A Unified Brain Theory?" *Nature Reviews Neuroscience* 11 (2): 127–38. doi:10.1038/nrn2787.

Gal, A., D. Eytan, A. Wallach, M. Sandler, J. Schiller, and S. Marom. 2010. "Dynamics of Excitability over Extended Timescales in Cultured Cortical Neurons." *Journal of Neuroscience* 30 (48): 16332–42. doi:10.1523/JNEUROSCI.4859-10.2010.

Gardner, Esther P., and Kenneth O. Johnson. 2013. "Sensory Coding." In *Principles of Neural Science*, 158:449–74. doi:10.1126/science.158.3799.399.

Georgopoulos, Apostolos P., Andrew B. Schwartz, and Ronald E. Kettner. 1986. "Neuronal Population Coding of Movement Direction." *Science* 233 (4771): 1416–19.

doi:10.1126/science.3749885.

- Gilbert., Scott F. 2000a. “Neural Crest Cells and Axonal Specificity.” In *Developmental Biology*, 6th ed. Sunderland (MA): Sinauer Associates.
- . 2000b. “The Central Nervous System and the Epidermis.” In *Developmental Biology*, 6th ed. Sunderland (MA): Sinauer Associates.
- Hardie, D. Grahame. 2011. “Sensing of Energy and Nutrients by AMP-Activated Protein Kinase.” In *American Journal of Clinical Nutrition*. doi:10.3945/ajcn.110.001925.
- Hardie, D. Grahame, Fiona A. Ross, and Simon A. Hawley. 2012. “AMPK: A Nutrient and Energy Sensor That Maintains Energy Homeostasis.” *Nature Reviews Molecular Cell Biology*. doi:10.1038/nrm3311.
- Harris, Julia J., Renaud Jolivet, and David Attwell. 2012. “Synaptic Energy Use and Supply.” *Neuron* 75 (5): 762–77. doi:10.1016/j.neuron.2012.08.019.
- Herculano-Houzel, Suzana. 2011. “Scaling of Brain Metabolism with a Fixed Energy Budget per Neuron: Implications for Neuronal Activity, Plasticity and Evolution.” *PLoS ONE*. doi:10.1371/journal.pone.0017514.
- Hofmeyr, Jan-Hendrik S., and Athel Cornish-Bowden. 2000. “Regulating the Cellular Economy of Supply and Demand.” *FEBS Letters* 476 (1–2): 47–51. doi:10.1016/S0014-5793(00)01668-9.
- Horton, April C., and Michael D. Ehlers. 2003. “Neuronal Polarity and Trafficking.” *Neuron*. doi:10.1016/S0896-6273(03)00629-9.
- Hovland, Carl I. 1960. “Computer Simulation of Thinking.” *American Psychologist* 15 (11): 687–93. doi:10.1037/h0044165.
- Huang, Chin-Wei, Chao-Ching Huang, Juei-Tang Cheng, Jing-Jane Tsai, and Sheng-Nan Wu. 2007. “Glucose and Hippocampal Neuronal Excitability: Role of ATP-Sensitive Potassium Channels.” *Journal of Neuroscience Research* 85 (7): 1468–77. doi:10.1002/jnr.21284.
- Hubel, D. H., and T. N. Wiesel. 1962. “Receptive Fields, Binocular Interaction and Functional Architecture in the Cat’s Visual Cortex.” *The Journal of Physiology* 160 (January): 106–54. <http://www.ncbi.nlm.nih.gov/pubmed/14449617>.
- Huttenlocher, P R. 1990. “Morphometric Study of Human Cerebral Cortex Development.” *Neuropsychologia* 28 (6): 517–27. doi:10.1016/0028-3932(90)90031-I.
- Hyder, Fahmeed, Anant B. Patel, Albert Gjedde, Douglas L. Rothman, Kevin L. Behar, and

- Robert G. Shulman. 2006. "Neuronal–Glial Glucose Oxidation and Glutamatergic–GABAergic Function." *Journal of Cerebral Blood Flow & Metabolism* 26 (7): 865–77. doi:10.1038/sj.jcbfm.9600263.
- Hyder, Fahmeed, Douglas L. Rothman, and Maxwell R. Bennett. 2013. "Cortical Energy Demands of Signaling and Nonsignaling Components in Brain Are Conserved across Mammalian Species and Activity Levels." *Proceedings of the National Academy of Sciences*. doi:10.1073/pnas.1214912110.
- Jaynes, E. T. 1965. "Gibbs vs Boltzmann Entropies." *American Journal of Physics* 33 (5): 391–98. doi:10.1119/1.1971557.
- Jekabsons, Mika B., and David G. Nicholls. 2004. "In Situ Respiration and Bioenergetic Status of Mitochondria in Primary Cerebellar Granule Neuronal Cultures Exposed Continuously to Glutamate." *Journal of Biological Chemistry*. doi:10.1074/jbc.M401540200.
- Jia, Zhengping, and Graham L. Collingridge. 2017. "Learning about Synaptic GluA3." *Neuron* 93 (2): 254–56. doi:10.1016/j.neuron.2017.01.004.
- Jimbo, Y., T. Tateno, and H. P.C. Robinson. 1999. "Simultaneous Induction of Pathway-Specific Potentiation and Depression in Networks of Cortical Neurons." *Biophysical Journal* 76 (2): 670–78. doi:10.1016/S0006-3495(99)77234-6.
- Kempler, Richard, Wulfram Gerstner, and J.L. Van Hemmen. 1999. "Hebbian Learning and Spiking Neurons." *Physical Review E* 59 (4): 4498–4514. doi:10.1103/PhysRevE.59.4498.
- Kostrzewa, Richard M., and Juan Segura-Aguilar. 2003. "Novel Mechanisms and Approaches in the Study of Neurodegeneration and Neuroprotection. A Review." *Neurotoxicity Research* 5 (6): 375–83. doi:10.1007/BF03033166.
- Lange, Sofie C., Ulrike Winkler, Lars Andresen, Mathilde Byhrø, Helle S. Waagepetersen, Johannes Hirrlinger, and Lasse K. Bak. 2015. "Dynamic Changes in Cytosolic ATP Levels in Cultured Glutamatergic Neurons During NMDA-Induced Synaptic Activity Supported by Glucose or Lactate." *Neurochemical Research* 40 (12): 2517–26. doi:10.1007/s11064-015-1651-9.
- Laughlin, S. B. 2001. "Energy as a Constraint on the Coding and Processing of Sensory Information." *Current Opinion in Neurobiology* 11 (4): 475–80. doi:10.1016/S0959-4388(00)00237-3.
- Lemak, Maria S., Oksana Voloshanenko, Andreas Draguhn, and Alexei V. Egorov. 2014.

- “KATP Channels Modulate Intrinsic Firing Activity of Immature Entorhinal Cortex Layer III Neurons.” *Frontiers in Cellular Neuroscience* 8 (August): 255.
doi:10.3389/fncel.2014.00255.
- Lennie, Peter. 2003. “The Cost of Cortical Computation.” *Current Biology* 13 (6): 493–97.
doi:10.1016/S0960-9822(03)00135-0.
- Lettvin, JY, HR Maturana, W. S. McCulloch, and W. H. Pitts. 1959. “What the Frog’s Eye Tells the Frog’s Brain.” *Proceedings of The*.
http://ieeexplore.ieee.org/xpls/abs_all.jsp?arnumber=4065609.
- Lorenz, Dirk M, Alice Jeng, and Michael W Deem. 2011. “The Emergence of Modularity in Biological Systems.” *Physics of Life Reviews* 8 (2). Elsevier B.V.: 129–60.
doi:10.1016/j.plrev.2011.02.003.
- Madhavan, Radhika, Zenas C. Chao, and Steve M. Potter. 2007. “Plasticity of Recurring Spatiotemporal Activity Patterns in Cortical Networks.” *Physical Biology* 4 (3): 181–93.
doi:10.1088/1478-3975/4/3/005.
- Magistretti, P. J. 2006. “Neuron – Glia Metabolic Coupling and Plasticity,” 2304–11.
doi:10.1242/jeb.02208.
- Magistretti, P. J., and Igor Allaman. 2015. “A Cellular Perspective on Brain Energy Metabolism and Functional Imaging.” *Neuron* 86 (4). Elsevier Inc.: 883–901.
doi:10.1016/j.neuron.2015.03.035.
- Magistretti, Pierre J., and Igor Allaman. 2018. “Lactate in the Brain: From Metabolic End-Product to Signalling Molecule.” *Nature Reviews Neuroscience*. doi:10.1038/nrn.2018.19.
- Marcaida, Goizane, María Dolores Miñana, Santiago Grisolia, and Vicente Felipo. 1995. “Lack of Correlation between Glutamate-Induced Depletion of ATP and Neuronal Death in Primary Cultures of Cerebellum.” *Brain Research*. doi:10.1016/0006-8993(95)00703-S.
- Marcaida, Goizane, María-Dolores Miñana, Santiago Grisolia, and Vicente Felipo. 1997. “Determination of Intracellular ATP in Primary Cultures of Neurons.” *Brain Research Protocols* 1 (1): 75–78. doi:10.1016/S1385-299X(96)00009-8.
- Marsland, Robert, and Jeremy England. 2018. “Limits of Predictions in Thermodynamic Systems: A Review.” *Reports on Progress in Physics* 81 (1): 016601. doi:10.1088/1361-6633/aa9101.
- Miller, Kenneth D., and David J. C. MacKay. 1994. “The Role of Constraints in Hebbian

- Learning.” *Neural Computation*. doi:10.1162/neco.1994.6.1.100.
- Moser, May-Britt, and Edvard I. Moser. 2015. “Where Am I? Where Am I Going?” *Scientific American* 314 (1). Nature Publishing Group: 26–33. doi:10.1038/scientificamerican0116-26.
- Moxon, Karen A., and Guglielmo Foffani. 2015. “Brain-Machine Interfaces beyond Neuroprosthetics.” *Neuron* 86 (1). Elsevier Inc.: 55–67. doi:10.1016/j.neuron.2015.03.036.
- Mulas, Marcello, and Paolo Massobrio. 2010. “A Simulated Neuro-Robotic Environment for Bi-Directional Closed-Loop Experiments.” *Communication* 1 (3): 179–86. doi:10.2478/s13230-011-0004-x.
- Nabavi, Sadegh, Rocky Fox, Christophe D. Proulx, John Y. Lin, Roger Y. Tsien, and Roberto Malinow. 2014. “Engineering a Memory with LTD and LTP.” *Nature* 511 (7509). Nature Publishing Group: 348–52. doi:10.1038/nature13294.
- Nicholls, David. 2013. *Bioenergetics*. 4th edn. Academic Press.
- Nicolelis, Miguel A L. 2003. “Brain??Machine Interfaces to Restore Motor Function and Probe Neural Circuits.” *Nature Reviews Neuroscience* 4 (5): 417–22. doi:10.1038/nrn1105.
- Novellino, A, P. D’Angelo, L. Cozzi, M. Chiappalone, V. Sanguineti, and S. Martinoia. 2007. “Connecting Neurons to a Mobile Robot: An In Vitro Bidirectional Neural Interface.” *Computational Intelligence and Neuroscience* 2007 (January): 1–13. doi:10.1155/2007/12725.
- Olshausen, Bruno a, and David J Field. 2005. “How Close Are We to Understanding V1?” *Neural Computation* 17 (8): 1665–99. doi:10.1162/0899766054026639.
- Otsu, Yo, Kiri Couchman, Declan G Lyons, Mayeul Collot, Amit Agarwal, Jean-Maurice Mallet, Frank W Pfrieger, Dwight E Bergles, and Serge Charpak. 2015. “Calcium Dynamics in Astrocyte Processes during Neurovascular Coupling.” *Nature Neuroscience* 18 (2): 210–18. doi:10.1038/nn.3906.
- Ouldridge, Thomas E. 2018. “The Importance of Thermodynamics for Molecular Systems, and the Importance of Molecular Systems for Thermodynamics.” *Natural Computing* 17 (1): 3–29. doi:10.1007/s11047-017-9646-x.
- Penfield, Wilder. 1965. “Speech, Perception and the Uncommitted Cortex.” In *Brain and Conscious Experience*, 217–37. Heidelberg: Springer, Berlin.
- Penfield, Wilder, and Edwin Boldrey. 1937. “Somatic Motor and Sensory Representation in

- Man.” *Brain*, 389–443. doi:10.1093/brain/60.4.389.
- Perry, E. K., I. McKeith, P. Thompson, E. Marshall, J. Kerwin, S. Jabeen, J. A. Edwardson, et al. 1991. “Topography, Extent, and Clinical Relevance of Neurochemical Deficits in Dementia of Lewy Body Type, Parkinson’s Disease, and Alzheimer’s Disease.” *Annals of the New York Academy of Sciences* 640: 197–202. doi:10.1111/j.1749-6632.1991.tb00217.x.
- Pino, Oscar, Georgina Guflera, Juana Gómez-Benito, Antonia Najas-García, Silvia Rufián, and Emilio Rojo. 2014. “Neurodevelopment or Neurodegeneration: Review of Theories of Schizophrenia.” *Actas Espanolas de Psiquiatria* 42 (4): 185–95.
- Pissadaki, Eleftheria K., and J. Paul Bolam. 2013. “The Energy Cost of Action Potential Propagation in Dopamine Neurons: Clues to Susceptibility in Parkinson’s Disease.” *Frontiers in Computational Neuroscience* 7. doi:10.3389/fncom.2013.00013.
- Potter, Wyatt B., Kenneth J. O’Riordan, David Barnett, Susan M.K. Osting, Matthew Wagoner, Corinna Burger, and Avtar Roopra. 2010. “Metabolic Regulation of Neuronal Plasticity by the Energy Sensor AMPK.” *PLoS ONE*. doi:10.1371/journal.pone.0008996.
- Ramamurthy, S., E. Chang, Y. Cao, J. Zhu, and G.V. Ronnett. 2014. “AMPK Activation Regulates Neuronal Structure in Developing Hippocampal Neurons.” *Neuroscience* 259 (February): 13–24. doi:10.1016/j.neuroscience.2013.11.048.
- Rangaraju, Vidhya, Nathaniel Calloway, and Timothy A. Ryan. 2014. “Activity-Driven Local ATP Synthesis Is Required for Synaptic Function.” *Cell*. doi:10.1016/j.cell.2013.12.042.
- Reid, R. Clay, and Robert M. Shapley. 1992. “Spatial Structure of Cone Inputs to Receptive Fields in Primate Lateral Geniculate Nucleus.” *Nature* 356 (6371): 716–18. doi:10.1038/356716a0.
- Riccomagno, Martin M., and Alex L. Kolodkin. 2015. “Sculpting Neural Circuits by Axon and Dendrite Pruning.” *Annual Review of Cell and Developmental Biology* 31 (1): 779–805. doi:10.1146/annurev-cellbio-100913-013038.
- Robbins, P. 2010. “Modularity of Mind.” Edited by Edward N. Zalta. *The Stanford Encyclopedia of Philosophy*. <http://plato.stanford.edu/entries/modularity-mind/>.
- Robinson, Michael B., and Joshua G. Jackson. 2016. “Astroglial Glutamate Transporters Coordinate Excitatory Signaling and Brain Energetics.” *Neurochemistry International* 98 (September): 56–71. doi:10.1016/j.neuint.2016.03.014.
- Ruth Clemo, H., and Barry E. Stein. 1982. “Somatosensory Cortex: A ‘new’ Somatotopic

- Representation.” *Brain Research* 235 (1): 162–68. doi:10.1016/0006-8993(82)90207-4.
- Saez, Isabel, Jordi Duran, Christopher Sinadinós, Antoni Beltran, Oscar Yanes, María F. Tevy, Carlos Martínez-Pons, Marco Milán, and Joan J. Guinovart. 2014. “Neurons Have an Active Glycogen Metabolism That Contributes to Tolerance to Hypoxia.” *Journal of Cerebral Blood Flow and Metabolism*. doi:10.1038/jcbfm.2014.33.
- Schroeder, Charles E, Donald A Wilson, Thomas Radman, Helen Scharfman, and Peter Lakatos. 2010. “Dynamics of Active Sensing and Perceptual Selection.” *Current Opinion in Neurobiology* 20 (2): 172–76. doi:10.1016/j.conb.2010.02.010.
- Searle, John R. 1980. “MINDS , BRAINS , AND PROGRAMS” 3: 1–19.
- Sengupta, Biswa, A. Aldo Faisal, Simon B. Laughlin, and Jeremy E. Niven. 2013. “The Effect of Cell Size and Channel Density on Neuronal Information Encoding and Energy Efficiency.” *Journal of Cerebral Blood Flow and Metabolism*. doi:10.1038/jcbfm.2013.103.
- Sheppard, G.M. 2016. *Foundations of the Neuron Doctrine: 25th Anniversary Edition*. Oxford: Oxford Univ Press.
- Shulman, Robert G., Douglas L. Rothman, Kevin L. Behar, and Fahmeed Hyder. 2004. “Energetic Basis of Brain Activity: Implications for Neuroimaging.” *Trends in Neurosciences* 27 (8): 489–95. doi:10.1016/j.tins.2004.06.005.
- Silbey, Robert J., Robert A. Alberty, and Bawendi G. Mounji. 2004. *Physical Chemistry*. 4th edn. Hoboken, NJ: Wiley.
- Sokoloff, Louis. 2008. “The Physiological and Biochemical Bases of Functional Brain Imaging.” *Cognitive Neurodynamics*. doi:10.1007/s11571-007-9033-x.
- Tarasov, Andrei I., Elinor J. Griffiths, and Guy A. Rutter. 2012. “Regulation of ATP Production by Mitochondrial Ca²⁺.” *Cell Calcium*. doi:10.1016/j.ceca.2012.03.003.
- Tateno, Takashi, and Yasuhiko Jimbo. 1999. “Activity-Dependent Enhancement in the Reliability of Correlated Spike Timings in Cultured Cortical Neurons.” *Biological Cybernetics* 80 (1): 45–55. doi:10.1007/s004220050503.
- Taylor, Cahlker; 1978. “Theory of Edge Detection” 207 (1167): 187–217. doi:10.1098/rspa.1966.0046.
- Tessadori, Jacopo, Marta Bisio, Sergio Martinoia, and Michela Chiappalone. 2012. “Modular Neuronal Assemblies Embodied in a Closed-Loop Environment: Toward Future Integration of Brains and Machines.” *Frontiers in Neural Circuits* 6 (December): 1–16.

doi:10.3389/fncir.2012.00099.

- Tessadori, Jacopo, Daniele Venuta, Sreedhar S. Kumar, Marta Bisio, Valentina Pasquale, and Michela Chiappalone. 2013. “Embodied Neuronal Assemblies: A Closed-Loop Environment for Coding and Decoding Studies.” *International IEEE/EMBS Conference on Neural Engineering, NER*, 899–902. doi:10.1109/NER.2013.6696080.
- Toloe, J., R. Mollajew, S. Kügler, and S. L. Mironov. 2014. “Metabolic Differences in Hippocampal ‘Rett’ Neurons Revealed by ATP Imaging.” *Molecular and Cellular Neuroscience*. doi:10.1016/j.mcn.2013.12.008.
- Trevisiol, Andrea, Aiman S. Saab, Ulrike Winkler, Grit Marx, Hiromi Imamura, Wiebke Möbius, Kathrin Kusch, Klaus-Armin Nave, and Johannes Hirrlinger. 2017. “Monitoring ATP Dynamics in Electrically Active White Matter Tracts.” *ELife* 6 (April). doi:10.7554/eLife.24241.
- Turrigiano, Gina. 2012. “Homeostatic Synaptic Plasticity: Local and Global Mechanisms for Stabilizing Neuronal Function.” *Cold Spring Harbor Perspectives in Biology*. doi:10.1101/cshperspect.a005736.
- Turrigiano, Gina, K. R. Leslie, N. S. Desai, L. C. Rutherford, and S. B. Nelson. 1998. “Activity-Dependent Scaling of Quantal Amplitude in Neocortical Neurons.” *Nature*. doi:10.1038/36103.
- Wang, Doris D., and Angélique Bordey. 2008. “The Astrocyte Odyssey.” *Progress in Neurobiology* 86 (4): 342–67. doi:10.1016/j.pneurobio.2008.09.015.
- Wang, Shenglan, Kimiko Kobayashi, Yoko Kogure, Hiroki Yamanaka, Satoshi Yamamoto, Hideshi Yagi, Koichi Noguchi, and Yi Dai. 2018. “Negative Regulation of TRPA1 by AMPK in Primary Sensory Neurons as a Potential Mechanism of Painful Diabetic Neuropathy.” *Diabetes* 67 (1): 98–109. doi:10.2337/db17-0503.
- Weber, Bruno, and L. Felipe Barros. 2015. “The Astrocyte: Powerhouse and Recycling Center.” *Cold Spring Harbor Perspectives in Biology*, February, a020396. doi:10.1101/cshperspect.a020396.
- Yuste, Rafael. 2015. “From the Neuron Doctrine to Neural Networks.” *Nature Reviews Neuroscience* 16 (8). Nature Publishing Group: 487–97. doi:10.1038/nrn3962.

Figure Legends




Figure 1. Homeostasis requires a balance between ATP production (metabolism) and ATP consumption (synaptic activity): **A**) Production of energy molecules by metabolism supports neuron activity, in addition to cell maintenance processes, most notably by ATP ($A(t)$). Neuronal homeostasis depends on a balance between production and consumption of high-energy molecules. **B**) Synaptic activity has been estimated to amount for half of total ATP consumption (diagram redrawn from Harris, Jolivet, and Attwell 2012). For cellular homeostasis to be achieved, neurons must regulate their activity and metabolism in response to changing external perturbations. We propose that regulatory mechanisms, responsible for changes synaptic plasticity, reflect the requirement for maintaining a constant level of energy resources available for neurons to function.




Figure 2. Neuron activity induces changes in metabolism and synaptic activity to maintain homeostatic energy levels: Schematic illustration of neuronal response to an increase in activity. Above, in scale of colors, ATP concentration indicating from low to high A, with A_H at the center. Below, step 1 depicts a resting state with $A(t_0)=A_H$. In time step 2, the increase in activity would result in additional ATP consumption, and therefore reduce ATP concentration. We propose that neurons would respond to this perturbation by decreasing ATP consumption ($C(t)$) and increasing its ATP production ($P(t)$), as represented in time step 3. Finally, neurons would return to homeostatic levels of ATP (A_H), which is illustrated in time step 4. During all these steps, $A(t)$ is colored following the above ATP concentration color scale. Some factors that contribute to ATP consumption are as follows: Synaptic post-excitatory currents (excitatory inputs), firing rate (excitatory outputs), ion channel density, size of soma, dendrite arborization and axonal length, neurotransmitter recycling and release, and cytoarchitecture adaptations. Whereas, factors

contributing to ATP production are: glycolysis supported by glucose or glycogen breakdown, oxidative phosphorylation supported by neuronal pyruvate, astrocyte-derived lactate, and ketone bodies. Neurons possess regulatory mechanisms that sense current energy levels (here represented by a rotating wheel with a floating balloon), and control production and consumption, to maintain homeostasis. Examples of these control mechanisms would include ATP-Sensitive K⁺ channels, and AMPK signaling.




Figure 3. Hypothetical experiment to evaluate the influence of energy availability on synaptic scaling: The influence of energy availability in synaptic plasticity could be evaluated empirically. Here we propose a simple hypothetical scenario and explain what outcomes we predict under each condition. **(A)** Cultured neurons are stimulated with bicuculine (Bic) for 48 hours (denoted by a black bar), which presumably induces transient changes in ATP concentration ($A(t)$). Neurons respond by increasing ATP production ($P(t)$) and reducing ATP consumption ($C(t)$) by reducing its firing rate, which leads to the reestablishment of homeostatic ATP concentration within the time window enclosed by dotted lines. **(B)** During stimulation, cultured neurons can be

pharmacologically treated to partially inhibit oxidative phosphorylation (i.e. reducing ATP synthesis) denoted by a black arrow. Following the Energy Homeostasis Principle, we propose this will result in a further reduction of ATP concentration, which will induce an accelerated reduction in ATP consumption through the reduction of synapse firing rates. Thus, we propose that in this scenario the time window required to return to homeostasis is shortened. **(C)** Almost an identical protocol to **(B)** is applied to neurons; however, using an ATP mimetic molecule (denoted by the black arrow). We assume that ATP mimetic molecules would delay the reduction of synapsis firing rate by allosterically inhibiting AMPK, resulting in an enlarged period of energy consumption. Thus, we propose a wider time window before reaching A_H . All graphics follow equations 4 and 5, with the additional assumption that the magnitude of the adjustment of $P(t)$ and $C(t)$ are proportional to the distance of ATP levels $A(t)$ to homeostatic levels A_H . Results from these kinds of experiments could advance the understanding (and potentially manipulate) of the mechanisms responsible for neural adaptations, uncovering the relevant role of metabolic elements, such as metabolic sensors and/or nutrient availability.




Figure 4. Energy homeostasis: An integrated view of neurons, networks and behavior schematic of the three nested levels of the energy homeostasis system. Each level represents one unit and its proximal operations. Neuron refers to one neuron which must manage its energy consumption, which will trigger neuron-plastic changes. Many of these neurons will build a network, which has connectivity properties and population energy demands. Many networks working together will deploy behavior through motor actions, while also receive the sensory input. All levels present a two-way energy interaction between behavior, neural networks, and neurons. The figure intends

to present how sensory input can be considered an energy demand at network and neuron levels, while motor output through behavior gives room to control part of the sensory input.




Figure 5. Depiction of Braitenberg's vehicle behavior as a controlled platform to study learning and network energy adaptation: In this figure we hypothesize, based on the Energy Homeostasis Principle, how a hybot would learn. **(A)** Different behaviors observed at different learning levels of the Braitenberg's vehicle (top panel), and the corresponding sensory input (bottom panel) from an obstacle (circle centered to x symbol) is detected by a sensor (dashed area), while the vehicle explores the environment. **(B)** Learning curve of the vehicle when learning to avoid the obstacles, passing from frequent collisions to full avoidance behavior. **(C)** Network energy adaptation triggered by the sensory input while minimizing the energy stress. Plasticity induced periods are marked with arrows.