

Perspective

Beyond Biodiversity: Can Environmental DNA (eDNA) Cut it as a Population Genetics Tool?

Clare I.M. Adams^{1,*}, Michael Knapp¹, Neil J. Gemmell¹, Gert-Jan Jeunen¹, Michael Bunce², Miles Lamare³, and Helen R. Taylor¹

¹ Department of Anatomy, University of Otago, 270 Great King Street, Dunedin, Otago, 9016, New Zealand; clare.adams@postgrad.otago.ac.nz (C.I.M.A.); michael.knapp@otago.ac.nz (M.K.); neil.gemmell@otago.ac.nz (N.J.G.); gjeunen@gmail.com (G.J.); helen.taylor@otago.ac.nz (H.R.T.)

² Trace and Environmental DNA (TrEnd) Laboratory, Department of Environment and Agriculture, Curtin University, Bentley, Perth, WA 6102, Australia; michael.bunce@curtin.edu.au (M.B.)

³ Department of Marine Science, University of Otago, 310 Castle Street, Dunedin, Otago, 9016, New Zealand; miles.lamare@otago.ac.nz (M.L.)

* Correspondence: clare.adams@postgrad.otago.ac.nz; Tel.: +64-022-012-9075

Abstract: Population genetic data underpin many studies of behavioral, ecological, and evolutionary processes in wild populations and contribute to effective conservation management. However, collecting genetic samples can be challenging when working with endangered, invasive, or cryptic species. Environmental DNA (eDNA) offers a way to sample genetic material non-invasively without requiring visual observation. While eDNA has been trialed extensively as a biodiversity and biosecurity monitoring tool with a strong taxonomic focus, it has yet to be fully explored as a means for obtaining population genetic information. Here, we review current research that employs eDNA approaches for the study of populations. We outline challenges facing eDNA-based population genetic methodologies, and suggest avenues of research for future developments. We advocate that with further optimizations, this emergent field holds great potential as part of the population genetics toolkit.

Keywords: Sampling methodology; mtDNA; mitochondrial DNA; conservation; biodiversity; populations; genetics

1. Introduction

The study of population genetics describes changes in the genetic diversity of populations through evolutionary mechanisms such as natural selection, genetic drift, mutation, and gene flow, the totality of which have evolutionary consequences and can lead to speciation [e.g., 1–6]. The direction, magnitude and mechanisms of population connectivity can be examined by quantifying gene flow [e.g., 4–6]. Furthermore, genetic data can be used to estimate inbreeding, which has potential fitness consequences [e.g., 12–14]. Understanding the genetics of populations can help us understand the past distributions [10], present status [11] and future prospects of species [12]. Genetic data are, therefore, useful for informing conservation management strategies.

Population genetic research requires genetic sampling from organisms of interest. For example, blood and tissue samples have traditionally been sources of genetic material for large marine mammals, e.g., flipper clips from seal populations [13]. However, sampling tissue has negative effects on the target organism, such as: lethal sampling [14], causing harm or discomfort [15], disfigurement [16], or imposing stress [17]. Sampling can also be expensive, dangerous and time consuming for researchers [18–20]. For example, studying venomous reptiles may put researchers at risk of a bite [21] and traditional sampling efforts may be more costly than reagents required for eDNA analysis [22]. More recently, non-invasive sampling techniques have been developed to reduce harm and



expense [23]. In marine environments, for example, fecal plumes, respiratory blow, parasites, and shed skin have proven to be useful DNA sources [24–28]. Such indirect samples have been used to amplify mitochondrial DNA (mtDNA) and nuclear DNA (nuDNA) markers, allowing measurement of allelic diversity, kinship, sex ratios, abundance and effective population size, density, evolutionary significant units, mating systems, and patterns of dispersal in species that might otherwise be difficult to sample [26,27,29–36]. The advent of promising new non-invasive technologies, such as environmental DNA (eDNA), creates further options for non-invasive population genetic sampling.

Here we adopt the definition of eDNA from Taberlet et al. [37] defining it as a “*complex mixture of genomic DNA from many different organisms found in an environmental sample.*” However, we take a narrower view of “environmental sample”; defining eDNA as DNA taken from soil, water, or air, excluding samples from direct individual remains such as fecal matter, hair, or feeding traces [18,34,38]. Direct organismal traces may allow for assignment of genetic data to specific individuals [38–40]. From these non-invasive samples, DNA may be used to describe abundance [32–34], effective population size [41], density [35], diet [42], and sex of individuals [43,44]. Additionally, estimates of genetic diversity (e.g., inbreeding within a population) [45,46], gene flow [47,48], evolutionary significant units [49] and mating systems (e.g., multiple paternity) [29] have been obtained. However, eDNA samples from soil, water, or air are useful when individual traces cannot easily be identified and sampled. For example, eDNA can be used to target sites of suspected occupancy before intensive, invasive sampling effort is carried out in difficult-to-sample habitat [50]. With eDNA metabarcoding of environmental samples, the use of traditional methodology in tandem with eDNA methodology may reveal more biodiversity than either method alone [51].

Environmental DNA is already being used to improve biodiversity estimates (e.g., species-level taxa), when a complete snapshot of biodiversity is difficult to obtain via traditional methods (e.g., transects, video, trawl nets, trapping) [52–55]. The sensitivity of eDNA methods makes them ideal for detecting the presence of endangered, low-density invasive, transient, and cryptic species [56–61], especially when sampling efforts to detect low density species would be unmanageable [19]. Environmental DNA methods are not without challenges, but because eDNA offers sampling advantages such as sensitivity, simplicity and reduced harm, it is already being employed across a variety of biomes as a biodiversity monitoring tool, frequently in tandem with traditional methodologies [62–65]. In addition to biodiversity monitoring, eDNA could potentially be used for population genetic studies, mitigating some of the issues around invasive sampling. Population genetic eDNA is a newly emerging field that offers exciting prospects, but at present comes with challenges [39,66–70]. These challenges, discussed below, include but are not limited to: distinguishing individuals, allelic dropout and false alleles related to amplification error, difficulty in relating eDNA to biologically relevant abundances, and expanding to nuclear markers.

2. Environmental DNA for population genetics

The use of eDNA in population genetics is in its infancy, but a handful of notable studies demonstrate the potential of eDNA to obtain within-species population genetic data. A recent landmark study used eDNA from seawater to examine the mtDNA haplotype variation of whale sharks (*Rhincodon typus*), to assess shark population structure, and their trophic interactions [66]. This work demonstrated that eDNA sampling in areas of known whale shark presence could detect some haplotypes that were also known from traditional tissue samples. Data from eDNA suggested a recently identified Qatar population of whale sharks was more closely related to the Indo-Pacific whale shark aggregation compared to the Atlantic aggregation [66]. In addition to haplotypic variation, this study reported a positive eDNA copy number correlation between quantities of whale shark eDNA and quantities of eDNA from their mackerel tuna prey (*Euthynnus affinis*) over two years, likely reflective of food web interactions [66]. This research suggests avenues for further eDNA mtDNA haplotype studies, especially that of examining population-specific haplotypic variation.

More recently, seawater eDNA work on North American killer whale (*Orcinus orca*) explored whether eDNA-obtained haplotypes could be assigned to known haplotypic variation, and how long genetic material can be detected after known target animal presence [69]. Using droplet digital PCR

(ddPCR), a sensitive PCR technique for the absolute quantification of DNA copy number in a sample, killer whale eDNA was detected up to two hours after individuals were observed in the sampled area [69]. Sequencing of these eDNA samples also identified a haplotype belonging to the southern resident killer whale ecotype, a genetically and usually geographically distinct variation of this species [69], showing that eDNA can be used to detect the genetic diversity of a vagile species like killer whales even after they have passed through an area. It may be possible to detect ecotype movements and interactions through eDNA, genetically confirming visual identification.

Beyond assignment to known haplotypes, eDNA has been used to uncover previously unknown intraspecific genetic diversity. For example, previously unknown harbor porpoise (*Phocoena phocoena*) mtDNA control region haplotypes have been detected via seawater eDNA [67]. These haplotypes differed from established haplotypes by one base pair, a small difference, and added additional support to the overall population structure previously described among the different sampling locations for this species [67]. The population structure as resolved with the assistance of eDNA derived haplotypes, supports managing the southeast Alaskan harbor porpoise as two separate populations [67]. Importantly, this study featured positive controls that helped lay the groundwork for strict quality filtering of the sequence data. This strict quality filtering has also been emphasized in other studies as well [61,66–68,70].

In addition to single-species targets, eDNA metabarcoding primer sets have been used to identify multiple haplotypes across multiple species simultaneously [68]. For example, multiple arthropod mtDNA haplotype ratios have been identified using eDNA from freshwater streams [68]. After strict data filtering and only considering haplotypes found in multiple replicates, between 177 to 200 operational taxonomic units (OTUs), were recovered with the average number of detected haplotypes for each OTU ranging between 2.40 to 3.30 [68].

Separately, 16S rDNA eDNA metabarcoding primers described multiple intraspecific haplotypes from the *Lethrinus* (snapper) genus from seawater samples, also using strict sequence read filtering [70]. Both studies required primers that gave enough resolution to discern intraspecific haplotypes while also being broad enough to capture multiple species [68,70]. These studies help build a framework for determining multi-taxa haplotypic diversity with eDNA metabarcoding technology, but more investigation is needed for other environments and taxa [68,70]. Importantly, the error profiles around each amplicon need to be carefully considered to differentiate rare haplotypes from sequencing error [71].

2.1. Why use eDNA over direct sampling for population genetics?

Some of the species in the aforementioned studies are of conservation and management interest and can be difficult to sample [66,67,69]. Sampling protected species (e.g. endangered or threatened species) and species of cultural significance requires permitting processes that cost time and effort [72–74]. Data deficient species may not have a good sampling framework in place [75]. Species such as cetaceans may travel widely, while stranded individuals (often a major source of tissue samples) may not correspond to a population's usual distribution [67]. Although studies using eDNA methods for population genetics to date have been heavily focused on the marine biome, eDNA could also be useful for other difficult-to-sample species in various habitats. For example, eDNA has been used to detect the presence of endangered species such as the great crested newt (*Triturus cristatus*) in freshwater habitat [56,76,77]. Terrestrial endangered species, such as the Bornean orangutan (*Pongo pygmaeus*) and Sunda pangolin (*Manis javanica*), may leave eDNA traces in locations such as salt licks [78]. Given the elusiveness of some imperiled organisms, eDNA may be one of the best options for describing genetic variation in specific cases. Understanding the population dynamics of these species is important for effective conservation management, especially if they fall under a protected status [79].

Not only can eDNA be useful for managing imperiled species, but invasive species may also be suitable targets. For example, freshwater invasive species such as bullfrogs (*Lithobates catesbeianus*) and Asian carp (*Hypophthalmichthys* sp.), among many others, have already been successfully detected with eDNA methods [19,80]. In addition to presence, population genetic eDNA methods may be

useful for identifying source populations and pathways of invasion spread, as previous population genetic studies have done [19,81]. We anticipate that developing a widely usable population genetic eDNA toolkit could facilitate monitoring for these invasive species in the future. By sensitively detecting an invasion front and source population, managers may further be able to prevent damage and associated economic costs [82,83]. However, going beyond proof-of-concept studies to answer questions at greater depth and scales about gene flow, genetic drift, mutation, and natural selection will require some significant challenges to be overcome.

3. Challenges facing environmental DNA population genetics

Despite recent success and future potential, eDNA population genetic methods require fine tuning before they can be widely applied. Perhaps one of the biggest obstacles in transitioning from species detection to population genetics using eDNA is assigning sequences to individuals. Previous non-invasive genetic techniques for terrestrial organisms have been able to assume different individuals from separate scats [34,45] or feeding sites [39,40,84]. Additionally, DNA extracted from invertebrate stomachs (iDNA) has also been used to obtain host DNA [28,85–87]. [18]. Indeed, whale shark population structure was confirmed using iDNA from a whale shark copepod (*Pandarus rhincodonicus*) parasite [28]. However, capture of whole, discrete fecal matter beyond initial deposition, feeding traces, or parasites may be challenging in environmental sampling scenarios. Currently, environmental samples cannot parse apart individuals. The same haplotype from a sample could indicate one individual or multiple individuals with the same haplotype. However obtained, confidence in individually-sorted genetic data is important for downstream analysis (e.g. *adegenet* for R, BAPS, or DADA2 [88–90]) as population genetic theory concerns the existence and change of genetic variation within individuals between and among populations.

Even when sequences are obtained in sufficient quantities and can be assigned to individuals, many non-invasive genetic sampling techniques, including eDNA, suffer from amplification challenges. Allelic dropout (the loss of allelic variation during sequencing), and false alleles (the apparent presence of non-existent alleles in samples), are notable examples [38,46,91]. Polymerase errors may inflate marker variation, leading to false alleles [92]. As in any population genetic study, regardless of DNA source, missing or false alleles will bias genetic diversity estimates with implications for further analyses [93]. Degraded DNA samples yielding low quality and/or template can contribute to these challenges, requiring increased replicates, increased PCR cycles, or increased sequencing depth [94,95]. Recent eDNA sampling has shown that allelic variation can be missed in multiple markers depending on extraction and capture methodologies [40]. Therefore, testing methodologies and using multiple markers will be important for reliable, reproducible eDNA capture and amplification in eDNA population genetic studies.

Some techniques for mitigating quantity challenges for eDNA samples, such as PCR, are not without biases which may impact allelic abundance [95–97]. For example, PCR may exponentially amplify common DNA, decreasing the signals of rare genetic variation [98]. When using next-generation sequencing (NGS), there is a fine line between NGS error rates and low-quantity alleles when filtering and analyzing data [68]. The inclusion of positive controls and stringent sequence read filtering could act as a guideline to determine real genetic variation [67,99]. Multiple sampling replicates could be used to further verify the legitimacy of low-abundance genetic variation [67,70]. The same sequencing errors likely will not occur over multiple sampling replicates, and these reads could be considered, even if present at low copy number. Additionally, some previous studies have not found read depth for species biodiversity to correlate with individuals, biomass, or density [59,100–102]. However, a more recent study indicates strong evidence that flow-corrected eDNA abundance reflects salmon count abundance in two species across two years and life stages [103]. Relating allelic abundance to eDNA genetic variation is necessary to understand allelic frequency differences between and within populations. Correlation correction factors may help, but for multicellular species obtained from environmental sampling, variation in individual rates of DNA shedding may obscure true haplotype ratios [104–106].

Furthermore, the number of reads from eDNA-obtained genetic data for target species must be sufficient enough to be distinguishable from background noise to ensure confidence that true genetic variation has been obtained. If initial DNA copy numbers are insufficient (eg. <200 copies/L, [59,107]), PCR may be too stochastic to amplify genetic variation reliably, or may not amplify it at all. This issue of low copy number (LCN) has been addressed previously in the forensic and ancient DNA (aDNA) fields with technological advancements and modified protocols (e.g. qPCR, [108]; increased reagents, [109]) and awareness of LCN difficulties [110]. Sufficient sampling may increase changes of capturing target eDNA, and limits of detection have been explored for eDNA at the species level [111–113]. Ensuring confidence in genetic assignments with sufficient copy number is necessary to have confidence in data.

Increasing the length of sequences and marker types that can be obtained from eDNA would also be advantageous for answering population genetic questions. Thus far, eDNA research has focused primarily on short (100-400 bp) mtDNA fragments. One reason for choosing mtDNA is its high copy number availability in environmental samples [114–116]. Further, as eDNA degrades over time and space, short (100-400 bp) mtDNA sequences are more likely to be available in environmental samples [117–120], although larger (>16kb) mtDNA fragments have recently been obtained from eDNA [121]. Additionally, commonly barcoded regions (e.g. *cytochrome oxidase I*) in mtDNA are widely available in public databases for many metazoans [122,123]. However, it would be useful to expand into other, longer molecular markers as the exclusive use of short, maternally-inherited mtDNA fails to incorporate non-maternal genetic variation for most organisms [124,125]. For a deeper understanding of population structure, nuDNA data should be considered [126,127]. For instance, nuclear variation may show evidence of introgression when mtDNA does not, which could potentially be important to understanding how species and populations are interacting [eg. 82]. Phylogenetically, nuDNA may show different patterns of divergence within populations compared to mtDNA data [124,127,129]. Obtaining more markers is easier with nuDNA (e.g., 10's of microsatellites or 10000's of SNPs) and can also help provide the ability to distinguish individuals with similar genetic signatures [40,130–132]. For example, multi-allelic microsatellites could be used, and they have the added advantage of establishing a minimum number of individuals present in an eDNA sample. For diploid species, if three alleles are found in a mixture, at least two individuals must exist in that environment. Furthermore, markers that identify sex in genetically sex-determined species could give insight into the presence of sexes present in non-sexually dimorphic species.

Due to the aforementioned challenges, and the emergent nature of the eDNA field, population genetic-oriented eDNA sampling methodologies currently lag behind those of contemporary population genetic methods. Population genetic studies now regularly use thousands of single nucleotide polymorphisms (SNPs) for multiple individuals [133]. Additionally, whole genome sequencing (WGS) is increasingly used to examine population-level genetic variation because it offers very high genomic resolution for detecting selection and the genetic basis of phenotypes (Figure 1) [134]. Already, population-level data collected via RADSeq and other reduced-representation sequencing approaches can compare exomes or SNP-based genotypes to answer population-level questions (Figure 1) [135–138]. It is likely that eDNA population genetics will continue to lag behind traditional population genetics while the aforementioned challenges are being addressed. However, exciting technological developments suggest that at least some of these challenges can be overcome.




Figure 1. Figure 1. Possible avenues of analyzing environmental DNA (eDNA) for population information. After eDNA is shed into the water column by target organisms of interest, collected, and extracted, multiple analyses can be used to obtain population data. These analyses have diverse applications, highlighted in the top yellow row, but each technique also presents its own challenges, highlighted in the bottom yellow row.

4. Further developments and future technologies

For all eDNA population genetic studies, a robust sampling design will help minimize LCN DNA profiles. Choosing the correct substrate for sampling a target organism, such as the position in the water column with the highest concentration of DNA of the target organism, will maximize potential copy number [139]. On this front, collecting eDNA may become easier with automated sampling systems for standardized, long-term sampling. After this, carefully considering the extraction methodology, assay, and bioinformatic approaches best for the biological question asked is key, as method choice influences genotypes obtained [95,140].

Several technologies that could help address the difficulty of assigning genetic information from eDNA samples to individuals, are already available or in development. One way forward might include single-cell sequencing. As eDNA contains sloughed cells, collecting and sequencing whole, individual cells could facilitate extraction of individual-level genetic variation [141]. Cells could be collected using flow cytometry and microfluidic technology [142]. From there, extracted DNA from these cells can be sequenced using a variety of sequencing tools (e.g., Illumina, Nanopore) [143]. Advances in droplet microfluidics could potentially facilitate the capture and sequencing of DNA from single cells in parallel [144,145]. For instance, microbial single cell sequencing has already been used to characterize the genomes of marine bacterioplankton after an oil spill [146]. In general, single-cell sequencing technology remains very much the domain of medical and model research systems and there are still issues with biases [147,148]. Small quantities of DNA can result in allelic dropout and low sequencing coverage [147,149]. However, bioinformatics tools to address single-cell sequencing challenges are growing and could be developed for eDNA techniques to facilitate the use of single-cell sequencing for obtaining individual variation from eDNA samples.

Good sampling design, in tandem with technological developments, could also help mitigate issues around obtaining allelic ratios and abundance [140,150]. Increasing sequencing depth also helps to guard against allelic dropout [151]. Additionally, the inclusion of positive controls during sequencing or PCRs can assist data processing, serving as a benchmark for separating true haplotypic diversity from sequencing noise where NGS is used [67]. Droplet digital PCR (ddPCR) could also be

used to address allelic and copy number abundance in a sensitive way [152,153]. This technology is already being explored for abundance estimations of species using eDNA techniques and could potentially quantify allelic diversity [59,69,152,153]. Other ways to get at quantification of allelic copy number include a correction factor depending on cell type, size, life-history stage and species, which could be integrated into single-cell methodology [104]. Lastly, the more that is understood about the ecology of eDNA in a field setting, the better we can understand and model eDNA abundance [103,141].

Another challenge for population genetic eDNA methods will be to fully develop nuDNA markers for the species of interest using eDNA sampling techniques. Capture methodology, where “bait” molecules bind to target DNA of interest, is currently being explored for eDNA with some success [39,154]. This methodology can target specific SNPs in nuDNA and may “fish out” extracellular, degraded eDNA [154]. Beyond current technologies, future sequencing developments may also enable longer reads to be obtained without sacrificing quality. Longer reads at greater depth may help to string haplotypes together, increasing genomic coverage. Once environmental nuDNA has been captured, statistical programs can be developed specifically to analyze environmental genetic data. It may be possible to use rarefaction methods to estimate how well genetic diversity has been sampled in a location, as has been used in eDNA metabarcoding studies to estimate taxa obtained [155]. Potentially, models can be built for total variation detection at a specific locus [111]. This has already been done for taxon biodiversity [100,111,156]. Additionally, estimates of relatedness between populations could be based on genetic similarities [39,66,67]. If nuDNA eDNA techniques are developed for eDNA samples, it would broaden the range of biological questions that eDNA could be used to answer.

These exciting technological advances paired with environmental genetics techniques offer promising new ways to address broad evolutionary genetics questions without direct sampling. Amazing progress is occurring all the time in the use of environmentally derived genetic material. For example, despite the much more rapid degradation of RNA in the environment, even environmental RNA (eRNA) samples are now being used to provide a snapshot of the biological diversity and processes present at the community level [157–159]. Additionally, eRNA has identified foraminiferal taxa not found with eDNA techniques [160]. It is feasible to compare these eDNA and eRNA snapshots between populations to understand differential expression of genes in macrofauna at different sites [159]. Studying community-level genetics may also be a promising research avenue [161]. “Universal” primers can already target haplotypes of multiple species [68,70]. Once these technologies are further refined, eDNA and eRNA may become tools for examining selection at the community level [161]. By examining targeted gene expression with eRNA, documenting community responses to common pressures, such as anthropogenic climate shifts, may be possible. Furthermore, combining eDNA with aDNA techniques may give researchers a lens into past communities [162]. With new technology and human ingenuity driving unprecedented datasets, we can answer questions about the patterns and processes of how populations are changing in new ways that may be less harmful.

Table 1. Potential challenges facing environmental DNA population genetic research with suggestions for tools to help mitigate some of the challenges. While previous research has applied a number of these tools to these challenges (as cited), research still remains to be done to fully address how each tool can help mitigate each challenge, and to identify advantages and drawbacks of each. Note that not all tools and techniques apply to all challenges.

	Challenges					
	Abundance	Allelic drop-out	Bioinformatic challenges	Identifying individuals	Long-term datasets	Obtaining nuclear markers
Tools and Techniques	Automated sampling	Reduce spatial and temporal variance, especially for difficult-to-sample areas. Standardized deployment may help detect abundance changes in regular intervals. [163,164]		Document individual presence repeated through time.	Precise, standardized capture across time and space.	
	Baited capture methodology	Targeted capture of specific allelic variation [165].	Identification of specific SNPs in a population.		Capture of specific allelic variation across time.	Target of nuDNA, especially SNP markers [165].
	Droplet digital PCR	Absolute quantification of target molecules [59,152,166].	Sensitively amplify alleles, could reduce drop-out.	Provides absolute quantification of specific molecule abundance.	Perhaps amplify and quantify single-cell eDNA.	Sensitively quantify changes in target molecules over time.
	eRNA	Increase temporal resolution [157].	Increase temporal resolution of transcribed alleles.	Identify gene expression at the population level.	Detection of live individuals.	Examine expressed gene changes within a population or community [158,160]
	Increased sequencing depth		Increased detection chance of low copy number alleles.	Increased chance of detecting genetic diversity in replicates, perhaps can apply stricter filtering.	Increased confidence in detection of individuals, especially if using single-cell techniques.	More robust datasets may show change throughout time at a finer scale.

	Increased sequencing read length (nanopore, long-range mtDNA)	Capture of long STRs or mtDNA genomes [121]	Longer reads may help compile individuals' mtDNA [121].	Links SNPs to form genomic or mtDNA haplotypes.	May see recombination patterns through time.	Increased genomic coverage [121].
Single-cell sequencing	Approximation of cells per sample.	Identify cells from the same individual for sequencing to pool samples and decrease the probability of allelic dropout.	Identification of individuals allows for information to be analyzed with traditional population-genetics methodology.	Identify individuals based on cell genome.	Identify changes in individual presence.	Target of nuDNA, perhaps even able to aid in sequencing of whole genome.
"Universal" primers specific enough for intraspecific variation		Alleles of multiple species identified in the same sample with same primer [70].		Possibility to identify multiple individuals of multiple species if individuals can be sorted.	Multiple species targeted for community composition snapshots [68].	Possibility to target nuclear markers in multiple species in the same sample.

5. Conclusions

The need for non-invasive sampling, especially for low-density and cryptic populations, will continue to drive technological development. Environmental DNA population genetic methods have the potential to fill this need if developed and tested rigorously. Limited conservation funding and a need to minimize stress to wild animals, coupled with increasing molecular expertise and declining sequencing cost make eDNA an attractive genetic sampling tool [167]. Sampling with eDNA will likely never completely replace invasive sampling because organism capture allows for the collection of additional and important non-genetic data, such as size, age, maturity, isotopic measurements, enzymatic activity, and hormone activity [168–171]. Population genetic eDNA methods can be a useful complement to traditional, direct-sampling studies; eDNA is another tool in the population genetic toolbox. While there are challenges facing eDNA population genetic methods, the field can profit from - and build upon - the framework of extensive technological development already associated with the fields of aDNA and forensics. Technological advances have facilitated studies of disease, migration, and population dynamics of past organisms with aDNA techniques that were not feasible even 20 years ago [172–174]. Environmental DNA research faces similar technological challenges to aDNA associated with the recovery and sequencing of degraded, low copy number target DNA. Improving conservation management tools is of critical importance in light of global environmental change [175,176]. We predict that eDNA technology will have a key role to play in

providing rapid and broad scale insights into the population genetics of imperilled and difficult-to-sample species around the globe.

Author Contributions: Conceptualization, C.I.M.A. and H.T.; writing—original draft preparation, C.I.M.A.; writing—review and editing, C.I.M.A., M.B., N.J.G., G.J.J., M.K., M.L., and H.T.; supervision, N.J.G., G.J.J., M.K., and H.T.; funding acquisition, M.K.

Funding: This research was funded by NATIONAL SCIENCE CHALLENGE SUSTAINABLE SEAS INNOVATION FUND, grant number C01X1515.

Acknowledgments: Thank you to Mateus Detoni, Natalie Forsdick, and Denise Martini for providing draft comments.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Lenormand, T. Gene flow and the limits to natural selection. *Trends Ecol. Evol.* **2002**, *17*, 183–189, doi:10.1016/S0169-5347(02)02497-7.
2. McGuire, J.M.; Scribner, K.T.; Congdon, J.D. Spatial aspects of movements, mating patterns, and nest distributions influence gene flow among population subunits of Blanding's turtles (*Emydoidea blandingii*). *Conserv. Genet.* **2013**, *14*, 1029–1042, doi:10.1007/s10592-013-0493-8.
3. Epps, C.W.; Palsbøll, P.J.; Wehausen, J.D.; Roderick, G.K.; Ramey, R.R.; McCullough, D.R. Highways block gene flow and cause a rapid decline in genetic diversity of desert bighorn sheep. *Ecol. Lett.* **2005**, *8*, 1029–1038, doi:10.1111/j.1461-0248.2005.00804.x.
4. McGuigan, K. Studying phenotypic evolution using multivariate quantitative genetics. *Mol. Ecol.* **2006**, *15*, 883–896, doi:10.1111/j.1365-294X.2006.02809.x.
5. Lande, R.; Arnold, S.J. The measurement of selection on correlated characters. *Evolution* **1983**, *37*, 1210–1226, doi:10.1111/j.1558-5646.1983.tb00236.x.
6. Wu, C.-I. The genic view of the process of speciation. *J. Evol. Biol.* **2001**, *14*, 851–865, doi:10.1046/j.1420-9101.2001.00335.x.
7. Keller, L.F.; Waller, D.M. Inbreeding effects in wild populations. *Trends Ecol. Evol.* **2002**, *17*, 230–241, doi:10.1016/S0169-5347(02)02489-8.
8. Keller, L.F. Inbreeding and its fitness effects in an insular population of song sparrows (*Melospiza melodia*). *Evolution* **1998**, *52*, 240–250, doi:10.1111/j.1558-5646.1998.tb05157.x.
9. Mills, L.S.; Smouse, P.E. Demographic consequences of inbreeding in remnant populations. *Am. Nat.* **1994**, *144*, 412–431, doi:10.1086/285684.
10. Drummond, A.J.; Rambaut, A.; Shapiro, B.; Pybus, O.G. Bayesian coalescent inference of past population dynamics from molecular sequences. *Mol. Biol. Evol.* **2005**, *22*, 1185–1192, doi:10.1093/molbev/msi103.
11. Dussex, N.; Robertson, B.C.; Salis, A.T.; Kalinin, A.; Best, H.; Gemmell, N.J. Low spatial genetic differentiation associated with rapid recolonization in the New Zealand Fur Seal *Arctocephalus forsteri*. *J. Hered.* **2016**, *107*, 581–592, doi:10.1093/jhered/esw056.
12. Shaffer, M.L. Minimum population sizes for species conservation. *Bioscience* **1981**, *31*, 131–134, doi:10.2307/1308256.
13. Gemmell, N.J.; Burg, T.M.; Boyd, I.L.; Amos, W. Low reproductive success in territorial male Antarctic fur seals (*Arctocephalus gazella*) suggests the existence of alternative mating strategies. *Mol. Ecol.* **2001**, *10*, 451–60.
14. Erwin, T.L. Beetles and other insects of tropical forest canopies at Manaus, Brazil, sampled by insecticidal fogging. *Trop. Rain For. Ecol. Manag.* **1983**, *2*, 59–75.
15. Bearzi, G. First report of a common dolphin (*Delphinus delphis*) death following penetration of a biopsy dart. *J. Cetacean Res. Manag.* **2000**, *2*, 217–221.
16. Cassone, B.J.; Boulding, E.G. Genetic structure and phylogeography of the lined shore crab, *Pachygrapsus crassipes*, along the northeastern and western Pacific coasts. *Mar. Biol.* **2006**, *149*, 213–226, doi:10.1007/s00227-005-0197-9.
17. Romero, L.M.; Reed, J.M. Collecting baseline corticosterone samples in the field: is under 3 min good enough? *Comp. Biochem. Physiol. Part A Mol. Integr. Physiol.* **2005**, *140*, 73–79,

- doi:10.1016/J.CBPP.2004.11.004.
- 18. Rodgers, T.W.; Janečka, J.E. Applications and techniques for non-invasive faecal genetics research in felid conservation. *Eur. J. Wildl. Res.* **2013**, *59*, 1–16, doi:10.1007/s10344-012-0675-6.
 - 19. Jerde, C.L.; Mahon, A.R.; Chadderton, W.L.; Lodge, D.M. “Sight-unseen” detection of rare aquatic species using environmental DNA. *Conserv. Lett.* **2011**, *4*, 150–157, doi:10.1111/j.1755-263X.2010.00158.x.
 - 20. Greenbaum, E. A practical and safe method for the capture and restraint of venomous snakes in the field. *Bull. Chicago Herpetol. Soc.* **2003**, *38*, 1–3.
 - 21. Ivanyi, C.; Altimari, W. Venomous reptile bites in academic research. *Herpetol. Rev.* **2004**, *35*, 49–50.
 - 22. Sigsgaard, E.E.; Carl, H.; Møller, P.R.; Thomsen, P.F. Monitoring the near-extinct European weather loach in Denmark based on environmental DNA from water samples. *Biol. Conserv.* **2015**, *183*, 46–52, doi:10.1016/j.biocon.2014.11.023.
 - 23. Lugg, W.H.; Griffiths, J.; van Rooyen, A.R.; Weeks, A.R.; Tingley, R. Optimal survey designs for environmental DNA sampling. *Methods Ecol. Evol.* **2017**, *9*, 1049–1059, doi:10.1111/2041-210X.12951.
 - 24. Parsons, K.M.; Dallas, J.F.; Claridge, D.E.; Durban, J.W.; Balcomb, K.C.; Thompson, P.M.; Noble, L.R. Amplifying dolphin mitochondrial DNA from faecal plumes. *Mol. Ecol.* **1999**, *8*, 1766–1768, doi:10.1046/j.1365-294x.1999.00723.x.
 - 25. Frère, C.H.; Krzyszczyk, E.; Patterson, E.M.; Hunter, S.; Ginsburg, A.; Mann, J. Thar she blows! A novel method for DNA collection from cetacean blow. *PLoS One* **2010**, *5*, e12299, doi:10.1371/journal.pone.0012299.
 - 26. Pierszalowski, S.P.; Ferrari, M.; Glockner-Ferrari, D.; Mizroch, S.; Clapham, P.J.; Dickerson, B.R. Investigating the feasibility of using DNA from sloughed skin for individual identification and kinship analysis in humpback whales (*Megaptera novaeangliae*). *Mar. Mammal Sci.* **2013**, *29*, 533–541, doi:10.1111/j.1748-7692.2012.00580.x.
 - 27. Swanson, B.J.; Kelly, B.P.; Maddox, C.K.; Moran, J.R. Shed skin as a source of DNA for genotyping seals. *Mol. Ecol. Notes* **2006**, *6*, 1006–1009, doi:10.1111/j.1471-8286.2006.01473.x.
 - 28. Meekan, M.; Austin, C.M.; Tan, M.H.; Wei, N.-W. V.; Miller, A.; Pierce, S.J.; Rowat, D.; Stevens, G.; Davies, T.K.; Ponzo, A.; Gan, H.M. iDNA at Sea: Recovery of whale shark (*Rhincodon typus*) mitochondrial DNA sequences from the whale shark copepod (*Pandarus rhincodonicus*) confirms global population structure. *Front. Mar. Sci.* **2017**, *4*, 420, doi:10.3389/fmars.2017.00420.
 - 29. Gottelli, D.; Wang, J.; Bashir, S.; Durant, S.M. Genetic analysis reveals promiscuity among female cheetahs. *Proceedings. Biol. Sci.* **2007**, *274*, 1993–2001, doi:10.1098/rspb.2007.0502.
 - 30. Parsons, K.M.; Durban, J.W.; Claridge, D.E.; Balcomb, K.C.; Noble, L.R.; Thompson, P.M. Kinship as a basis for alliance formation between male bottlenose dolphins, *Tursiops truncatus*, in the Bahamas. *Anim. Behav.* **2003**, *66*, 185–194, doi:10.1006/ANBE.2003.2186.
 - 31. Parsons, K.M.; Durban, J.W.; Claridge, D.E.; Herzing, D.L.; Balcomb, K.C.; Noble, L.R. Population genetic structure of coastal bottlenose dolphins (*Tursiops truncatus*) in the northern Bahamas. *Mar. Mammal Sci.* **2006**, *22*, 276–298, doi:10.1111/j.1748-7692.2006.00019.x.
 - 32. Lukacs, P.M.; Burnham, K.P. Review of capture-recapture methods applicable to noninvasive genetic sampling. *Mol. Ecol.* **2005**, *14*, 3909–3919, doi:10.1111/j.1365-294X.2005.02717.x.
 - 33. Ruell, E.W.; Riley, S.P.D.; Douglas, M.R.; Pollinger, J.P.; Crooks, K.R. Estimating bobcat population sizes and densities in a fragmented urban landscape using noninvasive capture–recapture sampling. *J. Mammal.* **2009**, *90*, 129–135, doi:10.1644/07-MAMM-A-249.1.
 - 34. Mondol, S.; Ullas Karanth, K.; Samba Kumar, N.; Gopalaswamy, A.M.; Andheria, A.; Ramakrishnan, U. Evaluation of non-invasive genetic sampling methods for estimating tiger population size. *Biol. Conserv.* **2009**, *142*, 2350–2360, doi:10.1016/J.BIOCON.2009.05.014.
 - 35. Janečka, J.E.; Munkhtsog, B.; Jackson, R.M.; Naranbaatar, G.; Mallon, D.P.; Murphy, W.J. Comparison of noninvasive genetic and camera-trapping techniques for surveying snow leopards. *J. Mammal.* **2011**, *92*, 771–783, doi:10.1644/10-MAMM-A-036.1.
 - 36. Sharma, R.; Stuckas, H.; Bhaskar, R.; Khan, I.; Goyal, S.P.; Tiedemann, R. Genetically distinct population of Bengal tiger (*Panthera tigris tigris*) in Terai Arc Landscape (TAL) of India. *Mamm. Biol.* **2011**, *76*, 484–490, doi:10.1016/j.mambio.2010.10.005.
 - 37. Taberlet, P.; Bonin, A.; Zinger, L.; Coissac, E. *Environmental DNA: For Biodiversity Research and Monitoring*; Oxford University Press, 2018; ISBN 0198767226.
 - 38. Gillett, R.M.; White, B.N.; Rolland, R.M. Quantification and genetic profiling of DNA isolated from free-

- floating feces of the North Atlantic right whale (*Eubalaena glacialis*). *Mar. Mammal Sci.* **2008**, *24*, 341–355, doi:10.1111/j.1748-7692.2008.00192.x.
39. Aylward, M.L.; Sullivan, A.P.; Perry, G.H.; Johnson, S.E.; Louis, E.E. An environmental DNA sampling method for aye-ayes from their feeding traces. *Ecol. Evol.* **2018**, *8*, 9229–9240, doi:10.1002/ece3.4341.
40. Monge, O.; Dumas, D.; Baus, I. Environmental DNA from avian residual saliva in fruits and its potential uses in population genetics. *Conserv. Genet. Resour.* **2018**, *1*–9, doi:10.1007/s12686-018-1074-4.
41. Perez, I.; Geffen, E.; Mokady, O. Critically endangered Arabian leopards *Panthera pardus nimr* in Israel: estimating population parameters using molecular scatology. *Oryx* **2006**, *40*, 295–301, doi:10.1017/S0030605306000846.
42. Farrell, L.E.; Roman, J.; Sunquist, M.E. Dietary separation of sympatric carnivores identified by molecular analysis of scats. *Mol. Ecol.* **2000**, *9*, 1583–90.
43. Janečka, J.E.; Jackson, R.; Yuquang, Z.; Diqiang, L.; Munkhtsog, B.; Buckley-Beason, V.; Murphy, W.J. Population monitoring of snow leopards using noninvasive collection of scat samples: a pilot study. *Anim. Conserv.* **2008**, *11*, 401–411, doi:10.1111/j.1469-1795.2008.00195.x.
44. Borthakur, U.; Barman, R.D.; Das, C.; Basumatary, A.; Talukdar, A.; Ahmed, M.F.; Talukdar, B.K.; Bharali, R. Noninvasive genetic monitoring of tiger (*Panthera tigris tigris*) population of Orang National Park in the Brahmaputra floodplain, Assam, India. *Eur. J. Wildl. Res.* **2011**, *57*, 603–613, doi:10.1007/s10344-010-0471-0.
45. Russello, M.A.; Gladyshev, E.; Miquelle, D.; Caccone, A. Potential genetic consequences of a recent bottleneck in the Amur tiger of the Russian far east. *Conserv. Genet.* **2004**, *5*, 707–713, doi:10.1007/s10592-004-1860-2.
46. Smith, O.; Wang, J. When can noninvasive samples provide sufficient information in conservation genetics studies? *Mol. Ecol. Resour.* **2014**, *14*, 1011–1023, doi:10.1111/1755-0998.12250.
47. Ernest, H.B.; Boyce, W.M.; Bleich, V.C.; May, B.; Stiver, S.J.; Torres, S.G. Genetic structure of mountain lion (*Puma concolor*) populations in California. *Conserv. Genet.* **2003**, *4*, 353–366, doi:10.1023/A:1024069014911.
48. Haag, T.; Santos, A.S.; Sana, D.A.; Morato, R.G.; Cullen JR, L.; Cranshaw JR, P.G.; De Angelo, C.; Di Bitetti, M.S.; Salzano, F.M.; Eizirik, E. The effect of habitat fragmentation on the genetic structure of a top predator: loss of diversity and high differentiation among remnant populations of Atlantic Forest jaguars (*Panthera onca*). *Mol. Ecol.* **2010**, *19*, 4906–4921, doi:10.1111/j.1365-294X.2010.04856.x.
49. Sharma, R.; Stuckas, H.; Bhaskar, R.; Khan, I.; Goyal, S.P.; Tiedemann, R. Genetically distinct population of Bengal tiger (*Panthera tigris tigris*) in Terai Arc Landscape (TAL) of India. *Mamm. Biol.* **2011**, *76*, 484–490, doi:10.1016/j.mambio.2010.10.005.
50. Goricki, Š.; Stankovic, D.; Snoj, A.; Kuntner, M.; Jeffery, W.R.; Trontelj, P.; Pavic, M.; Grzelj, Z.; Naparus-Aljancic, M.; Aljancic, G. Environmental DNA in subterranean biology: Range extension and taxonomic implications for *Proteus*. *Sci. Rep.* **2017**, *7*, 45054, doi:10.1038/srep45054.
51. Stat, M.; John, J.; DiBattista, J.D.; Newman, S.J.; Bunce, M.; Harvey, E.S. Combined use of eDNA metabarcoding and video surveillance for the assessment of fish biodiversity. *Conserv. Biol.* **2018**, doi:10.1111/cobi.13183.
52. Boussarie, G.; Bakker, J.; Wangensteen, O.S.; Mariani, S.; Bonnin, L.; Juhel, J.-B.; Kiszka, J.J.; Kulwicki, M.; Manel, S.; Robbins, W.D.; Vigliola, L.; Mouillot, D. Environmental DNA illuminates the dark diversity of sharks. *Sci. Adv.* **2018**, *4*, eaap9661, doi:10.1126/sciadv.aap9661.
53. Thomsen, P.F.; Møller, P.R.; Sigsgaard, E.E.; Knudsen, S.W.; Jørgensen, O.A.; Willerslev, E. Environmental DNA from seawater samples correlate with trawl catches of Subarctic, deepwater fishes. *PLoS One* **2016**, *11*, e0165252, doi:10.1371/journal.pone.0165252.
54. Smart, A.S.; Weeks, A.R.; van Rooyen, A.R.; Moore, A.; McCarthy, M.A.; Tingley, R. Assessing the cost-efficiency of environmental DNA sampling. *Methods Ecol. Evol.* **2016**, *7*, 1291–1298, doi:10.1111/2041-210X.12598.
55. Port, J.A.; O'Donnell, J.L.; Romero-Maraccini, O.C.; Leary, P.R.; Litvin, S.Y.; Nickols, K.J.; Yamahara, K.M.; Kelly, R.P. Assessing vertebrate biodiversity in a kelp forest ecosystem using environmental DNA. *Mol. Ecol.* **2016**, *25*, 527–541, doi:10.1111/mec.13481.
56. Rees, H.C.; Bishop, K.; Middleditch, D.J.; Patmore, J.R.M.; Maddison, B.C.; Gough, K.C. The application of eDNA for monitoring of the great crested newt in the UK. *Ecol. Evol.* **2014**, *4*, 4023–4032, doi:10.1002/ece3.1272.

57. Ardura, A.; Zaiko, A.; Martinez, J.L.; Samulioviene, A.; Semenova, A.; Garcia-Vazquez, E. eDNA and specific primers for early detection of invasive species – A case study on the bivalve *Rangia cuneata*, currently spreading in Europe. *Mar. Environ. Res.* **2015**, *112*, 48–55, doi:10.1016/J.MARENVRRES.2015.09.013.
58. Smith, M.A.; Woodley, N.E.; Janzen, D.H.; Hallwachs, W.; Hebert, P.D.N. DNA barcodes reveal cryptic host-specificity within the presumed polyphagous members of a genus of parasitoid flies (Diptera: Tachinidae). *Proc. Natl. Acad. Sci. U. S. A.* **2006**, *103*, 3657–62, doi:10.1073/pnas.0511318103.
59. Uthicke, S.; Lamare, M.; Doyle, J.R. eDNA detection of corallivorous seastar (*Acanthaster cf. solaris*) outbreaks on the Great Barrier Reef using digital droplet PCR. *Coral Reefs* **2018**, *37*, 1229–1239.
60. Yamanaka, H.; Minamoto, T. The use of environmental DNA of fishes as an efficient method of determining habitat connectivity. *Ecol. Indic.* **2016**, *62*, 147–153, doi:10.1016/j.ecolind.2015.11.022.
61. Gargan, L.M.; Morato, T.; Pham, C.K.; Finarelli, J.A.; Carlsson, J.E.L.; Carlsson, J. Development of a sensitive detection method to survey pelagic biodiversity using eDNA and quantitative PCR: a case study of devil ray at seamounts. *Mar. Biol.* **2017**, *164*, 112, doi:10.1007/s00227-017-3141-x.
62. McKelvey, K.S.; Young, M.K.; Knotek, W.L.; Carim, K.J.; Wilcox, T.M.; Padgett-Stewart, T.M.; Schwartz, M.K. Sampling large geographic areas for rare species using environmental DNA: A study of bull trout *Salvelinus confluentus* occupancy in western Montana. *J. Fish Biol.* **2016**, *88*, 1215–1222, doi:10.1111/jfb.12863.
63. Wilcox, T.M.; McKelvey, K.S.; Young, M.K.; Sepulveda, A.J.; Shepard, B.B.; Jane, S.F.; Whiteley, A.R.; Lowe, W.H.; Schwartz, M.K. Understanding environmental DNA detection probabilities: A case study using a stream-dwelling char *Salvelinus fontinalis*. *Biol. Conserv.* **2016**, *194*, 209–216, doi:10.1016/j.biocon.2015.12.023.
64. Snyder, D.E. Invited overview: Conclusions from a review of electrofishing and its harmful effects on fish. *Rev. Fish Biol. Fish.* **2004**, *13*, 445–453.
65. Kim, P.; Kim, D.; Yoon, T.J.; Shin, S. Early detection of marine invasive species, *Bugula neritina* (Bryozoa: Cheilostomatida), using species-specific primers and environmental DNA analysis in Korea. *Mar. Environ. Res.* **2018**, *139*, 1–10.
66. Sigsgaard, E.E.; Nielsen, I.B.; Bach, S.S.; Lorenzen, E.D.; Robinson, D.P.; Knudsen, S.W.; Pedersen, M.W.; Jaidah, M. Al; Orlando, L.; Willerslev, E.; Møller, P.R.; Thomsen, P.F. Population characteristics of a large whale shark aggregation inferred from seawater environmental DNA. *Nat. Ecol. Evol.* **2016**, *1*, 0004, doi:10.1038/s41559-016-0004.
67. Parsons, K.M.; Everett, M.; Dahlheim, M.; Park, L. Water, water everywhere: environmental DNA can unlock population structure in elusive marine species. *R. Soc. Open Sci.* **2018**, *5*, 180537, doi:10.1098/rsos.180537.
68. Elbrecht, V.; Vamos, E.E.; Steinke, D.; Leese, F. Assessing intraspecific genetic diversity from community DNA metabarcoding data. *PeerJ* **2018**, *6*, e4644, doi:https://doi.org/10.7717/peerj.4644.
69. Baker, C.S.; Steel, D.; Niekirk, S.; Klinck, H. Environmental DNA (eDNA) from the wake of the whales: droplet digital PCR for detection and species identification. *Front. Mar. Sci.* **2018**, *5*, 133, doi:10.3389/fmars.2018.00133.
70. Stat, M.; Huggett, M.J.; Bernasconi, R.; DiBattista, J.D.; Berry, T.E.; Newman, S.J.; Harvey, E.S.; Bunce, M. Ecosystem biomonitoring with eDNA: metabarcoding across the tree of life in a tropical marine environment. *Sci. Rep.* **2017**, *7*, 12240, doi:10.1038/s41598-017-12501-5.
71. Murray, D.C.; Coghlan, M.L.; Bunce, M. From benchtop to desktop: Important considerations when designing amplicon sequencing workflows. *PLoS One* **2015**, *10*, e0124671, doi:10.1371/journal.pone.0124671.
72. Ruru, J.; O’Lyver, P.B.; Scott, N.; Edmunds, D. Reversing the decline in New Zealand’s biodiversity: empowering Māori within reformed conservation law. *Policy Q.* **2017**, *13*, doi:10.26686/pq.v13i2.4657.
73. Tipa, G.; Nelson, K. Eco-cultural restoration across multiple spatial scales: a New Zealand case study. *Water Hist.* **2017**, *9*, 87–106, doi:10.1007/s12685-016-0175-9.
74. Witmer, G.W. Wildlife population monitoring: some practical considerations. *Wildl. Res.* **2005**, *32*, 259, doi:10.1071/WR04003.
75. Bland, L.M.; Collen, B.; Orme, C.D.L.; Bielby, J. Predicting the conservation status of data-deficient species. *Conserv. Biol.* **2015**, *29*, 250–259, doi:10.1111/cobi.12372.
76. Rees, H.C.; Baker, C.A.; Gardner, D.S.; Maddison, B.C.; Gough, K.C. The detection of great crested newts

- year round via environmental DNA analysis. *BMC Res. Notes* **2017**, *10*, 327, doi:10.1186/s13104-017-2657-y.
77. Grillas, C.; Ugnon-Coussioz, E.; Priol, P.; Renet, J. Low occurrence of the great crested newt *Triturus cristatus* at the limits of its range: an alarming preliminary study. *Herpetol. Bull.* **2018**, *145*, 14–18.
78. Ishige, T.; Miya, M.; Ushio, M.; Sado, T.; Ushioda, M.; Maebashi, K.; Yonechi, R.; Lagan, P.; Matsubayashi, H. Tropical-forest mammals as detected by environmental DNA at natural saltlicks in Borneo. *Biol. Conserv.* **2017**, *210*, 281–285, doi:10.1016/J.BIOCON.2017.04.023.
79. Allendorf, F.W. Genetics and the conservation of natural populations: allozymes to genomes. *Mol. Ecol.* **2017**, *26*, 420–430, doi:10.1111/mec.13948.
80. Dejean, T.; Valentini, A.; Miquel, C.; Taberlet, P.; Bellemain, E.; Miaud, C. Improved detection of an alien invasive species through environmental DNA barcoding: The example of the American bullfrog *Lithobates catesbeianus*. *J. Appl. Ecol.* **2012**, *49*, 953–959, doi:10.1111/j.1365-2664.2012.02171.x.
81. Goldstien, S.J.; Schiel, D.R.; Gemmell, N.J. Regional connectivity and coastal expansion: Differentiating pre-border and post-border vectors for the invasive tunicate *Styela clava*. *Mol. Ecol.* **2010**, *19*, 874–885, doi:10.1111/j.1365-294X.2010.04527.x.
82. Blackman, R.C.; Häneling, B.; Lawson-Handley, L. The use of environmental DNA as an early warning tool in the detection of new freshwater invasive non-native species. *CAB Rev.* **2018**, *13*, 1–15, doi:10.1079/PAVSNNR201813010.
83. Pimentel, D.; Zuniga, R.; Morrison, D. Update on the environmental and economic costs associated with alien-invasive species in the United States. *Ecol. Econ.* **2005**, *52*, 273–288, doi:10.1016/j.ecolecon.2004.10.002.
84. Nichols, R. V.; Cromsigt, J.P.G.M.; Spong, G. DNA left on browsed twigs uncovers bite-scale resource use patterns in European ungulates. *Oecologia* **2015**, *178*, 275–284, doi:10.1007/s00442-014-3196-z.
85. Bohmann, K.; Schnell, I.B.; Gilbert, M.T.P. When bugs reveal biodiversity. *Mol. Ecol.* **2013**, *22*, 909–911.
86. Calvignac-Spencer, S.; Leendertz, F.H.; Gilbert, M.T.P.; Schubert, G. An invertebrate stomach's view on vertebrate ecology: Certain invertebrates could be used as "vertebrate samplers" and deliver DNA-based information on many aspects of vertebrate ecology. *BioEssays* **2013**, doi:10.1002/bies.201300060.
87. Schnell, I.B.; Sollmann, R.; Calvignac-Spencer, S.; Siddall, M.E.; Yu, D.W.; Wilting, A.; Gilbert, M.T.P. iDNA from terrestrial haematophagous leeches as a wildlife surveying and monitoring tool - prospects, pitfalls and avenues to be developed. *Front. Zool.* **2015**, *12*, 24, doi:10.1186/s12983-015-0115-z.
88. Jombart, T.; Kamvar, Z.N.; Collins, C.; Lustrik, R.; Beugn, M.-P.; Knaus, B.J.; Jombart, M.T. Package 'adegenet.' **2018**.
89. Corander, J.; Waldmann, P.; Marttinen, P.; Sillanpää, M.J. BAPS 2: Enhanced possibilities for the analysis of genetic population structure. *Bioinformatics* **2004**, *20*, 2363–2369, doi:10.1093/bioinformatics/bth250.
90. Callahan, B.J.; McMurdie, P.J.; Rosen, M.J.; Han, A.W.; Johnson, A.J.A.; Holmes, S.P. DADA2: High-resolution sample inference from Illumina amplicon data. *Nat. Methods* **2016**, *13*, 581–583, doi:10.1038/nmeth.3869.
91. McKelvey, K.S.; Schwartz, M.K. Genetic errors associated with population estimation using non-invasive molecular tagging: problems and new solutions. *J. Wildl. Manage.* **2004**, *68*, 439–448, doi:10.2193/0022-541x(2004)068[0439:geawpe]2.0.co;2.
92. Oliver, A.K.; Brown, S.P.; Callaham, M.A.; Jumpponen, A. Polymerase matters: Non-proofreading enzymes inflate fungal community richness estimates by up to 15%. *Fungal Ecol.* **2015**, *15*, 86–89, doi:10.1016/j.funeco.2015.03.003.
93. Cooke, T.F.; Yee, M.C.; Muzzio, M.; Sockell, A.; Bell, R.; Cornejo, O.E.; Kelley, J.L.; Bailliet, G.; Bravi, C.M.; Bustamante, C.D.; Kenny, E.E. GBTools: A statistical method for estimating allelic dropout in reduced representation sequencing data. *PLoS Genet.* **2016**, *12*, e1005631, doi:10.1371/journal.pgen.1005631.
94. Ficetola, G.F.; Taberlet, P.; Coissac, E. How to limit false positives in environmental DNA and metabarcoding? *Mol. Ecol. Resour.* **2016**, *16*, 604–607, doi:10.1111/1755-0998.12508.
95. Alberdi, A.; Aizpurua, O.; Gilbert, M.T.P.; Bohmann, K. Scrutinizing key steps for reliable metabarcoding of environmental samples. *Methods Ecol. Evol.* **2018**, *9*, 134–147, doi:10.1111/2041-210X.12849.
96. Fonseca, V.G. Pitfalls in relative abundance estimation using eDNA metabarcoding. *Mol. Ecol. Resour.* **2018**, *18*, 923–926, doi:10.1111/1755-0998.12902.

97. Fernandez, S.; Sandin, M.M.; Beaulieu, P.G.; Clusa, L.; Martinez, J.L.; Ardura, A.; García-Vázquez, E. Environmental DNA for freshwater fish monitoring: insights for conservation within a protected area. *PeerJ* **2018**, *6*, e4486, doi:10.7717/peerj.4486.
98. Fernández, S.; Rodríguez, S.; Martínez, J.L.; Borrell, Y.J.; Ardura, A.; García-Vázquez, E. Evaluating freshwater macroinvertebrates from eDNA metabarcoding: A river Nalón case study. *PLoS One* **2018**, *13*, e0201741, doi:10.1371/journal.pone.0201741.
99. Eisenstein, M. Microbiology: making the best of PCR bias. *Nat. Methods* **2018**, *15*, 317–320, doi:10.1038/nmeth.4683.
100. Wangensteen, O.S.; Cebrian, E.; Palacín, C.; Turon, X. Under the canopy: Community-wide effects of invasive algae in Marine Protected Areas revealed by metabarcoding. *Mar. Pollut. Bull.* **2018**, *127*, 54–66, doi:10.1016/J.MARPOLBUL.2017.11.033.
101. Klobucar, S.L.; Rodgers, T.W.; Buday, P. At the forefront: evidence of the applicability of using environmental DNA to quantify the abundance of fish populations in natural lentic waters with additional sampling considerations. *Can. J. Fish. Aquat. Sci.* **2017**, *74*, 2030–2034, doi:10.1139/cjfas-2017-0114.
102. Tillotson, M.D.; Kelly, R.P.; Duda, J.J.; Hoy, M.; Kralj, J.; Quinn, T.P. Concentrations of environmental DNA (eDNA) reflect spawning salmon abundance at fine spatial and temporal scales. *Biol. Conserv.* **2018**, *220*, 1–11, doi:10.1016/j.biocon.2018.01.030.
103. Levi, T.; Allen, J.M.; Bell, D.; Joyce, J.; Russell, J.R.; Tallmon, D.A.; Vulsteek, S.C.; Yang, C.; Yu, D.W. Environmental DNA for the enumeration and management of Pacific salmon. *Mol. Ecol. Resour.* **2018**, *39*, 4445, doi:doi.org/10.1111/1755-0998.12987.
104. Vasselon, V.; Bouchez, A.; Rimet, F.; Jacquet, S.; Trobajo, R.; Cornuel, M.; Tapolczai, K.; Domaizon, I. Avoiding quantification bias in metabarcoding: Application of a cell biovolume correction factor in diatom molecular biomonitoring. *Methods Ecol. Evol.* **2018**, *9*, 1060–1069, doi:10.1111/2041-210X.12960.
105. Klymus, K.E.; Richter, C.A.; Chapman, D.C.; Paukert, C. Quantification of eDNA shedding rates from invasive bighead carp *Hypophthalmichthys nobilis* and silver carp *Hypophthalmichthys molitrix*. *Biol. Conserv.* **2015**, *183*, 77–84, doi:10.1016/j.biocon.2014.11.020.
106. Dunn, N.; Priestley, V.; Herraiz, A.; Arnold, R.; Savolainen, V. Behavior and season affect crayfish detection and density inference using environmental DNA. *Ecol. Evol.* **2017**, *7*, 7777–7785, doi:10.1002/ece3.3316.
107. Collins, R.A.; Wangensteen, O.S.; O’Gorman, E.J.; Mariani, S.; Sims, D.W.; Genner, M.J. Persistence of environmental DNA in marine systems. *Commun. Biol.* **2018**, *1*, 185, doi:10.1038/s42003-018-0192-6.
108. Alonso, A.; Martín, P.; Albarán, C.; García, P.; García, O.; Fernández De Simón, L.; García-Hirschfeld, J.; Sancho, M.; De La Rúa, C.; Fernández-Piqueras, J. Real-time PCR designs to estimate nuclear and mitochondrial DNA copy number in forensic and ancient DNA studies. *Forensic Sci. Int.* **2004**, *139*, 141–149, doi:10.1016/j.forsciint.2003.10.008.
109. Johnson, B.M.; Kemp, B.M. Rescue PCR: Reagent-rich PCR recipe improves amplification of degraded DNA extracts. *J. Archaeol. Sci. Reports* **2017**, *11*, 683–694, doi:10.1016/j.jasrep.2017.01.006.
110. Gill, P.; Curran, J.; Elliot, K. A graphical simulation model of the entire DNA process associated with the analysis of short tandem repeat loci. *Nucleic Acids Res.* **2005**, *33*, 632–643, doi:10.1093/nar/gki205.
111. Furlan, E.M.; Gleeson, D.; Hardy, C.M.; Duncan, R.P. A framework for estimating the sensitivity of eDNA surveys. *Mol. Ecol. Resour.* **2016**, *16*, 641–654, doi:10.1111/1755-0998.12483.
112. Hunter, M.E.; Dorazio, R.M.; Butterfield, J.S.S.; Meigs-Friend, G.; Nico, L.G.; Ferrante, J.A. Detection limits of quantitative and digital PCR assays and their influence in presence-absence surveys of environmental DNA. *Mol. Ecol. Resour.* **2017**, *17*, 221–229, doi:10.1111/1755-0998.12619.
113. Xia, Z.; Zhan, A.; Gao, Y.; Zhang, L.; Haffner, G.D.; MacIsaac, H.J. Early detection of a highly invasive bivalve based on environmental DNA (eDNA). *Biol. Invasions* **2018**, *20*, 437–447, doi:10.1007/s10530-017-1545-7.
114. Bogenhagen, D.; Clayton, D.A. The number of mitochondrial deoxyribonucleic acid genomes in mouse L and human HeLa cells. Quantitative isolation of mitochondrial deoxyribonucleic acid. *J. Biol. Chem.* **1974**, *249*, 7991–5.
115. Taberlet, P.; Luikart, G.; Waits, L.P. Noninvasive genetic sampling: Look before you leap. *Trends Ecol. Evol.* **1999**, *14*, 323–327.
116. Thomsen, P.F.; Willerslev, E. Environmental DNA - An emerging tool in conservation for monitoring

- past and present biodiversity. *Biol. Conserv.* **2015**, *183*, 4–18, doi:10.1016/j.biocon.2014.11.019.
117. Turner, C.R.; Barnes, M.A.; Xu, C.C.Y.; Jones, S.E.; Jerde, C.L.; Lodge, D.M. Particle size distribution and optimal capture of aqueous microbial eDNA. *Methods Ecol. Evol.* **2014**, *5*, 676–684, doi:10.1111/2041-210X.12206.
118. Nukazawa, K.; Hamasuna, Y.; Suzuki, Y. Simulating the advection and degradation of the environmental DNA of common carp along a River. *Environ. Sci. Technol.* **2018**, *52*, 10562–10570, doi:10.1021/acs.est.8b02293.
119. Bylemans, J.; Furlan, E.M.; Gleeson, D.M.; Hardy, C.M.; Duncan, R.P. Does size matter? An experimental evaluation of the relative abundance and decay rates of aquatic eDNA. *Environ. Sci. Technol.* **2018**, *52*, 6408–6416, doi:10.1021/acs.est.8b01071.
120. Dejean, T.; Valentini, A.; Duparc, A.; Pellier-Cuit, S.; Pompanon, F.; Taberlet, P.; Miaud, C. Persistence of environmental DNA in freshwater ecosystems. *PLoS One* **2011**, *6*, e23398, doi:10.1371/journal.pone.0023398.
121. Deiner, K.; Renshaw, M.A.; Li, Y.; Olds, B.P.; Lodge, D.M.; Pfrender, M.E. Long-range PCR allows sequencing of mitochondrial genomes from environmental DNA. *Methods Ecol. Evol.* **2017**, *8*, 1888–1898, doi:10.1111/2041-210X.12836.
122. Ratnasingham, S.; Hebert, P.D.N. bold: The Barcode of Life Data System (<http://www.barcodinglife.org>). *Mol. Ecol. Notes* **2007**, *7*, 355–364, doi:10.1111/j.1471-8286.2007.01678.x.
123. Hebert, P.D.N.; Cywinski, A.; Ball, S.L.; DeWaard, J.R. Biological identifications through DNA barcodes. *Proc. R. Soc. London B Biol. Sci.* **2003**, *270*, 313–321.
124. Rubinoff, D.; Cameron, S.; Will, K. A genomic perspective on the shortcomings of mitochondrial DNA for “barcoding” identification. *J. Hered.* **2006**, *97*, 581–594, doi:10.1093/jhered/esl036.
125. Hurst, G.D.D.; Jiggins, F.M. Problems with mitochondrial DNA as a marker in population, phylogeographic and phylogenetic studies: The effects of inherited symbionts. *Proc. R. Soc. B Biol. Sci.* **2005**, *272*, 1525–1534, doi:10.1098/rspb.2005.3056.
126. Rubinoff, D. Utility of mitochondrial DNA barcodes in species conservation. *Conserv. Biol.* **2006**, *20*, 1026–33.
127. Carranza, J.; Salinas, M.; de Andrés, D.; Pérez-González, J. Iberian red deer: paraphyletic nature at mtDNA but nuclear markers support its genetic identity. *Ecol. Evol.* **2016**, *6*, 905–922, doi:10.1002/ece3.1836.
128. Mitchell, S.M.; Muehlbauer, L.K.; Freedberg, S. Nuclear introgression without mitochondrial introgression in two turtle species exhibiting sex-specific trophic differentiation. *Ecol. Evol.* **2016**, *6*, 3280–3288, doi:10.1002/ece3.2087.
129. Pratlong, M.; Rancurel, C.; Pontarotti, P.; Aurelle, D. Monophyly of Anthozoa (Cnidaria): why do nuclear and mitochondrial phylogenies disagree? *Zool. Scr.* **2017**, *46*, 363–371, doi:10.1111/zsc.12208.
130. Knapp, S.M.; Craig, B.A.; Waits, L.P. Incorporating genotyping error into non-invasive DNA-based mark-recapture population estimates. *J. Wildl. Manage.* **2009**, *73*, 598–604, doi:10.2193/2007-156.
131. Mills, L.S.; Citta, J.J.; Lair, K.P.; Schwartz, M.K.; Tallmon, D.A. Estimating animal abundance using noninvasive DNA sampling: promise and pitfalls. *Ecol. Appl.* **2000**, *10*, 283–294, doi:10.1890/1051-0761(2000)010[0283:EAUAND]2.0.CO;2.
132. Waits, L.P.; Luikart, G.; Taberlet, P. Estimating the probability of identity among genotypes in natural populations: cautions and guidelines. *Mol. Ecol.* **2001**, *10*, 249–256, doi:10.1046/j.1365-294X.2001.01185.x.
133. Morin, P.A.; Luikart, G.; Wayne, R.K. SNPs in ecology, evolution and conservation. *Trends Ecol. Evol.* **2004**, *19*, 208–216, doi:10.1016/j.tree.2004.01.009.
134. Fuentes-Pardo, A.P.; Ruzzante, D.E. Whole-genome sequencing approaches for conservation biology: Advantages, limitations and practical recommendations. *Mol. Ecol.* **2017**, *26*, 5369–5406, doi:10.1111/mec.14264.
135. Davey, J.W.; Blaxter, M.L. RADSeq: next-generation population genetics. *Brief. Funct. Genomics* **2010**, *9*, 416–423, doi:10.1093/bfgp/elq031.
136. Schlotterer, C.; Tobler, R.; Kofler, R.; Nolte, V. Sequencing pools of individuals — mining genome-wide polymorphism data without big funding. *Nat. Rev. Genet.* **2014**, *15*, 749–763, doi:10.1038/nrg3803.
137. Luca, F.; Hudson, R.R.; Witonsky, D.B.; Di Rienzo, A. A reduced representation approach to population genetic analyses and applications to human evolution. *Genome Res.* **2011**, *21*, 1087–98, doi:10.1101/gr.119792.110.

138. Choi, M.; Scholl, U.I.; Ji, W.; Liu, T.; Tikhonova, I.R.; Zumbo, P.; Nayir, A.; Bakkaloğlu, A.; Ozen, S.; Sanjad, S.; Nelson-Williams, C.; Farhi, A.; Mane, S.; Lifton, R.P. Genetic diagnosis by whole exome capture and massively parallel DNA sequencing. *Proc. Natl. Acad. Sci. U. S. A.* **2009**, *106*, 19096–101, doi:10.1073/pnas.0910672106.
139. Kozoli, A.; Stat, M.; Simpson, T.; Jarman, S.; DiBattista, J.D.; Harvey, E.S.; Marnane, M.; McDonald, J.; Bunce, M. Environmental DNA metabarcoding studies are critically affected by substrate selection. *Mol. Ecol. Resour.* **2018**, doi:10.1111/1755-0998.12971.
140. Goldberg, C.S.; Turner, C.R.; Deiner, K.; Klymus, K.E.; Thomsen, P.F.; Murphy, M.A.; Spear, S.F.; McKee, A.; Oyler-McCance, S.J.; Cornman, R.S.; Laramie, M.B.; Mahon, A.R.; Lance, R.F.; Pilliod, D.S.; Strickler, K.M.; Waits, L.P.; Fremier, A.K.; Takahara, T.; Herder, J.E.; Taberlet, P. Critical considerations for the application of environmental DNA methods to detect aquatic species. *Methods Ecol. Evol.* **2016**, *7*, 1299–1307, doi:10.1111/2041-210X.12595.
141. Barnes, M.A.; Turner, C.R. The ecology of environmental DNA and implications for conservation genetics. *Conserv. Genet.* **2016**, *17*, 1–17, doi:10.1007/s10592-015-0775-4.
142. Kalisky, T.; Quake, S.R. Single-cell genomics. *Nat. Methods* **2011**, *8*, 311–314, doi:10.1038/nmeth0411-311.
143. Macaulay, I.C.; Ponting, C.P.; Voet, T. Single-cell multiomics: multiple measurements from single cells. *Trends Genet.* **2017**, *33*, 155–168, doi:10.1016/j.tig.2016.12.003.
144. Lan, F.; Demaree, B.; Ahmed, N.; Abate, A.R. Single-cell genome sequencing at ultra-high-throughput with microfluidic droplet barcoding. *Nat. Biotechnol.* **2017**, *35*, 640–646, doi:10.1038/nbt.3880.
145. Pellegrino, M.; Sciambi, A.; Treusch, S.; Durruthy-Durruthy, R.; Gokhale, K.; Jacob, J.; Chen, T.X.; Geis, J.A.; Oldham, W.; Matthews, J.; Kantarjian, H.; Futreal, P.A.; Patel, K.; Jones, K.W.; Takahashi, K.; Eastburn, D.J. High-throughput single-cell DNA sequencing of acute myeloid leukemia tumors with droplet microfluidics. *Genome Res.* **2018**, *28*, 1345–1352, doi:10.1101/gr.232272.117.
146. Mason, O.U.; Hazen, T.C.; Borglin, S.; Chain, P.S.G.; Dubinsky, E.A.; Fortney, J.L.; Han, J.; Holman, H.-Y.N.; Hultman, J.; Lamendella, R.; Mackelprang, R.; Malfatti, S.; Tom, L.M.; Tringe, S.G.; Woyke, T.; Zhou, J.; Rubin, E.M.; Jansson, J.K. Metagenome, metatranscriptome and single-cell sequencing reveal microbial response to Deepwater Horizon oil spill. *ISME J.* **2012**, *6*, 1715–1727, doi:10.1038/ismej.2012.59.
147. Yuan, G.-C.; Cai, L.; Elowitz, M.; Enver, T.; Fan, G.; Guo, G.; Irizarry, R.; Kharchenko, P.; Kim, J.; Orkin, S.; Quackenbush, J.; Saadatpour, A.; Schroeder, T.; Shivdasani, R.; Tirosh, I. Challenges and emerging directions in single-cell analysis. *Genome Biol.* **2017**, *18*, 84, doi:10.1186/s13059-017-1218-y.
148. Tanay, A.; Regev, A. Scaling single-cell genomics from phenomenology to mechanism. *Nature* **2017**, *541*, 331–338, doi:10.1038/nature21350.
149. Zhang, C.-Z.; Adalsteinsson, V.A.; Francis, J.; Cornils, H.; Jung, J.; Maire, C.; Ligon, K.L.; Meyerson, M.; Love, J.C. Calibrating genomic and allelic coverage bias in single-cell sequencing. *Nat. Commun.* **2015**, *6*, 6822, doi:10.1038/ncomms7822.
150. Ficetola, G.F.; Pansu, J.; Bonin, A.; Coissac, E.; Giguet-Covex, C.; De Barba, M.; Gielly, L.; Lopes, C.M.; Boyer, F.; Pompanon, F.; Rayé, G.; Taberlet, P. Replication levels, false presences and the estimation of the presence/absence from eDNA metabarcoding data. *Mol. Ecol. Resour.* **2015**, *15*, 543–556, doi:10.1111/1755-0998.12338.
151. Creer, S.; Deiner, K.; Frey, S.; Porazinska, D.; Taberlet, P.; Thomas, W.K.; Potter, C.; Bik, H.M. The ecologist's field guide to sequence-based identification of biodiversity. *Methods Ecol. Evol.* **2016**, *7*, 1008–1018, doi:10.1111/2041-210X.12574.
152. Nathan, L.M.; Simmons, M.; Wegleitner, B.J.; Jerde, C.L.; Mahon, A.R. Quantifying environmental DNA signals for aquatic invasive species across multiple detection platforms. *Environ. Sci. Technol.* **2014**, *48*, 12800–12806, doi:10.1021/es5034052.
153. Doi, H.; Uchii, K.; Takahara, T.; Matsuhashi, S.; Yamanaka, H.; Minamoto, T. Use of droplet digital PCR for estimation of fish abundance and biomass in environmental DNA surveys. *PLoS One* **2015**, *10*, e0122763, doi:10.1371/journal.pone.0122763.
154. Carpenter, M.L.; Buenrostro, J.D.; Valdiosera, C.; Schroeder, H.; Allentoft, M.E.; Sikora, M.; Rasmussen, M.; Gravel, S.; Guillén, S.; Nekhrizov, G.; Leshtakov, K.; Dimitrova, D.; Theodossiev, N.; Pettener, D.; Luiselli, D.; Sandoval, K.; Moreno-Estrada, A.; Li, Y.; Wang, J.; Gilbert, M.T.P.; Willerslev, E.; Greenleaf, W.J.; Bustamante, C.D. Pulling out the 1%: whole-genome capture for the targeted enrichment of ancient dna sequencing libraries. *Am. J. Hum. Genet.* **2013**, *93*, 852–864, doi:10.1016/j.ajhg.2013.10.002.
155. Günther, B.; Knebelberger, T.; Neumann, H.; Laakmann, S.; Martínez Arbizu, P. Metabarcoding of

- marine environmental DNA based on mitochondrial and nuclear genes. *Sci. Rep.* **2018**, *8*, 14822, doi:10.1038/s41598-018-32917-x.
156. Korpelainen, H.; Pietiläinen, M. Biodiversity of pollen in indoor air samples as revealed by DNA metabarcoding. *Nord. J. Bot.* **2017**, *35*, 602–608, doi:10.1111/njb.01623.
157. Pochon, X.; Zaiko, A.; Fletcher, L.M.; Laroche, O.; Wood, S.A. Wanted dead or alive? Using metabarcoding of environmental DNA and RNA to distinguish living assemblages for biosecurity applications. *PLoS One* **2017**, *12*, e0187636, doi:10.1371/journal.pone.0187636.
158. Zaiko, A.; Pochon, X.; Garcia-Vazquez, E.; Olenin, S.; Wood, S.A. Advantages and limitations of environmental DNA/RNA tools for marine biosecurity: Management and surveillance of non-indigenous species. *Front. Mar. Sci.* **2018**, *5*, 322, doi:10.3389/fmars.2018.00322.
159. Laroche, O.; Wood, S.A.; Tremblay, L.A.; Lear, G.; Ellis, J.I.; Pochon, X. Metabarcoding monitoring analysis: the pros and cons of using co-extracted environmental DNA and RNA data to assess offshore oil production impacts on benthic communities. *PeerJ* **2017**, *5*, e3347, doi:10.7717/peerj.3347.
160. Pochon, X.; Wood, S.A.; Keeley, N.B.; Lejzerowicz, F.; Esling, P.; Drew, J.; Pawlowski, J. Accurate assessment of the impact of salmon farming on benthic sediment enrichment using foraminiferal metabarcoding. *Mar. Pollut. Bull.* **2015**, *100*, 370–382, doi:10.1016/j.marpolbul.2015.08.022.
161. Wilson, D.S.; Swenson, W. Community genetics and community selection. *Ecology* **2003**, *84*, 586–588.
162. Bálint, M.; Pfenninger, M.; Grossart, H.-P.; Taberlet, P.; Vellend, M.; Leibold, M.A.; Englund, G.; Bowler, D. Environmental DNA time series in ecology. *Trends Ecol. Evol.* **2018**, *33*, 945–957, doi:10.1016/j.tree.2018.09.003.
163. Bohan, D.A.; Vacher, C.; Tamaddoni-Nezhad, A.; Raybould, A.; Dumbrell, A.J.; Woodward, G. Next-generation global biomonitoring: Large-scale, automated reconstruction of ecological networks. *Trends Ecol. Evol.* **2017**, *32*, 477–487.
164. Thyssen, M.; Tarran, G.A.; Zubkov, M. V.; Holland, R.J.; Gregori, G.; Burkhill, P.H.; Denis, M. The emergence of automated high-frequency flow cytometry: revealing temporal and spatial phytoplankton variability. *J. Plankton Res.* **2007**, *30*, 333–343, doi:10.1093/plankt/fbn005.
165. Wilcox, T.M.; Zarn, K.E.; Piggott, M.P.; Young, M.K.; McKelvey, K.S.; Schwartz, M.K. Capture enrichment of aquatic environmental DNA: A first proof of concept. *Mol. Ecol. Resour.* **2018**, *18*, 1–10, doi:10.1111/1755-0998.12928.
166. Hamaguchi, M.; Shimabukuro, H.; Hori, M.; Yoshida, G.; Terada, T.; Miyajima, T. Quantitative real-time polymerase chain reaction (PCR) and droplet digital PCR duplex assays for detecting *Zostera marina* DNA in coastal sediments. *Limnol. Oceanogr. Methods* **2018**, doi:10.1002/lom3.10242.
167. Sboner, A.; Mu, X.; Greenbaum, D.; Auerbach, R.K.; Gerstein, M.B. The real cost of sequencing: higher than you think! *Genome Biol.* **2011**, *12*, 125, doi:10.1186/gb-2011-12-8-125.
168. Bearhop, S.; Adams, C.E.; Waldron, S.; Fuller, R.A.; Macleod, H. Determining trophic niche width: A novel approach using stable isotope analysis. *J. Anim. Ecol.* **2004**, *73*, 1007–1012, doi:10.1111/j.0021-8790.2004.00861.x.
169. McKinney, M.A.; Arukwe, A.; De Guise, S.; Martineau, D.; Béland, P.; Dallaire, A.; Lair, S.; Lebeuf, M.; Letcher, R.J. Characterization and profiling of hepatic cytochromes P450 and phase II xenobiotic-metabolizing enzymes in beluga whales (*Delphinapterus leucas*) from the St. Lawrence River Estuary and the Canadian Arctic. *Aquat. Toxicol.* **2004**, *69*, 35–49, doi:10.1016/J.AQUATOX.2004.04.010.
170. Polich, R.L. Stress hormone levels in a freshwater turtle from sites differing in human activity. *Conserv. Physiol.* **2016**, *4*, cow016, doi:10.1093/conphys/cow016.
171. Greig, D.J.; Mashburn, K.L.; Rutishauser, M.; Gulland, F.M.D.; Williams, T.M.; Atkinson, S. Seasonal changes in circulating progesterone and estrogen concentrations in the California sea lion (*Zalophus californianus*). *J. Mammal.* **2007**, *88*, 67–72, doi:10.1644/06-MAMM-A-060R2.1.
172. Skoglund, P.; Posth, C.; Sirak, K.; Spriggs, M.; Valentin, F.; Bedford, S.; Clark, G.R.; Reepmeyer, C.; Petchey, F.; Fernandes, D.; Fu, Q.; Harney, E.; Lipson, M.; Mallick, S.; Novak, M.; Rohland, N.; Stewardson, K.; Abdullah, S.; Cox, M.P.; Friedlaender, F.R.; Friedlaender, J.S.; Kivisild, T.; Koki, G.; Kusuma, P.; Merriwether, D.A.; Ricaut, F.-X.; Wee, J.T.S.; Patterson, N.; Krause, J.; Pinhasi, R.; Reich, D. Genomic insights into the peopling of the Southwest Pacific. *Nature* **2016**, *538*, 510–513, doi:10.1038/nature19844.
173. Schuenemann, V.J.; Avanzi, C.; Krause-Kyora, B.; Seitz, A.; Herbig, A.; Inskip, S.; Bonazzi, M.; Reiter, E.; Urban, C.; Dangvard Pedersen, D.; Taylor, G.M.; Singh, P.; Stewart, G.R.; Velemínský, P.; Likovsky, J.;

- Marczik, A.; Molnár, E.; Pálfi, G.; Mariotti, V.; Riga, A.; Belcastro, M.G.; Boldsen, J.L.; Nebel, A.; Mays, S.; Donoghue, H.D.; Zakrzewski, S.; Benjak, A.; Nieselt, K.; Cole, S.T.; Krause, J. Ancient genomes reveal a high diversity of *Mycobacterium leprae* in medieval Europe. *PLoS Pathog.* **2018**, *14*, e1006997, doi:10.1371/journal.ppat.1006997.
174. Fellows Yates, J.A.; Drucker, D.G.; Reiter, E.; Heumos, S.; Welker, F.; Münzel, S.C.; Wojtal, P.; Lázničková-Galetová, M.; Conard, N.J.; Herbig, A.; Bocherens, H.; Krause, J. Central European Woolly Mammoth population dynamics: Insights from late Pleistocene mitochondrial genomes. *Sci. Rep.* **2017**, *7*, 17714, doi:10.1038/s41598-017-17723-1.
175. Pimm, S.L.; Russell, G.J.; Gittleman, J.L.; Brooks, T.M. The future of biodiversity. *Science*. **1995**, *269*, 347–350, doi:10.1126/science.269.5222.347.
176. Davis, M.; Faurby, S.; Svenning, J.-C. Mammal diversity will take millions of years to recover from the current biodiversity crisis. *Proc. Natl. Acad. Sci. U. S. A.* **2018**, *115*, 11262–11267, doi:10.1073/pnas.1804906115.