

Article

# Effects of Alcohol Hangover on Cognitive Performance: Findings from a Field/Internet Mixed Methodology Study

Andrew Scholey<sup>1\*</sup>, Sarah Benson<sup>1</sup>, Jordy Kaufman<sup>2</sup>, Chantal Terpstra<sup>1</sup>, Elizabeth Ayre<sup>1</sup>, Joris C Verster<sup>1,3</sup>, Cory Allen<sup>4</sup>, Grant Devilly<sup>5</sup>

<sup>1</sup> Centre for Human Psychopharmacology, Swinburne University, Melbourne, VIC 3122, Australia; andrew@scholeylab.com, cterpstra@swin.edu.au, eayre@swin.edu.au

<sup>2</sup> Swinburne BabyLab, Swinburne University, Melbourne, VIC 3122, Australia; jkaufman@swin.edu.au

<sup>3</sup> Institute for Risk Assessment Sciences, Utrecht University, The Netherlands; verster@neuroclinics.nl

<sup>4</sup> Queensland Police Service Academy, GPO Box 1110, Archerfield, QLD 4108, Australia; allencorey1966@hotmail.com

<sup>5</sup> School of Applied Psychology and Griffith Criminology Institute, Griffith University (Mt Gravatt Campus), QLD 4122, Australia; grant@devilly.org

\* Correspondence: andrew@scholeylab.com; Tel.: +44 (3)92148932

**Abstract:** Studies into the cognitive effects of alcohol have been mixed. They also present methodological challenges, often relying on self-report of alcohol consumption leading to hangover. The current study used BAC (obtained via breathalyser) and self-reported drinking behavior during a night out and related these to hangover severity and cognitive function measured over the internet in the same subjects the following morning. Volunteers were breathalysed and interviewed as they left a central entertainment district of an Australian state capital. They were provided with a unique identifier and, the following morning, logged on to a website. This included an online version of the Alcohol Hangover Severity Scale (AHSS), and number and type of drinks consumed the previous night and the eTMT-B - a validated, online analogue of the Trail Making Test B of executive function and working memory. Hangover severity was significantly correlated with one measure only, namely the previous night's BAC ( $r = .228$ ,  $p = .019$ ). Completion time on the eTMT-B was significantly correlated with hangover severity ( $r = .245$ ,  $p = .012$ ), previous night's BAC ( $r = .197$ ,  $p = .041$ ) and time spent drinking ( $r = .376$ ,  $p < .001$ ). These findings confirm that alcohol hangover negatively affects cognitive functioning and that poorer working memory and executive performance correlates with hangover severity. The results also support the utility of using online measures in hangover research.

**Keywords:** hangover; alcohol; internet; attention; executive function; working memory

## 1. Introduction

The alcohol hangover (AH) is defined as “the combination of mental and physical symptoms, experienced the day after a single episode of heavy drinking, starting when blood alcohol concentration approaches zero” [1]. It describes a feeling of malaise which follows a bout of drinking when blood alcohol levels are at or returning to zero [2-7]. The AH is variously characterized by somatic and behavioural symptoms including headache, thirst, stomach upsets, negative mood and cognitive problems.

There can be considerable inter-individual variability in the pattern, severity and temporal characteristics of hangover symptoms [8], with no clear relationship between AH severity and any single physiological process (although cytokine response to alcohol is emerging as a possible key factor [9, 10]). Other mechanisms that may contribute to AH include, but are not limited to, gut dysbiosis (including ghrelin-mediated), decreased blood glucose concentrations, poor sleep

architecture, dehydration (and concomitant electrolyte imbalances), oxidative stress and inflammatory responses [2, 11, 12]. These last two may in part be elevated in response to circulating ethanol metabolites.

The majority of previous research exploring the cognitive effects of alcohol has focused on acute intoxication, and the long-term neurocognitive consequences of alcohol dependence [7, 13, 14]. Acute intoxication impairs aspects of memory, attention and psychomotor performance [15-20]. Alcohol also produces a characteristic shift in the speed/accuracy trade-off. Unlike other impairing drugs which tend to slow responding, alcohol typically increases error rates with little effect on response speed [16, 17, 21].

Compared with alcohol intoxication, relatively little research has been directed at the specific cognitive effects of AH [22, 23]. A recent meta-analysis of next-day cognitive effects of heavy alcohol consumption included 19 studies published in 11 articles since 1970 [24]. It concluded that sustained attention, short- and long-term memory and psychomotor speed are the cognitive domains most susceptible to hangover.

Any hangover-related cognitive impairment could have major implications for everyday activities. For example hangover impairments to driving ability were in the range observed at BACs of 0.05-0.08% [25]. Such impairments have clear ramifications for safety-sensitive occupations, but also negatively impact on those that continue to engage in everyday activities while in a hangover state. In the context of absenteeism and presenteeism, it has been estimated that alcohol costs the US economy 179 billion annually in lost productivity [26]

There are a number of methodological approaches to the study of hangover effects on cognition (see Stephens et al., 2014, for a critical review). These include laboratory studies where controlled doses of alcohol are administered, usually in a relatively pure form (typically vodka) and cognitive outcomes are measured once BACs have returned to zero. Alcohol is either administered at fixed doses, or titrated to reach a target BAC. This approach has the advantage of providing relatively high levels of control, particularly regarding the timing of alcohol administration and measurement of physiological and functional endpoints. On the other hand, it may not have high ecological validity. For example in real-life drinking situations individuals may consume a variety of beverages over different lengths of time. Secondly, because Ethics Boards tend to err on the side of caution, laboratory studies typically use lower levels of alcohol than those observed in the field. Even when bar-like environments are used in the laboratory there tends to be a limit on target alcohol levels.

An alternative methodology in AH research is to use a so-called 'naturalistic' design where participants visit the laboratory on two mornings, one after a night's drinking and another after a sober night [27] (with order counterbalanced across participants). This method has the advantage of ecological validity, by not limiting participants' drinking. On the other hand, the approach relies on recollection of levels of alcohol consumed to generate an estimated Blood Alcohol Level (eBAC). Given that alcohol intoxication is associated with memory problems (as may hangover), this is problematic when trying to explore the relationship between alcohol consumed and functional consequences of AH.

The relative utility of these approaches is illustrated by findings regarding cognitive impairments, associated with hangover [28]. For example psychomotor deficits associated with AH were observed in naturalistic studies [23, 29, 30] but not in laboratory settings [31-33].

The current study took a somewhat different approach. Over the past decade or so, internet studies have been increasing used in psychological research, including to evaluate the impact of substance use [14, 34-40] and field [41] studies into the effects of recreational drugs including alcohol [14, 17, 41-43]. Here we employed mixed methodology, involving a subgroup of individuals who were taking part in ongoing field studies, SmartStart [44] and Last Drinks, into patterns and consequences of alcohol consumption. These studies included a cohort of patrons who were breathalysed while leaving the entertainment area of an Australian capital city and agreed to be contacted the following morning to complete an internet study of hangover. This included a version of the AHSS [1], and an online analogue of the Trail making B (TMT-B), named the eTMT-B, a test of psychomotor function, working memory and executive function. This task was chosen as it is

relatively brief, the task demands are easily understood and it captures elements of the major cognitive domains (psychomotor function, attention and executive function) affected by AH [24, 45].

This approach has several advantages. Next day symptomatology and performance can be linked to measured alcohol levels and drinking characteristics, collecting data via the internet is relatively convenient in that it does not require travelling to a testing location, and the method can address certain methodological questions – for example what factors contribute to attrition in hangover studies.

We hypothesized that previous night BAC would be related to both worse hangover severity and cognitive performance as measured using the eTMT-B. Further analyses explored the influence of beverage types and patterns of drinking on hangover severity and cognitive performance.

## 2. Experimental Section

### 2.1 Pilot study

A pilot study was conducted to evaluate the validity of the online eTMT-B. Twenty-four young adult volunteers (17 female, mean age 30.29 years, SD 5.03) completed both the pencil-and-paper and online versions of the test (with order counterbalanced across participants).

The traditional TMT-B requires participants to draw lines connecting 25 circles distributed on a page containing single digits and letters. Correct completion involves joining the stimuli alternating between ascending numbers and letters (e.g. 1-A-2-B-3-C... etc.), with completion time as the main outcome [29]. The TMT-B was administered according to the published protocol [46], with the exception that errors were not corrected (as this would not be possible in the online version).

6	A	B	11	1
F	8	2	7	4
H	10	5	D	J
G	3	9	E	C
I	L	K	12	13

**Figure 1.** Layout of the eTMT-B. The task requires participants to click ascending numbers and letters alternating between numbers and figure (i.e. 1-A-2-B-3-C etc.).

The eTMT-B was designed as an online analogue of the Pencil-and-Paper TMT-B. It consisted of a 5 × 5 grid of rectangular panels each labeled with a digit or number. Similar to the paper version, the task involved clicking on alternating ascending numbers and digits starting with 1-A and so on (see Figure 1). Once a panel was pressed it changed color from white to grey, after which it could clicking on it had no effect. If a participant attempted to press an incorrect button no response could be made and an incorrect response was recorded.

Both versions were scored for errors and completion time.

### 2.2. Main study

#### 2.2.1. Design

This study formed part of a larger series of studies aimed at determining patterns and drivers for drinking in and around an Australian state capital. This part of the study employed a mixed methodology approach, whereby individuals were approached as they left the Central Entertainment District (CED) and then contacted the following morning to conduct an internet study.

The study was approved by the Ethics committees of both Griffith University [2015/704] and Swinburne University [2016/167].

When first engaged with during the CED phase, participants were presented verbally with a summary of the nature of the study and what their involvement would entail. Survey and breathalyser measures were completed only if the participant provided verbal consent for both the concurrent and next morning measures (as approved by the Ethics committees) as is usual for this kind of research. Each participant was given a unique identifier card with a link to the study information sheet ([www.last-drinks.com.au](http://www.last-drinks.com.au)). They were informed that they would be allowed to withdraw their data (using the anonymous ID number given on the card) and obtain further information regarding the study if they so wished.

### 2.2.1. Participants

One hundred and five participants provided usable datasets. They were recruited at and around exit points (taxi ranks and train station) in the central entertainment area of Brisbane, Queensland, Australia.

### 2.2.3. Breath Alcohol

Breath alcohol levels were measured using an Alcolizer LE5. This device is used by law enforcement agencies throughout Australia and South East Asia, is Australia Standard 3547 certified, and has an accuracy of greater than 0.005 at 0.100 BAC g/100 ml. It has been demonstrated to have high reliability and validity for measuring intoxication in a sample of people attending nighttime entertainment districts [47].

### 2.2.4. Online measures

A website survey was constructed which included questions collecting demographic and morphometric information (birth year, gender, weight and height). There were asked specifically about the number of beers/ciders, glasses of wine, shots (unmixed) and alcohol mixed with either energy drinks or other beverages they had consumed the previous night and on a typical night out. This was followed by an 11-item version of the Alcohol Hangover Severity Scale (AHSS [28]), comprising of 11-point scales with endpoints (0 and 10) labelled as 'absent' and 'extreme'. Individual items were 'sweating', 'confusion', 'thirst', 'nausea', 'fatigue (being tired)', 'heart pounding', 'dizziness', 'shivering', 'clumsiness', 'apathy (lack of interest/concern)', and 'stomach pain'. Note that the 'difficulty concentrating' item was omitted to reduce expectancy effects while engaging in a task requiring an element of focused attention.

The next sections asked questions regarding the previous night's alcohol consumption (specifically numbers and types of beverages), sleep and qualitative hangover data (these will be reported elsewhere).

Participants were then directed to a website with the eTMT-B. Following completion of the online test, a debriefing page was displayed and participants were given details for entry to an online competition to win an iPad. They were also compensated with an 15 AUD iTunes voucher if they provided an Email address.

### 2.2.5. Procedure

Individuals were recruited by being approached as they left the central entertainment area of Brisbane, Queensland. If willing to be interviewed, they provided consent, were breathalysed, and gave details regarding their consumption of alcohol (including time drinking and number of drinks) that night. Those with BACs around 0.05% or above were asked if they agreed to be contacted the following morning. Those consenting (N = 346) provided contact details and estimated bedtime for that night. Participants were given a card with a unique identifier which allowed anonymous linking of their BAC and drinking data to the data collected over the internet.

The following morning individuals were contacted by text message 8 hours following their estimated bedtime and again 6 hours later if they had not completed the survey. If participating, they entered their unique identifier and then completed the webpages as described above.

### 2.2.6. Statistics and analysis

Number of drinks recorded were converted to standard drinks (one Australian standard drink is equivalent to 10 g alcohol). Initial analyses involved exploring the relationship between BAC, hangover severity and cognitive performance with various drinking factors. These included number of standard drinks, time drinking, and number of specific types of drink. The relationships between these factors were analysed using Pearsons correlations. To determine whether BAC affected participation in hangover part of the study, t-tests compared BACs of those who did and did not participate in the next day online phase of the study.

## 3. Results

### 3.1. Pilot Study

Response times were similar between the paper-and pencil and internet platforms. With a mean completion time of 40.27 sec (range 14.71 -71.59) for the paper version of the TMT-B and 41.49 sec (23.02 - 75.10) for the eTMT-B ( $t(23) = 0.116$ ,  $p = .909$ ). These were significantly correlated between individuals completing the two platforms ( $r = .499$ ,  $p = .013$ ).

Published norms for completion of the paper version of the task are typically around 50 sec [45, 48]. Considering our sample were predominantly students and the paper version involves error correction, our figures are in line with normative data. Errors were rare on both platforms with 4 people making errors on the paper version and 10 people on the eTMT-B. The vast majority of these were single errors.

### 3.1. Main Study

**Table 1.** Demographic and morphometric data of sample. Apart from gender figures are means with range in parentheses.

	N	105
Males/females (%)		51.4/48.6
Age		24.7 (17-49) years
Weight		74.0 (43-115) kg
BMI		24.15 (16.58-40.28)


Of 346 participants who were breathalysed and indicated that they may be prepared to participate in the next day phase of the study, 105 provided complete online datasets. Sample characteristics of this cohort are presented in Table 1.

Drinking characteristics of the cohort are presented in Table 2. The sample had a mean BAC of 0.11% (SD  $\pm 0.40$ ). They reported drinking for an average 7.5 ( $\pm 4.1$ ) hours and had consumed a mean number of 13.5 standard drinks. Analyses of next day reports revealed that the most consumed drink was alcohol mixed with non-energy drinks with a mean of  $5.7 \pm 6.0$  drinks, followed by beer/cider (4.0  $\pm 5.6$ ), wine (1.7  $\pm 3.6$ ), shots (1.5  $\pm 2.7$ ) then alcohol mixed with energy drinks (0.8  $\pm 1.7$ ).

The main focus of the study was to examine factors associated with hangover severity and cognitive performance (see Figure 2). Hangover severity was significantly related to one measure only, namely BAC ( $r = .228$ ,  $p = .019$ ). The correlation between total number of standard drinks and HSS gave a value of  $r = .184$ ,  $p = .064$ . Speed of completion of eTMT-B correlated with self-rated hangover severity ( $r = .245$ ,  $p = .0120$ , previous night's BAC ( $r = .197$ ,  $p = .04$ ) and drinking time ( $r = .376$ ,  $p < .001$ ).

**Table 2.** Drinking characteristics including types of drinks consumed reported for the night of data collection, BAC and hours drinking.

	N	Mean	SD	Max
<i>Drinks consumed</i>				
Beer/cider	59	7.16	5.86	30
Wine	31	5.68	4.56	20
Shots (unmixed)	43	3.51	3.21	16
Alcohol mixed with Energy Drink	30	2.87	1.98	8
Alcohol mixed with Other Beverage	83	7.27	5.80	40
Total	105	13.48	5.94	35
<i>Drinking measures</i>				
BAC (%)	-	0.110	0.040	0.25
Hours drinking	-	7.45	4.09	17

**Figure 2.** Graphs depicting significant associations with hangover severity (HSS score), previous night's blood alcohol content (BAC), drinking time and cognitive performance (eTMT-B).

Further analyses revealed that BAC correlated significantly with number of standard drinks consumed ( $r = .486, p < .001$ ) and time drinking ( $r = .376, p < .001$ ). Examination of the relationship between individual drinks and alcohol levels revealed significant correlations between BAC and amount of alcohol consumed as beer/cider ( $r = .361, p = .005$ ), wine ( $r = .398, p = .026$ ) and alcohol mixed with other beverages ( $r = .228, p = .038$ ) but not between BAC and shots alone or alcohol mixed with energy drink (Table 3).



**Table 3.** Correlations between Previous night's BAC, hangover severity and drinking characteristics.

	BAC (%)	Time drinking (h)	HSS score	Standard drinks (N)	Beer/cider (N)	Wine (N)	Shots (N)	AMED (N)	AMOB (N)
<i>N</i>	105	105	105	102	59	31	43	30	83
BAC (%)	-	.376***	.228*	.486***	.361**	.398*	-.028	-.099	.228*
Time drinking (h)	-	-	.148	.633***	.503***	.223	.166	.068	.257*
HSS score	-	-	-	.184	.171	.293	-.010	.019	.018
Standard drinks (N)	-	-	-	-	.623***	.437*	.249	.360	.369**
Beer/cider (N)	-	-	-	-	-	-.388	-.047	.033	-.255
Wine (N)	-	-	-	-	-	-	-.131	.162	-.284
Shots (N)	-	-	-	-	-	-	-	.102	-.067
AMED (N)	-	-	-	-	-	-	-	-	-.023

BAC = Blood Alcohol Content, HSS = Hangover Severity Scale, AMED = alcohol mixed with energy drinks, AMOB = alcohol mixed with other beverage. Significant correlations are indicated in bold (\*,  $p < .05$ ; \*\*,  $P < .01$ ; \*\*\*,  $P < .0001$ ). Drinking characteristics including types of drinks consumed reported for the night of data collection, BAC and hours drinking.

Time drinking significantly correlated with standard drinks consumed in total ( $r = .633$ ,  $p < .001$ ), and as beer/cider ( $r = .503$ ,  $p < .001$ ) and AMOB ( $r = .257$ ,  $p = .019$ ). Standard drinks significantly correlated with drinks consumed as beer/cider ( $r = .623$ ,  $p < .001$ ), wine ( $r = .437$ ,  $p = .014$ ) and AMOB ( $r = .369$ ,  $p = .001$ ). There were no other significant correlations.

There were no difference in the BACs of those who did ( $x = 0.11\% \pm SD 0.0408$ ) and did not ( $x = 0.11\% \pm SD 0.0405$ ) complete the online hangover phase of the study ( $t(345) = 0.240$ ,  $p = .81$ ).

#### 4. Discussion

Using mixed field and internet methodology, we found that hangover severity is significantly related to BAC and both are associated with worse performance on the eTMT-B test of attention and executive function. Drinking time was also associated with BAC and with worse performance on the task (though not with hangover score).

The current methodology allowed measurement of BACs, and next morning collection of data regarding types of alcohol consumed evaluation of severity of alcohol hangover and cognitive functioning. The study confirmed that previous night's BAC was significantly associated with hangover severity. No other measure was significantly correlated with hangover scores despite some intercorrelations with other measures and BAC, including type of beverage consumed.

It has been suggested that different types of alcoholic beverage may influence hangover severity. In particular it has been suggested that non-alcohol constituents of drinks, known as congeners, may differentially affect AH [49]. In particular it has been suggested that congener-rich drinks such as whiskey produce worse hangover symptoms than beverages with essentially no congeners, such as vodka, although there is little systematic research in this area. Our (albeit limited) data do not suggest that different alcohol types contribute differentially to AH symptomatology. Similarly, different types of mixers, had no differential effect on hangover (symptom) severity.

Conversely there was some evidence that certain drink types were more closely associated with BAC. As well as correlating with amount of time drinking, BAC was related to the amount of beer/cider, wine and AMOBs reportedly consumed (but not shots, or AMED). It seems unlikely that the nature of beverage consumed could differentially contribute to BAC. AMOBs were the most

commonly consumed drinks (by  $N = 83$  people, consuming  $\bar{x} = 7.27$  standard drinks), followed by beer/cider ( $N = 59$ ,  $\bar{x} = 7.16$ ), and shots ( $N = 43$ ,  $\bar{x} = 3.51$ ). AMEDs were the least consumed beverages ( $N = 30$ ,  $\bar{x} = 2.87$ ), and although wine was consumed by a similar number of people ( $N = 31$ ), on average it was consumed at a higher level ( $\bar{x} = 7.16$  drinks). Thus, our data confirm other findings using other methodology [50] showing that irrespective of the type of alcoholic drink or mixer, the most meaningful association was between the number of drinks consumed and BAC.

The error rate was generally low with more than half of the sample (54%) making no errors, and around one quarter (26%) making 1 or 2 errors only. This suggests that participants engaged with the eTMT-B and understood task demands of the online version. This confirms the results of the pilot study where there was high correspondence between the paper and online versions of the task. Thus impairment was largely manifest by slowed function. A pattern of slower reaction times with increasing hangover severity would differentiate alcohol hangover from intoxication. The latter is typified by a characteristic shift in the speed/accuracy trade-off (SATO) with intoxication leading to more errors but relatively little effect on reaction times [16, 17, 21, 51]. Since few errors were made on either version of the TMT-B, we cannot draw strong conclusion from this limited dataset, however. Nevertheless, the literature does suggest that slowing of response times during AH is more robust than increased errors [24, 52].

The current study had certain advantages over previous studies applying naturalistic methodology. For example, rather than relying on next day recall of previous night's drinks, we had an objective measure of BACs. Additionally the number of alcoholic beverages consumed and length of time drinking were recorded on the night, making the data less susceptible (though not invulnerable) to recall bias.

While we can be confident that measured BACs were accurate [47], we do not know which phase of drinking participants were when breathalysed. Specifically we do not know if BACs were measured at peak alcohol levels or during the rising or falling limb of the blood alcohol curve. Further, although the BAC measurement occurred at Central Entertainment District exit points (taxi ranks and train stations), we cannot preclude the possibility that some continued to consume alcohol, which would affect hangover and related functional consequences.

Another issue is that approximately one third of the sample did not attempt the next day measures. Drop-out rates in hangover studies are typically high (e.g. 70 % attrition in Grange et al., 2017), suggesting that the current methodology is as viable as others in this respect. One possibility is that those subjects who consume most alcohol and/or have the worse hangover symptoms are less likely to complete the next day measures. The current methodology allowed us to address this directly by comparing the BACs of those who did and did not complete the hangover part of the study. This showed that the BACs were more-or-less identical. Further our sample had a large range of BACs (up to 0.245%), suggesting that reaching a relatively high BAC did not affect the ability to complete the next-day measures. This may be an advantage of the current methodology, as next-day measures were completed online so were relatively convenient.

## 5. Conclusions

In conclusion, using novel methodology, this study has confirmed that higher BACs and associated measures result in worse hangover symptoms and poorer performance on a newly-validated online measure of working memory and executive function, the eTMT-B. Such hangover-related impairments are likely to have clear, real-life ramifications. For example they would impact on the ability to engage in complex behaviors, and may explain some of the hangover-associated impairment of driving [25]. They are also likely to impinge on fundamental aspects of cognitive functioning. Moreover, our findings suggest that a mixed field/internet approach provides a novel and viable methodology for hangover research.

**Author Contributions:** Conceptualization, AS, SB and GD; Data curation, SB, TP and EA; Formal analysis, AS, EA and GD; Funding acquisition, GD; Methodology, JV; Project administration, SB and CA; Resources, CA; Software, JK; Supervision, JV and GD; Validation, JK and TP; Visualization, AS; Writing – original draft, AS; Writing – review & editing, SB, JK, TP, EA, JV, CA and GD..



**Funding:** These data were drawn from a sample of a larger study funded by a grant to GD from the National Drug Strategy Law Enforcement Funding Committee (# 1314004). The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript, or in the decision to publish the results

**Conflicts of Interest:** The authors declare no conflict of interest.

## References

1. van Schrojenstein Lantman, M., et al., *Development of a definition for the alcohol hangover: consumer descriptions and expert consensus*. Current drug abuse reviews, 2016. **9**(2): p. 148-154.
2. Stephens, R., et al., *A critical analysis of alcohol hangover research methodology for surveys or studies of effects on cognition*. Psychopharmacology (Berl), 2014. **231**(11): p. 2223-36.
3. Prat, G., A. Adan, and M. Sanchez-Turet, *Alcohol hangover: a critical review of explanatory factors*. Hum Psychopharmacol, 2009. **24**(4): p. 259-67.
4. Penning, R., et al., *The pathology of alcohol hangover*. Current Drug Abuse Reviews, 2010. **3**: p. 68-75.
5. Swift, R.M.D. and D. Davidson, *Alcohol hangover: Mechanisms and mediators*. Alcohol Health & Research World 1998. **22**(1): p. 54-60.
6. Verster, J.C., *The alcohol hangover--a puzzling phenomenon*. Alcohol Alcohol, 2008. **43**(2): p. 124-6.
7. Wiese, J.G., M.G. Shlipak, and W.S. Browner, *The alcohol hangover*. Annals of Internal Medicine, 2000. **132**(11): p. 897-902.
8. Verster, J.C., et al., *Differences in the Temporal Typology of Alcohol Hangover*. Alcoholism: Clinical and Experimental Research, 2018. **42**(4): p. 691-697.
9. Van De Loo, A., et al., *Time-dependent changes in saliva cytokine concentrations during alcohol hangover: A comparison of two naturalistic studies*. Alcoholism: Clinical and Experimental Research, 2016. **40**(S1): p. 95A.
10. Van De Loo, A.J., et al., *Impact of mental resilience and perceived immune functioning on the severity of alcohol hangover*. BMC research notes, 2018. **11**(1): p. 526.
11. T Tipple, C., S. Benson, and A. Scholey, *A review of the physiological factors associated with alcohol hangover*. Current drug abuse reviews, 2016. **9**(2): p. 93-98.
12. Farokhnia, M., et al., *Pharmacological manipulation of the ghrelin system and alcohol hangover symptoms in heavy drinking individuals: Is there a link?* Pharmacology Biochemistry and Behavior, 2018.
13. Brust, J., *Ethanol and cognition: indirect effects, neurotoxicity and neuroprotection: a review*. International journal of environmental research and public health, 2010. **7**(4): p. 1540-1557.
14. Ling, J., et al., *Effects of alcohol on subjective ratings of prospective and everyday memory deficits*. Alcoholism: Clinical and Experimental Research, 2003. **27**(6): p. 970-974.
15. Ogden, E.J. and H. Moskowitz, *Effects of alcohol and other drugs on driver performance*. Traffic injury prevention, 2004. **5**(3): p. 185-198.
16. Scholey, A.B., et al., *Neurocognitive and mood effects of alcohol in a naturalistic setting*. Human Psychopharmacology: Clinical and Experimental, 2012. **27**(5): p. 514-516.
17. Tiplady, B., N. Franklin, and A. Scholey, *Effect of ethanol on judgments of performance*. British Journal of Psychology, 2004. **95**(1): p. 105-118.
18. Parrott, A., et al., *Understanding drugs and behaviour*. 2005, West Sussex, England: John Wiley & Sons Ltd.

19. Farquhar, K., et al., *Effect of ethanol on psychomotor performance and on risk taking behaviour*. J Psychopharmacol, 2002. **16**(4): p. 379-384.
20. Maylor, E.A. and P. Rabbitt, *Alcohol, reaction time and memory: a meta-analysis*. Brit J Psychol, 1993. **84**: p. 301-317.
21. Mackay, M., B. Tiplady, and A.B. Scholey, *Interactions between alcohol and caffeine in relation to psychomotor speed and accuracy*. Human Psychopharmacology: Clinical and Experimental, 2002. **17**(3): p. 151-156.
22. Schroyen Lantman, M., et al., *The impact of alcohol hangover symptoms on cognitive and physical functioning, and mood*. Human Psychopharmacology: Clinical and Experimental, 2017.
23. Verster, J.C., et al., *Alcohol hangover effects on memory functioning and vigilance performance after an evening of binge drinking*. Neuropsychopharmacology, 2003.
24. Gunn, C., et al., *A systematic review of the next-day effects of heavy alcohol consumption on cognitive performance*. Addiction, 2018.
25. Verster, J.C., et al., *Effects of alcohol hangover on simulated highway driving performance*. Psychopharmacology, 2014. **231**(15): p. 2999-3008.
26. Control, C.f.D., *The cost of excessive alcohol use*. 2015.
27. McKinney, A., et al., *Next day effects of naturalistic alcohol consumption on tasks of attention*. Human Psychopharmacology: Clinical and Experimental, 2012. **27**(6): p. 587-594.
28. Penning, R., et al., *Measurement of alcohol hangover severity: development of the Alcohol Hangover Severity Scale (AHSS)*. Psychopharmacology, 2013. **225**(4): p. 803-810.
29. McKinney, A. and K. Coyle, *Next-day effects of alcohol and an additional stressor on memory and psychomotor performance*. Journal of studies on Alcohol and Drugs, 2007. **68**(3): p. 446-454.
30. McKinney, A. and K. Coyle, *Next day effects of a normal night's drinking on memory and psychomotor performance*. Alcohol and Alcoholism, 2004. **39**(6): p. 509-513.
31. Collins, W.E. and W.D. Chiles, *Laboratory performance during acute alcohol intoxication and hangover*. Human Factors: The Journal of the Human Factors and Ergonomics Society, 1980. **22**(4): p. 445-462.
32. Myrsten, A., et al., *Acute effects and after-effects of alcohol on psychological and physiological functions*. Psychological lab. Univ. of Stockholm. 1970, Report.
33. Kruisselbrink, L.D., et al., *Physical and psychomotor functioning of females the morning after consuming low to moderate quantities of beer*. Journal of Studies on Alcohol, 2006. **67**(3): p. 416-420.
34. Heffernan, T.M., et al., *Prospective memory, everyday cognitive failure and central executive function in recreational users of Ecstasy*. Human psychopharmacology: Clinical and experimental, 2001. **16**(8): p. 607-612.
35. Parrott, A.C., et al., *Ecstasy/MDMA attributed problems reported by novice, moderate and heavy recreational users*. Human Psychopharmacology: Clinical and Experimental, 2002. **17**(6): p. 309-312.
36. Parrott, A.C., et al., *Dancing hot on Ecstasy: physical activity and thermal comfort ratings are associated with the memory and other psychobiological problems reported by recreational MDMA users*. Human Psychopharmacology: Clinical and Experimental, 2006. **21**(5): p. 285-298.
37. Rodgers, J., et al., *Differential effects of Ecstasy and cannabis on self-reports of memory ability: a web-based study*. Human Psychopharmacology: Clinical and Experimental, 2001. **16**(8): p. 619-625.

38. Rodgers, J., et al., *Patterns of drug use and the influence of gender on self-reports of memory ability in ecstasy users: a web-based study*. Journal of Psychopharmacology, 2003. **17**(4): p. 389-396.
39. Scholey, A.B., et al., *Increased intensity of Ecstasy and polydrug usage in the more experienced recreational Ecstasy/MDMA users: a WWW study*. Addictive behaviors, 2004. **29**(4): p. 743-752.
40. Verster, J.C., S. Benson, and A. Scholey, *Motives for mixing alcohol with energy drinks and other nonalcoholic beverages, and consequences for overall alcohol consumption*. International journal of general medicine, 2014. **7**: p. 285.
41. Scholey, A., et al., *Neurocognitive and mood effects of alcohol in a naturalistic setting*. Human Psychopharmacology, 2012. **27**: p. 514-516.
42. Scholey, A.B. and K.A. Fowles, *Retrograde enhancement of kinesthetic memory by alcohol and by glucose*. Neurobiology of learning and memory, 2002. **78**(2): p. 477-483.
43. Scholey, A., et al., *Effects of resveratrol and alcohol on mood and cognitive function in older individuals*. Nutrition and Aging, 2014. **2**(2): p. 133-138.
44. Devilly, G.J., C. Allen, and K. Brown, *SmartStart: Results of a large point of entry study into preloading alcohol and associated behaviours*. International Journal of Drug Policy, 2017. **43**: p. 130-139.
45. Tombaugh, T.N., *Trail Making Test A and B: normative data stratified by age and education*. Archives of clinical neuropsychology, 2004. **19**(2): p. 203-214.
46. Bowie, C.R. and P.D. Harvey, *Administration and interpretation of the Trail Making Test*. Nature protocols, 2006. **1**(5): p. 2277-2281.
47. Sorbello, J.G., et al., *Fuel-cell breathalyser use for field research on alcohol intoxication: an independent psychometric evaluation*. PeerJ, 2018. **6**: p. e4418.
48. Fernandez, A.L. and B.A. Marcopulos, *A comparison of normative data for the Trail Making Test from several countries: equivalence of norms and considerations for interpretation*. Scandinavian journal of psychology, 2008. **49**(3): p. 239-246.
49. J Rohsenow, D. and J. Howland, *The role of beverage congeners in hangover and other residual effects of alcohol intoxication: a review*. Current drug abuse reviews, 2010. **3**(2): p. 76-79.
50. Köchling, J., et al., *Grape or grain but never the twain? A randomized controlled multiarm matched-triplet crossover trial of beer and wine*. The American Journal of Clinical Nutrition, 2019. **109**(2): p. 345-352.
51. Benson, S., B. Tiplady, and A. Scholey, *Attentional and working memory performance following alcohol and energy drink: a randomised, double-blind, placebo-controlled, factorial design laboratory study*. PloS One, 2019.
52. Stephens, R., et al., *Review A review of the literature on the cognitive effects of alcohol hangover*. Alcohol & Alcoholism, 2008. **43**(2): p. 163-170.
53. Grange, J.A., et al., *The effect of alcohol hangover on choice response time*. Journal of Psychopharmacology, 2016. **30**(7): p. 654-661.