

Leishmania Proteomics: an in silico perspective [†]

Carlos A. Padilla,[‡] Maria J. Alvarez,[¶] and Aldo F. Combariza^{*,‡}

[‡]*Evolutionary Biology Research Group, in silico Molecular Modelling and Computational Simulation Research Group, Sciences and Education School, Biology and Chemistry Department, University of Sucre, Sincelejo, Colombia*

[¶]*Evolutionary Biology Research Group, in silico Molecular Modelling and Computational Simulation Research Group, Education and Sciences School, Biology and Chemistry Department, University of Sucre, Sincelejo, Colombia*

E-mail: aldo.combariza@unisucra.edu.co

Abstract

We report on the state of the art of proteins recognized as potential targets for the development of *Leishmania* treatments through the search of biologically active chemical species, either from experimental *in vitro*, *in vivo*, or *in silico* sources. We classify the gathered information, in several ways: vector taxonomy and geographical distribution, *Leishmania* parasite taxonomic and geographical distribution and enzymatic function (oxidoreductases, transferases, hydrolases, lyases, isomerases, ligases and cytokines). Our aim is to provide a much needed reference layout for research efforts aimed to understand the background of ligand-protein activation/inactivation processes, in this specific case, related with enzymes known to be part of biochemical cascade reactions initiated following a *Leishmania* infectious episode.

[†]To be corrected...

Introduction

Leishmaniasis is a tropical and subtropical group of zoonotic diseases, caused for different species of *Leishmania* genus.^{1,2} It mainly affects mammals and is transmitted through the bite of infected female sand flies.^{3,4} Sand fly taxonomic classification is presented in Figure 1. Actual molecular and phylogenetics analysis have allowed to build a rich and complex taxonomic classification of *Leishmania* genus.^{2,5} The division of *Leishmania* genus in euleishmania and paraleishmania was proposed as the result of molecular analysis.⁶ Euleishmania involves subgenus *L. (Viannia)*, *L. (Leishmania)* and *L. (Sauroleishmania)* (Fig. 2).⁶⁻⁹ Paraleishmania includes *L. (Endotrypanum)* subgenus, containing only *E. schaudinni* and *E. monterogeii* species (See Fig. 2).¹⁰⁻¹³ Colombia have reports of *L. amazonensis*, *L. braziliensis*, *L. mexicana*, *L. colombiensis*, *L. guyanensis*, *L. panamensis*, *L. chagasi*, *L. lainsoni* and *L. equatoriensis* as parasites of leishmaniasis transmitters.^{14,15}

The parasitization process involves a myriad of proteic agents, all of them playing preponderant roles in the development of the disease. Several proteins have been recognized as possible targets for the development of treatments.¹⁶ Currently, the first line drugs used to treat the disease are: Meglumine antimoniate (Glucantime), Sodium stibogluconate (Pentostam), Amphotericin B (Fungizone), Liposomal amphotericin B (AmBiosome), paromomycin and pentamidine.¹⁶⁻¹⁸ These drugs are not completely effective against *Leishmania*. There are many research efforts directed towards the development of new bioactive principles to control the disease, many of which turn to natural products (metabolites).^{19,20} as a source of inspiration

In this work, we set our goal as to shed light on the relationship between nature-inspired bioactive chemical structures and macromolecular proteic agents involved in any of the different levels of the *Leishmania* parasitization process, through the microscopic-atomistic lens of ab-initio and force-field mechanical-statistical based models, that is, to use hybrid in silico methodologies to model and simulate ligand-protein systems.




Figure 1: *Leishmania* vector taxonomic and classification and geographical distribution: Diptera order, Psychodidae family, phlebotominae subfamily, and, geographically, *Phlebotomus* genus in the old world and *Lutzomyia* genus for the new world.^{2,11}




Figure 2: *Leishmania* taxonomic classification and geographical distribution. *Leishmania* genus: euleishmania and paraleishmania. Paraleishmania is conformed by *L. (endotrypanum)* subgenus and euleishmania by *L. (viannia)*, *L. (leishmania)* and *L. (sauroleishmania)* subgenus.⁶⁻⁹

Leishmania life cycle and host immune response

Leishmania parasites invade, develop and replicate inside the host Mononuclear Phagocyte System (MPS), attacking macrophages and dendritic cells.²¹⁻²³ Incubation time of *Leishmania* parasite from promastigote to amastigote takes between two and three months, time range when the host immune system response activates and can be favorable or unfavorable.²² The parasite cycle begins with the bite of an infected female sandfly, carrier of promastigote *Leishmania* parasites in a meta-cyclic state (see Fig. 3).^{24,25} At this stage, neutrophils and macrophages are the first line immune cells activated, being neutrophils the initiators of the inflammatory process.^{25,26}

Leishmania parasite has two ways to enter the macrophage: A direct path, via the macrophage, and an indirect path, by attacking the neutrophils.²⁷ The direct path, occurs when the promastigote is directly endocited by the macrophage phagosome, or parasitophorous vacuole, which undergoes a biochemical transformation into phagolysosome.²⁸ The indirect path goes through the neutrophil mediated phagocytosis of the parasite, followed by a subsequent macrophage phagocytosis step (see Fig. 3).²⁷ *Leishmania* promastigote parasite survives inside the phagolysosome vacuole by producing Lipophosphoglycan (LPG), gp63 protein and glutathione transferase.^{24,29}

Following the initial macrophage attack, the biochemical response of the host immune system focus on the production of cytokines by specialized T_H $CD4^+$ cells, either T_H1 or T_H2 . Among the T_H1 secreted cytokines we found gamma interferon ($IFN-\gamma$), which activates and stimulates the macrophages, increasing its microbiocide activity. Interleucine (IL)-12 and IL-2, cytokines also secreted by T_H1 cells, help T_H $CD4^+$ transformation into T_H1 , which is the more suitable form to respond to the exogenous attack, in our case, the metacyclic promastigote form of *Leishmania*.^{30,31}).

Immune T_H2 response produces IL-4, IL-5, IL-10, Transforming Growth Factor Beta ($TGF-\beta$), among other cytokines. These compound are more effective facing allergenic diseases or helminthic infections, and therefore is more desirable the T_H1 response. Moreover,

the T_H2 response inhibits the T_H1 , favoring the propagation and survival of the *Leishmania* parasite.^{30,32}

***Leishmania* protein targets**

Crystal structures were retrieved from the Protein Data Bank (PDB) and classified accordingly.³³ The PDB search process was carried using the keywords "*Leishmaniasis*" and "*Leishmania*", providing 425 items, after which we classified the selected proteins according to the parasitization cycle. Other selection criteria considered were: proteins with different PDB-codes but same structure and proteins with equal structures but elucidated from different organisms were considered. An example of proteins that belong to different groups are: Dihydrofolate Reductase-Thymidylate Synthase (DHFR-TS) (PDB ID: 3INV), which is an oxidoreductase and a transferase. A total of 49 proteins comprised the final study population as possible drug-targets (see Fig. 4).

From the selected proteins, the most abundant group belongs to oxidoreductases (Enzyme Commission Number - EC 1) and transferases (EC 2), with 15 subjects each one, 6 hydrolases (EC 3), 5 lyases (EC 4), 3 isomerases (EC 5), 3 ligases (EC 6), one cytokine and one Lipid Binding Protein was classified too (see Fig. 4).

Oxidoreductases (EC 1)

This group is comprised by enzymes that catalyzes oxidation-reduction reactions.³⁴ Glyceraldehyde 3-phosphate dehydrogenase *Leishmania major* (PDB ID: 1GYP and 1A7K) belongs to this group. 1GYP is a homotetrameric enzyme of 156 kDa that catalyzes Glyceraldehyde-3-phosphate (GAP) oxidative phosphorylation to 1,3 Bisphosphoglycerate (BPG) in the glycolysis pathway, through the NAD^+ cofactor.³⁵

Malate dehydrogenase (PDB ID: 4H7P) participates in the glycolysis process by reducing oxaloacetate (OAA) to malate.³⁶ Dihydrofolate Reductase-Thymidylate Synthase (DHFR-




Figure 3: *Leishmania* life cycle can be divided in five steps. **Step 1:** Virulent metacyclic promastigotes are egested when a female sandfly carrier bites a possible host. **Step 2:** Here there are two possible paths. Direct path: promastigotes are phagocytized by macrophage cells. Indirect path: promastigotes are phagocytized by neutrophil cells, subsequently, neutrophils are phagocytized by macrophage. At the end of step 2, promastigotes end up inside a macrophage phagolysosome. **Step 3:** The promastigote parasite produces glutathione transferase, a protein that protects it from the acidic conditions of the phagolysosome, then initiating the replication process. **Step 4:** Transformation of vector promastigote into amastigote takes place. *Leishmania* amastigotes are experts at exploiting host cell machinery to thrive. **Step 5:** Amastigotes are then taken out when a sand fly, either infected or not, bites the host, closing the cycle.

TS) (PDB ID: 3INV) and Pteridine reductase (PTR1) (PDB ID: 1E92) from *L. major* and *T. cruzi*, respectively, are responsible of pterin salvaging in parasitic trypanosomatids. Specifically, this happens because *Leishmania* parasites are auxotrophics for folate and other pterins required in critical pathways, including nucleic acid and protein biosynthesis.^{37,38} Dihydroorotate dehydrogenase (DHODH) (PDB ID: 3GYE) is a flavoprotein enzyme involved in the *de novo* pyrimidine biosynthesis pathway. 3GYE catalyzes (S)-dihydroorotate oxidation to orotate, in a redox reaction. DHODH is located within the cytosol (Class 1) and inside the mitochondria (Class 2) in eukaryotes and some prokaryotes.³⁹

Thymine dehydrogenase JBP1 (PDB ID: 2XSE) is essential for J-Base DNA (β -D-glucosyl-hydroxymethyluracil) biosynthesis and maintenance.⁴⁰ Tryparedoxin (PDB ID: 3S9F) and Tryparedoxin peroxidase I (PDB ID: 3TUE) (TXN/TXNPx) reduces macrophages-hydroperoxides produced as the infection progresses.⁴¹ These proteins stay in a cytosolic form. The detoxification pathway is essential for parasite survival.⁴¹ Pseudoperoxidase (PP) (PDB ID: 5VIA) *L. major* is a heme protein, expressed in humans during pathogen attack.⁴² It is an important protein involved in pathogen defense mechanisms against reactive nitrogen species, such as peroxynitrite, produced as host immune response.⁴² Ascorbate peroxidase (APX) (PDB ID: 3RIV) is yet another enzyme involved in this metabolic pathway, and is considered as a potential drug specific target.⁴³

Enzyme superoxide dismutase - FeSODA (PDB ID: 4F2N) protects the parasite against macrophage toxic radicals. When the amastigote is phagocytized, macrophage cells produce a respiratory burst generating reactive oxygen intermediates, such as hydrogen peroxide, OH^- , O_2^- radicals and peroxynitrite, as part of the macrophage mechanism to fight invasive microorganisms.⁴⁴⁻⁴⁶ Thus, 4F2N acts as the first line of defense against those Reactive Oxygen Species (ROS). This fact, makes 2F2N a suitable enzymatic target for *Leishmania* controlling drug development.⁴⁴

Trypanothione reductase *Leishmania infantum* (PDB ID: 2YAU) is a key enzyme from the *Leishmania infantum* polyamine-dependent redox metabolism.⁴⁷ Prostaglandin F syn-

thase from *L. major* (PGF; PDB ID: 4F40) is involved in the lipid metabolic pathway, acting through a NADP cofactor.⁴⁸ Sterol 14 α -demethylase (CYP51) *L. infantum* is an enzyme that catalyzes the removal of the 14 α -methyl group from sterol precursors. This reaction is essential for membrane cell biosynthesis, specifically, CYP51 it relates to the ergosterol pathway, and is believed to be crucial for the survival of infectious *Leishmania* parasite.⁴⁹




Figure 4: *Leishmania* protein classification. Oxidoreductases (orange), transferases (blue), hydrolases (red), lyases (violet), isomerases (green), ligases (dark blue), cytokines (yellow) and Lipid Binding protein (light orange). The chart at the bottom right of the figure shows the number of proteins found for each species. *T. cruzi* and *T. brucei* do not belong at the *Leishmania* protein group, but they have been used in some studies as homologous proteins.^{38,50}

Transferases group (EC. 2)

Transferases catalyze reactions in which a chemical group is transferred from a electron/proton donor substrate to an electron/proton acceptor substrate.³⁴ These proteins catalyze key cellular processes in all kingdoms of life, such as, DNA repair, RNA editing, and activation of metabolites used in catabolic pathways.⁵¹ Adenine phosphoribosyltransferase (APRT) (PDB ID: 1QB7 and 1MZV) belongs to the phosphoribosyltransferase family type I (PRTs) and is involved in the purine-salvaging process, catalyzing adenines to adenosine-5-monophosphate (AMP).^{52,53} It is a key enzyme, because, protozoan parasites are auxotrophic for the purine *de novo* biosynthesis pathway. N-myristoyltransferase (NMT) (PDB ID: 4UCM) catalyzes the co-translational transfer of myristate from myristoyl-CoA to the N-terminal glycine in a large number of proteins. Modifications have been implicated in localization and/or substrate activation.⁵⁴ OMP decarboxylase (PDB ID: 3QW3) and UMP synthase (PDB ID: 3QW4) are two enzyme involved in the last two steps of *de novo* Uridine 5'-monophosphate (UMP) biosynthesis, for pyrimidine production (as opposed to the purine-salvaging pathway).⁵⁵ Both enzymes are recognized as possible drug targets.

Cysteine synthase (PDB ID: 4AIR) is involved in cysteine biosynthesis. Cysteine is required for protein biosynthesis in *Leishmania* species, also, it is a source of reduced sulfur for important metabolites, such as coenzyme A (CoA), enzyme cofactors and iron-sulfur clusters.⁵⁶ DNA polymerase Beta (PDB ID: 4P4M) is required for maintenance, replication and recombination of the DNA.⁵⁷ This protein is specially required for the *Leishmania* parasite amastigote form.⁵⁷ Thymidine kinase (PDB ID: 4UXH) catalyzes ATP γ -phosphate transfer to 2'-deoxythymidine (dThd), forming thymidine monophosphate (dTMP). It is a important enzyme because plays a key role in parasitization process.⁵⁸ Farnesyl pyrophosphate synthase (FPPS) (PDB ID: 4JZX) is involved in ergosterol synthesis too, acting in the early steps of isoprene synthesis and maintainance of lipid bilayer integrity.^{59,60} It is a potential enzymatic target, because, it was successful inhibited with bisphosphonate previously.⁵⁹

UDP-glucose pyrophosphorylase *L. major* (PDB ID: 4J18) catalyzes the reversible con-

version of glucose-1-phosphate (Glc-1-P) and uridine 5'-triphosphate (UTP) to UDP-Glc and inorganic pyrophosphate (PPi) in the presence of Mg^{2+} , into glycolytic pathway.⁵¹ Aspartate aminotransferase (AAT) (PDB ID: 4H51) catalyzes reversible transfer of the α -amino group of aspartate and glutamate, converting L-aspartate and 2-oxoglutamate to oxaloacetate and L-glutamate (<http://brenda-enzymes.info>).⁶¹ This enzyme is pyridoxal phosphate (PLP) co-factor dependent and a potential drug target.⁶² 3-mercaptopyruvate sulfurtransferase (PDB ID: 1OKG) is involved in cysteine metabolism, having three domains: N-terminal, core and C-terminal. The active site of the protein is localized along the N-terminal and core domains, containing the Cys-253, Arg-74 and 185 as sulfite inhibition amino acids. Asp-61, His-75 and Ser-255 are close to Cys-253, and polarize the carboxyl group of 3-mercaptopyruvate through a nucleophilic attack.⁶³ Transketolase (PDB ID: 1R9J) is a key enzyme in the nonoxidative branch of the pentose phosphate pathway (PPP), which transfers two-carbon glyceraldehyde units from ketose-donors to aldose-acceptor sugars.⁶⁴ Finally, pyruvate kinase (PDB ID: 1PKL) catalyzes the phosphoenolpyruvate-phosphate group transfer to adenosine diphosphate. This enzyme is involved in the glycolytic pathway.

Hydrolases group (EC. 3)

Hydrolases catalyze reactions in which a bond in a substrate is hydrolyzed to produce two fragments.³⁴ Purine nucleoside hydrolase *L. major* enzyme (PDB ID: 1EZR) is the main responsible of nucleotide salvaging from host, due *Leishmania* parasite lack *de novo* biosynthesis pathway of purines.⁶⁵ The *Leishmania* parasite promastigote form, express glycoproteins in its surface; these enzymes are known as Leishmanolysin (*gp63* gene) (PDB ID: 1LML).⁶⁶ It has an important role in macrophages infection process.⁶⁶

Oligopeptidase B (OPB) *L. major* (PDB ID: 2XE4) catalyzes the Arg- and Lys- bond formation in proteins, and is a potential drug target, according to several studies.⁶⁷ Tyrosine phosphatase (PRL-1) (PDB ID: 3S4O) is mainly secreted by the *Leishmania* parasite promastigote form, however, the parasite amastigote form produces PRL-1 more efficient and

abundantly during macrophage infection process.⁶⁸ It is important for the parasite survival. Putative aminopeptidase (LAP) *L. major* (PDB ID: 5NTH) is involved in N-terminus catalysis of proteins. This protein has multiple functions; in host infection processes it provides essential amino acid, and is a major transcription regulator, among other functions.⁶⁹ Finally, the arginase protein (PDB ID: 4ITY) catalyzes the first step polyamine biosynthesis. This process entails cellular growth as well as parasite survival.⁷⁰

Lyase group (EC. 4)

Group of enzymes that catalyzes non-hydrolytic reactions, in which a chemical group is cleaved and removed from a substrate leaving a double bond.³⁴ Adenylosuccinate lyase (ASL) (PDB ID: 4MX2) is a lyase protein, and have been identified as vital component of purine salvaging in *Leishmania donovani*.⁷¹ Fumarate hydrolase (FH) class 1 enzyme is a cytosolic protein with two isoforms: a mitochondrial (PDB ID: 6MSO) and a cytosolic ((PDB ID: 5L2R)). 5L2R produces fumarate substrate for the dihydroorotate dehydrogenase. Additionally, this enzyme migrate to cellular nucleus, where play a key rol in DNA repair.⁷² 6MSO catalyzes the stereospecific reversible conversion of fumarate to S-malate. This reaction is part of the tricarboxylic acid (TCA) cycle, participates in the succinic fermentation pathway, DNA repair and is proposed to provide fumarate for the de novo pyrimidine biosynthetic pathway.⁷³ Finally, aldolase *L. mexicana* (PDB ID: 1EPX) enzyme, is involves in glycolytic pathway and catalyzes the Fructose-1,6-bisphosphate conversion to glyceraldehyde-3-phosphate and dihydroxyacetone phosphate.⁷⁴

Isomerases group (EC. 5)

Isomerase proteins catalyzes one-substrate/one-product reactions that can be regarded as isomerization reactions.³⁴ Triosephosphate isomerase (TIM) (PDB ID: 1AMK) plays a central rol in the glycolysis process, as catalyst of dihydroxyacetone phosphate (DHAP) and D-glyceraldehyde-3-phosphate (GAP).⁷⁵ Currently, there exists a TIM E65Q mutant (PDB

ID: 1QDS), being more stable than the TIM wild-type.⁷⁶ Peptidyl-prolyl cis-trans isomerase (PDB ID: 4S1E), accelerates the folding process of proteins. It catalyzes the cis-trans isomerization of proline imidic peptide bonds in oligopeptides <https://www.uniprot.org/uniprot/Q9U9R3>.⁷⁷

Ligases group (EC. 6)

Ligase proteins catalyzes the joining together of two or more molecules coupled to hydrolysis of ATP or an analogous molecule.³⁴ This group has three proteins classified as possible enzymatic targets. First and second enzyme are Methionyl-tRNA synthetase (PDB ID: 3KFL) and Tyrosyl-tRNA synthetase (PDB ID: 3P0I), which were elucidated with MgATP as substrate and methionine as solvent, and are recognized as essential for biological processes, such as gene translation.^{78,79} The three enzyme is Asparagine synthetase A (ASNA) (PDB ID: 4LNS), which is an ammonium and glutamine dependent enzyme. In experimental (*in vivo* and *in vitro*) studies, this protein-deletion-gene caused parasite growth delay, therefore is recognized as a possible target.⁵⁰

Cytokines group

The Migration Inhibitory Factor (MIF) from *L. major* (PDB ID: 3B64) was classified as possible enzymatic target. This cytokine is an ortholog of human MIF, also known as Lm1740MIF. 3B64 interacts with MIF receptors, HLA class II histocompatibility antigen gamma chain (also called invariant chain or CD74) and exhibits an antiapoptotic activity that may facilitate the intracellular persistence of *Leishmania* into macrophages.⁸⁰

Lipid Binding Protein group

At this group belong the Apo- Acyl Carrier Protein (PDB ID: 5ZWS), where play crucial roles in biosynthesis of fatty acids, non-ribosomal polypeptides and polyketides. The fatty acids pathway and their more complex forms recently gained attention in *Leishmania*, because

it play crucial roles in the virulences of protozoan parasites for survival inside the host.⁸¹ Therefore, 5ZWS is a potential enzymatic target.

Proteomics analysis: *Leishmania* parasite metabolic pathways

Some of the previously classified proteins are involved in critically important metabolic pathways such as Glycolysis, the pentose-phosphate and citrate cycles. These pathways are essential for normal metabolic parasite processes, therefore, are essential to *Leishmania* parasite survival.⁸² As an example, Arginase participates in the following metabolic pathways: arginine biosynthesis, proline and arginine metabolism, secondary metabolite biosynthesis and antibiotic biosynthesis.⁸³ (see fig. 5).

The proteins with PDB code 1E92, 2XSE, 3VIA, 4F2N, 4F40, 4UCM, 2XE4, 3S4O, 5NTH, 4S1E, 3P0I, 3KFL y 5ZWS, do not appear in the metabolic pathway databases. Other proteins as 3B64, 1LML and 4P4M, are structural proteins of *Leishmania* parasite, but are not included into a metabolic pathway.

Leishmaniasis drugs

Pentavalent antimonials (Sb(V)) were firsts leishmaniasis drugs for along time, however, development of *Leishmania* resistance made it drug inefficient. New drug and treatment investigations have been development.² Currently, being developed much effective drugs, but it generates secondary effects as high toxicity, also, these treatments and drugs are expensive and generates resistance for *Leishmania* parasites.^{84,85}

Leishmaniasis disease is a Neglected Tropical Disease (NTD), therefore, it does not have necessary economic funds and strict epidemiological controls from governmental organizations, made difficult drugs and treatments development.⁸⁶

Standard drugs

Currently, leishmaniasis is treated using the following types of chemical substances: antimonials (Sb(V)), amphotericin B, Pentamidine, Miltefosine (hexadecylphosphocholine), paromomycin (aminosidine), sitamaquine and pamidronate (see Fig. 6).^{17,87} These drugs are used for treats three main leishmaniasis forms: cutaneous leishmaniasis (CL), mucocutaneous leishmaniasis (MCL) and visceral leishmaniasis (VL).⁸⁴ Antimonials were the first antileishmania compounds, introduced in the 40s decade.^{2,87} They are available as meglumine antimoniate (Glucantime) and sodium stibogluconate (Pentostam), and these are standard first line drugs for treatment, but emergence of resistance has limited their use.^{16,85} Antimonials are used for VL treatment, but, different studies found that *L. donovani* and *L. braziliensis* were more sensitive to sodium stibogluconate than *L. major*, *Leishmania tropica* and *L. mexicana*.^{2,17} Amphotericin B is a macrolide antibiotic isolated from *Streptomyces nodosus* in 1956 and widely used since the 80s as amphotericin B deoxycholate.^{2,87} It selectively inhibits the membrane synthesis of the parasite and causes holes in the membrane, leading to parasite death.⁸⁷ It is used as a second-line treatment, and is present in five formulations: amphotericin B deoxycholate (Fungizone), Emulsification of Fungizone in intralipid 20 %, liposomal amphotericin B (AmBisome), amphotericin B lipid complex (ABLC; Abelcet) and amphotericin B cholesterol dispersion (ABCD; Amphotec) (see Fig. 6).^{17,87} Amphotericin B and its lipid formulations are used as alternative chemotherapeutic treatments.¹⁶ Lipid formulations of amphotericin B have been gaining importance, becoming the fist-choice established leishmaniasis treatment by the US Food and Drug Administration (FDA).²

Pentamidine is widely recognised, its antileishmanial activity is on *Leishmania* parasite polyamines biosynthesis and mitochondrial membrane. They are considered second-line treatment drugs for VL, because its toxicity (myalgia, nausea, headache, hypoglycemia, irreversible insulin dependent diabetes mellitus and death), declining its efficacy. Also, its hig2 monetary costs renders them prohibitive.⁸⁷ India and East Africa used paromomycin as




Figure 6: Drugs used against leishmaniasis can be divided in two main groups: standard drugs and alternative drugs (metabolites). Graphics show molecular structures of drug active compounds. Structures were obtained from ChemSpider (<http://www.chemspider.com/>).

a cheap alternative treatment, despite its toxicity.² Paromomycin remained neglected until the 80s, when topical formulations for VL were developed.⁸⁷ One World Health, the Bill and Melinda Gates Foundation, Gland Pharma Limited, IDA Solutions and WHO/TDR partnered to develop Paromomycin as a public health tool to be sold on a not-for-profit basis, at a very low price.⁸⁷ Miltefosine, initially developed as an anticancer drug, currently is the first effective oral treatment of VL.^{2,87} Variation in the sensitivities of both, promastigote and amastigote stages of *L. donovani*, *L. major*, *L. tropica*, *Leishmania aethiopica*, *L. mexicana* and *L. panamensis*, were investigated *in vitro*. From these assays, *L. donovani* was recognized as the most sensitive species to this treatment.¹⁷ Miltefosine ED₅₀ against *L. donovani* was measured in the range of 0.12 to 1.32 μ M. Sitamaquine is rapidly metabolized, forming desethyl and 4-CH₂OH derivatives, which might be responsible for its activity. Toxicity appears to be relatively mild, as it causes mild methemglobinaemia.^{87,88} Finally, pamidronate is a bisphosphonate drug with significant activity against *Leishmania donovani in vitro*. FPPS protein (PDB ID: 4JZX) is potently inhibited by bisphosphonates in the trypanosomatid parasite.⁸⁷ Other two type of drugs considered for leishmaniasis treatment are azoles and nucleoside analogues.^{17,89} Within the azole group are, for example, ketoconazole and itraconazole, which inhibits the C14 α -demethylase. Nucleoside Analogues, such as allopurinol and pyrazolopyrimidine, are known to inhibit enzymatic processes of the purine salvaging pathway in *Leishmania*.¹⁷

Metabolomics: Non standard drugs from plant metabolites

Leishmania resistance against antileishmanial drugs suggests that the current use of monotherapies has to be reviewed. The rationale behind combination therapies is to prevent the risk of parasite resistance, provide increased efficacy resulting from synergistic effects, lower dose requirements and reduced duration of therapy, eventually leading to reduced secondary toxic

effects of the drugs.⁸⁴ Natural compounds, such as secondary plant metabolites, are another type of alternative treatment. We have retrieved information of 222 compounds, isolated from plants, with recognized antileishmanial activity (Supplementary Material). The search process was performed through revision of the state of the art.^{19,20,90} These compounds were isolated from several parts of the plants, and tested in experimental assays (*in vitro* and *in vivo*) under different physical chemistry conditions and *Leishmania* species.

References

- (1) Torres-Guerrero, E.; Quintanilla-Cedillo, M. R.; Ruiz-Esmenjaud, J.; Arenas, R. Leishmaniasis: a review. *F1000Research* **2017**, *6*, 750.
- (2) Bruschi, F.; Gradoni, L. In *The Leishmaniases: Old Neglected Tropical Diseases*; Bruschi, F., Gradoni, L., Eds.; Springer International Publishing: Cham, 2018; pp 1–245.
- (3) Killick-Kendrick, R. The biology and control of Phlebotomine sand flies. *Clinics in Dermatology* **1999**, *17*, 279–289.
- (4) (OMS), O. M. d. l. S. *Control de las leishmaniasis*; 2010; p 200.
- (5) Kostygov, A. Y.; Yurchenko, V. Revised classification of the subfamily Leishmaniinae (Trypanosomatidae). *Folia Parasitologica* **2017**, *64*, 1–5.
- (6) Cupolillo, E.; Medina-Acosta, E.; Noyes, H.; Momen, H.; Grimaldi, G. A revised classification for *Leishmania* and *Endotrypanum*. *Parasitology Today* **2000**, *16*, 142–144.
- (7) Lainson, R.; Shaw, J. In *The Leishmaniases in Biology and Epidemiology*; Peters, W., Killick-Kendrick, R., Eds.; Academic Press, 1987; Vol. 1; Chapter 1, pp 1 – 120.
- (8) Lainson, R.; Shaw, J. In *Biology of the Kinetoplastida*, 2nd ed.; In WHR Lumsden, D. E., Ed.; Academic Press: London and New York, 1979; p 116.

- (9) Corrêa, J. R.; Brazil, R. P.; Soares, M. J. *Leishmania* (*Viannia*) *lainsoni* (Kinetoplastida: Trypanosomatidae), a divergent *Leishmania* of the *Viannia* subgenus - A mini review. *Memorias do Instituto Oswaldo Cruz* **2005**, *100*, 587–592.
- (10) Franco, A. M.; Grimaldi, G. Characterization of *Endotrypanum* (Kinetoplastida: Trypanosomatidae), a Unique Parasite Infecting the Neotropical Tree Sloths (Edentata). *Memorias do Instituto Oswaldo Cruz* **1999**, *94*, 261–268.
- (11) Akhoundi, M.; Kuhls, K.; Cannet, A.; Votýpka, J.; Marty, P.; Delaunay, P.; Sereno, D. A Historical Overview of the Classification, Evolution, and Dispersion of *Leishmania* Parasites and Sandflies. *PLoS Neglected Tropical Diseases* **2016**, *10*, 1–40.
- (12) Mesnil F, B. E. Sur un hematozoaire nouveau (*Endotrypanum* n.gen.) d'un edente de la Guyane. *C R Soc Biol* **1908**, *65*: 587.
- (13) JJ, S. *London School of Hygiene and Tropical Medicine Memoir*, 13th ed.; H. K. Lewis (1969): London, 1969; p 132.
- (14) WHO/PAHO, *Leishmaniases: Epidemiological Report in the Americas*; 2018; pp 1 – 7.
- (15) Universidad del Rosario, Colombia, el país con más especies de parásitos de *Leishmania*. 2017; <http://www.urosario.edu.co/UCD/Leishmania/Documento/pdf-fasciculo-leishmaniasis-universidad-del-rosari/>.
- (16) Chawla, B.; Madhubala, R. Drug targets in *Leishmania*. *Journal of Parasitic Diseases* **2010**, *34*, 1–13.
- (17) Croft, S. L.; Sundar, S.; Fairlamb, A. H. Drug Resistance in Leishmaniasis. *Society* **2006**, *19*, 111–126.
- (18) OPS/OMS, O. P. d. l. S. *Leishmaniasis: Informe Epidemiológico de las Américas*; 2018; Vol. 6; pp 1 – 7.

- (19) Tiuman, T. S.; Santos, A. O.; Ueda-Nakamura, T.; Filho, B. P. D.; Nakamura, C. V. Recent advances in leishmaniasis treatment. *International Journal of Infectious Diseases* **2011**, *15*.
- (20) Sen, R.; Chatterjee, M. Plant derived therapeutics for the treatment of Leishmaniasis. *Phytomedicine* **2011**, *18*, 1056–1069.
- (21) Catta-Preta, C. M. C.; Mottram, J. C. Drug candidate and target for leishmaniasis. *Nature* **2018**, *560*, 171–172.
- (22) Chappuis, F.; Sundar, S.; Hailu, A.; Ghalib, H.; Rijal, S.; Peeling, R. W.; Alvar, J.; Boelaert, M. Visceral leishmaniasis: What are the needs for diagnosis, treatment and control? *Nature Reviews Microbiology* **2007**, *5*, 873–882.
- (23) Rittig, M. G.; Bogdan, C. Leishmania-host-cell interaction: Complexities and alternative views. *Parasitology Today* **2000**, *16*, 292–297.
- (24) Cunningham, A. C. Parasitic Adaptive Mechanisms in Infection by Leishmania. *Experimental and Molecular Pathology* **2002**, *72*, 132–141.
- (25) Nathan, C. Neutrophils and immunity: challenges and opportunities. *Nature Reviews Immunology* **2006**, *6*, 173–182.
- (26) Charmoy, M.; Auderset, F.; Allenbach, C.; Tacchini-Cottier, F. The Prominent Role of Neutrophils during the Initial Phase of Infection by Leishmania Parasites. *Journal of Biomedicine and Biotechnology* **2010**, *2010*, 1–8.
- (27) Laskay, T.; Zandbergen, G. v.; Solbach, W. Neutrophil granulocytes as host cells and transport vehicles for intracellular pathogens : Apoptosis as infection-promoting factor. *Immunology* **2008**, *213*, 183–191.
- (28) Naderer, T.; McConville, M. J. The Leishmania-macrophage interaction: a metabolic perspective. *Cellular Microbiology* **2008**, *10*, 301–308.

- (29) Naderer, T.; Vince, J.; McConville, M. Surface Determinants of Leishmania Parasites and their Role in Infectivity in the Mammalian Host. *Current Molecular Medicine* **2004**, *4*, 649–665.
- (30) Kindt, T. J.; Goldsby, R. A.; Osborne, B. A. In *INMUNOLOGÍA*, 6th ed.; Fraga, J. d. L., Ed.; Mc Graw Hill, interamericana, 2007.
- (31) Parham, P. In *The Immune System, Fourth Edition*, 4th ed.; Science, G., Ed.; Taylor and Francis Group, 2014; Vol. 39; p 624.
- (32) Kima, P. E. Leishmania molecules that mediate intracellular pathogenesis. *Microbes and Infection* **2014**, *16*, 721–726.
- (33) Berman, H. M. The Protein Data Bank. *Nucleic Acids Research* **2000**, *28*, 235–242.
- (34) Cornish-Bowden, A. Current IUBMB recommendations on enzyme nomenclature and kinetics. *Perspectives in Science* **2014**, *1*, 74–87.
- (35) Kim, H.; Feil, I. K.; Verlinde, C. L. M. J.; Petra, P. H.; Hol, W. G. J. Crystal Structure of Glycosomal Glyceraldehyde-3-phosphate Dehydrogenase from Leishmania mexicana: Implications for Structure-Based Drug Design and a New Position for the Inorganic Phosphate Binding Site. **1995**, *34*, 14975–14986.
- (36) Pelley, J. W. *Elsevier's Integrated Biochemistry*; Elsevier, 2007; pp 65–71.
- (37) Gourley, D. G.; Schüttelkopf, A. W.; Leonard, G. A.; Luba, J.; Hardy, L. W.; Beverley, S. M.; Hunter, W. N. Pteridine reductase mechanism correlates pterin metabolism with drug resistance in trypanosomatid parasites. *Nature Publishing Group* **2001**, *8*, 521 – 525.
- (38) Vadloori, B.; Sharath, A. K.; Prabhu, N. P.; Maurya, R. Homology modelling, molecular docking, and molecular dynamics simulations reveal the inhibition of Leishmania

- donovani dihydrofolate reductase-thymidylate synthase enzyme by Withaferin-A. *BMC Research Notes* **2018**, *11*, 1–7.
- (39) Cordeiro, A. T.; Feliciano, P. R.; Pinheiro, M. P.; Nonato, M. C. Crystal structure of dihydroorotate dehydrogenase from *Leishmania major*. *Biochimie* **2012**, *94*, 1739–1748.
- (40) Heidebrecht, T.; Christodoulou, E.; Chalmers, M. J.; Jan, S.; Ter Riet, B.; Grover, R. K.; Joosten, R. P.; Littler, D.; Van Luenen, H.; Griffin, P. R.; Wentworth, P.; Borst, P.; Perrakis, A. The structural basis for recognition of base J containing DNA by a novel DNA binding domain in JBP1. 2011.
- (41) Fiorillo, A.; Colotti, G.; Boffi, A.; Baiocco, P.; Ilari, A. The Crystal Structures of the Tryparedoxin-Tryparedoxin Peroxidase Couple Unveil the Structural Determinants of *Leishmania* Detoxification Pathway. *PLoS Neglected Tropical Diseases* **2012**, *6*, e1781.
- (42) Chreifi, G.; Dejam, D.; Poulos, T. L. Crystal structure and functional analysis of *Leishmania major* pseudoperoxidase. *Journal of Biological Inorganic Chemistry* **2017**,
- (43) Jasion, V. S.; Polanco, J. A.; Meharena, Y. T.; Li, H.; Poulos, T. L. Crystal structure of *Leishmania major* peroxidase and characterization of the compound I tryptophan radical. *Journal of Biological Chemistry* **2011**,
- (44) Phan, I. Q. H.; Davies, D. R.; Moretti, N. S.; Shanmugam, D.; Cestari, I.; Anupama, A.; Fairman, J. W.; Edwards, T. E.; Stuart, K.; Schenkman, S.; Myler, P. J. Iron superoxide dismutases in eukaryotic pathogens: new insights from Apicomplexa and Trypanosoma structures. *Acta Crystallographica Section F Structural Biology Communications* **2015**, *71*, 615–621.
- (45) GHOSH, S.; GOSWAMI, S.; ADHYA, S. Role of superoxide dismutase in survival of *Leishmania* within the macrophage. *Biochemical Journal* **2003**, *369*, 447–452.

- (46) Slauch, J. M. How does the oxidative burst of macrophages kill bacteria? Still an open question. *Molecular Microbiology* **2011**, *80*, 580–583.
- (47) Ilari, A.; Baiocco, P.; Messori, L.; Fiorillo, A.; Boffi, A.; Gramiccia, M.; Di Muccio, T.; Colotti, G. A gold-containing drug against parasitic polyamine metabolism: the X-ray structure of trypanothione reductase from *Leishmania infantum* in complex with auranofin reveals a dual mechanism of enzyme inhibition. *Amino Acids* **2012**, *42*, 803–811.
- (48) Moen, S. O.; Fairman, J. W.; Barnes, S. R.; Sullivan, A.; Nakazawa-Hewitt, S.; Van Voorhis, W. C.; Staker, B. L.; Lorimer, D. D.; Myler, P. J.; Edwards, T. E. Structures of prostaglandin F synthase from the protozoa *Leishmania major* and *Trypanosoma cruzi* with NADP. *Acta Crystallographica Section F Structural Biology Communications* **2015**, *71*, 609–614.
- (49) Hargrove, T. Y.; Wawrzak, Z.; Liu, J.; Nes, W. D.; Waterman, M. R.; Lepesheva, G. I. Substrate preferences and catalytic parameters determined by structural characteristics of sterol 14 α -demethylase (CYP51) from *Leishmania infantum*. *Journal of Biological Chemistry* **2011**,
- (50) Manhas, R.; Tripathi, P.; Khan, S.; Sethu Lakshmi, B.; Lal, S. K.; Gowri, V. S.; Sharma, A.; Madhubala, R. Identification and Functional Characterization of a Novel Bacterial Type Asparagine Synthetase A. *Journal of Biological Chemistry* **2014**, *289*, 12096–12108.
- (51) Fühling, J.; Cramer, J. T.; Routier, F. H.; Lamerz, A.-C.; Baruch, P.; Gerardy-Schahn, R.; Fedorov, R. Catalytic Mechanism and Allosteric Regulation of UDP-Glucose Pyrophosphorylase from *Leishmania major*. *ACS Catalysis* **2013**, *3*, 2976–2985.
- (52) Phillips, C. L.; Ullman, B.; Brennan, R. G.; Hill, C. P. Crystal structures of adenine

- phosphoribosyltransferase from *Leishmania donovani*. *EMBO Journal* **1999**, *18*, 3533–3545.
- (53) Silva, M.; Silva, C. H.; Iulek, J.; Oliva, G.; Thiemann, O. H. Crystal structure of adenine phosphoribosyltransferase from *Leishmania tarentolae*: Potential implications for APRT catalytic mechanism. *Biochimica et Biophysica Acta - Proteins and Proteomics* **2004**,
- (54) Robinson, D. A.; Wyatt, P. G. Identification and structure solution of fragment hits against kinetoplastid N -myristoyltransferase. *Acta Crystallographica Section F Structural Biology Communications* **2015**, *71*, 586–593.
- (55) French, J. B.; Yates, P. A.; Soysa, D. R.; Boitz, J. M.; Carter, N. S.; Chang, B.; Ullman, B.; Ealick, S. E. The *Leishmania donovani* UMP synthase is essential for promastigote viability and has an unusual tetrameric structure that exhibits substrate-controlled oligomerization. *Journal of Biological Chemistry* **2011**,
- (56) Fyfe, P. K.; Westrop, G. D.; Ramos, T.; Müller, S.; Coombs, G. H.; Hunter, W. N. Structure of *Leishmania major* cysteine synthase. *Acta Crystallographica Section F: Structural Biology and Crystallization Communications* **2012**,
- (57) Mejia, E.; Burak, M.; Alonso, A.; Larraga, V.; Kunkel, T. A.; Bebenek, K.; Garcia-Diaz, M. Structures of the *Leishmania infantum* polymerase beta. *DNA Repair* **2014**,
- (58) Timm, J.; Bosch-Navarrete, C.; Recio, E.; Nettleship, J. E.; Rada, H.; González-Pacanowska, D.; Wilson, K. S. Structural and Kinetic Characterization of Thymidine Kinase from *Leishmania major*. *PLOS Neglected Tropical Diseases* **2015**, *9*, e0003781.
- (59) Aripirala, S.; Gonzalez-Pacanowska, D.; Oldfield, E.; Kaiser, M.; Amzel, L. M.; Gabelli, S. B. Structural and thermodynamic basis of the inhibition of *Leishmania major* farnesyl diphosphate synthase by nitrogen-containing bisphosphonates. *Acta Crystallographica Section D: Biological Crystallography* **2014**,

- (60) de Mattos Oliveira, L.; Araújo, J. S. C.; Bacelar Costa Junior, D.; Santana, I. B.; Duarte, A. A.; Leite, F. H. A.; Benevides, R. G.; Coelho dos Santos Junior, M. Pharmacophore modeling, docking and molecular dynamics to identify Leishmania major farnesyl pyrophosphate synthase inhibitors. *Journal of Molecular Modeling* **2018**, *24*, 314.
- (61) Schomburg, I.; Jeske, L.; Ulbrich, M.; Placzek, S.; Chang, A.; Schomburg, D. The BRENDA enzyme information system—From a database to an expert system. *Journal of Biotechnology* **2017**, *261*, 194–206.
- (62) Abendroth, J.; Choi, R.; Wall, A.; Clifton, M. C.; Lukacs, C. M.; Staker, B. L.; Van Voorhis, W.; Myler, P.; Lorimer, D. D.; Edwards, T. E. Structures of aspartate aminotransferases from Trypanosoma brucei, Leishmania major and Giardia lamblia. *Acta Crystallographica Section F: Structural Biology Communications* **2015**,
- (63) Alpey, M. S.; Williams, R. A.; Mottram, J. C.; Coombs, G. H.; Hunter, W. N. The crystal structure of leishmania major 3-mercaptopyruvate sulfurtransferase. A three-domain architecture with a serine protease-like triad at the active site. *Journal of Biological Chemistry* **2003**, *278*, 48219–48227.
- (64) VEITCH, N. J.; MAUGERI, D. A.; CAZZULO, J. J.; LINDQVIST, Y.; BARRETT, M. P. Transketolase from Leishmania mexicana has a dual subcellular localization. *Biochemical Journal* **2004**, *382*, 759–767.
- (65) Shi, W.; Schramm, V. L.; Almo, S. C. Nucleoside Hydrolase from Leishmania major. *The Journal of Biological Chemistry* **1999**, *274*, 21114–21120.
- (66) Schlagenhauf, E.; Etges, R.; Metcalf, P. The crystal structure of the Leishmania major surface proteinase leishmanolysin (gp63). *Structure* **1998**, *6*, 1035–1046.
- (67) McLuskey, K.; Paterson, N. G.; Bland, N. D.; Isaacs, N. W.; Mottram, J. C. Crystal Structure of Leishmania major Oligopeptidase B Gives Insight into the Enzymatic Prop-

- erties of a Trypanosomatid Virulence Factor. *Journal of Biological Chemistry* **2010**, *285*, 39249–39259.
- (68) Leitherer, S.; Clos, J.; Liebler-Tenorio, E. M.; Schleicher, U.; Bogdan, C.; Soulat, D. Characterization of the protein tyrosine phosphatase LmPRL-1 secreted by *Leishmania major* via the exosome pathway. *Infection and Immunity* **2017**,
- (69) Timm, J.; Valente, M.; García-Caballero, D.; Wilson, K. S.; González-Pacanowska, D. Structural Characterization of Acidic M17 Leucine Aminopeptidases from the *Trypanosoma brucei* and Evaluation of Their Role in Nutrient Starvation in *Trypanosoma brucei*. *mSphere* **2017**,
- (70) D'Antonio, E. L.; Ullman, B.; Roberts, S. C.; Dixit, U. G.; Wilson, M. E.; Hai, Y.; Christianson, D. W. Crystal structure of arginase from *Leishmania mexicana* and implications for the inhibition of polyamine biosynthesis in parasitic infections. *Archives of Biochemistry and Biophysics* **2013**, *535*, 163–176.
- (71) Boitz, J. M.; Strasser, R.; Yates, P. A.; Jardim, A.; Ullman, B. Adenylosuccinate synthetase and adenylosuccinate lyase deficiencies trigger growth and infectivity deficits in *Leishmania donovani*. *Journal of Biological Chemistry* **2013**, *288*, 8977–8990.
- (72) Feliciano, P. R.; Drennan, C. L.; Nonato, M. C. Crystal structure of an Fe-S cluster-containing fumarate hydratase enzyme from *Leishmania major* reveals a unique protein fold. *Proceedings of the National Academy of Sciences* **2016**, *113*, 9804–9809.
- (73) Feliciano, P. R.; Drennan, C. L.; Nonato, M. C. Crystal Structures of Fumarate Hydratases from *Leishmania major* in a Complex with Inhibitor 2-Thiomalate. *ACS Chemical Biology* **2019**, *14*, 266–275.
- (74) Chudzik, D. M.; Michels, P. A.; De Walque, S.; Hol, W. G. Structures of type 2 peroxisomal targeting signals in two trypanosomatid aldolases. *Journal of Molecular Biology* **2000**, *300*, 697–707.

- (75) Williams, J. C.; Zeelen, J. P.; Neubauer, G.; Vriend, G.; Backmann, J.; Michels, P. a.; Lambeir, a. M.; Wierenga, R. K. Structural and mutagenesis studies of leishmania triosephosphate isomerase: a point mutation can convert a mesophilic enzyme into a superstable enzyme without losing catalytic power. *Protein engineering* **1999**, *12*, 243–250.
- (76) Lambeir, A. M.; Backmann, J.; Ruiz-Sanz, J.; Filimonov, V.; Nielsen, J. E.; Kursula, I.; Norledge, B. V.; Wierenga, R. K. The ionization of a buried glutamic acid is thermodynamically linked to the stability of *Leishmania mexicana* triose phosphate isomerase. *European Journal of Biochemistry* **2000**, *267*, 2516–2524.
- (77) Bateman, A. et al. UniProt: the universal protein knowledgebase. *Nucleic Acids Research* **2017**, *45*, D158–D169.
- (78) Larson, E. T.; Kim, J. E.; Zucker, F. H.; Kelley, A.; Mueller, N.; Napuli, A. J.; Verlinde, C. L.; Fan, E.; Buckner, F. S.; Van Voorhis, W. C.; Merritt, E. A.; Hol, W. G. Structure of *Leishmania major* methionyl-tRNA synthetase in complex with intermediate products methionyladenylate and pyrophosphate. *Biochimie* **2011**, *93*, 570–582.
- (79) Larson, E. T.; Kim, J. E.; Castaneda, L. J.; Napuli, A. J.; Zhang, Z.; Fan, E.; Zucker, F. H.; Verlinde, C. L. M. J.; Buckner, F. S.; Van Voorhis, W. C.; Hol, W. G. J.; Merritt, E. A. The double-length tyrosyl-tRNA synthetase from the eukaryote *leishmania major* forms an intrinsically asymmetric pseudo-dimer. *Journal of Molecular Biology* **2011**,
- (80) Kamir, D. et al. A *Leishmania* Ortholog of Macrophage Migration Inhibitory Factor Modulates Host Macrophage Responses. *The Journal of Immunology* **2008**, *180*, 8250–8261.
- (81) Arya, R.; Sharma, B.; Dhembala, C.; Pal, R. K.; Patel, A. K.; Sundd, M.; Ghosh, B.; Makde, R. D.; Kundu, S. A conformational switch from a closed apo- to an open

- holo-form equips the acyl carrier protein for acyl chain accommodation. *Biochimica et Biophysica Acta - Proteins and Proteomics* **2019**, 1867, 163–174.
- (82) McConville, M. J.; Naderer, T. Metabolic Pathways Required for the Intracellular Survival of Leishmania. *Annual Review of Microbiology* **2011**, 65, 543–561.
- (83) Kanehisa, M. From genomics to chemical genomics: new developments in KEGG. *Nucleic Acids Research* **2006**, 34, D354–D357.
- (84) Zulfiqar, B.; Shelper, T. B.; Avery, V. M. Leishmaniasis drug discovery: recent progress and challenges in assay development. *Drug Discovery Today* **2017**, 22, 1516–1531.
- (85) Ghorbani, M.; Farhoudi, R. Leishmaniasis in humans: Drug or vaccine therapy? *Drug Design, Development and Therapy* **2018**, 12, 25–40.
- (86) den Boer, M.; Argaw, D.; Jannin, J.; Alvar, J. Leishmaniasis impact and treatment access. *Clinical Microbiology and Infection* **2011**, 17, 1471–1477.
- (87) Kumar, A. *Leishmania and Leishmaniasis*; 2013; Vol. 3.
- (88) Soto, J.; Soto, P. Miltefosina oral para el tratamiento de la leishmaniasis. *Biomédica* **2012**, 26, 207.
- (89) Ramón Azanza, J.; García-Quetglas, E.; Sádaba, B. Farmacología de los azoles. *Rev Iberoam Micol* **2007**, 24, 223–227.
- (90) Gutiérrez-Rebolledo, G. A.; Drier-Jonas, S.; Jiménez-Arellanes, M. A. Natural compounds and extracts from Mexican medicinal plants with anti-leishmaniasis activity: An update. *Asian Pacific Journal of Tropical Medicine* **2017**, 10, 1105–1110.