

*Article*

# Breeding Biofortified Pearl Millet Varieties and Hybrids to Enhance Millet Markets for Human Nutrition

M. Govindaraj \*, KN Rai, Binu Cherian, W H Pfeiffer, A Kanatti and H Shivade

International Crops Research Institute for the Semi-Arid Tropics (ICRISAT), Patancheru 502 324, Telangana, India.

\* Correspondence: m.govindaraj@cgiar.org; Tel.: (optional; include country code; if there are multiple corresponding authors, add author initials) +xx-xxxx-xxx-xxxx (F.L.)

**Abstract:** Pearl millet is an important food crop in the arid and semi-arid tropical regions of Africa and Asia. These regions are home to millions of poor smallholder's households living in harsh agro-ecology and reported higher prevalence of malnutrition. . Indeed, pearl millet is one of the food crops they continue to grow for their food and nutritional security. Considering inherent high nutritional values and climate resilient nature (drought and heat), demand for pearl millet as food, beside valued for its Stover as a source of livestock fodder, is projected to grow strongly in Asia (India) and Africa (West and Central Africa). Iron (cause anemia) and zinc (cause stunting) deficiencies are widespread and serious public health problems worldwide, including India and Africa. Biofortification is a cost-effective and sustainable agricultural strategy to address this problem. Aim of this review is to provide current status and future directions of pearl millet biofortification for growing nutrition markets. Research on pearl millet has shown that large genetic variability (30-140 mg kg<sup>-1</sup> Fe and 20-90 mg kg<sup>-1</sup> Zn) available in this crop can be effectively utilized to develop high-yielding cultivars with high iron and zinc densities. Both Open –pollinated varieties (Dhanshakti and Chakti) and hybrids (ICMH 1202, ICMH 1203 and ICMH 1301) of pearl millet with high grain yield (>3.5 tons/ha in hybrids) and high levels of iron (70-75 mg kg<sup>-1</sup>) and zinc (35-40 mg kg<sup>-1</sup>) densities have been developed and released in India. Currently, India growing >70,000 ha of biofortified pearl millet, besides more pipeline hybrids and varieties are under various stage of testing at the national (India) and international (west Africa) trials for possible release. Biofortification program utilized the existing high-Fe lines identified from the mainstream program to serve the immediate objective of developing high-yielding and high-Fe hybrids. Genomic tools will be an integral part of biofortification breeding program particularly to use diagnostic markers and genomic selection for micronutrients. Several biofortified varieties and hybrids released in pearl millet. Till today, no markets to promote biofortified varieties and hybrids as no incentive price and such products aims to address food and nutritional security challenges simultaneously. The demand is likely to increase only after investment in crop breeding and integration into modern public distribution system, nutritional intervention schemes, private seed and food companies with strong mainstreaming nutritional policies. In the non-traditional regions, this will contribute to livestock and poultry feed industry as spill-over benefits to improve nutrition. The following sections of this review describing various aspects of breeding for increased market opportunity of this crop towards addressing micronutrient malnutrition.

**Keywords:** Biofortification; Pearl millet; Malnutrition; Iron; Zinc; Market.

## 1. Introduction

Inadequate intake of energy-providing organic macronutrients (largely carbohydrate, followed by protein and fat, in that order), leads to under-nutrition, with a consequent feeling of hunger. The result of chronic hunger, reflected in such apparent physical manifestations as underweight (being too thin for one's age), wasting (being too thin for one's height) and stunting (being too short for one's age), has long been debated as a food security issue in various global fora [1]. Unlike the macronutrients mentioned above, which are consumed in larger quantities for proper growth and development, there are several micronutrients, which are needed in trace amounts, but they play vital roles in various physiological functions. The deficiencies of these micronutrients, do not lead to obvious hunger effects, lead to all adverse physical manifestations as those arising from the deficiencies of macronutrients. Thus, these micronutrient deficiencies are termed as hidden hunger. Deficiencies of iron (Fe) and zinc (Zn) have been reported to be most widespread, affecting more than two billion people worldwide, mostly in the low- and middle-income countries.

Micronutrient malnutrition, primarily the result of poor quality of diets or poor intake of vitamins and minerals, affects more than 2 billion peoples in developing countries, especially women and preschool children [2,3]. The costs of these micronutrient deficiencies in terms of lives lost, adverse effect on economic growth and poor quality of life are very huge and more staggering in developing countries, including India. Therefore, considerable global development efforts are underway to improve the health of poor people by breeding staple food crops enriched with essential micronutrients, so called biofortification. It is a multidisciplinary, sustainable and cost-effective approach to bring the full potential of crop improvement and nutrition science to bear on the persistent problem of micronutrient malnutrition. Pearl millet is highly nutritious, largely grown under rainfed conditions in India (8 m ha) and Africa (18 m ha) [4]. It supplies 80-90% of the calories for several millions of the poor people in the world [5]. In India, highest per capita consumption by rural population, especially in the western Rajasthan and Gujarat, this contributes more than 50% of the cereal consumption in these regions [6] and also consumed other parts of Gujarat, Rajasthan, Maharashtra and Haryana [7].

HarvestPlus is a CGIAR Challenge Program, leading a global effort to breed and disseminate micronutrient-rich crop varieties inducing pearl millet through partnership in developing countries. While development of OPVs (100%) continues to be the thrust research area in Africa, development of hybrids (70%) is the primary focus in India. Pearl millet as such is a high-iron crop with a fairly high Zn content than rice and wheat; however, not all available cultivars have high Fe and Zn content. So far, crop breeding programs in ICRISAT and National Agricultural Research System (NARS) in India and Africa have largely been focused on grain yield and components traits (including biotic stress tolerance) and less emphasis was given to nutritional quality traits (iron and zinc) in the core line/cultivar development process. Consequently, narrow range of such micronutrients exhibited in most of the released cultivars (46-56 mg kg<sup>-1</sup> Fe) that are being used for consumption in poor households. For instance, all the commercial hybrids had 42 mg kg<sup>-1</sup> mean iron and 31 mg kg<sup>-1</sup> mean zinc content [8]. Therefore, thrust of the pearl millet biofortification research targeted for India is on the development of high-yielding and high-Fe hybrids since the entire efforts in public and private sector are geared towards hybrid development. This effort has been underway in ICRISAT biofortification program by utilizing the existing high-Fe hybrid parents and advanced breeding lines identified from the mainstream breeding program to serve the immediate objective of developing high-yielding and high-Fe hybrids. The gradual shift of biofortification research from detection of variability through breeding high-Fe cultivars using existing high-Fe lines to testing and delivery of biofortified hybrids realized in India. Whereas in Africa, 100% OPV cultivation, fast-track development of OPV using high-Fe populations are in progress. all these screening, breeding and market opportunities are briefly discussed in the following sections.

## 2. High-Throughput Micronutrient Phenotyping

The success of a breeding program depends on the precision phenotyping efficiency through high throughput tools. This is primary needs of the biofortification research and is a key to speedy progress in identifying high-Fe/Zn lines from large set of germplasm and gene pools. Atomic Absorption Spectrophotometry (AAS) and Inductively Coupled Plasma Optical Emission Spectrometry (ICP-OES) techniques (both are destructive method) are highly used by researchers and its results are reproducible for grain Fe and Zn densities. However, breeding for micronutrient dense cultivars needs screening large number of genetic material, such as germplasm collections, elite lines, segregating populations, hybrids etc. and phenotyping for micronutrients though destructive techniques involves high analytical cost and breeding resources. The recent study using advanced technique X-ray Fluorescence Spectrometry (XRF) estimates of Fe and Zn densities from diverse material of pearl millet showed highly significant and positive correlation with ICP values ( $r =$  up to 0.97;  $p < 0.01$  for Fe and  $r =$  up to 0.98;  $p < 0.01$  for Zn) [9]. Hence, high-throughput, non-destructive and low-cost quantitative technique XRF are now being used in several laboratories for initial screening of large genetic material. For final validation of those top-ranking entries based on XRF value can be further re-evaluated using ICP to obtain precision estimates for wider reporting and documentation [10].


## 3. Genetic Variability for Micronutrients

Genetic variability is very critical asset to breeders to start with any trait specific breeding. With the high throughput screening facilities, several thousands of pearl millet samples were analyzed for grain Fe and Zn density. This is largely contributed to scale available genetic variability for these two micronutrients in working germplasm and breeding materials at ICRISAT. Several studies at ICRISAT showed wide range of variability for grain Fe and Zn densities in diverse breeding materials such as, *Iniari* germplasm accessions (51-121 mg kg<sup>-1</sup> Fe; 46-87 mg kg<sup>-1</sup> Zn) population progenies (18.0-135.0 mg kg<sup>-1</sup> Fe; 22.0-92.0 mg kg<sup>-1</sup> Zn), inbred parents (30.3 – 102.0 mg kg<sup>-1</sup> Fe; 27.4 to 84.0 mg kg<sup>-1</sup> Zn), hybrids derived from diverse inbreds (25.8-80.0 mg kg<sup>-1</sup> Fe; 22.0-70 mg kg<sup>-1</sup> Zn) and commercial hybrids (31.0-61.0 mg kg<sup>-1</sup> Fe, 32.0-54.0 mg kg<sup>-1</sup>). Most of these genetic materials with high Fe and Zn levels were entirely or largely based on *Iniari* germplasm [11-13]. *Iniari* refers to early-maturing and large-seeded land races found in adjoining parts of Togo, Ghana, Benin and Burkina Faso [14]. Core collection will be evaluated to explore if *non-iniari* sources of high Fe can be identified to diversify the genetic based of high-Fe materials. A recent validation of advanced breeding lines (seed-parents and restorer parents' progenies) showed that about 128 progenies (81 seed-parents and 47 restorer-parents) were highly adapted to field condition with the Fe density 64-133 mg kg<sup>-1</sup> in seed-parents progenies and 56 - 139 mg kg<sup>-1</sup> in restorer-parents progenies. Of these, 53 seed-parent progenies exceeded 85 mg kg<sup>-1</sup> Fe density and had 46-80 mg kg<sup>-1</sup> Zn density. Commercially available cultivars (not bred for micronutrients) had mean 42 mg kg<sup>-1</sup> Fe [8]. Variability for grain minerals in this crop open the opportunity for breeding high iron breeding lines and hybrid parents and thereby high-iron cultivars for improved human nutrition in millet consuming populations.

## 4. Biofortification Breeding Approach

Pearl millet biofortification breeding program at ICRISAT has taken a three-pronged breeding phase – I, II and III. First phase is short-term approach dealing with traits genetics, germplasm screening and creating genetic variability. Second phase is medium-term approach consisted of validating identified high-iron and zinc breeding lines and hybrid parents from regular breeding program for developing fast-track biofortified variety/hybrids. Third phase consisted of long-term objective- development high-Fe/Zn breeding lines and hybrid parents and its genetic diversification through steady mainstreaming micronutrient traits at ICRISAT and NARS breeding programs. In brief, these micronutrients largely governed by additive genes thus biofortification breeding approaches are same as for as any other quantitative traits. Previous reports also indicate the greater

importance of additive gene action for grain Fe and Zn in pearl millet [12,15,16]. Heritability of these micronutrients was very high and not much influenced by environments [17-19]. Pedigree breeding method is a most common method in pearl millet breeding which is entirely deals with the progenies derived from bi-parental crosses and this method well described by Andrews et al. (1997) [20]. This method also utilizes composites as a base population that has potential to accelerate the genetic gains for yield in hybrids and widen the genetic base of hybrid-parents and cultivars. However, composites will be used for pedigree breeding of hybrid parents only when they fulfil the basic traits required of hybrid parents. Pearl millet has different cytoplasmic sterility systems (CMS), however, A1, A4, A5 are widely used. The adoption of cytoplasmic genetic male sterility under the standard three-line system (A, B, and R) to produce hybrid seed. Currently A1 and A4 exploited under commercial breeding programs in India. All the pearl millet hybrids so far developed in India is based on A1 CMS while biofortification program adventure both A1 and A4 CMS to diversify the new cultivar base. Currently, pearl millet biofortification is in gradual transition from fast-track breeding (breeding phase II) to genetic diversification and mainstreaming (breeding phase III). The development of biofortified fast-track hybrids/varieties at ICRISAT briefly described in **Figure 1**. Currently ICRISAT demonstrated the combining micronutrients and grain yield by conventional breeding approaches [Rai et al. The fast-track approach explained in Figure 1 uses existing moderate to higher Fe/Zn content breeding lines and population which is not bred for micronutrient content. Breeding parental lines (seed- and restorer-parents) bred for high Fe density as a target trait will address the long-term objective of breeding hybrids with even high-Fe levels. In this direction, 174 high-Fe early-generation progenies (B  $\times$  B progenies and R  $\times$  R progenies) have been developed, which in trials conducted at Patancheru had shown  $> 90 \text{ mg kg}^{-1}$  Fe density and 36 to  $72 \text{ mg kg}^{-1}$  Zn density. Based on preliminary study, identified common and overlapping Quantitative Trait Loci (QTL) for Fe and Zn densities in LG3 (chromosome 3) [21]. Since, substantial variation for Fe and Zn available in elite breeding populations, more emphasis is required on developing diagnostic makers for screening segregating materials in future.




**Figure 1.** Fast-Track breeding approach followed at ICRISAT for biofortified hybrid development in India and OPV development in West Africa.

## 5. Biofortified Cultivar Release and Adoption:


In continuation of germplasm screening, identified high-Fe progenies and breeding lines were used to develop initial crosses for building hybrid parents and OPV development, respectively. In the short and medium-term objectives, identified high-Fe hybrid parents and populations contributed to develop high-Fe cultivars for rapid delivery. A first high-Fe pearl millet variety 'Dhanashakti' was developed by utilizing the intra-population variability within ICTP 8203, an early-maturing, large-seeded, disease resistant and high-yielding open-pollinated variety that has been under cultivation in India since 1990. Dhanashakti is officially released and notified by Central Variety Releasing Committee in 2014 for cultivation in all pearl millet growing states of India [22]. Based on national testing trials, Dhanashakti had 71 mg kg<sup>-1</sup> Fe (9% higher) and 2.2 t ha<sup>-1</sup> grain yield (11% higher) compared to original. ICMV 221, another largely grown OPV also under improvement for Fe. The improved version (ICMV221 Fe 11-2) had 70 mg kg<sup>-1</sup> Fe density (11% higher), 58 mg kg<sup>-1</sup> Zn density (9% higher), and grain yield of 3.2 t/ha (5% higher) compared to original variety. This variety reached nearly 90,000 farmers in India and introduction of this biofortified variety in West Africa is in progress [23].

The current breeding efforts in the public and private sector are towards hybrid development and indeed supported by ICRISAT through the development and dissemination of a large number and diverse range of improved breeding lines and hybrid parents through Pearl Millet Hybrid Parent Research Consortia (PMHPRC). By utilizing identified high-Fe hybrid-parents from amongst those initially not bred for high-Fe as a target trait, several high-yielding and high-Fe hybrids has been developed, which are at various stages of testing. One such hybrid has been identified and designated as ICMH 1201, which in 48 field trials during 2011-2013 gave 75 mg kg<sup>-1</sup> Fe density (similar to ICTP 8203) but had 3.6 t ha<sup>-1</sup> grain yields (38% higher than ICTP 8203). ICMH 1201 flowered only 3 days later than ICTP 8203, so it falls in the early-maturity group and fits in the same production system. ICMH 1201 and eight other biofortified hybrids developed at ICRISAT and identified by our partners as promising hybrids have been tested over two years in All India Coordinated Research Project on Pearl Millet (AICRP-PM) trials [24]. At the same time, the commercial production of ICMH 1201 was undertaken by Shaktivardhak Seed Company since 2014 for marketing under its brand name Shakti-1201 [25]. Shakti-1201 has been adopted by 35,000 farmers, in Maharashtra and Rajasthan [24]. Several hybrids with high-Fe density comparable to high-Fe experimental hybrid ICMH 1201, with higher grain yield were identified at various stages of testing. Grain yield of these hybrids were not comparable but close (85-90%) to highest-yielding and high-Fe (56 mg kg<sup>-1</sup> Fe) commercial hybrid 86M86 (from Pioneer), but most of these biofortified hybrids flowered a week earlier than 86M86, hence suitable for medium maturity zone. The first wave of biofortified hybrids are AHB 1200 Fe (ICMH 1202), HHB 299 (ICMH 1203), and Phule Maha Shakti (ICMH 1301) officially released for national level in collaboration with agricultural universities of Maharashtra (VNMKV and MPKV), Haryana (CCSHAU). These biofortified hybrids contain more than 70 mg kg<sup>-1</sup> Fe and 35 mg kg<sup>-1</sup> Zn (Figure 2 and Figure 3).





**Figure 2.** Released and notified biofortified high-yielding hybrids in India.




**Figure 3.** Initiation of biofortified pearl millet cultivars cultivation spread in India (Adapted form HarvetPlus-India).

This biofortification products marks the first milestone in the mainstreaming the grain mineral traits in the cultivar development process and expected to have policy in place for grain mineral bench mark to enable long-term biofortification at all level of hybrid testing and release in India and Africa. With the great support of partners, especially MPKV-Dhule, AICRP-PM, Mondore and Nirmal seeds Pvt Ltd., first high-iron variety Dhanashakti reached regular seed systems that intend annually 200-300 kg of breeder seed to national partners; so far and reached >60,000 farmers in peninsular India. Truthfully labeled seed (TLS) production of Shakti-1201 is being undertaken by Shaktivardhak Seed Company for commercialization and it has been adopted by 20,000 ha, mostly in Maharashtra and Rajasthan. Much greater progress in adopting high-Fe hybrids with high-grain yield is expected in the near future.

## 6. Trade-Off Between Yield and Micronutrients:

The knowledge of relationship between micronutrient and yield traits is prerequisite to combining these two market traits into improved genetic background. Highly positive and significant correlation between Fe and Zn encourage concurrent genetic improvement [9,26-28]. Whereas both micronutrients are negatively and non-significantly associated with grain yield due to genetic architecture of breeding lines and cultivar as they were coming from targeted selections for yield and not targeted for micronutrients [Kanatti et al. 2015; Govindaraj et al Govindaraj et al 2018. Interestingly, the biofortified hybrids which are bred for both traits are reflected no negative association with grain yield [Figure 4]. Indication of this no relationship between the productivity level of the environments and grain yield and micronutrients imply that higher productivity not necessarily lower the grain micronutrients. Further, seed weight is positively correlated with grain micronutrients indicate indirect way of grain nutrients contributing to yield potential of a line or hybrid. Therefore, careful selection should be followed in initial breeding cycles to achieve high-yielding and higher micronutrients.



**Figure 4.** Relationship between Fe/Zn density and grain yield in biofortified hybrid

## 7. Market opportunities:

In India, strong pearl millet breeding programs exist in both the public and private sectors. The commercial market is largely driven by hybrids (90%) with the private sector playing a key role; hence, biofortification initiatives engages both actors in mainstreaming iron pearl millet. Private sector seed companies have strong pearl millet research and development programs for hybrids. To ensure long-term sustainability, seed companies must engage in iron pearl millet breeding and establish their own high-iron product lines. Therefore, the HarvestPlus strategy engages seed companies in genotype-by-environment (GxE) testing of hybrids and inbred lines developed at ICRISAT and encourage companies to develop their own high-iron hybrids for commercialization. Over the past 10 years of crop development with NARS and seed companies, many breeding and product testing challenges are addressed. However, involvement of seed companies is not up to the target to mainstream micronutrients since no policy in place to support biofortified varieties for public pull and profit marketing (**Figure 5**). Therefore, revisiting some of the mainstreaming challenges (yellow status) and addressing lack of policy to pull the nutritious varieties into market are high priority in the future. Private seed companies operate a two-tier distribution system, supplying seed directly to distributors who in turn sell to retailers [29]. Generally, each distributor sells to 40–50 retailers, depending on location and crops/products. These retailers ultimately sell the seeds to farmers. Pearl millet seeds are sold commercially in 1.5 and 3 kg packs, in attractive primary packaging with mandatory labeling (according to India's Seed Act). Demand for iron pearl millet seed is created by farmer demonstrations, field days, and promotions at points of sales. Nirmal Seed's channel partners and personnel have been trained in nutrition messaging for iron pearl millet, a crucial component in the delivery process [23]. Building product acceptance is further facilitated by the agronomic superiority of a recently released high-iron cultivars. Now biofortification programs required to collaborate with various actors in food and retail for demand creation of iron pearl millet grains, flour, and value-added products. This includes testing specific promotional messages and product benefits, communication channels and their effectiveness, and the selection of the brand name and advertising. With this backgrounds, proposed market channel given and will be channelized at the country level (**Figure 6**).

| Task | Key milestone             | Status | Timeline | Remarks                                       |
|------|---------------------------|--------|----------|---|
| 01   | Partnership establishment | Green  | 2011     | Complete strengthening <b>continued-test</b>  |
| 02   | Initial Product testing   | Green  | 2012     | Key resource has been <b>shared</b>           |
| 03   | Advanced product testing  | Yellow | 2013     | Project <b>funding</b> is not sufficient.     |
| 04   | National trial            | Yellow | 2014     | Improved <b>testing and sampling</b>          |
| 05   | Integration and Testing   | Green  | 2015     | Need <b>partners involvement</b>              |
| 06   | Initiate Mainstreaming    | Yellow | 2016     | Need <b>implementing strategy</b>             |
| 07   | Hybrid Release            | Green  | 2017     | Need more <b>products pipelines</b>           |
| 08   | Release & Accept. Stage   | Red    | 2018     | Need <b>policy driven</b> products            |
| 09   | Accept. & consumption     | Red    | ≥2020    | Need <b>policy and market</b> driven products |

**Legend:**


= On schedule/ completed

= problems exist, or re-visiting required

= Schedule has not been, or cannot be sustained (policy)

**Figure 5.** Snap-shot of pearl millet biofortification program challenges and opportunities in developing countries (India and Africa).





**Figure 6.** Proposed scaling up pathway for biofortified grains from varieties and hybrid in India (from production to consumption).

### 7.1. Mainstreaming cultivar policy:

Indian Council of Agricultural Research (ICAR) has endorsed to include the minimum levels of iron and zinc that have to be bred into the future varieties of pearl millet across the country. All-India Coordinated Research Project on Pearl Millet, decided a minimum of 42 mg kg<sup>-1</sup> of iron and zinc 32 mg kg<sup>-1</sup>. Any variety or hybrids to be sold to farmers by public or private seed producers should follow these micronutrient standards apart from giving a higher yield [30]. Dissemination of these biofortified breeding lines and hybrid parents to and their utilization by user-research organizations (both public and private sector) on a continuing basis, as done so far for the non-biofortified materials, will make biofortified hybrid development a matter of routine and hence significantly contribute to improved nutrition. Mainstreaming should be done

### 7.2. Market policy

Unlike other cereals, pearl millet consumed as whole grain and very little processing may require for new products. Processing technologies for pearl millet is yet to be commercialized for improving the edible and nutritional characteristics of millet as well as the grain supply challenges to promote both urban and rural household's utilization. Various forms of food products from industries is key to upscale underutilized nutritious crops for better human health. Climate and nutrition smart millet brings the prospects for millet processing entrepreneurs and consumers' willingness irrespective of the purchase power of rural and urban community.

### 7.3. Smart food

Pearl millet is climate smart crop by itself - dryland resilient with high metabolizable energy, high gluten-free protein, and more of balanced amino acids. So, this crop has potential role to play in the smart food initiative taken by ICRISAT [for more detail visit <https://www.icrisat.org/smartfood/>] which aims to build food systems where the food is good for you (highly nutritious), good for the planet (climate resilient) and good for the smallholder farmer. In general, several products of millets are available in various countries [[http://www.icrisat.org/wp-content/uploads/2018/05/Selected-Millet-and-sorghum-products\\_1.pdf](http://www.icrisat.org/wp-content/uploads/2018/05/Selected-Millet-and-sorghum-products_1.pdf)]. The same market linkage can be tapped for exploring nutri-dense grain and products markets in future.

## 8. Conclusion and Way Forward

Biofortification is scientifically proven sustainable and cost-effective approach to address malnutrition. This approach targets the root cause of the malnutrition. Combining yield and micronutrients highly feasible like other heritable traits provided mainstreaming these traits in the national and international breeding programs. Public and private partners need to take this opportunity to elevate the importance of this nutri-cereal at national and international level. Efficacy studies provides evidences that, consumption of grains from biofortified varieties would provide bioavailable Fe to meet full recommended daily allowance (RDA) in children, adult men and 80% of the RDA in women. In the modern era, genetic biofortification is a way of improving nutrition in staple crops for which enhanced food processing and markets will be the backbone to nutritional security in future. There is no regulatory mechanism in place support biofortified crop varieties and markets is the key for enhancing processed foods as well as related industries. This will also generate market opportunities for farmers. While rice and wheat will continue to play major roles in addressing the problem of food security, biofortified pearl millet have potential to make significant contributions to food-cum-nutritional security in dryland poor households. Biofortified cultivars are being developed through conventional breeding thus grains and foods produced from such cultivars do not face the possible challenge of food regulations and consumer acceptance.

**Author Contributions:** Conceptualization, MG; Writing original draft, MG, AK, BC; Data collection and tables, HS; Review & editing, MG, KN, BC and WHP.

**Acknowledgments:** This work was financially supported by HarvestPlus Challenge Program of the CGIAR.

**Conflicts of Interest:** The authors declare no conflict of interest.

### References:

1. Webb, P.; Stordalen, G.A.; Singh, S.; Wijesinha-Bettoni, R.; Shetty, P.; Lartey, A. Hunger and malnutrition in the 21st century. *BMJ*. **2018**, 361, k2238
2. Kramer, C.V.; Allen, S. Malnutrition in developing countries. *Paediatr. Child Health* **2015**, 25, 422–427.

3. Saltzman, A.; Birol, E.; Oparinde, A.; Andersson, M.S.; Asare-Marfo, D.; Diressie, M.T.; Gonzalez, C.; Lividini, K.; Moursi, M.; Zeller, M. Availability, production, and consumption of crops biofortified by plant breeding: current evidence and future potential. *Ann. N. Y. Acad. Sci.* **2017**, *1390*, 104-114.
4. Yadav, O.P.; Rai, K.N. Genetic Improvement of Pearl Millet in India. *Agric. Res.* **2013**, 1-18.
5. Burton, G.W.; Wallace, A.T.; Rachie, K.O. Chemical Composition and Nutritive Value of Pearl Millet (*Pennisetum typhoides* (Burm.) Stapf and EC Hubbard) Grain 1. *Crop Sci.* **1972**, *12*, 187-188.
6. Parthasarathy Rao, P.; Birthal, P.S.; Reddy, B.V.; Rai, K.N.; Ramesh, S. Diagnostics of sorghum and pearl millet grains-based nutrition in India. *Int. Sorghum Millets Newsl.* **2006**, *47*, 93-96.
7. Basavaraj, G.; Rao, P.P.; Bhagavatula, S.; Ahmed, W. Availability and utilization of pearl millet in India. *SAT eJournal* **2010**, *8*, 1-6.
8. Rai, K.N.; Yadav, O.P.; Govindaraj, M.; Pfeiffer, W.H.; Yadav, H.P.; Rajpurohit, B.S.; Patil, H.T.; Kanatti, A.; Rathore, A.; Rao, A.S.; Shivade, H. Grain iron and zinc densities in released and commercial cultivars of pearl millet (*Pennisetum glaucum*). *Indian J. Agric. Sci.* **2016**, *86*, 291-6.
9. Govindaraj, M.; Rai, K.N.; Pfeiffer, W.H.; Kanatti, A.; Shivade, H. Energy-Dispersive X-ray Fluorescence Spectrometry for Cost-Effective and Rapid Screening of Pearl Millet Germplasm and Breeding Lines for Grain Iron and Zinc Density. *Comm. Soil Sci. Plant Anal.* **2016**, *47*, 2126-2134.
10. Paltridge, N.G.; Milham, P.J.; Ortiz-Monasterio, J.I.; Velu, G.; Yasmin, Z.; Palmer, L.J.; Guild, G.E.; Stangoulis, J.C. Energy-dispersive X-ray fluorescence spectrometry as a tool for zinc, iron and selenium analysis in whole grain wheat. *Plant soil* **2012**, *361*, 261-269.
11. Velu, G.; Rai, K.N.; Muralidharan, V.; Kulkarni, V.N.; Longvah, T.; Raveendran, T.S. Prospects of breeding biofortified pearl millet with high grain iron and zinc content. *Plant Breed.* **2007**, *126*, 182-185.
12. Govindaraj, M.; Rai, K.N.; Shanmugasundaram, P.; Dwivedi, S.L.; Sahrawat, K.L.; Muthaiah, A.R.; Rao, A.S. Combining ability and heterosis for grain iron and zinc densities in pearl millet. *Crop Sci.* **2013**, *53*, 507-517.
13. Rai, K.N.; Gupta, S.K.; Sharma, R.; Govindaraj, M.; Rao, A.S.; Shivade, H.; Bonamigo, L.A. Pearl millet breeding lines developed at ICRISAT: A reservoir of variability and useful source of non-target traits. *SAT eJ.* **2014**, *1*, 1-13.
14. Andrews, D.J.; Kumar, K.A. Use of the West African pearl millet landrace Iniadi in cultivar development. *Plant Genet. Resour. Newsl.* **1996**, *105*, 15-22.
15. Velu, G.; Rai, K.N.; Muralidharan, V.; Longvah, T.; Crossa, J. Gene effects and heterosis for grain iron and zinc density in pearl millet (*Pennisetum glaucum* (L.) R. Br). *Euphytica* **2011**, *180*, 251-259.
16. Kanatti, A.; Rai, K.N.; Radhika, K.; Govindaraj, M.; Sahrawat, K.L.; Rao, A.S. Grain iron and zinc density in pearl millet: combining ability, heterosis and association with grain yield and grain size. *SpringerPlus* **2014**, *3*, 763.
17. Kanatti, A.; Radhika, K.; Padma, V.; Janila, P.; Govindaraj, M.; Rai, K.N. Intra-population variability for grain iron and zinc densities in commercial open-pollinated varieties of pearl millet. *Intern. J. Economic Plants*, **2015**, 171-174.
18. Govindaraj, M.; Rai, K.N.; Shanmugasundaram, P. Intra-population genetic variance for grain iron and zinc contents and agronomic traits in pearl millet. *Crop J.* **2016**, *4*, 48-54.
19. Gupta, S.K.; Velu, G.; Rai, K.N.; Sumalini K. Association of grain iron and zinc content with grain yield and other traits in pearl millet (*Pennisetum glaucum* (L.) R.Br.). *Crop Imp.* **2009**, *36*, 4-7.
20. Andrews, D.J.; Ejeta, G.; Gilbert, M.; Goswami, P.; Anand Kumar, K.; Maunder, A.B.; Porter, K.; Rai, K.N.; Rajewski, J.F.; Reddy, B.V.S.; Stegmeier, W.; Talukdar, B.S. Breeding hybrid parents. In *Proceedings of an International Conference on the Genetic Improvement of Sorghum and Pearl Millet*, Lubbock, Texas, 22-27 September 1996. International Sorghum and Millet Research (INTSORMIL) – International Crops Research Institute for the Semi-arid Tropics (ICRISAT), 1997; pp.173-187.
21. Kumar, S.; Hash, C.T.; Thirunavukkarasu, N.; Singh, G.; Rajaram, V.; Rathore, A.; Senapathy, S.; Mahendrakar, M.D.; Yadav, R.S.; Srivastava, R.K. Mapping quantitative trait loci controlling high iron and zinc content in self and open pollinated grains of pearl millet [*Pennisetum glaucum* (L.) R. Br.]. *Front. Plant Sci.* **2016**, *7*, 1636.
22. Rai, K.N.; Patil, H.T.; Yadav, O.P.; Govindaraj, M.; Khairwal, I.S.; Cherian, B.; Rajpurohit, B.S.; Rao, A.S.; Shivade, H.; Kulkarni, M.P. Variety Dhanashakti (Pearlmillet). *Indian J. Genet.* **2014**, *74*, 405-406.

23. Pfeiffer, W.; Andersson, M.; Govindaraj, M.; Parminder, V.; Cherian, B.; Illona, P.; Magezi, S.; Mulambu, J. Biofortification in Underutilized Staple Crops for Nutrition in Asia and Africa. In *Regional Expert Consultation on Underutilized Crops for Food and Nutritional Security in Asia and the Pacific – Thematic, Strategic Papers and Country Status Reports*. Asia-Pacific Association of Agricultural Research Institutions (APAARI), Bangkok, Thailand, 2018; pp. 70-81.
24. Govindaraj, M.; Rai, K.N. Breeding Biofortified Pearl Millet Cultivars with high iron density. *Indian Farm*. **2016**,65,53-55.
25. Purushottam Singh, S.K.; Uddeen, R. Nutri-farms for mitigating malnutrition in India. In *Biofortification of Food Crops*. Singh, U.; Praharaj, C.S.; Singh, S.S.; Singh, N.P. Springer, India, 2016; pp. 461-477.
26. Velu, G.; Rai, K.N.; Sahrawat, K.L.; Sumalini, K. Variability for grain iron and zinc contents in pearl millet hybrids. *J. SAT Agric. Res.* **2008**,6:1-4.
27. Rai, K.N.; Govindaraj, M.; Rao, A.S. Genetic enhancement of grain iron and zinc content in pearl millet. *Qual. Assur. Safety Crops Foods* **2012**,4,119-125
28. Kanatti, A.; Rai, K.N.; Radhika, K.; Govindaraj, M.; Sahrawat, K.L.; Srinivasu, K.; Shivade, H. Relationship of grain iron and zinc content with grain yield in pearl millet hybrids. *Crop Imp.* **2014**,41,91-96.
29. Karandikar, B.; Smale, M.; Birol, E.; Tedla-Diressie, M. India's Pearl Millet Seed Industry: Prospects for High-Iron Hybrids. HarvestPlus Working Paper 2018, 28,1-41.
30. AICRP Pearl Millet. 2018. *Proceedings of the 53rd Annual Group Meeting of ICAR - All India Coordinated Research Project on Pearl Millet*. <http://www.webcitation.org/76fWm4e89> (accessed on 6-March-2019)