

Original Research Article

Self-Reported Exposure to ETS (Environmental Tobacco Smoke), Urinary Cotinine and Oxidative Stress Parameters in Pregnant Women – the Pilot Study

Lubica Argalasova^{1*}, Ingrid Zitnanova², Diana Vondrova¹, Monika Dvorakova², Lucia Laubertova², Jana Jurkovicova¹, Juraj Stofko³, Michael Weitzman⁴, Martin Simko⁵

¹ Institute of Hygiene, Faculty of Medicine, Comenius University, Bratislava, Slovakia; lubica.argalasova@fmed.uniba.sk (L.A.); diana.vondrova@fmed.uniba.sk (D.V.)

² Institute of Medical Chemistry, Biochemistry and Clinical Biochemistry, Faculty of Medicine, Comenius University, Bratislava, Slovakia; ingrid.zitnanova@fmed.uniba.sk (I.Z.); monika.dvorakova@fmed.uniba.sk; lucia.laubertova@fmed.uniba.sk (L.L.)

³ Institute of Physiotherapy, Balneology and Medical Rehabilitation, University of Ss. Cyril and Methodius in Trnava, Slovakia; juraj.stofko@gmail.com (J.S.)

⁴ Department of Pediatrics, New York University, New York, NY, 10016, USA; Michael.Weitzman@nyulangone.org (M.W.)

⁵ IInd Gynecology and Obstetrics Clinic, Faculty of Medicine, Comenius University Bratislava, Slovakia; cyklomartin@gmail.com (M.S.)

* Correspondence: lubica.argalasova@fmed.uniba.sk; Tel.: (optional; include country code; if there are multiple corresponding authors, add author initials) +421-905-209-114 (L.A.)

Abstract: Background: Exposure to ETS (Environmental Tobacco Smoke) is one of the most toxic environmental exposures. Objective: To investigate the impact of ETS on physiological, biochemical, psychological indicators, on the urine antioxidant capacity (AC) and oxidative damage to lipids in a pilot sample of healthy pregnant women. Methods: The exposure to ETS was investigated by a validated questionnaire, urine cotinine and the marker of oxidative damage to lipids - 8-isoprostanate concentrations using an ELISA kit. Urine AC was determined by the spectrophotometric TEAC method. From the sample of pregnant women (n=319, average age 30.84 ± 5.09 years) in 80 the levels of cotinine and oxidative stress markers were analyzed. Results: From our sample, 5 % individuals (7.4 % objectified by cotinine) were current smokers and 25 % reported passive smoking in the household (18.8 % objectified by cotinine). The Kappa was 0.78 for smokers and 0.22 for ETS exposed non-smokers. Smokers as well as non-smokers had significantly higher ($p<0.05$) urine AC than ETS exposed non-smokers. Non-smokers had significantly lower levels of 8-isoprostanate than smokers ($p<0.01$) and ETS-exposed non-smokers ($p<0.05$). Correlations between urine levels of cotinine and AC were positive in ETS exposed non-smokers. Conclusion: The harmful effect of active and passive smoking on oxidative stress parameters has been indicated.

Keywords: Environmental Tobacco Smoke (ETS); pregnant women; questionnaire; urinary cotinine; oxidative stress parameters

1. Introduction

Exposure to environmental risk factors has a negative impact on health, especially in vulnerable population groups, which include the children, mothers and pregnant women. Exposure to tobacco smoke is one of the most toxic environmental exposures. Globally, more than a third of all people are regularly exposed to the harmful effects of smoke. This exposure is responsible for about 600,000 deaths per year, and about 1% of the global burden of diseases worldwide [1]. Around the world, 40% of children, 33% of male non-smokers, and 35% of female non-smokers were exposed to ETS in 2004 [2]. According to the Global Adult Tobacco Survey (GATS) (2008-2010), which investigated the prevalence of smoking and passive smoking among women aged 15-49 years in 14 low- and middle-income countries, the prevalence was 0.4% in Egypt, 30.8% in Russia, 17.8% in Mexico and 72.3% in Vietnam. In Poland 26.9% of women smoke, 45.4% are exposed to ETS at home and 24.3% at work. Slovakia and the Czech Republic did not take part in this survey [3]. According to the WHO, the prevalence of daily adult tobacco smokers in Slovakia in 2016 was 29%, 24% of women and 34% of men [4].

Diseases arising from smoking are referred to as „smoking-related diseases“. These include tumors (lips, throat, esophagus, colon, kidney, bladder, liver, lung), non-cancerous respiratory system diseases, cardiovascular diseases and many other diseases affecting a wide variety of organ systems that increase the morbidity, mortality, shorter life expectancy, and worse quality of life [5-10]

Smoking, however, also affects non-smokers in households and public places, where smoking is allowed [11-13]. Environmental tobacco smoke (ETS) exposure, defined as smoke emitted from the burning end of a cigarette or cigar or exhaled by a smoker, represents a well-established and significant health risk [11, 12]. Recent studies demonstrate that ETS is composed not only of second-hand smoke (SHS) but also of third-hand smoke (THS). Third-hand smoke is a complex phenomenon resulting from residual tobacco smoke pollutants that adhere to the clothing and hair of smokers and to surfaces, furnishings, and dust in indoor environments. Exposure can even take place long after smoking has ceased, through the close contact with smokers and in indoor environments in which tobacco is regularly smoked [14, 15].

There have been many studies pointing to the harmful effects of passive smoking on exposed groups of adults, children, pregnant women and their fetuses [13, 16-20].

The most serious complications of ETS in pregnancy include spontaneous abortion, preterm birth fetal developmental anomalies, ectopic pregnancy, preterm labor, intrauterine growth retardation of the fetus (IUGR), fetal death, sudden infant death syndrome (SIDS) [11, 12, 21-24]. Newborns exposed to cigarette smoke during pregnancy are more affected by neurological disorders with long-term deterioration in behavioral, emotional and cognitive functions at a later age [25-27].

Tobacco smoke contains toxic, carcinogenic and mutagenic chemicals as well as free radicals and reactive oxygen species with the potential of oxidative damage to biomolecules. The increased production of reactive oxygen species is related to the depletion of antioxidants and the formation of oxidative stress in the organism [28, 29]. As a result, lipid oxidation, cell membrane damage, DNA strand breaks and the inactivation of some enzymes may occur [30]. Exposure of pregnant women to tobacco smoke causes oxidative stress not only in pregnant women but also in their fetuses [31, 32]. Nicotine and its major metabolite cotinine (the most common biomarker for exposure to cigarette smoke assessed in hair, serum or urine) have high lipid solubility; therefore, they pass rapidly through the placenta into the fetal circulation, with higher levels of cotinine recorded in the fetus than in the mother's plasma [33-37].

The aim of this international and interdisciplinary project is to assess the degree of ETS exposure and its impact on physiological, biochemical and psychological indicators and on the urine antioxidant capacity and oxidative damage to lipids in a pilot sample of healthy pregnant women. The specific aim is to determine the extent to which self-reported smoking and exposure to ETS are in agreement with the levels of nicotine metabolite (urinary cotinine). The benefit of the study will be the development of the basis for primary preventive interventions in clinical and preventive practice.

2. Materials and Methods

Researchers from the Comenius University's Obstetrics and Gynecology (OB/Gyn) Department and Institute of Hygiene in Bratislava, Slovakia distributed surveys to pregnant women in the 36th – 41st week of pregnancy being seen for the follow-up at the OB/Gyn Department of the Faculty Hospital and Clinic. This survey is the continuation and re-analysis of the previous survey that was designed to evaluate environmental, behavioral, and psychosocial factors in the lives of women [20]. The results of the study have shown that ETC exposure is an independent risk factor associated with the worse physical health of non-smoking mothers in the reproductive age and the worse mental health in the smaller sample of pregnant women [20]. In the present study, we have enlarged the sample of the pregnant women and objectified the self-reported smoking and ETS exposure by the levels of urinary cotinine. The study was approved by the Ethical Committee of the Faculty of Medicine, Comenius University Bratislava, Slovakia and by the Institutional Review Board of New York University School of Medicine, New York, U.S.A (IRB number: 09-0331).

In the present study 319 (average age 30.84 ± 5.09) healthy pregnant women without any medical treatment were included and in 80 of them (average age 30.24 ± 4.92 years) the levels of cotinine and oxidative stress markers in urine specimens were analyzed from March to June 2018. Exposure to tobacco smoke as well as the analysis of the lifestyle and demographic determinants of passive smoking were assessed by the validated Questionnaire for mothers used in the previous study [20]. Based on the obtained data we have evaluated the environmental, behavioral and psychosocial factors in the mother's life. For the verification and objectification of women's exposure to tobacco smoke, the levels of urinary cotinine were evaluated [38].

Urine specimens were taken at the routine control into plastic containers that were subsequently frozen at -20°C . In the urine samples, levels of cotinine and oxidative stress marker (8-isoprostanates) were analyzed within 3 months of sampling.

2.1 Sample

In the sample of healthy pregnant women (n=319) 79.9 % were younger than 35 years old, most of Slovak nationality (94.3 %), 78.2 % were married or in a relationship, 50.5 % graduated from the university, 60.6 % were employed, 57.4 % had children under 18 years of age in their household (Table 1). In the sample of healthy pregnant women in whom we analyzed the levels of cotinine and oxidative stress markers in urine specimens (n=80), 81.20 % were younger than 35 years old, most of Slovak nationality (93.8 %), 78.8 % were married or in a relationship, 68.80 % graduated from the college, 66.3 % of mothers employed, 31.3 % had children under 18 years of age in their household (Table 2).

2.2 Questionnaire

The validated "Questionnaire for Mothers" administered by a trained person, contained questions on environmental, behavioral and psychosocial factors in the life of pregnant women. Besides questions on personal (age, nationality, marital status, education, employment, children), behavioural (smoking, lifestyle, nutrition), housing (residence) and economic characteristics (household income), it also included questions on mothers' smoking and ETS exposure in the

household (smoking spouse or other members of the family, number of cigarettes and number of years of smoking). In the case of a former smoker, there was a question for how many years she/he has not smoked. Former smokers were considered non-smokers.

Table 1. Characteristics of a sample of pregnant women (n=319)

Indicator*	N	%
Age group		
≤ 35	255	79.9
> 35	64	20.1
Nationality		
Slovak	299	94.3
other	18	5.7
Marital status		
married/in a relationship	248	78.2
single/divorced	69	21.8
Number of children under 18		
no	100	42.6
1	107	45.5
2	24	10.2
≥ 3	4	1.7
Mother's education		
secondary school or lower	42	13.2
high school graduate	116	36.4
university degree	161	50.5
Employment status of the mother		
employed	191	60.6
unemployed	124	39.4
Father's education		
secondary school or lower	63	20.0
high school graduate	118	37.5
university degree	134	42.5
Employment status of the father		
employed	307	98.4
unemployed	5	1.6
Household income		
≤ 700 €	62	20.1
> 700 €	246	79.9
Residence		
urban-metropolitan area	229	72.2
rural-non-metropolitan area	88	27.8
Physical activity		
regular	129	41.1
irregular	185	58.9

Healthy lifestyle		
yes	207	65.9
no/not sure	107	34.1
Number of daily meals		
≤ 4	192	60.4
> 4	126	39.6
Smoking status (self-reported)		
non-smoker	187	58.6
ex-smoker	103	32.3
current smoker	29	9.1
Exposure to tobacco smoke (self-reported)^a		
not exposed	156	62.2
exposed	95	37.8

^a If somebody living in the household is smoking. ^b There are some data missing in each variable category

Table 2. Characteristics of a sample of pregnant women with urinary cotinine and oxidative stress parameters (n=80)

Indicator*	N	%
Age		
≤ 35	65	81.2
> 35	15	18.8
Nationality		
Slovak	75	93.8
other	5	6.2
Marital status		
married/in a relationship	63	78.8
single	17	21.3
Number of children under 18		
no	55	68.7
1	20	25.0
2	5	6.3
≥ 3	0	0.0
Mother's education		
secondary school or lower	6	7.5
high school graduate	19	23.7
college graduate and higher	55	68.8
Employment status of the mother		
employed	53	66.3
unemployed	27	33.7
Father's education		
secondary school or lower	10	12.6
high school graduate	21	26.6

college graduate and higher	48	60.8
Employment status of the father		
employed	78	98.7
unemployed	1	1.3
Household income		
≤ 700 €	5	6.5
> 700 €	72	93.5
Residence		
urban-metropolitan area	57	72.2
rural-non-metropolitan area	22	27.8
Physical activity		
regular	40	50.6
irregular	39	49.4
Healthy lifestyle		
yes	54	67.5
no/not sure	26	32.5
Number of daily meals		
≤ 4	43	54.5
> 4	36	45.5
Smoking status (self-reported)		
non-smoker	59	73.8
ex-smoker	17	21.2
smoker	4	5.0
Smoking status (cotinine objectified)		
no	74	92.6
yes	6	7.4
Exposure to tobacco smoke (self-reported)^a		
not exposed	56	70.0
exposed	20	25.0
Exposure to tobacco smoke (cotinine objectified)^a		
not exposed	59	73.8
exposed	15	18.8

^a If somebody living in the household is smoking. * There are some data missing in each variable category

2.3 Chemical analyses

2.3.1 Cotinine

The level of cotinine was measured in urine samples using a competitive ELISA kit (MyBioSource, San Diego, CA, USA) according to the manufacturer's instructions. Obtained results were expressed in mg/mol of creatinine. Pregnant women were assigned into three experimental groups based on the urine cotinine levels: 58 women with cotinine levels above 2 mg/mol creatinine were included in the smoker group (S), 15 women with cotinine levels between 0.06 - 2 mg/mol

creatinine into the ETS group (environmental tobacco smoke) and 7 women with cotinine levels below 0.06 mg/mol creatinine were included into the non-smoker group (NS).

2.3.2 Antioxidant capacity of urine (TEAC)

Trolox equivalent antioxidant capacity (TEAC) decolorization assay (Re et al. 1999) is a decolorization method applicable for both the lipophilic and hydrophilic antioxidants. A cation radical 2,2'-azino-bis-3-ethyl benzothiazoline-6-sulfonic acid (ABTS⁺) is produced by the oxidation of ABTS with potassium persulfate (K₂S₂O₈). Added antioxidants reduce the cation radical in a dose- and time-response manner. Decolorization of the cation radical is related to the standard trolox (synthetic, water-soluble form of vitamin E). Results are expressed in mmol of trolox/L/mol of creatinine.

2.3.3 8-isoprostane

Isoprostane (8-iso prostaglandin F₂ α) levels in urine were determined by the commercial competitive ELISA kit (Cayman Chemical, USA) following manufacturer's instructions. Results are expressed in ng/mL/mmol of creatinine.

2.3.4 Creatinine

Urine creatinine was determined in the certified laboratory (Medirex, a.s., Bratislava, Slovakia)

2.4. Statistical analysis

To evaluate the results, we used the methods of descriptive and analytical statistics (categorical data analysis) to identify mutual associations between factors assessed in the questionnaire and self-reported ETS exposure. Kappa statistics, sensitivity, specificity and correlations were used to determine the extent to which self-reported smoking and exposure to ETS are in agreement with the degree of ETS exposure determined by the levels of urinary cotinine (i.e. to determine the accuracy of self-reported smoking status). Kappa is the percentage of cases in which the two measures are in agreement after accounting for chance agreement [39]. It does not take into account which measure is considered the gold standard. Sensitivity is the percentage of true positive (the percentage of respondents who reported being smokers or ETS exposed non-smokers among those classified as smokers or ETS exposed non-smokers based on cotinine concentrations). Specificity is the percentage of true negatives (the percentage of respondents who reported being non-smokers among those classified as non-smokers based on cotinine concentrations). The predictive value positive (PVP) and predictive value negative (PVN) are the complements of the percent false positive and percent false negative, respectively [38, 40, 41]. Statistical package SPSS, version 24 (International Business Machines Corp.; New Orchard Road; Armonk, New York, USA) was used for the data analysis.

To evaluate the results of chemical analysis the statistical package SPSS ver. 18 (SPSS Inc., Chicago, IL, USA) was used. The results are expressed as mean \pm standard deviation (SD) for normally distributed data, or median (lower quartile – upper quartile) for data not normally distributed. The Student's unpaired t-test or non-parametric Mann-Whitney test were used for the comparison between groups of continuous parameters as appropriate. To quantify the association between two variables, Pearson or Spearman correlations were used.

The significance level was set at $p < 0.05$.

3. Results

In the sample of 319 healthy pregnant women, there were 58.6% (187) self-reported non-smokers, 32.3 % (103) ex-smokers and 9.1% (29) current smokers smoking from one to 15 cigarettes a day. The average number of cigarettes was 6.66 ± 4.16 per day; median 5 (lower quartile 3 – upper quartile 10); the average duration of smoking was 8.90 ± 5.46 years; median 10 (4– 12). Current smokers were excluded from the analysis. ETS exposure (somebody living in the household is smoking) reported 37.8 % (95) non-smoking respondents. The average number of cigarettes smoked by the partner/person living in the same household was 13.12 ± 8.12 per day; median 11.5 (7– 20). The average duration of smoking was 12.81 ± 6.38 years (Table 1). In the analysis of the lifestyle and demographic determinants of passive smoking in the household significant negative relationships between ETS and the level of mother's and father's education ($p < 0.001$) and household income ($p < 0.05$) were found. ETS exposed non-smoking pregnant women live mostly in the urban/metropolitan area, have reportedly worse healthy lifestyle ($p < 0.05$) and indicated lower physical activity ($p = 0.057$) (Table 3).

Table 3. The relation between demographic factors and mother's exposure to tobacco smoke (self-reported)

Indicator ^a	ETS-		ETS+		p-value
	N	%	N	%	
Age group					
≤ 35	117	75.0	78	82.1	n.s.
> 35	39	25.0	17	17.9	
Nationality					
Slovak	148	94.9	91	95.8	n.s.
other	8	5.1	4	4.3	
Marital status					
married/in a relationship	127	81.9	72	76.6	n.s.
single/divorced	28	18.1	22	23.4	
Number of children					
any	50	40.0	31	40.8	n.s.
1-2	73	58.4	43	56.6	
≥ 3	2	1.6	2	2.6	
Mother's education					
secondary school or lower	12	7.7	21	22.1	<0.001
high school graduate	48	30.8	44	46.3	
university degree	96	61.5	30	31.6	
Employment status of the mother					
employed	100	65.8	56	58.9	n.s.
unemployed	52	34.2	39	41.1	
Father's education					
secondary school or lower	19	12.3	32	33.7	<0.001

high school graduate	58	37.7	42	44.2	
university degree	77	50.0	21	22.1	
Employment status of the father					
employed	149	98.0	93	98.9	
unemployed	3	2.0	1	1.1	n.s.
Household income					
≤ 700 €	23	15.3	24	26.1	
> 700 €	127	84.7	68	73.9	<0.05
Residence					
urban-metropolitan area	114	73.1	56	60.2	
rural-non-metropolitan area	42	26.9	37	39.8	<0.05
Physical activity					
regular	70	44.9	30	32.6	
irregular	86	55.1	62	67.4	0.057
Healthy lifestyle					
yes	112	72.3	54	57.4	
no/not sure	43	27.7	40	42.6	<0.05
Number of daily meals					
≤ 4	97	62.2	60	63.8	
> 4	59	37.8	34	36.2	n.s.

ETS+ exposed to tobacco smoke (self-reported); ETS- not exposed to tobacco smoke (self-reported); p<0.05 is considered as statistically significant *There are some data missing in each variable category

In the sample of healthy pregnant women in whom we analyzed the levels of cotinine and oxidative stress markers in urine specimens (n=80), there were 5 % (4) self-reported smokers, 73.8 % (59) non-smokers and 21.2 % (17) ex-smokers. The average number of cigarettes was 5.67 ± 4.04 per day; median 5 (2–5), the average duration of smoking was 14.50 ± 7.78 years; median 14.50 (9.0–14.5). ETS exposure (somebody living in the household is smoking) reported 25 % (20) non-smoking respondents. The presence of ETS exposure objectified by cotinine was confirmed in 18.8 % (15) respondents. The average number of cigarettes smoked by the partner/person living in the same household was 10.39 ± 6.50 per day; median 10 (5–16.3). The average duration of smoking was 13.43 ± 5.90 years (Table 2). There were 5 % (4) self-reported current smokers and 7.4 % (6) current smokers objectified by the level of cotinine in the urine sample and 25 % (20) self-reported ETS exposed non-smokers and 18.8 % (15) ETS exposed non-smokers confirmed by the level of cotinine in the urine sample. The sensitivity for self-reported smoking status was 66.7 %, specificity 100 %, positive predictive value 100 % and negative predictive value 95.8 %. Kappa was 0.78 indicating the substantial agreement [42] or excellent agreement [39]. The sensitivity for self-reported ETS exposure was 46.7 %, specificity 78 %, positive predictive value 35 %, negative predictive value 85.2 %. Kappa was 0.22 indicating the fair agreement [42] or the poor agreement [39]. The agreement for self-reported ETS exposure was better for women from the younger age group (≤ 35 yrs) and with lower education reaching to moderate or fair to good agreement (Kappa=0.44) [39, 42].

Table 4. Measures of agreement to determine the accuracy of self-reported smoking status and exposure to ETS in the sample of pregnant women (n=80)

Measures of agreement	Smoking status			
	Non-smoker vs. current smoker		ETS- vs. ETS+	
	Total	Total	Younger age group (≤ 35 yrs)	Lower education
Kappa	0.78	0.22	0.30	0.45
Spearman correlation	0.80	0.22	0.29	0.44
Sensitivity	66.7%	46.7%	54.6%	66.7%
Specificity	100.0%	78.0%	79.6%	80.0%
Positive predictive value	100.0%	35.0%	37.5%	57.1%
Negative predictive value	95.8%	85.2%	88.6%	85.7%
Diagnostic accuracy	96.2%	71.6%	75.0%	76.2%

ETS+ exposed to tobacco smoke; ETS- not exposed to tobacco smoke

The median value of cotinine in ETS exposed pregnant women was 0.22 (0.129-0.338) and in currently smoking pregnant women 253.19 (181.82-498.31) mg/mol creatinine. The urine antioxidant capacity (TEAC) mean value in ETS exposed pregnant women was 0.91 ± 0.28 and 1.3 ± 0.43 mmol trolox/L/mol creatinine in current smokers; median values of isoprostanes 258.41 (112.26-411.88) in ETS exposed and 293.74 (250.17-377.51) ng/mL/mmol creatinine in currently smoking pregnant women (Table 5).

Pregnant women in the ETS+ group had significantly reduced urine antioxidant capacity (TEAC) compared to both the non-smoker (ETS-) and the smoker groups (Tables 5 and 6). There was no significant difference in urine antioxidant capacity between the non-smokers and the smokers. The marker of oxidative damage to lipids - 8-isoprostanes were significantly increased in the ETS+ and the smoker group compared to the non-smoker group. 8-isoprostane levels were the highest in the smoker group; however, there was no significant difference between ETS+ and smoker groups.

Significant positive correlation between urine cotinine levels and urine antioxidant capacity (TEAC) in the ETS exposed group was found (Table 7). The same correlation was negative in the non-smoker group; however, this correlation was marginally significant.

Table 5. Cotinine levels, TEAC and levels of 8-isoprostanes in the analyzed groups

	Cotinine	TEAC	8-isoprostanes
Smoking status	mg/mol creatinine	mmol trolox/L/mol creatinine	ng/mL/mmol creatinine
ETS-	0.00±0.00	1.2±0.4	143.6(73.91-197.54)
ETS+	0.22(0.129-0.338)	0.91±0.28	238.41(112.26-411.88)
current smoker	253.19(181.82-498.31)	1.3±0.43	293.74(250.17-377.51)

ETS+ exposed to tobacco smoke; ETS- not exposed to tobacco smoke (cotinine objectified). Results are expressed as the mean ± SD or the median (lower quartile – upper quartile).

Table 6. Statistical significance (p-value) of TEAC and levels of 8-isoprostanes between the analyzed groups

Smoking status	TEAC	8-isoprostanes
ETS+ vs. ETS-	0.0105*	0.0487*
ETS+ vs. current smoker	0.0199*	0.4702
current smoker vs. ETS-	0.7374	0.0055*

ETS+ exposed to tobacco smoke; ETS- not exposed to tobacco smoke (cotinine objectified); * significant at p<0.05

Table 7. Correlations between cotinine levels in urine and oxidative stress parameters in the analyzed groups

Antioxidant parameters	Rho	p-value
ETS-		
TEAC	-0.2036	0.0642
isoprostanes	0.0676	0.3097
ETS+		
TEAC	0.7607	0.0007*
8-isoprostanes	-0.2179	0.2171
Current smoker		
TEAC	-0.0857	0.4014
8-isoprostanes	-0.5429	0.1208

ETS+ exposed to tobacco smoke; ETS- not exposed to tobacco smoke (cotinine objectified); Rho - Spearman's rank correlation coefficient; * significant at p<0.05

4. Discussion

The results of our previous studies show that ETS is one of the most important health hazards influencing the physical and mental health of the exposed non-smoking partners [20]. The study published by the members of our research team on a nationally representative data from the year 2000 to 2004 Medical Expenditure Panel Survey in the USA [18] showed a relationship between living with smokers and worsened maternal physical and mental health in non-smoking mothers with children. The risk was discernible with the presence of a single adult smoker in a household and increased with the number of smokers [18]. The limitation of our previous studies was the fact that the smoking status was ascertained via self-reporting. Since there is a considerable public awareness about the effects of cigarette smoking and ETS exposure on humans, participants might be motivated

to under-report their smoking status although there is evidence in some studies to show that self-report is an accurate way to measure smoking behaviors [38, 43, 44].

The problem might be the ETS exposure of pregnant respondents and the motivation to under-report [45, 46, 47] or over-report their exposure (we have not found a study on pregnant women over-reporting their ETS exposure).

The strength of our present study is the determination of the accuracy of self-reported smoking and ETS exposure status by urinary cotinine and investigation of the impact of ETS besides physiological, biochemical, and psychological indicators on the urine antioxidant capacity (AC) and oxidative damage to lipids. Active smoking of pregnant women or ETS exposure results in several problems such as intrauterine growth retardation, an increased risk of spontaneous abortion, reduction of pulmonary function in healthy neonates or a higher risk of sudden infant death syndrome [22]. One of the mechanisms explaining these effects is the presence of the smoke-induced oxidative stress leading to the oxidative damage to molecules and to the inflammatory response [48]. The cigarette smoke contains a large number of free radicals as well as metals such as copper, mercury and zinc [49], which may catalyze the production of the very reactive hydroxyl radical by the Fenton reaction [50]. Smoking may increase oxidative stress not only through the generation of free radicals but also through the depletion of the antioxidant systems protecting the organism against deleterious effects of oxygen radicals.

In our study we have examined the effect of the ETS exposure and the active smoking on the oxidative damage to lipids and on the antioxidant capacity of urine in pregnant women. In the past decades, numerous studies have shown that 8-isoprostanes are extremely accurate markers of lipid peroxidation [51, 52, 53, 54]. 8-isoprostanes are compounds produced by the non-enzymatic oxidation of arachidonic acid. We have found that pregnant women exposed to ETS had significantly higher oxidative damage to lipids and significantly lower urine antioxidant capacity than non-smokers. These results indicate that ETS-exposed pregnant women are under increased oxidative stress which is in accord with other studies [55, 56, 57]. Smoking pregnant women had 8-isoprostanes level similar to the ETS group and antioxidant capacity similar to the non-smokers. In the smoker group compared to the ETS and the non-smoker groups women are exposed to the higher load of oxidants, which may stimulate compensatory mechanisms leading to the increased antioxidant capacity. Results of other studies on oxidative stress of smoking pregnant women are ambiguous. Similar results were reported also in plasma and saliva by other studies [28, 32, 58]. In contrast, Fayol et al (2005) have detected higher plasma antioxidant activity in ETS exposed pregnant women than in controls [59].

In addition, we have observed a strong, significant, positive correlation between the urine antioxidant capacity and the urine cotinine levels only in the ETS+ group. ETS exposed pregnant women might be sensitive to tobacco smoke and able to correspondingly stimulate their antioxidant system. However, in the smoker group this correlation was non-significantly negative which might be the consequence of the higher use of antioxidant compounds by the fetus in order to counteract the increased oxidative burden in active smokers.

ETS exposure or active smoking of pregnant women can have negative effects on their fetuses. There are several reports providing evidence of increased oxidative damage to lipids, DNA and proteins in the blood of such neonates (Kurt et al, 2016) and the correlations between oxidative stress parameters of pregnant women and their neonates [60]. Increased oxidative damage to important biomolecules in fetus caused by cigarette smoke has been implicated in the etiopathogenesis of over

100 disorders [59]. Increased consumption of dietary antioxidants might be a potential therapeutic means against increased oxidative stress in ETS exposed pregnant women and actively smoking pregnant women.

The validity of self-reported smoking in population surveys remains an important question [44]. In our study self-report seems to be in the best agreement with the self-reported smoking status (78 % agreement). Sensitivity was 66.7% and specificity 100 %. There are studies with higher sensitivities, but using larger samples [38, 44, 61]. Self-reported non-smokers who seem to be smokers based on biochemical measurements are generally considered “deceivers” of their true smoking status [43]. In a summary of studies in which questionnaire responses regarding smoking status were compared with cotinine measurements, the estimated misclassification rates (proportion of self-reported non-smokers with increased cotinine levels indicative of active smoking) ranged from 0.9% to 9.8% [43, 44, 62]. Misclassification rates reported among pregnant women may be as different as 3% in a population-based survey and 26.2% in a smoking cessation trial [43, 46, 63]. In our study the misclassification rate for pregnant smokers was 3.9 %.

The agreement for ETS exposed pregnant non-smokers is much lower (22 % agreement, 46.7 % sensitivity and 78 % specificity). The misclassification rate for under-reported ETS exposure was 10.81 %, but for over reported ETS exposure 17.57 %. The older and more educated respondents seem to overestimate their ETS exposure (Table 4). The pregnancy itself may also play a role in overestimation of ETS exposure. The analysis of the lifestyle and demographic determinants on a larger sample of 319 pregnant non-smokers revealed negative relationships between ETS and the level of mother’s and father’s education ($p<0.001$) and household income ($p<0.05$) similar to the other studies [18, 20].

The main conclusion of several studies on large population samples is that the validity of self-reported smoking is consistently high in population-based studies and therefore the extended use of cotinine measurements for validation purposes may not be justified [38, 43, 44]. Nevertheless, further research may focus on assessing the optimal cut off point for validating smoking status among specific groups, such as pregnant women. These studies will also improve our understanding of the effects of gender, social conditions, and pregnancy status on the metabolism of nicotine and on smoking behaviors that may affect nicotine intake.

Findings of several studies suggest that most pregnant women disclose their smoking and ETS exposure as well. Universal urinary cotinine screening of pregnant women could aid in appropriately counseling women about second-hand exposure, as well as monitoring women at high risk for adverse pregnancy outcomes [64, 65]. In contrast, there is a substantial within-person fluctuation in pregnancy smoking, as women try repeatedly to quit or cut down. In this case, cotinine measures may be of limited use for validation of amount smoked, as they are informative only about a recent exposure, vary with individual smoking topography and are dependent on the time elapsed since the last cigarette smoked [61]. The results of the study by Xiao, 2018 on rural pregnant women indicated that, regardless of trimester, more than 15% of pregnant women with actual exposure to ETS might not perceive themselves as passive smokers in prenatal care, especially in the first trimester [66]. The third trimester of pregnancy was the proper period to follow our respondents.

The limitation of our study is the small sample size, the cross-sectional design, and self-reported ETS exposure in the larger sample. In our study, we did not use the medical outcomes short form-12 (SF-12) to quantify the mental and physical health of mothers because pregnancy itself could

influence mental and physical health as well. Especially, physical health is very much influenced by pregnancy and its analysis by SF-12 could be biased [20, 66]. The strength of our study is the separation of current smokers, ETS exposed and ETS not exposed non-smokers and the investigation of the harmful effects of active and passive smoking on detected oxidative stress parameters.

5. Conclusions

Data from our study show that maternal cigarette smoking and ETS exposure during pregnancy may compromise the balance between reactive oxygen species (ROS) and antioxidant defense and can cause potent oxidative stress with all negative consequences in pregnancy.

Combining the maternal self-report of smoking with the level of urine cotinine concentration could improve the precision of the exposure to tobacco smoke. Urine testing for cotinine may be useful in reducing the nondisclosure surrounding prenatal tobacco use. This screening could be a valuable tool for counseling to help pregnant women in tobacco smoking cessation. The presented results might be used in clinical practice and in campaigns for smoke-free environments and in the promotion of community-based smoke-free programs. Furthermore, they represent an important argument for intervention in families. A complete smoking ban at home should be considered to avoid potential adverse effects on pregnancy outcomes due to ETS.

Author Contributions: L.A., I. Z., M.S., M.W. conceived and designed the project, L.A., I. Z., M.S. wrote the manuscript, L.A., D.V., I.Z. analyzed the data, I.Z., M.D., L.L., did the chemical laboratory analysis, J.J., M.W. provided critical revision of the manuscript, M.S., I.Z., L.A., J.J. performed literature search and drafted sections of the manuscript, J.S. was responsible for funding acquisition and project administration, M.W. provided critical revision of the manuscript and did the language, style and spell check.

All authors have approved the submitted version.

Funding: This manuscript was partially supported by the grant Y.A.B.S. (Youth and Parents Behavioral Survey in Slovakia) O-15-101-/0001-00, by the ESFOPV project, MPH study program development at Comenius University in Bratislava in English language (Master of Public Health) ITMS code of the project: 261402300093.

Acknowledgments: We thank the staff of IInd Gynecology and Obstetrics Clinic, Medical Faculty, Comenius University Bratislava, Slovakia helping us with the performance of the project.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. World Health Organization. *Global Health Observatory Data. Second-Hand Smoke*. 2017. Available online: http://www.who.int/gho/phe/secondhand_smoke/en/ (accessed on 22 January 2019).
2. Oberg, M.; Jaakkola, M.S.; Woodward, A.; Peruga, A.; Pruess-Ustuen, A. Worldwide burden of disease from exposure to second-hand smoke: a retrospective analysis of data from 192 countries. *Lancet* **2011**, *377*, 139–146, doi:10.1016/s0140-6736(10)61388-8.
3. Centers for Disease Control and Prevention. Current tobacco use and secondhand smoke exposure among women of reproductive age-14 countries, 2008-2010. *MMWR Morb. Mortal. Wkly. Rep.* **2012**, *2*, 877–882.
4. World Health Organization. *WHO Report on the Global Tobacco Epidemic, 2017: Monitoring Tobacco Use and Prevention Policies*; WHO: Geneva, Switzerland, 2017.
5. Ezzati, M.; Lopez, A.D. Estimates of global mortality attributable to smoking in 2000. *Lancet* **2003**, *362*, 847–852, doi:10.1016/s0140-6736(03)14338-3.

6. Teo, K.K.; Ounpuu, S.; Hawken, S.; Pandey, M.R.; Valentin, V.; Hunt, D.; Diaz, R.; Rashed, W.; Freeman, R.; Jiang, L., et al. Tobacco use and risk of myocardial infarction in 52 countries in the INTERHEART study: a case-control study. *Lancet* **2006**, *368*, 647–658, doi:10.1016/s0140-6736(06)69249-0.
7. Jacob, V.; Vellappally, S.; Smejkalova, J. The influence of cigarette smoking on various aspects of periodontal health. *Acta Medica (Hradec Kralove)* **2007**, *50*, 3–5.
8. Jang, A.-Y.; Lee, J.-K.; Shin, J.-Y.; Lee, H.-Y. Association between Smoking and Periodontal Disease in Korean Adults: The Fifth Korea National Health and Nutrition Examination Survey (2010 and 2012). *Korean J. Fam. Med.* **2016**, *37*, 117–122, doi:10.4082/kjfm.2016.37.2.117.
9. Kasajova, P.; Holubekova, V.; Mendelova, A.; Lasabova, Z.; Zubor, P.; Kudela, E.; Biskupska-Bodova, K.; Danko, J. Active cigarette smoking and the risk of breast cancer at the level of N-Acetyltransferase 2 (NAT2) gene polymorphisms. *Tumor Biol.* **2016**, *37*, 7929–7937, doi:10.1007/s13277-015-4685-3.
10. Santoro, A.; Prinzi, G.; Lamonaca, P.; Cardaci, V.; Fini, M.; Russo, P. Tobacco Smoking: Risk to Develop Addiction, Chronic Obstructive Pulmonary Disease, and Lung Cancer. *Recent Pat. Anticancer Drug Discov.* **2019**, doi:10.2174/1574892814666190102122848.
11. U.S. Department of Health and Human Services. The health consequences of smoking – 50 years of progress: a report of the surgeon general. Atlanta. GA: U.S. Department of Health and Human Services. Centers for Disease Control and Prevention. National Center for Chronic Disease Prevention and Health Promotion. Office on Smoking and Health, 2014.
12. U.S. Department of Health and Human Services. The health consequences of involuntary exposure to tobacco smoke: a report of the surgeon general. Atlanta. GA: U.S. Department of Health and Human Services. Centers for Disease Control and Prevention. Coordinating Center for Health Promotion. National Center for Chronic Disease Prevention and Health Promotion. Office on Smoking and Health, 2006.
13. Besaratinia, A.; Pfeifer, G.P. Second-hand smoke and human lung cancer. *Lancet Oncol.* **2008**, *9*, 657–666, doi:10.1016/s1470-2045(08)70172-4.
14. Protano, C.; Vitali, M. The New Danger of Thirdhand Smoke: Why Passive Smoking Does Not Stop at Secondhand Smoke. *Environ. Health Perspect.* **2011**, *119*, a422.
15. Matt, G.E.; Quintana, P.J.E.; Destaillats, H.; Gundel, L.A.; Sleiman, M.; Singer, B.C.; Jacob, P.; Benowitz, N.; Winickoff, J.P.; Rehan, V.; et al. Thirdhand tobacco smoke: Emerging evidence and arguments for a multidisciplinary research agenda. *Environ Health Perspect.* **2011**, *119*, 1218–1226.
16. Forastiere, F.; Mallone, S.; Lo Presti, E.; et al. Characteristics of nonsmoking women exposed to spouses who smoke: Epidemiologic study on environment and health in women from four Italian areas. *Environ. Health Perspect.* **2000**, *108*, 1171–1177.
17. Wu, F.Y.; Wu, H.D.I.; Yang, H.L.; Kuo, H.W.; Ying, J.C.; Lin, C.J.; Yang, C.C.; Lin, L.Y.; Chiu, T.H.; Lai, J.S. Associations among genetic susceptibility, DNA damage, and pregnancy outcomes of expectant mothers exposed to environmental tobacco smoke. *Sci. Total Environ.* **2008**, *404*, 218–219, doi:10.1016/j.scitotenv.2008.04.015.
18. Sobotova, L.; Liu, Y.H.; Burakoff, A.; Sevcikova, L.; Weitzman, M. Household Exposure to Secondhand Smoke is Associated with Decreased Physical and Mental Health of Mothers in the USA. *Matern. Child Health J.* **2011**, *15*, 128–137, doi:10.1007/s10995-009-0549-z.

19. Zhou, S.; Rosenthal, D.G.; Sherman, S.; Zelikoff, J.; Gordon, T.; Weitzman, M. Physical, behavioral, and cognitive effects of prenatal tobacco and postnatal secondhand smoke exposure. *Curr. Probl. Pediatr. Adolesc. Health Care* **2014**, *44*, 219–241, doi:10.1016/j.cppeds.2014.03.007
20. Argalasova, L.; Sevcikova, L.; Jurkovicova, J.; Babjakova, J.; Janekova, E.; Totka, A.; Simko, M.; Weitzman, M. Determinants of ETS exposure in a sample of Slovak pregnant women. *Rev. Environ. Health* **2017**, *32*, 201–205, doi:10.1515/reveh-2016-0029.
21. Hrubá, D.; Kachlík, P. The influence of maternal active and passive smoking during pregnancy on birth weight in newborns. *Cent. Eur. J. Public Health* **2000**, *8*, 249–252.
22. DiFranza, J.; Aline, C.; Weitzman, M. Prenatal and postnatal environmental tobacco smoke exposure and children's health. *Pediatrics* **2004**, *113*, 1007–1015.
23. Polanska, K.; Hanke, W.; Ronchetti, R.; van den Hazel, P.; Zuurbier, M.; Koppe, J.; Bartonova, A. Environmental tobacco smoke exposure and children's health. *Acta Paediatr.* **2006**, *95*, 86–92, doi:10.1080/08035320600886562.
24. Cui, H.; Gong, T.-T.; Liu, C.-X.; Wu, Q.-J. Associations between Passive Maternal Smoking during Pregnancy and Preterm Birth: Evidence from a Meta-Analysis of Observational Studies. *Plos One* **2016**, *11*, doi:10.1371/journal.pone.0147848.
25. Button, T.; Maughan, B.; McGuffin, P. The relationship of maternal smoking to psychological problems in the offspring. *Early Hum. Dev.* **2007**, *83*, 727–732, doi:10.1016/j.earlhumdev.2007.07.006.
26. Kukla, L.; Hrubá, D.; Tyrlík, M. Maternal smoking during pregnancy, behavioral problems and school performances of their school-aged children. *Cent. Eur. J. Public Health* **2008**, *16*, 71–76.
27. Bublitz, M.H.; Stroud, L.R. Maternal smoking during pregnancy and offspring brain structure and function: review and agenda for future research. *Nicotine Tob. Res.* **2012**, *14*, 388–397, doi:10.1093/ntr/ntr191.
28. Aycicek, A.; Erel, O.; Kocyigit, A. Decreased total antioxidant capacity and increased oxidative stress in passive smoker infants and their mothers. *Pediatr. Int.* **2005**, *47*, 635–639, doi:10.1111/j.1442-200x.2005.02137.x.
29. Aycicek, A.; Erel, O.; Kocyigit, A. Increased oxidative stress in infants exposed to passive smoking. *Eur. J. Pediatr.* **2005**, *164*, 775–778, doi:10.1007/s00431-005-1720-1.
30. Seet, R.; Lee, C.; Loke, W.; Huang, S.; Huang, H.; Looi, W.; Chew, E.; Quek, A.; Lim, E.; Halliwell, B. Biomarkers of oxidative damage in cigarette smokers: Which biomarkers might reflect acute versus chronic oxidative stress? *Free Radic. Biol. Med.* **2011**, *50*, 1787–1793, doi:10.1016/j.freeradbiomed.2011.03.019.
31. Aycicek, A. Tobacco Smoking and Oxidative Stress in Pregnancy. In *Perinatal and Prenatal Disorders, Oxidative Stress in Applied Basic Research and Clinical Practice*, Dennery, P.A. et al. (eds.), Springer Science+Business Media New York, USA, 2014; DOI 10.1007/978-1-4939-1405-0_4
32. Chelchowska, M.; Ambroszkiewicz, J.; Gajewska, J.; Laskowska-Klita, T.; Leibschang, J. The effect of tobacco smoking during pregnancy on plasma oxidant and antioxidant status in mother and newborn. *Eur. J. Obstet. Gynecol. Reprod. Biol.* **2011**, *155*, 132–136, doi:10.1016/j.ejogrb.2010.12.006.

33. Sastry, B.V.; Chance, M.B.; Hemontolor, M.E.; Goddijn-Wessel, T.A. Formation and retention of cotinine during placental transfer of nicotine in human placental cotyledon. *Pharmacology* **1998**, *57*, 104–116.
34. Al-Delaimy, W.K.; Crane, J.; Woodward, A. Is the hair nicotine level a more accurate biomarker of environmental tobacco smoke exposure than urine cotinine? *J. Epidemiol. Community Health* **2002**, *56*, 66–71.
35. Behera, D.; Uppal, R.; Majumdar, S. Urinary levels of nicotine & cotinine in tobacco users. *Indian J. Med. Res.* **2003**, *118*, 129–133.
36. Wipfli, H.; Avila-Tang, E.; Navas-Acien, A.; Kim, S.; Onicescu, G.; Yuan, J.; Breysse, P.; Samet J.M. Secondhand Smoke Exposure Among Women and Children: Evidence From 31 Countries. *Am. J. Public Health* **2008**, *98*, 672–679.
37. Berlin, I.; Heilbronner, C.; Georgieu, S.; Meier, C.; Spreux-Varoquaux, O. Newborns cord blood plasma cotinine concentrations are similar to that of their delivering smokers mothers. *Drug Alcohol Depend.* **2010**, *107*, 250–252, doi:10.1016/j.drugalcdep.2009.10.008
38. Wong, S.L.; Shields, M.; Leatherdale, S.; Malaison, E.; Hammond, D. Assessment of validity of self-reported smoking status. *Health Rep.* **2012**, *23*, 47–53.
39. Fleiss, J.L. *Statistical methods for rates and proportions*, 2nd ed.; John Wiley, New York, USA, 1981; pp. 38–46.
40. Hennekens, C.H.; Buring, J.E. *Epidemiology in Medicine*. 1st ed.; Little, Brown and Company, Boston, USA, 1987; pp. 327–335.
41. Rothmann, K.J.; Greenland, S.; Lash, T.L. *Modern Epidemiology*. 3rd ed.; Lippincott Williams & Wilkins, Philadelphia, USA, 2008; pp. 642–646.
42. Landis, J.R.; Koch, G.G. The measurement of observer agreement for categorical data. *Biometrics* **1977**, *33*, 159–174, doi:10.2307/2529310
43. Rebagliato, M. Validation of self-reported smoking. *J. Epidemiol. Community Health* **2002**, *56*, 163–164.
44. Vartiainen, E.; Seppälä, T.; Lillsunde, P.; Puska, P. Validation of self-reported smoking by serum cotinine measurement in a community-based study. *J. Epidemiol. Community. Health* **2002**, *56*, 167–170.
45. de Chazeron, I.; Llorca, P.M.; Ughetto, S.; Coudore, F.; Boussiron, D.; Perriot, J.; Vendittelli, F.; Sapin, V.; Lemery, D. Occult maternal exposure to environmental tobacco smoke exposure. *Tob. Control* **2007**, *16*, 64–65.
46. Paek, Y.J.; Kang, J.B.; Myung, S.K.; Lee, D.H.; Seong, M.W.; Seo, H.G.; Cho, J.J.; Song, H.J.; Park, K.H.; Kim, CH.; Ko, J.A. Self-reported exposure to second-hand smoke and positive urinary cotinine in pregnant nonsmokers. *Yonsei Med. J.* **2009**, *50*, 345–351, doi:10.3349/ymj.2009.50.3.345.
47. Xiao, X.; Li, Y.; Song, X.; Xu, Q.; Yang, S.; Wu, J.; Seto, E. Discrepancy between Self-Reported and Urine Cotinine-Verified Environmental Tobacco Smoke Exposure among Rural Pregnant Women in China. *Int. J. Environ. Res. Public Health* **2018**, *15*, pii:E1499. doi:10.3390/ijerph15071499.
48. Biri, A.; Bozkurt, N.; Turp, A.; Kavutcu, M.; Himmetoglu, O.; Durak, I. Role of oxidative stress in intrauterine growth restriction. *Gynecol. Obstet. Invest.* **2007**, *64*, 187–192.
49. Pappas, R.S. Toxic elements in tobacco and in cigarette smoke: inflammation and sensitization. *Metallomics* **2011**, *11*, 1181–1198, doi:10.1039/c1mt00066g.

50. Kanti Das, T.; Wati, M.R.; Fatima-Shad, K.J.A.N. Oxidative Stress Gated by Fenton and Haber Weiss Reactions and its Association with Alzheimer's disease. *Arch. Neurosci.* **2015**, *2*, e20078, doi:10.5812/archneurosci.20078

51. Janicka, M.; Kot-Wasik, A.; Kot, J.; Namieśnik, J. Isoprostanes-Biomarkers of Lipid Peroxidation: Their Utility in Evaluating Oxidative Stress and Analysis. *Int. J. Mol. Sci.* **2010**, *11*, 4631–4659, doi:10.3390/ijms11114631

52. Fam, S.S.; Morrow, J.D. The isoprostanes: unique products of arachidonic acid oxidation—a review. *Curr Med Chem* **2003**, *10*, 1723–1740.

53. Cracowski, J.L.; Durand, T.; Bessard, G. Isoprostanes as a biomarker of lipid peroxidation in humans: physiology, pharmacology and clinical implications. *Trends Pharmacol. Sci.* **2002**, *23*, 360–366.

54. Lawson, J.A.; Rokach, J.; FitzGerald, G.A. Isoprostanes: formation, analysis and use as indices of lipid peroxidation *in vivo*. *J. Biol. Chem.* **1999**, *274*, 24441–24444.

55. Park, E.Y., Hong, Y.C.; Lee, K.H.; Im, M.W.; Ha, E.; Kim, Y.J.; Ha, M. Maternal exposure to environmental tobacco smoke, GSTM1/T1 polymorphisms and oxidative stress. *Reprod. Toxicol.* **2008**, *26*, 197–202, doi:10.1016/j.reprotox.2008.08.010.

56. Aydogan, U.; Durmaz, E.; Ercan, C.M.; Eken, A.; Ulutas, O.K.; Kavuk, S.; Gursel, O.; Alanbay, I.; Akay, C.; Kurekci, A.E.; Aydin, A.; Sayal, A.; Saglam, K.; Cok, I. Effects of smoking during pregnancy on DNA damage and ROS level consequences in maternal and newborns' blood. *Arh. Hig. Rada Toksikol.* **2013**, *64*, 35–46, doi:10.2478/10004-1254-64-2013-2232

57. Rua, E.A.O.; Porto, M.L.; Ramos, J.P.L.; Nogueira, B.V.; Meyrelles, S.S.; Vasquez, E.C.; Pereira, T.M.C. Effects of tobacco smoking during pregnancy on oxidative stress in the umbilical cord and mononuclear blood cells of neonates. *J. Biomed. Sci.* **2014**, *21*, 105, doi:10.1186/s12929-014-0105-z

58. Mottalebnejad, M.; Pouramir, M.; Jenabian, N.; Ranjbar Omrani, M.; Bijani, A.; Yarmand, F. Evaluating the association between passive smoking with total antioxidant capacity and salivary lipid peroxidation levels in 12 to 15 year old adolescents. *J. Res. Dent. Sci.* **2014**, *11*, 40–44.

59. Fayol, L.; Gulian, J.M.; Dalmasso, C.; Calaf, R.; Simeoni, U.; Millet, V. Antioxidant status of neonates exposed in utero to tobacco smoke. *Biol. Neonate* **2005**, *87*, 121–126.

60. Kurt, A.; Kurt, A.N.C.; Benzer, D.; Aygün, A.D.; Ustündag, B.; Dogan, Y.; Erel, O. Exposure to Environmental Tobacco Smoke during Pregnancy Restrains the Antioxidant Response of their Neonates. *J. Neonatal Biol.* **2016**, *5*, 1, doi:10.4172/2167-0897.1000210

61. Pickett, K.E.; Rathouz, P.J.; Kasza, K.; Wakschlag, L.S.; Wright, R. Self-reported smoking, cotinine levels, and patterns of smoking in pregnancy. *Paediatr. Perinat. Epidemiol.* **2005**, *19*, 368–376, doi:10.1111/j.1365-3016.2005.00660.x.

62. Seccareccia, F.; Zuccaro, P.; Pacifici, R.; Meli, P.; Pannozzo, F.; Freeman, K.M.; Santaquilani, A.; Giampaoli, S. Research Group of the MATISS Project. Serum cotinine as a marker of environmental tobacco smoke exposure in epidemiological studies: the experience of the MATISS project. *Eur. J. Epidemiol.* **2003**, *18*, 487–492.

63. Boyd, N.R.; Windsor, R.A.; Perkins, L.L.; Lowe, J.B. Quality of measurement of smoking status by self-report and saliva cotinine among pregnant women. *Matern. Child. Health J.* **1998**, *2*, 77–83.

64. Swamy, G.K.; Reddick, K.L.B.; Brouwer, R.J.N.; Pollak, K.I.; Myers, E.R. Smoking prevalence in early pregnancy: comparison of self-report and anonymous urine cotinine testing. *J. Matern. Fetal Neonatal Med.* **2011**, *24*, 86–90, doi:10.3109/14767051003758887.
65. Bobb-Semple, A.A.; Williams, A.F.; Boggs, M.E.; Gold, K.J. Prenatal Point-of-Care Tobacco Screening and Clinical Relationships. *Ann. Fam. Med.* **2018**, *16*, 507–514, doi:10.1370/afm.2290.
66. Xiao, X.; Li, Y.; Song, X.; Xu, Q.; Yang, S.; Wu, J.; Seto, E. Discrepancy between Self-Reported and Urine Cotinine-Verified Environmental Tobacco Smoke Exposure among Rural Pregnant Women in China. *Int. J. Environ. Res. Public Health* **2018**, *15*, pii:E1499, doi:10.3390/ijerph15071499.
67. Ware, J. Jr; Kosinski, M.; Keller, S.D. A 12-item short-form health survey: construction of scales and preliminary tests of reliability and validity. *Med. Care* **1996**, *34*, 220–233.