

**Assessment of Residual Feed Intake and its Relevant Measurements in Two  
Varieties of Japanese Quail (*Coturnix coturnix japonica*) under High Environmental  
Temperature**

**Moataz M. Fathi<sup>1,2,\*</sup>, Ibrahim Al-Homidan<sup>1</sup>, Tarek A. Ebeid<sup>1,3</sup>, Ahmed Galal<sup>2</sup> and  
Osama K. Abou-Emera<sup>1</sup>**

<sup>1</sup>Department of Animal Production and Breeding, College of Agriculture and Veterinary  
Medicine, Qassim University, Al-Qassim, 51452, Saudi Arabia;  
mmfathi@fulbrightmail.org

<sup>2</sup>Department of Poultry Production, Faculty of Agriculture, Ain Shams  
University, Hadayek Shoubra 11241, Cairo, Egypt; galaly2k\_2005@yahoo.com

<sup>3</sup>Department of Poultry Production, Faculty of Agriculture, Kafrelsheikh  
University, 33516, Kafr El-Sheikh, Egypt; tarkamin@gmail.com

\* Correspondence: [mmfathi@fulbrightmail.org](mailto:mmfathi@fulbrightmail.org)

**Simple Summary:** Residual feed intake (RFI) is an important factor in improving poultry production and laying performance, particularly for those raised under heat stress. An experiment was conducted to assess RFI and its related measurements in Japanese quails of two varieties (gray and white) reared under high environmental temperatures. Results confirmed that raising gray quails for egg production under high environmental temperature is recommended. Multiple regression analysis clearly identified a significant effect of metabolic body weight and egg mass in computing expected feed intake rather than body weight gain in both varieties of Japanese quails.

**Abstract:** Three hundred ten 12-week-old laying quails (155 each) were randomly selected from the initial population and kept in individual battery cages. The measurements of growth and egg production were determined to derive RFI. The

relationship between RFI and egg quality, blood parameters and carcass characteristics 31  
was also determined. The results indicated that the gray quails had significantly higher 32  
egg mass and lower broken eggs compared to the white quails. A significant increase for 33  
eggshell strength and shell % was found in eggs produced from gray quails compared to 34  
white counterparts, although the shell thickness was the same. The results of multiple 35  
regression analysis clearly identified a significant effect of metabolic body weight and 36  
egg mass in computing expected feed intake rather than body weight gain in both 37  
varieties of Japanese quails. Strongly positive correlation between RFI and feed intake in 38  
both gray and white quail varieties was found. The same trend was also observed for feed 39  
conversion ratio (FCR). Therefore, including RFI in selection criteria of Japanese quail to 40  
improve FCR under high environmental temperature is highly recommended. 41

**Keywords:** residual feed consumption; quail; high environmental temperature 42

## 1. Introduction

Feed expenses represent almost 70% of the gross cost of poultry production. Lowering costs of maintenance processes would leave more energy remaining for higher output. Minimizing residual feed intake (RFI) and, in turn improving feed efficiency would be beneficial for more efficient quail hens particularly under high environmental temperatures. However, the bird's ability to convert consumed feed to produce eggs and/or meat is greatly influenced by genotype and environmental factors. Birds that require less feed than expected for maintenance and production requirements have a negative RFI and are desirable in poultry breeding programs to reduce feed costs. RFI has increasingly become a critical factor for measuring feed efficiency and is considered to be one of the target traits in animal breeding programs [1]. However, many selection programs take RFI into consideration to improve economic productive traits of synthetic or commercial strains. It has already been reported that RFI could be used in selection programs in laying hens and quails. Altan et al. [2] indicated that selection for RFI in Japanese quails might provide a tool to improve efficiency of feed utilization without significant negative changes in egg production and egg quality traits and with a decreased susceptibility to stress. Most researchers concluded that a four-week recording period provides sufficient information for genetic evaluation of RFC in many species of poultry [3-7].

Direct selection for more efficient birds is becoming one of the primary goals in breeding programs of laying hens [7]. Improving feed efficiency is of great economic concern for commercial egg producers to maximize project outcomes. Traditionally, feed efficiency has been improved by selection for increased egg mass and decreased body weight and getting a correlated response in feed efficiency [2]. Identifying birds that require less feed for body maintenance could improve feed efficiency. However, selection for feed conversion ratio can lead to unfavorable changes in the component traits. Additionally, direct selection for feed efficiency requires measurement of individual feed intake which is time consuming and very expensive and needs to use well designed feeders to prevent feed wastage. On the other hand, additional criteria for feed utilization should be involved. Residual feed consumption (RFC) may be used as selection criteria to attain these goals [2]. To present, there are no reports available on RFI traits in varied

varieties of Japanese quail [4]. To our knowledge, there are no previous reports on the residual feed intake of different lines or varieties for Japanese quails raised under hot ambient temperature. Due to adjusting pattern of feed consumption according to ambient temperature, the present study was carried out to estimate RFI as well as its relationship with productive traits in two varieties of Japanese quails under high environmental temperature.

## 2. Materials and methods

### 2.1. Birds, Housing and Management

A total of 310 laying quails of two varieties (gray and white) (155 each) were randomly selected from an initial population of 800 day olds that were reared to point of lay and transferred to individual battery cages. Each hen was individually housed in a wire cage (20 × 20 × 20 cm) supplied with individual feed trough in the front and a nipple drinker. The quail received a laying ration containing 18% CP and 2850 kcal/kg ME during the experimental period. Throughout the experiment, feed and water were available *ad libitum*. Birds were exposed to a lighting period of 16 h per day. All quails received uniform care and management practices throughout the whole experimental period. The average high and low ambient temperatures recorded during the experimental period were 38.9°C and 24.3°C, respectively. No vaccination or medication was performed. The use and handling of quails were approved by the Ethical Committee of Qassim University.

### 2.2. Egg Production Parameters

Starting from 12-wk of age, egg production (weight and number), feed intake (FI), and body weight (initial and final) were determined for each hen over a four-week experimental period. Egg production (%) was calculated as total laid eggs divided by the total number of days (28 days). Feed intake was measured on a cage basis and combined with egg production data to calculate feed conversion ratio (FCR).

### 2.3. Egg Quality Measurements

A total of 460 eggs (230 eggs from each variety) were collected at 16 weeks of age. 105  
External and internal egg quality measurements of each variety were assessed according 106  
to Fathi et al. [8]. Each egg was weighed to the nearest 0.1 g. Egg width (equatorial axis) 107  
and egg length (longitudinal axis) were measured using Vernier caliper in 0.1 mm. Egg 108  
shape was calculated according to the following formula: 109

Egg shape = [egg width / egg length] × 100 110

The breaking strength for intact eggs was determined in Kg/cm<sup>2</sup> using Egg Force 111  
Reader™, Orka Food Technology Ltd, USA. The liquid contents were put aside and the 112  
shell plus membranes were washed under running water to remove the adherent albumen. 113

The wet eggshell was left for 24h at room temperature for drying and then weighed to the 114  
nearest 0.01g. The relative weight of dry eggshell was calculated on the basis of egg 115

weight. To measure shell thickness, pieces from three different regions (two poles and 116  
equator) of each eggshell with intact membranes were measured with a dial gauge 117

micrometer to the nearest 0.01 mm. The height of thick albumen and egg yolk was 118  
measured by placing the liquid content on a balanced surface using a tripod micrometer. 119

Then, the yolk was separated and rolled on tissue papers to remove the residual albumen. 120

Albumen weight was calculated by subtracting the yolk and shell weight from egg 121

weight. The weight of eggshell, yolk and albumen were expressed as a percentage of egg 122

weight. Haugh units (HU) were calculated according to the following formula: 123

$HU = 100 \log (H - 1.7W^{0.37} + 7.57)$ . Where H is the albumen height (mm) and W is the 124

egg weight (g). Yolk colour was measured by comparing yolk colour to the Roche yolk 125  
colour fan. 126

#### 2.4. Carcass and Internal Organs 127 128

At the end of the experiment, 50 quails (25 from each variety) were randomly assigned 129

for carcass yield assessment. After a pre-slaughter fasting period for 4 h, the quails were 130

weighed and slaughtered by cutting their jugular veins. Following a 2-min bleeding time, 131

each quail was dipped in a hot water bath at 60 °C for 60 sec. and manually defeathered. 132

Head and feet were removed. The carcass was eviscerated manually and weighed. Upon 133

evisceration, the weight of eviscerated carcass, liver, heart and gizzard was recorded and 134

expressed as a percentage of live body weight. To minimize variations in the carcass procedure, all dissections were carried out by the same person.

### 2.5. Blood Hematology and Plasma Biochemistry

During slaughter, 25 blood samples were collected from each variety in heparinized tubes. The hematological parameters were determined by using Automatic Fully Digital Hematology Analyzer (BC-3000 Plus, Shenzhen Mindray, Bio-Medical Electronics Co., LTD). These parameters were total count of red blood cells (RBC), hemoglobin (HGB), hematocrit (HT) and thrombocytes. The collected blood samples were centrifuged at 4000×rpm for 15 min. The resulting plasma samples were frozen at -20 C for further analysis. The plasma concentrations of total protein, albumen, total cholesterol and triglycerides were spectrophotometrically determined using commercial reagent kits (Stanbio Laboratory, Boerne, TX). The globulin was calculated as the difference between the total protein and albumen.

### 2.6. Calculation of Residual Feed Intake and Statistical Analysis

Expected feed intake was computed using mid-metabolic body weight ( $BW^{0.75}$ ), body weight gain ( $\Delta BW$ ) and total egg mass (EM) for a given time considered by multiple regression analysis. Residual feed intake (RFI) was calculated as the difference between observed (OFI) and expected feed intake (EFI) for each experimental hen using the PROC REG procedure using JMP Ver. 11 [9]. Each variety had its own partial regression coefficients according to the following multiple regression equation:

$$EFI = aBW_i^{0.75} + bEM_i + c\Delta BW_i + d$$

where:

EFI = Expected feed intake of hen i (g).

$BW_i^{0.75}$  = Mean metabolic body weight of hen i ( $g^{0.75}$ ).

EM<sub>i</sub> = Egg mass production of hen i (g).

$\Delta BW_i$  = Body weight gain (g).

a,b and c = Partial regression coefficients.

d = Intercept.

Student *t* test analysis was applied to separate between means. All data were presented as means and the pooled SEM. Correlation coefficient was computed between RFI and some studied traits within each variety using PROC CORR procedure.

### 3. Results and discussion

Productive performance of two varieties of Japanese quail is shown in Table 1. No significant difference for body weight (initial and final), weight gain, FI and FCR was identified between the varieties. Gray quails had significantly higher ( $P<0.02$ ) egg mass and egg production percentage than that of white quails. A superiority of egg production in brown variety compared to both gray and white ones was detected [10]. Broken eggs was significantly ( $P<0.05$ ) affected by variety of quail. The gray quails recorded the lower value (1.13%) compared to the white one (2.06%). Mortality levels fell within the normal range and there was no significant difference between quail varieties (data not shown).

**Table 1.** Productive performance of two varieties of Japanese quails.

Parameter	Variety		SEM	P-value
	Gray	White		
Initial body weight, g	203.9	201.1	1.72	0.41
Final body weight, g	213.8	211.5	1.74	0.51
Body weight gain, g	9.8	10.4	1.32	0.84
Egg mass, g	289.5 <sup>a</sup>	277.8 <sup>b</sup>	2.43	0.02
FI, g	719.9	722.6	3.75	0.87
FCR	2.49	2.60	0.07	0.11
Egg production, %	89.9 <sup>a</sup>	86.4 <sup>b</sup>	0.73	0.02
Broken eggs, %	1.13 <sup>b</sup>	2.06 <sup>a</sup>	0.02	0.05
Egg weight, g	11.5	11.4	0.023	0.93

N= 155 quails/ variety

Internal and external egg quality characteristics are presented in Table 2. Shape index was significantly ( $P<0.01$ ) higher in egg produced from white quails compared to gray counterparts. Consistent with our results, Yilmaz et al. [11] and Sari et al. [12] reported

that the egg shape index depends on plumage colour of the quails. They found that the mean shape index obtained from gray plumage colour was significantly lower than that of white plumage colour. In contrast to our results, Bagh et al. [10] did not find a significant difference between gray and white lines for all physical properties of egg quality. A numerical increase ( $p=0.08$ ) in HU was found in gray quails when compared with white birds. However, Bagh et al. [10] reported that there was no significance difference among the quail varieties for HU. In terms of yolk properties, it could be noticed that there were no significant differences between quail varieties for yolk colour, yolk index and yolk percentage. Significant ( $P<0.04$ ) higher albumen percentage was associated with eggs produced from white variety compared to the gray one. In regard to eggshell quality, a significant ( $P<0.01$ ) increase in eggshell breaking strength was found in eggs produced from gray quails ( $1.43 \text{ kg/cm}^2$ ) compared to the white quails ( $1.34 \text{ kg/cm}^2$ ). Also, gray quails had a significantly ( $P<0.01$ ) higher relative weight of eggshell (9.4%) compared to that of white counterparts (9.0%). However, shell thickness did not exhibit a significant difference due to variety effect. This advantage in eggshell strength associated with gray quails may be due to better ultrastructural feature compared to eggshell of white variety. However, changes in external and internal quality characteristics of eggs obtained from quails with different plumage colours have previously been reported [10-12]. However, literature on the external and internal quality characteristics of eggs obtained from quail varieties with different plumage colour under high environmental conditions is very limited.

**Table 2.** Internal and external egg quality of two varieties of Japanese quails.

Parameter	Variety		SEM	P-value
	Gray	White		
Shape index, %	75.95 <sup>b</sup>	76.90 <sup>a</sup>	0.15	<0.01
HU	88.7	88.2	0.13	0.08
Yolk colour <sup>1</sup>	6.96	6.98	0.03	0.70
Yolk index	47.67	47.31	0.12	0.15



Yolk %	33.5	33.3	0.12	0.39
Albumen %	57.1 <sup>b</sup>	57.7 <sup>a</sup>	0.14	0.04
Shell%	9.4 <sup>a</sup>	9.0 <sup>b</sup>	0.05	<0.01
Shell thickness, $\mu$	254.0	255.2	0.73	0.40
Breaking strength, kg/cm <sup>2</sup>	1.43 <sup>a</sup>	1.34 <sup>b</sup>	0.01	<0.01

---

N= 230 intact eggs/variety

209

<sup>1</sup>Roche yolk colour fan

210

211

Plasma biochemical and hematological parameters in gray and white feathered Japanese quails are summarized in Table 3. As shown in, no significant effect on the blood biochemical variables was detected due to variety except for cholesterol level. White quails had a significant ( $P<0.01$ ) increase in cholesterol level (198.5 mg/dL) compared to gray variety (152.5 mg/dL). In terms of blood hematology, the white feathered quails had significantly higher levels of RBC, HGB and HT compared to the gray quails. Generally, hematological parameter fell within the normal range for quails [13]. These results indicate that different genotypes in the present study were in normal physiological status.

212

213

214

215

216

217

218

219

220

221

**Table 3.** Biochemical and hematological blood parameters of two varieties of Japanese quails.

222

223

Parameter	Variety		SEM	P-value
	Gray	White		
Total Protein, g/dL	5.34	5.43	0.17	0.82
Albumin, g/dL	3.13	3.42	0.09	0.27
Globulin, g/dL	2.21	2.01	0.18	0.69
Cholesterol, mg/dL	152.50 <sup>b</sup>	198.47 <sup>a</sup>	8.98	<0.01

Triglyceride, mg/dL	121.75	120.67	4.69	0.93
RBC	3.28 <sup>b</sup>	3.51 <sup>a</sup>	0.05	0.02
HGB	19.36 <sup>b</sup>	20.59 <sup>a</sup>	0.25	<0.01
HT	44.37 <sup>b</sup>	47.21 <sup>a</sup>	0.63	0.02
Thrombocytes	15.84	15.83	0.85	0.99

---

N= 25 quails/variety

224

Results of carcass traits studied as affected by variety of quails are shown in Table 4. 225  
 It was found that either carcass percentage or giblets (liver, heart and gizzard) did not 226  
 significantly differ between varieties. However, insignificant increase ( $p=0.15$ ) in in 227  
 dressed carcass % was found in white quails (63.3%) compared to gray counterparts 228  
 (62.4%). Like in the present study, Charati and Esmailzadeh [14] found that genotype 229  
 had no significant effect on carcass percentage in white and wild (gray) Japanese quail. In 230  
 contrast, several previous studies reported that feather colours had a significant effect on 231  
 live weight, and carcass characteristics in Japanese quail. The white feathered quail had 232  
 less body weight than that of the wild-type [15-17]. Similarly, Vali et al. [18] found 233  
 significant differences in two quail strains for carcass weight, carcass percentage, and the 234  
 relative weight of breast and femur. 235

236

**Table 4.** Carcass traits of two varieties of Japanese quails. 237

Trait	Variety		SEM	P-value
	Gray	White		
Live body weight, g	223.9	223.5	3.18	0.93
Dressed carcass, %	62.4	63.3	0.30	0.15
Liver, %	2.02	2.12	0.08	0.51
Heart, %	1.20	1.14	0.02	0.19
Gizzard, %	1.66	1.56	0.04	0.13

---

N= 25 quails/variety

238

239

The results of multiple regression analysis are listed in Table 5. Below the table, the prediction equations for expected feed consumption for each variety were provided. RFI is defined as the difference between the realized feed consumption and the expected feed consumption estimated based on metabolic BW, body weight gain and EM [3,19]. As shown, the partial regression coefficients for metabolic body weight and egg mass had a significantly effect in computing expected feed intake in both quail varieties. Intercept values had also a significant effect. On the other hand, body weight gain ( $\Delta$  BWT) did not significantly affect computing RFI either in gray variety ( $p= 0.08$ ) or in white one ( $p= 0.63$ ). Estimates of regression coefficients in the models for gray and white quails were close. Similarly, Badawe et al. [3] found that the prediction of feed intake and residual feed consumption computing from multiple regression analysis were significantly affected by metabolic body weight and egg mass in laying hen chickens.

240

241

242

243

244

245

246

247

248

249

250

251

252

**Table 5.** Partial regression coefficients for factors affecting expected feed intake of two varieties of Japanese quails.

253

254

Parameter estimate	Partial regression coefficient		Prob.	
	Gray	White	Gray	White
Intercept	914.6**	396.2*	<0.001	0.02
$\Delta$ BWT	-0.96	-0.36	0.08	0.63
(BWT) <sup>0.75</sup>	5.21*	11.84**	0.04	<0.001
Egg mass (EM)	-1.64**	-1.11**	<0.001	<0.001

N= 155 individual records/ variety, \*P< 0.05, \*\*P< 0.01

255

Prediction equations:

256

$Y = 914.6 - 0.96 \Delta \text{BWT} + 5.21 (\text{BWT})^{0.75} - 1.64 \text{EM}$  (Gray variety)

257

$Y = 396.2 - 0.36 \Delta \text{BWT} + 11.84 (\text{BWT})^{0.75} - 1.11 \text{EM}$  (White variety)

258

Where: Y stands for expected feed consumption,  $\Delta$  BWT = body weight gain, (BWT)<sup>0.75</sup>= metabolic body weight, EM = total egg mass.

259

260

261

262

Phenotypic correlations between RFI and some studied traits are presented in Table 6. Strongly positive correlation between RFI and FI in gray and white quail varieties (0.89

263

264

and 0.91, respectively) was found. Notably, correlation between RFI and FI in our study was much higher than those estimated in previous work on laying chickens [7,20,21]. Selection for low RFI could reduce FI without significant changes in EM [7]. FCR is widely used but not a suitable selection trait because of its complex correlations with growth and production traits [6,7]. As shown in Table 6, a significantly high correlation was recorded between RFI and FCR (0.55 and 0.49, respectively). These strong relationships have indicated that selection for negative RFI would genetically improve feed efficiency and reduce feed intake. These results are consistent with those of Zhang et al.[1], who found high phenotypic correlation between RFI and FCR (0.55). Moreover, RFI was strongly correlated with FI (0.82) in a random population of Pekin duck. Similarly, RFC was positively correlated with FI in laying hens of chicken [20-22] and Japanese quail [2]. It is worthy to note that there was a low or neglected correlation between RFI and both egg weight and egg production% for the quail varieties. A low correlation between RFI with body weight gain was found (close to zero) in both quail varieties. Our results are in accordance with the findings of Luiting and Urff [23] and Altan et al. [2], who described also that the phenotypic correlations of RFI with egg mass and body weight goes to zero. Likewise, these results are in agreement with the findings of Varkoohi et al. [5] reflecting the fact that RFI is phenotypically independent of weight gain (WG), which tends to make the genetic correlation between RFI and WG low as well. On the other hand, phenotypic correlations between RFI and both blood parameters and carcass traits were found to be rather low and insignificant. No significant relationship was observed between RFI with live body weight and eviscerated carcass weight [24].

**Table 6.** Phenotypic correlations between RFI and some studied traits in Japanese quails.

Trait	Phenotypic correlation	
	Gray	White
Δ Body wt	-0.003	0.002
Feed intake	0.89**	0.91**
FCR	0.55*	0.49*

Prod%	-0.07	-0.05
Egg wt	0.11	0.18
Total protein	-0.21	0.22
Albumin	-0.38	0.38
Globulin	0.01	0.03
Cholesterol	0.23	-0.15
Triglyceride	0.51	0.20
Hemoglobin	-0.18	-0.40
Red blood cells	0.32	-0.49
Hematocrit	0.20	-0.50
Thrombocytes	-0.15	-0.14
Carcass	0.09	0.09
Liver	0.04	-0.22
Heart	-0.38	-0.03
Gizzard	0.09	-0.22

\*P< 0.05, \*\*P< 0.01

290

291

#### 4. Conclusion

292

The current results indicate that egg mass significantly increased in gray variety compared to white one. Additionally, gray quails had a significantly lower percentage of broken eggs. Additionally, colour variations in Japanese quails should be considered when selecting for type of production. We recommended raising gray quails for egg production under high environmental temperature, while white variety may be more suitable for meat type. Results derived from multiple regression analysis clearly identified a significant effect of metabolic body weight and egg mass in computing expected feed intake rather than body weight gain in both varieties of Japanese quails. Including RFI in selection criteria of Japanese quail to improve FCR under high environmental temperature is highly recommended.

301

302

303

304

**Author Contributions:** Designing M.F.; performing the experiment and data collecting, 305  
M.F., I.A., T.E. and O.A.; Data processing and statistical analysis, M.F. and A.G.; Final 306  
revision, M.F. All authors have been involved in developing, writing and commenting on 307  
the manuscript. All authors read and approved the final manuscript. 308

**Funding:** This research was funded by Qassim University. 309

**Acknowledgments:** This work was supported by Qassim University, Al-Qassim, KSA. 310  
Also, Mr. A. Al-Moshawah must be thanked for his technical assistance during the course 311  
of the study. 312

**Conflicts of Interest:** The authors declare no conflict of interest. 313

314

315

316

317

318

319

320

<b>References</b>	321
1. Zhang, Y.; Guo, Z.B.; Z. Zhang, M. Xie; Hou, S. Genetic parameters for residual feed intake in a random population of Pekin duck. <i>Asian-Australas. J. Anim. Sci.</i> <b>2017</b> , <i>30</i> , 167-170.	322 323 324 325
2. Altan, O.; Oguz, I.; Akbas, Y.; Aksit, M. Genetic variability of residual feed consumption (RFC) and its relationships with some production traits and fear response in Japanese quail hens ( <i>Coturnix coturnix japonica</i> ). <i>Arch. Geflügelk.</i> <b>2004</b> , <i>68</i> ,223–229.	326 327 328 329 330
3. Badawe, M.I.; Abd El-Hameid, E.F.; Fathi, M.M.; Zein El-Dein, A. Prediction of feed intake and residual feed consumption in laying hen chickens. <i>Egypt. Poult. Sci.</i> <b>2005</b> , <i>25</i> ,1003-1016.	331 332 333
4. Varkoohi, S.; Babak, M.; Pakdel, A.; Javaremi, A.; Zaghari, M.; Kause, A. Response to selection for feed conversion ratio in Japanese quail. <i>Poult. Sci.</i> <b>2010</b> , <i>89</i> ,1590-1598.	334 335 336
5. Varkoohi, S.; Pakdel, A.; Babak, M.S.; Javaremi, A.N.; Kause, A.; Zaghari, M. Genetic parameters for feed utilization traits in Japanese quail. <i>Poult. Sci.</i> <b>2011</b> , <i>90</i> , 42–47.	337 338 339
6. Foomani, N.N.; Zerehdaran, S.; Azari M. A.; Lotfi, E. Genetic parameters for feed efficiency and body weight traits in Japanese quail. <i>Br. Poult. Sci.</i> <b>2014</b> , <i>55</i> :298-304.	340 341 342
7. Yuan, J.; Wang, K.; Yi, G.; Ma, M.; Dou, T.; Sun, C.; Qu, L.J.; Shen, M.; Qu, L.; Yang, N. Genome-wide association studies for feed intake and efficiency in two laying periods of chickens. <i>Genet. Sel. Evol.</i> <b>2015</b> , <i>47</i> ,82-95.	343 344 345
8. Fathi, M.M.; El-Dlebshany, A.E.; Bahie El-Deen, M.; Radwan, L.M.; Rayan G.N. Effect of long-term selection for egg production on eggshell quality of Japanese quail ( <i>Coturnix japonica</i> ). <i>Poult. Sci.</i> <b>2016</b> , <i>95</i> ,2570–2575.	346 347 348
9. SAS Institute. JMP Version 11. <b>2013</b> , User’s Guide, SAS.	349
10. Bagh, J.; Panigrahi, B.; Panda, N.; Pradhan, C.R.; Mallik, B.K.; Majhi, B.; Rout. S.S. Body weight, egg production, and egg quality traits of gray, brown, and white	350 351

- varieties of Japanese quail (*Coturnix coturnix japonica*) in coastal climatic condition of Odisha. *Vet. World.* **2016**, *9*,832-836.
11. Yilmaz, A.; Tepeli, C.; Caglayan, T. External and internal egg quality characteristics in Japanese quails of different plumage color lines. *J. Food Agric. Environ.* **2011**, *9*,375-379.
12. Sari, M.; Işık, S.; Önk, K.; Tilki, M.; Kirmizibayrak, T. Effects of layer age and different plumage colors on external and internal egg quality characteristics in Japanese quails (*Coturnix coturnix japonica*). *Arch. Geflügelk.* **2012**, *76*, 254–258.
13. Nazifi, S.; Asasi, K. Hematological and serum biochemical studies on Japanese quails (*Coturnix coturnix japonica*) fed different levels of furazolidone. *Revue Méd. Vét.* **2001**, *152*, 705-708.
14. Charati, H.; Esmailizadeh, A.K. Carcass traits and physical characteristics of eggs in Japanese quail as affected by genotype, sex and hatch. *J. Livest. Sci. Technol.* **2013**, *1*,57-62.
15. Petek, M.; Ozen, Y.; Karakas, E. Effects of recessive white plumage color mutation on hatchability and growth of quail hatched from breeders of different ages. *Br. Poult. Sci.* **2004**, *45*,769-774.
16. Minvielle, F.; Gourichon, D.; Moussu, C. Two new plumage mutations in the Japanese quail: “curly” feather and “rusty” plumage. *BMC Genet.* **2005**, *6*,14.
17. Inci, H.; Sogut, B.; Sengul, T.; Sengul, A.Y.; Taysi, M.R. Comparison of fattening performance, carcass characteristics, and egg quality characteristics of Japanese quails with different feather colors. *R. Bras. Zootec.* **2015**, *44*, 390-396.
18. Vali, N.; Edriss, M.A.; Rahmani, H.R. Genetic parameters of body and some carcass traits in two quail strains. *Poult. Sci.* **2005**, *45*, 296-300.
19. Katle, J.; Kolstad, N. Selection for efficiency of food utilisation in laying hens: Direct response in residual food consumption and correlated responses in weight gain, egg production and body weight. *Br. Poult. Sci.* **1991**, *32*, 939–953.
20. Schulman, N.; Tuiskula-Haavisto, M.; Siitonen L.; Mantysaari, E.A. Genetic variation of residual feed consumption in a selected Finnish egg-layer population. *Poult. Sci.* **1994**, *73*, 1479–1484.



21. Tixier-Boichard, M.; Boichard, D.; Groeneveld, E.; Bordas, A. Restricted maximum likelihood estimates of genetic parameters of adult male and female Rhode Island Red Chickens divergently selected for residual feed consumption. *Poult. Sci.* **1995**, *74*, 1245–1252. 383  
384  
385  
386
22. Bordas, A.; Minvielle, F. Patterns of growth and feed intake in divergent lines of laying domestic fowl selected for residual feed consumption. *Poult. Sci.* **1999**, *78*, 317–323. 387  
388  
389
23. Luiting, P.; Urff, E.M. Residual feed consumption in laying hens. 2. Genetic variation and correlations. *Poult. Sci.* **1991**, *70*, 1663–1672. 390  
391
24. Beglia, H.E.; Torshizi, R.V.; Masoudi, A.A.; Ehsani, A.; Jensen, J. Relationship between residual feed intake and carcass composition, meat quality and size of small intestine in a population of F2 chickens. *Livest. Sci.* **2017**, *205*, 10–15. 392  
393  
394