

Review

Use of a novel enhanced DNA vaccine vector for preclinical virus vaccine investigation

Rosamund Chapman¹ and Edward P Rybicki^{1,2,*}

¹ Institute of Infectious Disease and Molecular Medicine, Faculty of Health Sciences, University of Cape Town, Observatory 7925, Cape Town, South Africa; ros.chapman@uct.ac.za

² Biopharming Research Unit, Department of Molecular & Cell Biology, University of Cape Town, PB X3 Rondebosch 7701, Cape Town, South Africa; ed.rybicki@uct.ac.za

* Correspondence: ed.rybicki@uct.ac.za; Tel.: +27 21 6503265

Abstract: DNA vaccines are stable, safe, cost effective to produce and relatively quick and easy to manufacture. However, to date DNA vaccines have shown relatively poor immunogenicity in humans despite promising preclinical results. Consequently, a number of different approaches have been investigated to improve the immunogenicity of DNA vaccines. These include the use of improved delivery methods, adjuvants, stronger promoters and enhancer elements to increase antigen expression, and codon optimization of the gene of interest. This review describes the creation and use of a DNA vaccine vector containing a porcine circovirus (PCV-1) enhancer element that significantly increases recombinant antigen expression and immunogenicity and allows for dose sparing. A 172bp region containing the PCV-1 capsid protein promoter (Pcap) and a smaller element (PC; 70 bp) within this were found to be equally effective. DNA vaccines containing the Pcap region expressing various HIV-1 antigens were found to be highly immunogenic in mice, rabbits and macaques, at 4 to 10-fold lower doses than normally used and to be highly effective in heterologous prime-boost regimens. By lowering the amount of DNA used for immunization, safety concerns over injecting large amounts of DNA into humans can be overcome.

Keywords: DNA vaccine; [HIV-1](#); enhancer element; circovirus; [immunogenicity](#)

1. Introduction


DNA vaccines were hailed as long ago as the 1990s as the next best thing in vaccines: plasmid-based DNA vaccines are relatively easy and affordable to produce, sharing a common production method for all vaccines; they are thermostable, safe with no risk of virulence or apparently of anti-vector immunity, can be administered to immunocompromised individuals, and multiple plasmids can be mixed and used as a broad spectrum combination vaccine. DNA vaccines elicit mainly cell-mediated immune responses due to presentation of expressed antigens via MHC-I presentation, which is similar to viral pathogens, and a desirable feature of a vaccine [1]. One important drawback to DNA vaccines, however, is their lack of immunogenicity compared to protein-based or whole virus vaccines: humoral responses are generally weak if not lacking altogether, and high, repeated doses of DNA are needed in order to obtain reasonable response rates in animal models. Additionally, results in small experimental animals have not translated well into human clinical trial results, and there are concerns over the safety of injecting large amounts of DNA (milligrams) into humans [2].

2. A novel enhancer sequence for DNA vaccine antigen expression

Our group therefore previously investigated the potential of short enhancer sequences derived from a mammalian single-stranded DNA virus – porcine circovirus type I (PCV-1) – for dose-sparing potential and immunogenicity enhancement in a clinically-trialed HIV-1 subtype C DNA vaccine [3]. The plasmid vector (pTH) has been well used in preclinical and clinical studies [4-6] and is regarded

as being a high potency vaccine antigen vector for HIV and other agents. It relies on the human cytomegalovirus immediate/early promoter/enhancer element (CMV I/E) constituting the promoter Pcmv [7], one of the strongest known promoters in mammalian expression systems, driving in vivo antigen expression with the help of the CMV intron A and the bovine growth hormone polyadenylation signal. It has been used to vector the synthetic HIV-1 subtype C vaccine antigen GrttnC, a polyprotein incorporating Gag, reverse transcriptase (RT), Tat and Nef sequences, in studies in mice, guinea pigs, monkeys and humans [8-12].

PCV-1, like all circoviruses, has a compact, genetically dense, bi-directionally transcribed genome of 1759 bp that encodes only a viral capsid protein (*cap* gene) and the replication associated proteins Rep and Rep', which derive by alternative splicing from one ORF (*rep*) (Fig. 1). Bidirectional transcription of the two genes originates in the origin of replication (Ori) for *rep*, and in an intron within *rep* for *cap* [13]. In vitro expression studies in HEK293 cells with various constructs derived from the PCV-1 genomes showed that enhancement activity resided in a 70 bp "core sequence" (PC) of the 172 base pair capsid promoter, Pcap, that includes a putative composite transcription factor binding site comprising C/EBPb, GATA-1 and CREB sites, as well as a 47-bp conserved late element, or CLE. Inclusion of the 70-bp sequence in the reverse orientation immediately upstream of the Pcmv sequence in pTHgrttnC (yielding pTHCRgrttnC) resulted in 2.4-fold enhancement of polyprotein expression level in vitro following transfection of HEK293 cells, as assessed by Gag p24 ELISA. The cognate sequence from the related PCV-2 was equally effective. The 172 base Pcap sequence also enhanced luciferase expression in HEK293 cells by 3-fold when inserted in reverse orientation upstream of the SV40 promoter in the commercial pGL vector [3]. Accordingly, we tested enhancement of immunogenicity in vivo by intramuscular injection of mice with a variety of pTHgrttnC constructs with additives from PCV-1: the best enhancement over pTHgrttnC, as assayed by IFN- γ ELISPOT responses to a RT CD8⁺ peptide, was obtained using the Pcap (172 bp) insert, after two intramuscular inoculations of 100 μ g of pTHPcapRgrttnC DNA (5-fold increase in sfu/10⁶ splenocytes). Moreover, two inoculations of 10 μ g of pTHPcapgrttnC DNA was significantly more immunogenic (3.5-fold) than pTHgrttnC and boosting with 10⁴ pfu of modified vaccinia Ankara (MVA) vectoring Grttn showed the same trend (Fig. 2). The response to the 10 μ g of pTHPcapgrttnC DNA alone was also equivalent to or higher than to 100 μ g of pTHgrttnC, indicating that significant dose-sparing (10-fold) was possible for the same priming effect for a vaccine-relevant antigen. This proof that a simple enhancement could dramatically improve the functionality of a DNA vaccine vector led to its being employed in subsequent studies in our HIV vaccine research programme.



B

PCV-1 172 bp PcapR fragment (1218 – 1389)

```

                                AP2                c/EBPb                GATA-1
GAGCAGTTC CCTGTAACGTATGTGAGAAATTTCCGCGGGCTGGCTGAAC TTTGAAAGTGAGCGGGAAGATGCAGCAGCGTGATTGGAA

                                CREB                SP1                AP3
GACAGCTGTACACGTCATAGTGGCCCGCCCGGTTGTGGGAAGAGCCAGTGGGCCCGTAATTTGCTGAGCCTAGCGACACC

```

Figure 1. PCV-1 genome arrangement. (A) Diagram of the linearized PCV-1 genome, depicted in the orientation cloned into pTHCapgrttnC. The *rep* intron is enlarged and the capsid gene promoter (Pcap) indicated. The core and Conserved Late Elements (CLE) components of Pcap are shown. *rep* = replication associated protein gene, *cap* = capsid protein gene, Prep = *rep* gene promoter, Ori = origin of replication, core = composite host transcription factor binding site. (B) DNA sequence of 172 bp PcapR fragment. Putative host transcription factor binding sites are indicated and underlined, CLE motifs are in bold and the minimal PcapR sequence (1252-1238; as identified by Mankertz & Hillenbrand [13]) is highlighted in grey. PCV-1 Accession Number Y09921. Figure reproduced from Tanzer *et al.* [3] under the Creative Commons Attribution (CC-BY) licence as specified by BioMed Central.




Figure 2. HIV-1 specific IFN- γ ELISPOT responses to pTHgrttnC DNA vaccines containing portions of the PCV-1 genome. Groups of mice were vaccinated intramuscularly with DNA vaccines on days 0 and 28. Two groups of mice were subsequently boosted with 10⁴ pfu of MVA on day 56. A separate group of mice was vaccinated with 10 μ g pTH (empty vector) on days 0 and 28 and subsequently boosted with 10⁴ pfu of MVA on day 56. * $p < 0.001$; ** $p < 0.05$ Student t-test. Figure reproduced from Tanzer *et al.* [3] under the Creative Commons Attribution (CC-BY) licence as specified by BioMed Central.

3. Testing the enhanced DNA vector with HIV-1 subtype C pr55Gag

Strong polyfunctional CD8⁺ T cell responses to HIV-1 Gag or Gag-derived antigens have been found to be important for controlling viraemia in HIV+ people who are termed “elite controllers”. Accordingly, Gag should be and often is included in candidate HIV vaccination regimes, so as to allow early clearance of infected cells at the initial sites of infection, as well as control of spread from these sites and later control of viraemia [14]. A subtype C mosaic Gag sequence was chosen to increase the coverage of both CD8⁺ and CD4⁺ T cell epitopes from that of natural sequences with the hope of reducing the HIV-1 escape pathways [15-18]. Subtype C (HIV-1C) was chosen as it is the most prevalent subtype in the world, accounting for over 50% of all global infections and is the dominant subtype in southern Africa. In a study carried out by our group, the pTHPcapR plasmid backbone [3] was used to construct a DNA vaccine containing an HIV-1 subtype C mosaic *gag* gene, DNA-Gag^M [19, 20].

HEK293T cells transfected with DNA-Gag^M expressed high levels of Gag (up to 26ng/ml in the media). The immune responses to the DNA vaccine were evaluated in mice using homologous and heterologous prime boosts with MVA vaccine expressing the matching HIV-1 subtype C mosaic Gag antigen (MVA-Gag^M). To confirm that the DNA vaccine was immunogenic at a low dose, mice were vaccinated with 10 μ g of the DNA vaccine. Mice vaccinated with two doses of DNA-Gag^M had mean cumulative Gag-specific IFN- γ ELISPOT responses of 882 sfu/10⁶ splenocytes (Fig. 3). These responses were higher to CD8⁺ rather than CD4⁺ Gag peptides (604 and 278 sfu/10⁶ respectively). Mice that received a heterologous prime boost consisting of two doses of DNA-Gag^M followed by a single dose of MVA-Gag^M had mean cumulative Gag-specific IFN- γ ELISPOT responses of 2675 sfu/10⁶, that was evenly balanced to both Gag CD4⁺ and CD8⁺ peptides. Both the homologous and heterologous vaccination regimen elicited a higher proportion of CD8⁺ T cells expressing cytokines than CD4⁺ T cells. All the cytokine-positive CD8⁺ T cells had an effector memory phenotype. This study confirmed that the pTHPcapR DNA vector backbone containing the porcine circovirus enhancer elicits high magnitude, Gag-specific T cell responses in BALB/c mice at a low dose.




Figure 3. DNA vaccine elicits high Gag-specific IFN- γ ELISPOT responses both alone and in a heterologous prime boost with MVA. (A) Vaccination schedule. DNA-Gag^M = pTHPcapR containing mosaic gag; DNA^E = pTHPcapR empty vector; MVA-Gag^M = MVA containing mosaic gag (B) Cumulative IFN- γ ELISPOT CD8⁺ and CD4⁺ responses of vaccinated mice to HIV-1 Gag peptides. ** p<0.01 Student t-test of unpaired data.

4. Testing the enhanced DNA vector with HIV-1 subtype C Env immunogens

The kinds of immune responses that an effective HIV-1 vaccine would need to elicit include non-neutralizing antibody responses as well as broadly neutralizing antibody responses, together with polyfunctional cytotoxic T cell responses to a variety of epitopes from the HIV-1 proteome [21-23]. One of the main targets of recent HIV-1 vaccine candidates is broadly neutralizing antibody (bNAb) responses: bNAbs that can neutralize diverse primary HIV-1 subtype isolates protect against viral challenge in nonhuman primates (NHP) with Env-pseudotyped simian-human immunodeficiency viruses (SHIVs), suggesting that infection in humans could be similarly prevented [24, 25]. Ranking of HIV-1 isolates according to their sensitivities to neutralizing antibodies allows identification of viruses as Tier 1 (sensitive), Tier 2 (moderately resistant), and Tier 3 (resistant) [26]. The circulating viruses that vaccines will need to protect against are largely Tier 2-type: accordingly, HIV vaccines should elicit responses that neutralize laboratory Tier 2 virus isolates. We showed previously that using a DNA prime / MVA boost immunisation regime in mice with vaccines expressing HIV-1 subtype C mosaic Gag resulted in strong cellular immune responses directed against Gag [19]. We wished to extend these results by improving the vaccine regimen to allow the elicitation of Env-specific neutralizing antibodies in a rabbit model.

The pTHPcapR vector was used to construct a DNA vaccine expressing HIV-1 envelope (DNA Env). The envelope sequence (CAP256SU) used in this study was selected as it elicited broadly neutralising antibodies (bNAbs) in the patient [27] and is sensitive to several prototype broadly neutralizing monoclonal antibodies [28]. Several modifications were made to the envelope sequence, these included replacing the native leader sequence with the tissue plasminogen activator leader sequence, replacing the furin cleavage site with a flexible linker, introducing an I548P mutation equivalent to the I559P in the SOSIP trimers to improve the trimerization of gp41 [29] and truncating the sequence from gp160 to gp150 [30]. A second plasmid expressing the soluble envelope protein (gp140) with the same modifications was also constructed using the pTHPcapR backbone [31]. This plasmid was used to generate a stable cell line expressing high levels of the soluble HIV-1 envelope protein which was subsequently purified and utilised as a protein boost in rabbit immunogenicity studies. MVA vaccines expressing the matching gp150 Env and Env plus mosaic Gag were also constructed.

Rabbits were inoculated with different combinations of vaccines in different regimens, in order to ascertain the overall effects on immunogenicity of the Env component. The first test group was injected with 100 µg of each of DNA Env and DNA-Gag^M-encoding plasmids at weeks 0 and 4, boosted with doses of 10⁸ PFU of rMVA Env+Gag^M at weeks 8 and 12, and further boosted with gp140Env protein at weeks 20 and 28 (regime designated asDDMMPP). The other group received 10⁸ PFU of rMVA Env+Gag^M intramuscularly at weeks 0 and 4, followed by three protein boosts at weeks 12, 20, and 28 (MMPPP) (Fig. 4).

Both the DDMMPP and MMPPP vaccination regimens elicited NAbs to the autologous Tier 2 CAP256SU pseudovirion. Moreover, high titres of antibodies that bound to the homologous CAP256 Env and a CAP256 V1V2 loop scaffold were also elicited [30]. It was noticeable that the DDMMPP regimen elicited higher mean peak titres of Tier 2 NAbs than did the MMPPP regimen: this suggests that priming with a DNA vaccine (DDMMPP) gives a better, wider anti-Env immune response than the MMPP regime (Fig. 4). The DDMMPP regimen rabbits also apparently developed a slight increase in breadth of the response as they had low levels of NAbs to clade A pseudovirus 398F1. Our findings that DNA primes a good humoral response agree with others: for example, adding DNA-C priming in the EV01 phase I trial resulted in increased anti-Env IgG responses (from 27% for NYVAC alone to 75% for DNA + NYVAC [32]). Priming with DNA also resulted in significantly boosted T cell responses.




Figure 4. Rabbits primed with DNA produce higher autologous Tier 2 neutralising antibodies than those receiving MVA and protein alone. (A) Longitudinal, Tier 2 neutralising antibody responses to autologous CAP256.SU pseudovirion from serum of individual rabbits. (B) Neutralizing antibody titres at week 30. ** $p < 0.01$ Mann-Whitney U Test.

5. Comparison of DNA vaccines between two initiatives in South Africa

In 2000, a UCT-based consortium headed by Prof Anna-Lise Williamson was awarded funds by the South African AIDS Vaccine Initiative (SAAVI) for the development of HIV-1C vaccines for South Africa. Two vaccines - designated SAAVI DNA-C2 and SAAVI MVA-C - were deemed suitable for human clinical trials [9, 10]. The vaccines expressed a HIV-1 subtype C truncated envelope protein Du151 (gp150) and the polyprotein designated Grttn described above, consisting of translational fusions of HIV-1 subtype C Gag Du422, and modified reverse transcriptase (RT), Tat and Nef-encoding ORFs. The vector backbone utilised for the DNA vaccines contained the regulatory R region from the 5' long terminal repeat (LTR) of human T-cell leukemia virus type 1, which acts as a transcriptional and post-transcriptional enhancer [33]. Rhesus macaques were inoculated at weeks 0, 4 and 8 with 4mg of SAAVI DNA-C2. No HIV-specific ELISPOT responses were detected following the DNA vaccinations (unpublished data). In a more recent study DNA vaccines expressing the SIV Gag and HIV-1 subtype C truncated envelope ZM109F.PB4 were constructed utilising the pTHPcapR vector backbone (unpublished data). Rhesus macaques were inoculated, at weeks 0 and 4 with 1mg of the DNA vaccines (4-fold lower dose). Four out of five macaques developed IFN- γ ELISPOT responses following stimulation with SIV Gag and HIV-1 subtype Env peptides. It should be noted that the antigens used in the SHIP vaccines have been designed to be more immunogenic than those used in the SAAVI vaccines and thus the improvement in the immune response cannot be solely attributed to the increased expression of the Gag and Env due to the inclusion of the porcine circovirus in the DNA vaccines. However, the SHIP DNA vaccines elicited an HIV-specific T cell response despite being administered at a four-fold lower dose than the SAAVI DNA-C2 vaccine.

6. Future possibilities for enhanced DNA vaccine or expression vectors based on circoviruses

Our group has recently published an investigation of the possibility of using circovirus-derived replication control elements to create replicons, or replicating dsDNA plasmid-like molecules, in plants and in mammalian cells [34]. This followed our extensive success with use of a plant ssDNA geminivirus-derived expression vector in plants as an enhanced expression vector [35]: geminiviruses are very similar to circoviruses in having small circular ssDNA genomes that replicate via a Rep-mediated rolling circle mechanism, and very similar sequences for their nonanucleotide origins of replication (TAATATT/AC vs. TAGTATT/AC). In this study we used a synthetic, partially dimeric clone of the genome of beak and feather disease virus (BFDV), a circovirus generically related to PCV,

to investigate cross-potential of replication between the plant and animal viruses in plants, and replication of the BFDV genome alone in HEK293TT cells.

Initial experiments where both the geminivirus-derived vector (BeYDV) and BFDV genome were introduced into *Nicotiana benthamiana* plants via *Agrobacterium tumefaciens*-mediated DNA transfer showed that replication of BeYDV facilitated the co-replication of BFDV, albeit to levels only 100x less than for BeYDV replicons. More importantly, however, transfection of HEK293TT cells with the BFDV construct resulted in a ten-fold increase in genome copy number after three days. This was the first time that BFDV genomes have been shown to replicate in any animal-derived cell culture, in contrast to PCVs which readily infect a variety of cells [36]. Improvement in replicon copy number could be achieved by expressing BFDV *in trans* from another co-transfected vector with a strong promoter: this is not surprising, considering the native *rep* promoter is quite weak, and is probably not well recognized in mammalian cells, meaning expression *in trans* could mean a far higher availability of Rep.

These results open up a number of fascinating possibilities for using BFDV-derived sequences as replication-competent DNA expression and vaccine vectors, several of which we are currently investigating (W de Moor, G Regnard, A-L Williamson, EP Rybicki, unpublished results and ongoing work). There are currently no small DNA virus-derived vectors in use in vaccinology, other than recombinant adeno-associated viruses (rAAV), and AAV2 has recently been implicated in insertional mutagenesis in human hepatocellular carcinomas [37]. Papilloma- and polyomaviruses are also known to be associated with cancers, which may preclude their use as replicating vectors. The essentially ubiquitous ssDNA torque teno viruses are potentially associated with some human disease conditions, although causation is not proven [38, 39].

Circoviruses have never been implicated in any human disease: although PCV-1 and PCV-2 were famously discovered in live rotavirus vaccines given to millions of children [40], and PCV-1 was shown to be able to infect a human hepatocellular carcinoma cell line [41], there was no evidence that PCV-1 infected the infants given the contaminated rotavirus vaccine [42]. There have been concerns, however, over ssDNA viruses of pigs associated with xenotransplantation in humans [43], and swine-human contacts are frequent and worldwide in agriculture. Thus, use of a circovirus such as BFDV as the source of elements for a replicating DNA expression vector, when the virus is host-restricted to one type of birds and has never been associated with human disease, is probably more likely to be regarded as safe. Our preliminary investigations have revealed considerable promise in this regard; however, these will be reported elsewhere (W de Moor, G Regnard, A-L Williamson, EP Rybicki, unpublished results).

7. Conclusions

It has been over 25 years since DNA vaccines were first introduced and many advances have been made in the field. However, despite showing promise in small animals, with some DNA vaccines being licensed for veterinary use [44, 45], no DNA vaccines have been licensed for human use as immunogenicity is still relatively poor. Thus, a great deal of research has gone into improving the immunogenicity of DNA vaccines. Some of the strategies that have been shown to be effective are: i) RNA optimisation to remove mRNA structures that inhibit ribosomal loading and sequences that inhibit nuclear export of mRNA [46, 47]; ii) codon optimisation [46, 48, 49]; iii) use of Kozak sequences [50]; iv) use of leader sequences to improve stability, translation and secretion [46]; v) use of 3' UTR such as polyadenylation signals and posttranscriptional response elements which are important for nuclear export, translation and mRNA stability [51]; vi) use of different promoters & enhancers [52-54]; vii) the inclusion of genes expressing immunomodulatory molecules in the plasmid vector such as GM-CSF or IL-2 [55, 56]; viii) formulation of DNA vaccines in lipids and polymers [57]; ix) use of better delivery systems [58-60] and x) use of suitable adjuvants [58, 59].

In this review we have only focused on a single method of improving DNA vaccine immunogenicity. This was the use of a short enhancer sequence derived from the circovirus PCV-1 capsid gene promoter to increase recombinant antigen expression. This enhancer element led to increased antigen expression and immunogenicity of HIV-1 subtype C candidate DNA vaccines and

allowed for the use of 10-fold lower doses. The improved performance of the DNA vaccines with these candidates, compared to non-enhanced vectors that went into human clinical trial, has prompted the inclusion of the enhancer into all DNA vaccines under investigation in our research group, with excellent results. Future use of replicating circovirus-derived DNA expression and vaccine vectors may yet open up even more exciting possibilities.

Author Contributions: Both authors contributed equally to the review article.

Funding: This work is based upon research supported by the South African Medical Research Council with funds received from the South African Department of Science and Technology and the South African Research Chairs Initiative of the Department of Science and Technology and National Research Foundation.

Acknowledgments: We acknowledge the pioneering work of Dr Fiona Tanzer, and the gift of the original PCV-1 genome by Dr Kenneth Palmer.

Conflicts of Interest: The authors declare no conflict of interest. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript, or in the decision to publish the results.

References

- [1] Doria-Rose NA, Haigwood NL. DNA vaccine strategies: candidates for immune modulation and immunization regimens. *Methods*. **2003**;31:207-16.
- [2] Forde GM. Rapid-response vaccines--does DNA offer a solution? *Nat Biotechnol*. **2005**;23:1059-62.
- [3] Tanzer FL, Shephard EG, Palmer KE, Burger M, Williamson AL, Rybicki EP. The porcine circovirus type 1 capsid gene promoter improves antigen expression and immunogenicity in a HIV-1 plasmid vaccine. *Virology journal*. **2011**;8:51-422X-8-51.
- [4] Hanke T, Schneider J, Gilbert SC, Hill AV, McMichael A. DNA multi-CTL epitope vaccines for HIV and Plasmodium falciparum: immunogenicity in mice. *Vaccine*. **1998**;16:426-35.
- [5] Im EJ, Nkolola JP, di Gleria K, McMichael AJ, Hanke T. Induction of long-lasting multi-specific CD8+ T cells by a four-component DNA-MVA/HIVA-RENTA candidate HIV-1 vaccine in rhesus macaques. *Eur J Immunol*. **2006**;36:2574-84.
- [6] Sharpe S, Hanke T, Tinsley-Bown A, Dennis M, Dowall S, McMichael A, et al. Mucosal immunization with PLGA-microencapsulated DNA primes a SIV-specific CTL response revealed by boosting with cognate recombinant modified vaccinia virus Ankara. *Virology*. **2003**;313:13-21.
- [7] Stinski MF, Isomura H. Role of the cytomegalovirus major immediate early enhancer in acute infection and reactivation from latency. *Med Microbiol Immunol*. **2008**;197:223-31.
- [8] Chege GK, Burgers WA, Muller TL, Gray CM, Shephard EG, Barnett SW, et al. DNA-MVA-protein vaccination of rhesus macaques induces HIV-specific immunity in mucosal-associated lymph nodes and functional antibodies. *Vaccine*. **2017**;35:929-37.
- [9] Churchyard G, Mlisana K, Karuna S, Williamson AL, Williamson C, Morris L, et al. Sequential Immunization with gp140 Boosts Immune Responses Primed by Modified Vaccinia Ankara or DNA in HIV-Uninfected South African Participants. *PLoS One*. **2016**;11:e0161753.
- [10] Gray GE, Mayer KH, Elizaga ML, Bekker LG, Allen M, Morris L, et al. Subtype C gp140 Vaccine Boosts Immune Responses Primed by the South African AIDS Vaccine Initiative DNA-C2 and MVA-C HIV Vaccines after More than a 2-Year Gap. *Clin Vaccine Immunol*. **2016**;23:496-506.
- [11] Shephard E, Burgers WA, Van Harmelen JH, Monroe JE, Greenhalgh T, Williamson C, et al. A multigene HIV type 1 subtype C modified vaccinia Ankara (MVA) vaccine efficiently boosts immune responses to a DNA vaccine in mice. *AIDS Res Hum Retroviruses*. **2008**;24:207-17.

- [12] Williamson AL, Rybicki E, Shephard E, Gray G, Bekker LG, Downing K, et al. South African HIV-1 vaccine candidates - the journey from the bench to clinical trials. *S Afr Med J*. **2012**;102:452-5.
- [13] Mankertz A, Hillenbrand B. Analysis of transcription of Porcine circovirus type 1. *J Gen Virol*. **2002**;83:2743-51.
- [14] Williamson AL, Rybicki EP. Justification for the inclusion of Gag in HIV vaccine candidates. *Expert Rev Vaccines*. **2016**;15:585-98.
- [15] Abdul-Jawad S, Ondondo B, van Hateren A, Gardner A, Elliott T, Korber B, et al. Increased Valency of Conserved-mosaic Vaccines Enhances the Breadth and Depth of Epitope Recognition. *Mol Ther*. **2016**;24:375-84.
- [16] Barouch DH, O'Brien KL, Simmons NL, King SL, Abbink P, Maxfield LF, et al. Mosaic HIV-1 vaccines expand the breadth and depth of cellular immune responses in rhesus monkeys. *Nat Med*. **2010**;16:319-23.
- [17] Barouch DH, Stephenson KE, Borducchi EN, Smith K, Stanley K, McNally AG, et al. Protective efficacy of a global HIV-1 mosaic vaccine against heterologous SHIV challenges in rhesus monkeys. *Cell*. **2013**;155:531-9.
- [18] Santra S, Liao HX, Zhang R, Muldoon M, Watson S, Fischer W, et al. Mosaic vaccines elicit CD8+ T lymphocyte responses that confer enhanced immune coverage of diverse HIV strains in monkeys. *Nat Med*. **2010**;16:324-8.
- [19] Chapman R, Jongwe TI, Douglass N, Chege G, Williamson AL. Heterologous prime-boost vaccination with DNA and MVA vaccines, expressing HIV-1 subtype C mosaic Gag virus-like particles, is highly immunogenic in mice. *PLoS One*. **2017**;12:e0173352.
- [20] Fischer W, Perkins S, Theiler J, Bhattacharya T, Yusim K, Funkhouser R, et al. Polyvalent vaccines for optimal coverage of potential T-cell epitopes in global HIV-1 variants. *Nat Med*. **2007**;13:100-6.
- [21] Bricault CA, Kovacs JM, Badamchi-Zadeh A, McKee K, Shields JL, Gunn BM, et al. Neutralizing Antibody Responses following Long-Term Vaccination with HIV-1 Env gp140 in Guinea Pigs. *J Virol*. **2018**.
- [22] Cohen KW, Frahm N. Current views on the potential for development of a HIV vaccine. *Expert Opin Biol Ther*. **2017**;17:295-303.
- [23] Plotkin SA. Complex correlates of protection after vaccination. *Clin Infect Dis*. **2013**;56:1458-65.
- [24] Hessel AJ, Poignard P, Hunter M, Hangartner L, Tehrani DM, Bleeker WK, et al. Effective, low-titer antibody protection against low-dose repeated mucosal SHIV challenge in macaques. *Nat Med*. **2009**;15:951-4.
- [25] Julg B, Sok D, Schmidt SD, Abbink P, Newman RM, Broge T, et al. Protective Efficacy of Broadly Neutralizing Antibodies with Incomplete Neutralization Activity against Simian-Human Immunodeficiency Virus in Rhesus Monkeys. *J Virol*. **2017**;91.
- [26] Kim M, Qiao ZS, Montefiori DC, Haynes BF, Reinherz EL, Liao HX. Comparison of HIV Type 1 ADA gp120 monomers versus gp140 trimers as immunogens for the induction of neutralizing antibodies. *AIDS Res Hum Retroviruses*. **2005**;21:58-67.
- [27] Bhiman JN, Anthony C, Doria-Rose NA, Karimanzira O, Schramm CA, Khoza T, et al. Viral variants that initiate and drive maturation of V1V2-directed HIV-1 broadly neutralizing antibodies. *Nat Med*. **2015**;21:1332-6.
- [28] Moore PL, Sheward D, Nonyane M, Ranchope N, Hermanus T, Gray ES, et al. Multiple pathways of escape from HIV broadly cross-neutralizing V2-dependent antibodies. *J Virol*. **2013**;87:4882-94.
- [29] Sanders RW, Vesanen M, Schuelke N, Master A, Schiffner L, Kalyanaraman R, et al. Stabilization of the soluble, cleaved, trimeric form of the envelope glycoprotein complex of human immunodeficiency virus type 1. *J Virol*. **2002**;76:8875-89.
- [30] van Diepen MT, Chapman R, Douglass N, Galant S, Moore PL, Margolin E, et al. Prime boost immunisations with DNA, MVA and protein-based vaccines elicit robust HIV-1, Tier 2 neutralizing antibodies against the CAP256 superinfecting virus. *J Virol*. **2019**.

- [31] van Diepen MT, Chapman R, Moore PL, Margolin E, Hermanus T, Morris L, et al. The adjuvant AlhydroGel elicits higher antibody titres than AddaVax when combined with HIV-1 subtype C gp140 from CAP256. *PLoS One*. **2018**;13:e0208310.
- [32] Asbach B, Kliche A, Kostler J, Perdiguero B, Esteban M, Jacobs BL, et al. Potential To Streamline Heterologous DNA Prime and NYVAC/Protein Boost HIV Vaccine Regimens in Rhesus Macaques by Employing Improved Antigens. *Journal of Virology*. **2016**;90:4133-49.
- [33] Barouch DH, Yang ZY, Kong WP, Koriath-Schmitz B, Sumida SM, Truitt DM, et al. A human T-cell leukemia virus type 1 regulatory element enhances the immunogenicity of human immunodeficiency virus type 1 DNA vaccines in mice and nonhuman primates. *J Virol*. **2005**;79:8828-34.
- [34] Regnard GL, de Moor WRJ, Hitzeroth, II, Williamson AL, Rybicki EP. Xenogenic rolling-circle replication of a synthetic beak and feather disease virus genomic clone in 293TT mammalian cells and *Nicotiana benthamiana*. *J Gen Virol*. **2017**;98:2329-38.
- [35] Regnard GL, Halley-Stott RP, Tanzer FL, Hitzeroth, II, Rybicki EP. High level protein expression in plants through the use of a novel autonomously replicating geminivirus shuttle vector. *Plant Biotechnol J*. **2010**;8:38-46.
- [36] Karuppannan AK, Kwang J. ORF3 of porcine circovirus 2 enhances the in vitro and in vivo spread of the of the virus. *Virology*. **2011**;410:248-56.
- [37] Nault JC, Datta S, Imbeaud S, Franconi A, Mallet M, Couchy G, et al. Recurrent AAV2-related insertional mutagenesis in human hepatocellular carcinomas. *Nat Genet*. **2015**;47:1187-93.
- [38] Eibach D, Hogan B, Sarpong N, Winter D, Struck NS, Adu-Sarkodie Y, et al. Viral metagenomics revealed novel betatorquevirus species in pediatric inpatients with encephalitis/meningoencephalitis from Ghana. *Sci Rep*. **2019**;9:2360.
- [39] Herrmann A, Sandmann L, Adams O, Herrmann D, Dirks M, Widera M, et al. Role of BK polyomavirus (BKV) and Torque teno virus (TTV) in liver transplant recipients with renal impairment. *J Med Microbiol*. **2018**;67:1496-508.
- [40] Gilliland SM, Forrest L, Carre H, Jenkins A, Berry N, Martin J, et al. Investigation of porcine circovirus contamination in human vaccines. *Biologicals*. **2012**;40:270-7.
- [41] Beach NM, Cordoba L, Kenney SP, Meng XJ. Productive infection of human hepatocellular carcinoma cells by porcine circovirus type 1. *Vaccine*. **2011**;29:7303-6.
- [42] Dubin G, Toussaint JF, Cassart JP, Howe B, Boyce D, Friedland L, et al. Investigation of a regulatory agency enquiry into potential porcine circovirus type 1 contamination of the human rotavirus vaccine, Rotarix: approach and outcome. *Hum Vaccin Immunother*. **2013**;9:2398-408.
- [43] Karuppannan AK, Opriessnig T. Possible risks posed by single-stranded DNA viruses of pigs associated with xenotransplantation. *Xenotransplantation*. **2018**;25:e12453.
- [44] Anderson ED, Mourich DV, Fahrenkrug SC, LaPatra S, Shepherd J, Leong JA. Genetic immunization of rainbow trout (*Oncorhynchus mykiss*) against infectious hematopoietic necrosis virus. *Mol Mar Biol Biotechnol*. **1996**;5:114-22.
- [45] Davis BS, Chang GJ, Cropp B, Roehrig JT, Martin DA, Mitchell CJ, et al. West Nile virus recombinant DNA vaccine protects mouse and horse from virus challenge and expresses in vitro a noninfectious recombinant antigen that can be used in enzyme-linked immunosorbent assays. *J Virol*. **2001**;75:4040-7.
- [46] Megati S, Garcia-Hand D, Cappello S, Roopchand V, Masood A, Xu R, et al. Modifying the HIV-1 env gp160 gene to improve pDNA vaccine-elicited cell-mediated immune responses. *Vaccine*. **2008**;26:5083-94.
- [47] Zhou W, Cook RF, Cook SJ, Hammond SA, Rushlow K, Ghabrial NN, et al. Multiple RNA splicing and the presence of cryptic RNA splice donor and acceptor sites may contribute to low expression levels and poor

immunogenicity of potential DNA vaccines containing the env gene of equine infectious anemia virus (EIAV). *Vet Microbiol.* **2002**;88:127-51.

[48] Fath S, Bauer AP, Liss M, Spriestersbach A, Maertens B, Hahn P, et al. Multiparameter RNA and codon optimization: a standardized tool to assess and enhance autologous mammalian gene expression. *PLoS One.* **2011**;6:e17596.

[49] Ko HJ, Ko SY, Kim YJ, Lee EG, Cho SN, Kang CY. Optimization of codon usage enhances the immunogenicity of a DNA vaccine encoding mycobacterial antigen Ag85B. *Infect Immun.* **2005**;73:5666-74.

[50] Kozak M. Point mutations define a sequence flanking the AUG initiator codon that modulates translation by eukaryotic ribosomes. *Cell.* **1986**;44:283-92.

[51] Barrett LW, Fletcher S, Wilton SD. Regulation of eukaryotic gene expression by the untranslated gene regions and other non-coding elements. *Cell Mol Life Sci.* **2012**;69:3613-34.

[52] Garg S, Oran AE, Hon H, Jacob J. The hybrid cytomegalovirus enhancer/chicken beta-actin promoter along with woodchuck hepatitis virus posttranscriptional regulatory element enhances the protective efficacy of DNA vaccines. *J Immunol.* **2004**;173:550-8.

[53] Vanniasinkam T, Reddy ST, Ertl HC. DNA immunization using a non-viral promoter. *Virology.* **2006**;344:412-20.

[54] Wang S, Farfan-Arribas DJ, Shen S, Chou TH, Hirsch A, He F, et al. Relative contributions of codon usage, promoter efficiency and leader sequence to the antigen expression and immunogenicity of HIV-1 Env DNA vaccine. *Vaccine.* **2006**;24:4531-40.

[55] Hellerstein M, Xu Y, Marino T, Lu S, Yi H, Wright ER, et al. Co-expression of HIV-1 virus-like particles and granulocyte-macrophage colony stimulating factor by GEO-D03 DNA vaccine. *Hum Vaccin Immunother.* **2012**;8:1654-8.

[56] Henke A, Rohland N, Zell R, Wutzler P. Co-expression of interleukin-2 by a bicistronic plasmid increases the efficacy of DNA immunization to prevent influenza virus infections. *Intervirology.* **2006**;49:249-52.

[57] O'Hagan DT, Singh M, Ulmer JB. Microparticle-based technologies for vaccines. *Methods.* **2006**;40:10-9.

[58] Burton S, Spicer LM, Charles TP, Gangadhara S, Reddy PBJ, Styles TM, et al. Clade C HIV-1 Envelope Vaccination Regimens Differ in Their Ability To Elicit Antibodies with Moderate Neutralization Breadth against Genetically Diverse Tier 2 HIV-1 Envelope Variants. *J Virol.* **2019**;93.

[59] Li L, Petrovsky N. Molecular Adjuvants for DNA Vaccines. *Curr Issues Mol Biol.* **2017**;22:17-40.

[60] Porter KR, Raviprakash K. DNA Vaccine Delivery and Improved Immunogenicity. *Curr Issues Mol Biol.* **2017**;22:129-38.