

Article

Response relating factors of bone marrow-derived mesenchymal stem cells transplantation in patients with alcoholic cirrhosis

HariPriya Gupta¹, Gi Soo Youn¹, Sang Hak Han², Min Jea Shin¹, Sang Jun Yoon¹, Dae Hee Han¹, Na Young Lee¹, Dong Joon Kim¹, Soon Koo Baik³, Ki Tae Suk^{1*}

¹ Institute for Liver and Digestive Diseases, Hallym University College of Medicine, Chuncheon, 24253, South Korea; phr.haripriya13@gmail.com (H.G.); gisu0428@gmail.com (G.S.Y.); wehome3@hallym.ac.kr (M. J. S.); ysjtlhuman@gmail.com (S.J.Y.); eogmlgks@gmail.com (D.H.H.); na0lee0218@gmail.com (N.Y.L.); djkim@hallym.ac.kr (D.J.K.)

² Department of Pathology, Hallym University College of Medicine, Chuncheon, 24253, South Korea; drhsh74@hallym.or.kr (S.H.H.)

³ Department of Pathology, Wonju Severance Christian Hospital, Yonsei University, Wonju College of Medicine, Wonju 26426, South Korea; baiksk@yonsei.ac.kr (S.K.B.)

* Correspondence: ktsuk@hallym.ac.kr, H.G., G.S.Y., and S.H.H. equally contributed to this study

Abstract: Liver cirrhosis leads to hepatic dysfunction and life-threatening conditions. Though clinical efficacy of autologous bone marrow-derived mesenchymal stem cells (BM-MSC) transplantation in alcoholic cirrhosis (AC) was demonstrated, the relevant mechanism has not been elucidated. We aimed to identify predictive factors and gene/pathways for responders after autologous BM-MSC transplantation. Fifty-five patients with biopsy-proven AC underwent autologous BM-MSC transplantation. The characteristics of responders who showed improvement in Laennec score (≥ 1) after transplantation were compared with those of non-responders. BM-MSCs were analyzed with cDNA microarrays to identify gene/pathways that were differentially expressed in responder. Thirty-three patients (66%) were responders. Initial high Laennec score ($p=0.007$, odds ratio 3.73) and performance of BM-MSC transplantation ($p=0.033$, odds ratio 5.75) were predictive factors for responder. Three genes (olfactory receptor2L8, microRNA4520-2, and chloride intracellular channel protein3) were upregulated in responders and CD36 and retinol-binding protein 4 are associated with biologic processes dominant in non-responder. Eleven pathways (inositol phosphate, ATP-binding cassette transporters, protein-kinase signaling, extracellular matrix-receptor interaction, endocytosis, phagosome, hematopoietic cell lineage, adipocytokine, peroxisome proliferator-activated receptor, fat digestion/absorption, and insulin resistance) were upregulated in non-responders ($p<0.05$). BM-MSC transplantation is warranted treatment for AC patients with high Laennec score. Cell-based therapy utilizing response-relating genes or pathway can be treatment candidate.

Keywords: Cirrhosis; Bone marrow; Mesenchymal stem cells; Characteristics; Liver regeneration

1. Introduction

Cirrhosis is the end stage of liver injury with chronic necrosis and inflammation of cells combined with fibrogenic processes, followed by transformation of normal liver tissue into regenerative nodules [1,2]. Currently, the only definitive treatment for advanced liver cirrhosis is liver transplantation [3]. However, transplantation has limitations such as deficiency in donors, surgical complications, immunosuppression after transplantation, and high cost [4]. New therapeutic options, such as regeneration, bioartificial liver, and cell-based transplantation are performing investigation to improve the prognosis of patients with liver cirrhosis [5].

With recent advancements and research-related cell therapies, mesenchymal stem cell (MSC) have demonstrated improved clinical applications because of their accessibility, expandability, and multipotentiality [6]. Various MSCs have been transplanted for the treatment of liver cirrhosis [7,8]. Recently, the positive influence of MSCs transplantation has been reaffirmed against alcoholic cirrhosis (AC), including clinical trials for autologous bone marrow-derived mesenchymal stem cell (BM-MSC) transplantation to patients with AC, that have shown the affirmative response of MSCs to reduce liver cirrhosis and improve liver function. [9-11].

Therapeutic potentials of MSCs include their immunosuppressive properties, their ability to secrete various trophic factors and hepatic regeneration [8,12]. After the homing of MSCs to injured tissue sites for repair, when in contact with local stimuli, such as inflammatory cytokines and ligands of toll-like receptors (TLRs), these stimuli can trigger MSCs to release a broad spectrum of bioactive molecules related with tissue remodeling and regeneration [10]. Since liver regeneration is mainly regulated by various cytokines and growth factors, such as insulin-like factor 1, fibroblast growth factors, hepatocyte growth factor, and chemokine ligand 2, they are potent mediators of angiogenesis and restrain cell apoptosis [13,14]. However, these mechanisms have been mostly investigated in preclinical studies and application in clinical studies is still ambiguous.

In the current scenario, cell therapies, including hepatocytes or the direct infusion of cells in cirrhotic liver have been shown to be ineffective because obtaining human hepatocytes and maintaining their viability and function are difficult in clinical setting [10]. Despite the keen interest of using stem cells as opposed to liver transplantation, the therapeutic mechanisms have not yet been characterized [15,16]. To better understand the degree of responded and secreted factors by undifferentiated BM-MSCs, we tried to unravel the predictive factors for responder, response relating genes, and signal pathway after BM-MSC transplantation. In addition, we tried to explore molecular, biological and cellular functions of BM-MSCs in AC by microarray analysis so that this study may intercept novel insights into the gene related therapeutic approaches in liver disease that can be utilized for further research.

2. Materials and Methods

2.1. Patients

From January 2013 to November 2015, a randomized open-label trial was prospectively conducted at 12 university-affiliated hospitals in Korea (clinicaltrial.gov: NCT01875081). Patients with biopsy-proven AC were recruited and all patients discontinued alcohol intake for at least 6 months before participating in this study. AC was diagnosed on the basis of imaging studies, history of alcohol consumption, and pathologic examinations of liver biopsy samples [17]. Abstaining from alcohol consumption for a 6-month period has been shown to improve inflammation and to facilitate stable results from liver histology, thereby increasing the power of the sample [18]. The eligibility criteria included age between 20 to 70 years and Child-Pugh scores of B or C. The institutional review boards of all participating hospitals and the Korea Food and Drug Administration (KFDA) approved the protocol (KFDA clinical trial No. 27), and all participating patients provided written informed consent to participate in this study. All protocol used in the study conformed to the ethical guidelines of the 2013 Declaration of Helsinki.

We conducted a baseline evaluation, that included a family history, BMI calculation, abdominal ultrasound, a complete blood count, liver function test, and screening for viral markers. The serum biochemical parameters included total bilirubin, alanine aminotransferase, aspartate aminotransferase, gamma-glutamyl transpeptidase, alkaline phosphatase, albumin, blood urea nitrogen, creatinine, α -fetoprotein, prothrombin time, blood glucose, triglycerides, total protein, and total cholesterol. All patients were tested for viral hepatitis, and their Child-Pugh scores, and model for end-stage liver disease (MELD) scores were calculated.

During the clinical trial, we collected BM-MSCs from passage 4-5 and media from sub-cultured samples. Patients were randomly assigned to a control group or one of two autologous BM-MSC groups, in which patients received one or two hepatic arterial injections of 5×10^7 BM-MSCs 30 days

after bone marrow aspiration. Patients in whom the METAVIR fibrosis score improved (by more than 1 point) post-transplantation were defined as responders. We compared BM-MSCs from passages 4-5 that showed significant good responses ($n=3$) or no response ($n=3$) after transplantation with BM-MSCs from the normal healthy control group ($n=3$). In the analysis of BM-MSCs, cDNA microarrays and bio-mathematical analyses were used to identify the secreted genes and related pathways that were differentially expressed in specific stem cell populations in AC. (Fig. 1)




Figure 1. Study design. AC, alcoholic cirrhosis; BM-MSCs, bone marrow driven mesenchymal stem cells; n, number.

2.2. Bone marrow aspiration, isolation, and culture

All manufacturing and product-testing procedures for the production of clinical-grade autologous MSCs (Livercellgram, Pharmicell Co., Ltd., Seongnam, Korea) were performed in a manner consistent with good manufacturing practices and the regulatory guidelines of the Ministry of Food and Drug Safety. With patients under local anesthesia, roughly 10-20 mL of bone marrow was aspirated from the posterior iliac crest. Mononuclear BM cells were isolated using density-gradient centrifugation (Histopaque-1077, Sigma-Aldrich, St. Louis, MO, USA) and plated in 75 cm² flasks (Falcon, Franklin Lakes, NJ, USA) with low-glucose Dulbecco's modified Eagle's medium (DMEM; Gibco, Grand Island, NY, USA) containing 10% fetal bovine serum (FBS, Gibco) and 20 g/mL gentamicin (Gibco), after which they were cultured at 37°C in a 5% CO₂ atmosphere. After 5-7 days, nonadherent cells were removed through replacement of the medium, and the adherent cells were cultured for 2-3 additional days. After detaching the colonized cells using a trypsin/EDTA solution (Gibco), they were re-plated in 175 cm² flasks. When the cultures reached 70-80% confluence, serial subcultures were performed up to passages 4 or 5 for injection. For the second injection in the two-injection BM-MSC group, some cells from passage -1 were harvested during the cell culture process and cryopreserved in 10% dimethyl sulfoxide (Sigma-Aldrich) and 90% FBS. These cells were thawed according to the injection schedule and sub-cultured to passage 4 or 5.

2.2. Viability, immunophenotype, and differential potential of stem cells

A trypan blue exclusion assay was used to assess the viability of the BM-MSCs. The immunophenotypes of the final harvested BM-MSCs were analyzed utilizing CD14, CD34, CD45, CD73, and CD105 as markers. To determine their immunophenotype, MSCs were stained with the following antibodies conjugated with phycoerythrin or fluorescein isothiocyanate: anti-CD14-FITC, anti-CD34-FITC, anti-CD45-FITC, anti-CD73-PE, and anti-CD105-PE (BD Biosciences, San Jose, CA, USA). Briefly, 5×10^5 cells were re-suspended in 0.2 mL of phosphate-buffered saline and incubated with antibodies at room temperature for 20 minutes. As isotypes control, fluorescein isothiocyanate

- or phycoerythrin-conjugated mouse IgGs were used. The intensity of the fluorescence of the cells was evaluated using flow cytometry (Navios™; Beckman Coulter, Fullerton, CA, USA)(Fig. 1).

Furthermore, the potential of BM-MSCs to differentiate into osteoblasts and adipocytes was tested following procedures published in an earlier study [7]. Osteogenic differentiation was assessed by plating the cells at 2×10^4 cells/cm² in six-well plates and culturing them for 2-3 weeks in osteogenic medium composed of low-glucose DMEM medium supplemented with 10% FBS, 10 mM β -glycerophosphate, 10^{-7} M dexamethasone and 0.2 mM ascorbic acid (Sigma-Aldrich) [19]. To evaluate adipogenic differentiation, BM-MSCs were plated at 2×10^4 cells/cm² in six-well plates and cultured for 1 week. Differentiation was induced through an adipogenic medium composed of 10% FBS, 1 μ M dexamethasone, 0.5 mM 3-isobutyl-1-methylxanthine, 10 μ g/mL insulin, and 100 μ M indomethacin in high-glucose DMEM for 3 additional weeks. The differentiated cells were fixed in 4% paraformaldehyde for 10 minutes, after which they were stained with fresh Oil Red-O solution (Sigma-Aldrich) to detect lipid droplets [7].

2.3. Whole transcript expression arrays and data preparation

RNA extraction procedures were performed according to the manufacture's guidelines. A ND-1000 Spectrophotometer (NanoDrop, Wilmington, DE, USA) and a Agilent 2100 Bioanalyzer (Agilent Technologies, Palo Alto, CA, USA) were used to evaluate purity and integrity. The Affymetrix Whole transcript expression array process was performed following the manufacturer's protocol (GeneChip Whole Transcript PLUS reagent Kit). cDNA was synthesized using the GeneChip WT (Whole Transcript) amplification kit, according to the manufacturer's instructions. Fragmentation and biotin-labeling of the sense cDNA were performed with terminal deoxynucleotidyl transferase (TdT) using the GeneChip WT Terminal labeling kit. Roughly 5.5 μ g of the labeled DNA target was hybridized to the Affymetrix GeneChip Human 2.0 ST Array at 45°C for 16 hours. The hybridized arrays were washed and stained using a GeneChip Fluidics Station 450 and scanned on a GCS3000 Scanner (Affymetrix). The Affymetrix® GeneChip™ Command Console software was used to compute signal values.

Raw data were automatically obtained by means of the Affymetrix data extraction protocol in the Affymetrix GeneChip® Command Console® software. Once the CEL files were imported, the data were summarized and normalized using the robust multi-average method implemented in the Affymetrix® Expression Console™ software. The results of the gene-level analysis were exported and differentially expressed gene (DEG) analysis were performed. A comparative analysis between the test and control samples was carried out using the LPE test to calculate fold changes, and the null hypothesis was demonstrated supported by the absence of any significant difference between the two groups. The false discovery rate was controlled by using the Benjamini-Hochberg algorithm to adjust the p-value. For each DEG set, a hierarchical cluster analysis was performed using complete linkage and Euclidean distance as a similarity measure.

2.4. Statistical analysis

The chi-square test and Student's t-test were used to evaluate baseline characteristics. Categorical variables were analyzed by the chi-square test, and continuous variables were assessed by Student's t-test and paired t-test. Multivariate logistic regression was used to detect relating factors for responder BM-MSC transplantation. Data were analyzed with statistical software (SPSS, version 19.0, SPSS, Inc., Chicago, IL, USA) and GraphPad Prism version 6.0 for Windows (GraphPad Software, San Diego, CA, USA). For all tests, p-values < 0.05 were considered significant.

Gene enrichment and functional annotation analyses for significant probe lists were performed using Gene Ontology (<http://geneontology.org/>) and KEGG (<http://kegg.jp>). All statistical tests and visualization of differentially expressed genes were conducted using R v. 3.1.2. statistical software (www.r-project.org).

3. Results

3.1. Patients

A total of 72 patients with AC were randomly assigned to the three groups, and 17 patients were excluded after randomization. In the end, 55 patients (18 in the control group and 37 in the BM-MSC group) completed this study. After 6 month of the BM-MSC transplantation, liver biopsy was performed for histological analysis of treated AC patients. Thirty-three patients (66%) were identified as responders in this study. In the comparison between responders and non-responders, the initial high Laennec score was significantly increased only in the responder group ($p < 0.05$). Other variables did not show differences in baseline characteristics between responders and non-responders (Table 1 and Table 2).

Table 1. Baseline characteristics of patients.

Variables	AC control		P-value	BM-MSC transplantation		P-value
	Responder (n=8)	Non-responder (n=10)		Responder (n=25)	Non-responder (n=12)	
Male (n, %)	8 (100)	9 (90)	NS	21 (84)	9 (82)	NS
Age ^a	51 (8)	54 (8)	NS	54 (8)	55 (9)	NS
Functional analysis ^a						
Child-Pugh score	8.5 (1.4)	8.0 (1.5)	NS	7.8 (1.1)	7.8 (1.1)	NS
MELD score	14.9 (4.9)	12.8 (3.2)	NS	10.3 (3.9)	8.3 (4.7)	NS
Histologic analysis ^a						
Laennec score	8.9 (1.4)	7.5 (1.1)	0.029	7.7 (1.2)	6.7 (1.2)	0.021
Biochemical analysis ^a						
AST (IU/L)	61 (52)	42 (9)	NS	39 (18)	43 (15)	NS
ALT (IU/L)	37 (28)	28 (9)	NS	23 (10)	22 (13)	NS
Albumin (g/dL)	3.2 (0.6)	3.6 (0.5)	NS	3.7 (0.8)	3.3 (0.6)	NS
Bilirubin (mg/dL)	3.5 (2.9)	2.4 (1.5)	NS	1.6 (1.2)	1.5 (1.0)	NS
ALP (IU/L)	169 (66)	97 (35)	0.009	118 (68)	169 (88)	NS
GGT (IU/L)	74 (97)	61 (46)	NS	69 (68)	78 (53)	NS

a Continuous variables are expressed as mean values (standard deviation)

n, number; MELD, AST, aspartate aminotransferase; ALT, alanine aminotransferase; ALP, alkaline phosphatase; GGT, gamma glutamyl transferase; NS, not significant

In the multivariate analysis, the initial high Laennec score ($p = 0.007$, odds ratio 3.73) and performance of BM-MSC transplantation ($p = 0.033$, odds ratio 5.75) were predictive factors for the responders (Supplementary Table 1).

Table 2. Characteristics of responder and non-responder.

Variables	Responder (n=33)		P-value	Non-responder (n=22)		P-value
	Pre	Post		Pre	Post	
Functional analysis ^a						
Child-Pugh score	7.7 (1.1)	6.7 (1.7)	<0.001	8.1 (1.4)	7.3 (2.1)	NS
Histologic analysis ^a						
Laennec score	8.0 (1.3)	6.3 (1.3)	<0.001	7.1 (1.2)	7.4 (1.5)	0.031
Biochemical analysis ^a						
AST (IU/L)	44 (31)	38 (21)	0.012	47 (15)	49 (23)	0.036
ALT (IU/L)	25 (17)	21 (11)	NS	28 (12)	30 (14)	NS

Albumin (g/dL)	3.7 (0.7)	4.1 (1.9)	NS	3.3 (0.6)	3.3 (0.6)	NS
Bilirubin (mg/dL)	1.8 (1.2)	1.8 (1.5)	NS	2.7 (2.3)	2.1 (0.9)	NS
ALP (IU/L)	118 (59)	104 (35)	NS	142 (74)	126 (58)	NS
GGT (IU/L)	76 (70)	75 (50)	NS	57 (52)	77 (75)	NS

a Continuous variables are expressed as mean values (standard deviation)

n, number; MELD, AST, aspartate aminotransferase; ALT, alanine aminotransferase; ALP, alkaline phosphatase; GGT, gamma glutamyl transferase; NS, not significant

3.2. Viability, immunophenotype, and differentiation potential

The mean viability percentage of the BM-MSCs, assessed by the trypan blue exclusion assay was approximately 86-89% in all groups (approved viability is greater than 70%). In the immunophenotyping analysis, BM-MSCs were positive for CD73 and CD105 and negative for CD14, CD34, and CD45 [7]. At the second passage, it was observed that BM-MSCs had successfully differentiated into osteoblasts and adipocytes as evidenced by 5-bromo-4-chloro-3-indolyl phosphate/intro blue tetrazolium staining for alkaline phosphatase activity and Oil Red-O staining for lipids.

3.3. Response relating biological processes and signal pathway

In the analysis between the AC and control groups, a total of 1,028 terms were identified. Among all detected terms, 823 in biological processes, 109 in cellular components, and 98 in molecular functions were identified. The most significant terms are listed in Supplementary Table 2, 3, 4, and 5. In the analysis between non-responders and responders, all related terms were divided into different processes, resulting in the identification of 1,160 terms. Among all detected terms, there were 947 in biological processes, 90 in cellular components, and 123 in molecular functions. The most significant of all the terms were found to be associated with biological processes, which are listed in Table 3.

Table 3. Significant biological process between non-responder and responder.

Term	P-value	Genes
Lipid transport	0.012	ABCA9, RBP4, CD36
Lipid localization	0.015	ABCA9, RBP4, CD36
Secretion	0.022	MEG3, RBP4, STC1, CD36
System process	0.024	EYA4, OR2L8, MEG3, RBP4, STC1
Positive regulation of production of molecular mediator of immune response	0.037	RBP4, CD36
Establishment of localization	0.040	ABCA9, GEM, MEG3, RBP4, STC1, CLIC3, CD36
Anion transport	0.042	STC1, CLIC3, CD36

ABCA9, ATP binding cassette subfamily A member 9; RBP4, retinol binding protein-4; CD36, CD36 Molecule; MEG3, maternally expressed gene 3; STC1, stanniocalcin-1; EYA4, eyes absent homolog 4; OR2L8, olfactory receptor family 2 subfamily L member 8; GEM, GTP-binding protein GEM; CLIC3, chloride intracellular channel 3

Among all evaluated pathways, 15 pathways showed statistical significance between the AC and normal control groups (Fig. 2). The Ras signaling, Wnt signaling, antigen processing and presentation, Axon guidance, and graft-versus-host disease pathways were upregulated, whereas metabolic, arginine and proline metabolism, TGF (transforming growth factor)-beta signaling, HIF (hypoxia-inducible factor) signaling, FoxO signaling, cytokine interaction, stem cell pluripotency, natural killer cell mediated cytotoxicity, and central carbon metabolism were all downregulated.




Figure 2. Microarray analysis between AC and normal control group. A. Heat map depicting the expression of differentially expressed genes encoding various markers in BM-MSCs. B. the KEGG Pathway mapped for significantly dysregulated pathways in AC and normal control group. TGF, transforming growth factor; HIF, hypoxia-inducible factor

In the comparison between the non-responder and responder groups, 12 pathways showed statistical significance which included the metabolic, inositol phosphate metabolism, ABC (ATP-binding cassette) transporters, AMPK (AMP-activated protein kinase) signaling, ECM (extracellular matrix)-receptor interaction, endocytosis, phagosome, hematopoietic cell lineage, adipocytokine signaling, PPAR (peroxisome proliferator-activated receptor) signaling, fat digestion and absorption, olfactory transduction, and insulin resistance (Fig. 3).




Figure 3. Microarray analysis between non-responder and responder. A. Heat map depicting the expression of differentially expressed genes encoding various markers in BM-MSCs. B. the KEGG Pathway mapped for significantly dysregulated pathways in non-responder and responder.

ABC, ATP-binding cassette transporters; AMKP, AMP-activated protein kinase; ECM, extracellular matrix; PPAR, peroxisome proliferator-activated receptor

3.4. Differentially expressed genes

To detect the most dysregulated genes of significance, we showed the top 10 DEGs between the AC and normal control groups. Among the top 10 DEGs, 6 DEGs (GDNF family receptor alpha-1 [GFRA1], fibrillin-2 precursor [FBN2], RAS P21 protein activator 4B [RASA4B], killer cell immunoglobulin like receptor, three Ig domains, and long cytoplasmic tail 2, microRNA 4436a [KIR3DL2], and SERPINF1) were increased and stanniocalcin-1 (STC1), RNA, U5F small nuclear 1 (RNU5F-1), tektin 4 pseudogene 2 (TEKT4P2) and zinc finger protein (ZFY) were downregulated in AC group (Supplementary Table 5).

In the comparison the between non-responder and responder groups, 10 DEGs (maternally expressed gene 3 [MEG3], retinol binding protein 4 [RBP4], LOC101928395, sortilin related VPS10 domain containing receptor 2 [SORCS2], MIR2355, MIR520D, zinc finger protein Y-Linked platelet glycoprotein 4 [ZFY], chloride intracellular channel 3 [CLIC3], MIR4520-2 and olfactory receptor family 2 subfamily L member 8 [OR2L8]) were significantly upregulated and CLIC3, MIR4520-2 and OR2L8 were downregulated in non-responder (Table 4).

Table 4. Profiling of up or down regulated gene expression in bone marrow-driven mesenchymal stem cells between non-responder and responder group.

Gene	Fold change	Function
MEG3	2.93	Growth suppressor in tumor cells and cell growth Activator for p53 and cell apoptosis
RBP4	2.38	Delivers retinol from the liver stores to the peripheral tissues
LOC101928395	2.22	NA
SORCS2	2.02	Represents the only module of the luminal/extracellular moiety Exhibit disparate functions depending on its proteolytic processing
LINC01111	1.97	NA
MIR2355	1.91	NA
MIR520D	1.86	Regulation of malignancy and maintaining p53 upregulation
ZFY	1.80	Function as a transcription factor
CD36	1.80	Function as receptors engaged in platelet adhesion to collagen Initiates signal transduction and internalization of receptor-ligand complexes.
STC1	1.79	Regulation of renal and intestinal calcium and phosphate transport, cell metabolism, or cellular calcium/phosphate homeostasis.
CLIC3	-2.02	Regulator of a recycling pathway and regulates cell migration and invasion
MIR4520-2	-1.70	NA
OR2L8	-1.61	Responsible for the recognition/G protein-mediated transduction of odorant signals.

MEG3, maternally expressed gene 3; RBP4, retinol binding protein 4; SORCS2, sortilin related VPS10 domain containing receptor 2; LINC01111, long intergenic non-protein coding RNA 1111; MIR, microRNA; ZFY, zinc finger protein Y-Linked platelet glycoprotein 4; STC, stanniocalcin1; CLIC3, chloride intracellular channel 3; OR2L8, olfactory receptor family 2 subfamily L member 8

4. Discussion

From our previous lab data, it was corroborated that transplantation of BM-MSs to AC patients has proved to be new perspective in this area of research. Improved Child-Pugh score ($p < 0.05$) after 6 months of BM-MS transplantation showed changes in the liver fibrosis which provided insight of regeneration of the liver tissue. Although other characteristics of the cirrhosis patients were statistically insignificant, but mild improvement was observed which provided valuable insight into the efficacy of BM-MS transplantation in AC patients [9]. The related clinical factors and precise mechanism of response after BM-MS transplantation have not been elucidated. Thus, this intriguing function of BM-MSs led us to perform the microarray analysis so as to unfold possible genes and their signaling pathways involved in BM-MSs function.

In this study, the initial high Laennec score (odds ratio 3.73) and BM-MS transplantation (odds ratio 5.7) were predictive factors for the responders. In a previous *in vivo* study, MSC sheets prevented the growth of hepatocellular carcinoma and progression of advanced stage of cirrhosis [20]. Regarding the initial high Laennec score, the exact mechanism of why the advanced stage in histology is prone to a good response is not clear. In this study, we strictly selected stable patients that abstained from alcohol consumption for 6 months. Because patients with a high Laennec score have increased potential for reversal of fibrosis, an initial high Laennec score might be supposed to be the predictor for good responders. Additional clinical studies on the mechanism are needed in the future.

In this study, 33 patients (66%) were identified as responders after transplantation, and MSC transplantation is another predictive factor for good responders. As we discussed in the previous trial, MSC transplantation has been identified as a promising treatment for liver regeneration in patients with AC [21,22]. As a result, BM-MS transplantation effectively improved the fibrosis stage and liver testing in patients with AC.

In the present study, we identified specific genes and evaluated their pharmacological activity could affect the regulation of liver cirrhosis. By using a cDNA microarray and immunophenotyping assays, we focused on identifying of MSC functions and related trophic factors for secreted genes and their pathways that are differentially expressed in specific stem cell populations in patients with liver cirrhosis.

Through microarray analysis, 466 DEGs were upregulated and 449 DEGs were downregulated in all the groups. This finding suggests that the effect of BM-MS transplantation therapy in the AC group was a multigene-regulated complex process. In addition, the functions of the identified differentially expressed genes were involved in various biological processes, cellular components and molecular functions and various related signaling pathways. Six genes were significantly upregulated, and 9 genes were downregulated in the AC group compared with the control group. In the comparison between responders and non-responders, 3 genes (OR2L8, MIR4520-2, and CLIC3, $p < 0.05$) were functionally upregulated and 28 genes are significantly downregulated in the responders.

Of biologic process and relating genes, lipid associated process is upregulated and CD36 is involved in most biologic process which dominant in non-responder. CD36 has also been implicated in hemostasis, thrombosis, malaria, inflammation, lipid metabolism and atherogenesis [23,24]. CD36 is expressed on multiple cell types and has numerous functions. In a previous study, patient who got hematogenous stem cells transplantation with CD36 positive showed transfusion refractoriness which was significantly improved after transplantation from CD36 deficiency donor [25]. Taken together, CD36 is related with non-responder after transplantation and can be defined as a biomarker for expecting responsiveness in AC patients.

RBP4 is secreted mainly by hepatocytes and acts as the major transporter for vitamin A and retinol acid in serum [26,27]. RBP4 was overexpressed in non-responders in this study. Overexpression of RBP4 resulted in cancer cell migration, and RBP4 was upregulated in ovarian cancer cells. MMP-2 and MMP-9, which are key factors in cancer metastasis, were affected by RBP4 overexpression. This suggested that RBP4 stimulates cancer cell migration along with cell proliferation [28]. Overall, RBP4 could be a negative factor in the response to MSC transplantation, although it is related to cell migration in tumors.

CD36 and RBP4 are upregulated in non-responder and commonly associated genes in all biologic processes. CD36 and RBP4 can be used to as predictive markers after transplantation as the expression is negatively associated with poor outcome in various human signal pathway. So far, the mechanism has not been elucidated, so further researches evaluating the role of CD36 and RBP4 are needed in the future.

Long noncoding RNAs, such as the tumor-suppressing MEG3, have been identified as regulators of various types of human tumors. Previous reports have indicated a positive correlation with human diseases showing abnormal differentiation of BM-MSCs [29]. In one report, MEG3 inhibits of cell proliferation and down regulation of MEG3 activates the Wnt/beta-catenin signaling pathway via epigenetically regulating Wnt genes promotor [30]. Another study demonstrated that down regulation of MEG3 promotes BM-MSCs differentiation [31]. In this study, MEG3 expression was significantly increased in non-responders compared with responders. Taken together, MEG3 is down regulated in responder and further study using MEG3 gene are needed in the future.

CLIC3 is clearly increased in the responder group in this result. CLIC3 acts as a key regulator of cell migration in tissues and as an independent prognostic indicator in pancreatic cancer [32]. Additionally, CLIC3 regulates the recycling of membrane-type 1 matrix metalloproteinase, and transport of endosomes and/or lysosomes back to the plasma membrane and demonstrates potential for the development of new therapies for more efficient mobilization and homing [33]. Moreover, CLIC3, which is expressed in mesenchymal stromal cells, demonstrated a lineage decision-making process and is overexpressed during osteogenic differentiation [34]. This finding suggests that CLIC3 is a lineage-specific gene and could be a potential factor for the differentiation of MSCs to hepatocytes. Additional investigations should be performed to determine the exact mechanism of CLIC3 in the future.

Human miR-520d is a minor miRNA, and its upregulation was observed to induce tumor-suppressive effects and inhibit metastasis [35]. miR-520d-5p is suspected to act in the transformation of hepatoma cells to noncancerous cells in vivo via p53 upregulation [36]. The signaling pathway induced by miR-520d-5p has been shown to inhibit cell motility and invasiveness [37]. Another study demonstrated that miR-520d inhibits c-Myc, Cyclin D1, and MMP9 expression and their inhibitory role demonstrated a the reduction in proliferation, invasion, and migration of cancer cells [38]. Our results are similar to those reported in previous studies. Additionally, nonresponder showed increased expression of miR-520d.

STC1, glycoprotein, having pleiotropic effects may show autocrine and paracrine functions. In two separate studies, STC1 mediates anti-apoptotic effects of MSCs in Lung epithelial cells [39], and MSCs exhibited anti-fibrotic impact via STC1 secretion in pulmonary fibrosis that manifest paracrine function by the reduction in TGF- β 1 production from alveolar macrophages [40]. Thus these facts suggest that STC1 is directly related to MSCs and down-regulation of STC1 disrupt MSCs function which can be clearly seen in our study in the AC group. GFRA1 and FBN2 upregulation in the AC group can also be related with the previous studies that show implication in carcinogenesis [41,42].

Biological process analysis indicated that DEGs in the BM-MSC treated group were mainly involved in the processes of biological regulation, stress development, immune response, cellular localization, cellular response to hypoxia and secretions. It has been found that HIF-1 regulates the expression of STC1 as well as BNIP3 and shows a relation to the Warburg effect in the progression of tumorigenesis under hypoxic conditions [43]. Both BNIP3 and STC1 are hypoxia responsive genes and their overexpression actuates tumor progression. Furthermore, downregulation of these genes alleviates their effects and promotes cancer cell apoptosis. Additionally, MCSs under hypoxic conditions have been shown to downregulate the proapoptotic protein BNIP3 by exerting paracrine effects [44]. Metabolic reprogramming has been shown to increase tumorigenesis in harsh cellular environments, such as hypoxia and oxidative stress. A previous study also inferred that high expression in the liver of PDK1, a downstream target of HIF-1 α factor, regulates metabolism by metabolic reprogramming, and it has been shown to promote liver metastasis [45]. Therefore, these studies, together with ours, specify the conclusive effect of BM-MSC transplantation in AC compared

with the normal control group. BM-MSC treatment of AC patients was mediated by the interplay between multiple genes involved in the regulation of diversified biological systems.

Via of KEGG pathway analysis, pathways that reduce liver injury and promote its growth were identified. BM-MSC transplantation upregulated the protein that negatively regulates the Ras signaling pathway and RASA4b. RASA4b is a GTPase-activating protein that subdues the RAS/MAPK pathway, resulting in the inactivation of the signaling pathway [46]. Since aberrant activation of the RAS/MAPK signaling pathway is implicated in liver cancer progression [47], its inactivation can prevent downstream pathways and contribute to the survival of hepatocytes. However, RASA4b does not have a direct relationship with BM-MSCs, and further study is needed to establish a direct relationship with BM-MSCs. Thus, BM-MSC transplantation can improve cirrhosis in association with the participation of multiple genes and signaling pathways.

5. Conclusions

BM-MSC transplantation is an effective treatment in patients with AC, and responders revealed different gene expression and signaling pathways such as CD36, RBP4, and lipid associated process compared with non-responders. Therefore, cell-based therapy utilizing response-related genes is warranted in patients with AC, and this study may act as a guide for future fundamental and clinical research.

Supplementary Materials: The following are available online: Table S1, S2, S3, and S4

Author Contributions: conceptualization, H.G. S.H.H. and K.T.S.; methodology, H.G.; software, G.S.Y.; investigation, K.T.S.; writing—original draft preparation, S.J.Y.; writing—review and editing, H.G. S.H.H. and K.T.S.; supervision, M.J.S.; project administration, D.J.K. and S.K.B; funding acquisition, K.T.S..

Funding: This research was funded by Hallym University Research Fund, Korea National Research Foundation (NRF-2017R1A5A2015369 and NRF-2018M3A9F3020956), and Hallym University Research Fund 2016 and 2017 (HURF-2016-60 and HURF-2017-18)(K.T.S.).

Acknowledgments: In this section you can acknowledge any support given which is not covered by the author contribution or funding sections. This may include administrative and technical support, or donations in kind (e.g., materials used for experiments).

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Gao, B.; Bataller, R. Alcoholic liver disease: Pathogenesis and new therapeutic targets. *Gastroenterology* **2011**, *141*, 1572-1585.
2. Suk, K.T.; Baik, S.K.; Yoon, J.H.; Cheong, J.Y.; Paik, Y.H.; Lee, C.H.; Kim, Y.S.; Lee, J.W.; Kim, D.J.; Cho, S.W., *et al.* Revision and update on clinical practice guideline for liver cirrhosis. *Korean J Hepatol* **2012**, *18*, 1-21.
3. Burra, P.; Senzolo, M.; Adam, R.; Delvart, V.; Karam, V.; Germani, G.; Neuberger, J.; Elita; Centers, E.L.T. Liver transplantation for alcoholic liver disease in europe: A study from the eltr (european liver transplant registry). *American journal of transplantation : official journal of the American Society of Transplantation and the American Society of Transplant Surgeons* **2010**, *10*, 138-148.
4. Lucey, M.R. Liver transplantation in patients with alcoholic liver disease. *Liver transplantation : official publication of the American Association for the Study of Liver Diseases and the International Liver Transplantation Society* **2011**, *17*, 751-759.
5. Khan, J.A.; Mendelson, A.; Kunisaki, Y.; Birbrair, A.; Kou, Y.; Arnal-Estape, A.; Pinho, S.; Ciero, P.; Nakahara, F.; Ma'ayan, A., *et al.* Fetal liver hematopoietic stem cell niches associate with portal vessels. *Science* **2016**, *351*, 176-180.
6. Ullah, I.; Subbarao, R.B.; Rho, G.J. Human mesenchymal stem cells - current trends and future prospective. *Biosci Rep* **2015**, *35*.
7. Jang, Y.O.; Kim, Y.J.; Baik, S.K.; Kim, M.Y.; Eom, Y.W.; Cho, M.Y.; Park, H.J.; Park, S.Y.; Kim, B.R.; Kim, J.W., *et al.* Histological improvement following administration of autologous bone marrow-derived mesenchymal stem cells for alcoholic cirrhosis: A pilot study. *Liver international : official journal of the International Association for the Study of the Liver* **2014**, *34*, 33-41.

8. Alfaifi, M.; Eom, Y.W.; Newsome, P.N.; Baik, S.K. Mesenchymal stromal cell therapy for liver diseases. *J Hepatol* **2018**, *68*, 1272-1285.
9. Suk, K.T.; Yoon, J.H.; Kim, M.Y.; Kim, C.W.; Kim, J.K.; Park, H.; Hwang, S.G.; Kim, D.J.; Lee, B.S.; Lee, S.H., *et al.* Transplantation with autologous bone marrow-derived mesenchymal stem cells for alcoholic cirrhosis: Phase 2 trial. *Hepatology* **2016**, *64*, 2185-2197.
10. Eom, Y.W.; Kim, G.; Baik, S.K. Mesenchymal stem cell therapy for cirrhosis: Present and future perspectives. *World J Gastroenterol* **2015**, *21*, 10253-10261.
11. Shim, K.Y.; Eom, Y.W.; Kim, M.Y.; Kang, S.H.; Baik, S.K. Role of the renin-angiotensin system in hepatic fibrosis and portal hypertension. *Korean J Intern Med* **2018**, *33*, 453-461.
12. Li, D.L.; He, X.H.; Zhang, S.A.; Fang, J.; Chen, F.S.; Fan, J.J. Bone marrow-derived mesenchymal stem cells promote hepatic regeneration after partial hepatectomy in rats. *Pathobiology* **2013**, *80*, 228-234.
13. Wei, X.; Yang, X.; Han, Z.P.; Qu, F.F.; Shao, L.; Shi, Y.F. Mesenchymal stem cells: A new trend for cell therapy. *Acta Pharmacol Sin* **2013**, *34*, 747-754.
14. Kim, D.; Cho, G.S.; Han, C.; Park, D.H.; Park, H.K.; Woo, D.H.; Kim, J.H. Current understanding of stem cell and secretome therapies in liver diseases. *Tissue Eng Regen Med* **2017**, *14*, 653-665.
15. Higashiyama, R.; Inagaki, Y.; Hong, Y.Y.; Kushida, M.; Nakao, S.; Niioka, M.; Watanabe, T.; Okano, H.; Matsuzaki, Y.; Shiota, G., *et al.* Bone marrow-derived cells express matrix metalloproteinases and contribute to regression of liver fibrosis in mice. *Hepatology* **2007**, *45*, 213-222.
16. Jang, Y.O.; Jun, B.G.; Baik, S.K.; Kim, M.Y.; Kwon, S.O. Inhibition of hepatic stellate cells by bone marrow-derived mesenchymal stem cells in hepatic fibrosis. *Clinical and molecular hepatology* **2015**, *21*, 141-149.
17. European Association for the Study of, L. Easl clinical practical guidelines: Management of alcoholic liver disease. *J Hepatol* **2012**, *57*, 399-420.
18. D'Amico, G.; Garcia-Tsao, G.; Pagliaro, L. Natural history and prognostic indicators of survival in cirrhosis: A systematic review of 118 studies. *J Hepatol* **2006**, *44*, 217-231.
19. Hu, Y.; Liao, L.; Wang, Q.; Ma, L.; Ma, G.; Jiang, X.; Zhao, R.C. Isolation and identification of mesenchymal stem cells from human fetal pancreas. *The Journal of laboratory and clinical medicine* **2003**, *141*, 342-349.
20. Alshareeda, A.T.; Alsowayan, B.; Almubarak, A.; Alghuwainem, A.; Alshawakir, Y.; Alahmed, M. Exploring the potential of mesenchymal stem cell sheet on the development of hepatocellular carcinoma in vivo. *J Vis Exp* **2018**.
21. Si-Tayeb, K.; Noto, F.K.; Nagaoka, M.; Li, J.; Battle, M.A.; Duris, C.; North, P.E.; Dalton, S.; Duncan, S.A. Highly efficient generation of human hepatocyte-like cells from induced pluripotent stem cells. *Hepatology* **2010**, *51*, 297-305.
22. Chamberlain, G.; Fox, J.; Ashton, B.; Middleton, J. Concise review: Mesenchymal stem cells: Their phenotype, differentiation capacity, immunological features, and potential for homing. *Stem cells* **2007**, *25*, 2739-2749.
23. Collins, R.F.; Touret, N.; Kuwata, H.; Tandon, N.N.; Grinstein, S.; Trimble, W.S. Uptake of oxidized low density lipoprotein by cd36 occurs by an actin-dependent pathway distinct from macropinocytosis. *J Biol Chem* **2009**, *284*, 30288-30297.
24. Stewart, C.R.; Stuart, L.M.; Wilkinson, K.; van Gils, J.M.; Deng, J.; Halle, A.; Rayner, K.J.; Boyer, L.; Zhong, R.; Frazier, W.A., *et al.* Cd36 ligands promote sterile inflammation through assembly of a toll-like receptor 4 and 6 heterodimer. *Nat Immunol* **2010**, *11*, 155-161.
25. Zhou, Y.; Li, L.L.; Zhong, Z.L.; Liu, X.J.; Liu, J.L.; Shen, W.D.; Wu, G.G. [anti-cd36 mediated platelet transfusion refractoriness and related cases after stem cell transplantation]. *Zhongguo Shi Yan Xue Ye Xue Za Zhi* **2018**, *26*, 541-546.
26. Yang, Q.; Graham, T.E.; Mody, N.; Preitner, F.; Peroni, O.D.; Zabolotny, J.M.; Kotani, K.; Quadro, L.; Kahn, B.B. Serum retinol binding protein 4 contributes to insulin resistance in obesity and type 2 diabetes. *Nature* **2005**, *436*, 356-362.
27. Blaner, W.S. Retinol-binding protein: The serum transport protein for vitamin a. *Endocr Rev* **1989**, *10*, 308-316.
28. Li, X.; Fei, T.; Wang, Y.; Zhao, Y.; Pan, Y.; Li, D. Wheat starch with low retrogradation properties produced by modification of the gtfb enzyme 4,6-alpha-glucanotransferase from streptococcus thermophilus. *J Agric Food Chem* **2018**, *66*, 3891-3898.
29. Zhuang, W.; Ge, X.; Yang, S.; Huang, M.; Zhuang, W.; Chen, P.; Zhang, X.; Fu, J.; Qu, J.; Li, B. Upregulation of lncrna meg3 promotes osteogenic differentiation of mesenchymal stem cells from multiple myeloma patients by targeting bmp4 transcription. *Stem cells* **2015**, *33*, 1985-1997.

30. Deng, L.; Hong, H.; Zhang, X.; Chen, D.; Chen, Z.; Ling, J.; Wu, L. Down-regulated lncrna meg3 promotes osteogenic differentiation of human dental follicle stem cells by epigenetically regulating wnt pathway. *Biochem Biophys Res Commun* **2018**, *503*, 2061-2067.
31. Sun, X.; Luo, L.H.; Feng, L.; Li, D.S. Down-regulation of lncrna meg3 promotes endothelial differentiation of bone marrow derived mesenchymal stem cells in repairing erectile dysfunction. *Life Sci* **2018**, *208*, 246-252.
32. Dozynkiewicz, M.A.; Jamieson, N.B.; Macpherson, I.; Grindlay, J.; van den Berghe, P.V.; von Thun, A.; Morton, J.P.; Gourley, C.; Timpson, P.; Nixon, C., *et al.* Rab25 and clic3 collaborate to promote integrin recycling from late endosomes/lysosomes and drive cancer progression. *Dev Cell* **2012**, *22*, 131-145.
33. Macpherson, I.R.; Rainero, E.; Mitchell, L.E.; van den Berghe, P.V.; Speirs, C.; Dozynkiewicz, M.A.; Chaudhary, S.; Kalna, G.; Edwards, J.; Timpson, P., *et al.* Clic3 controls recycling of late endosomal mt1-mmp and dictates invasion and metastasis in breast cancer. *J Cell Sci* **2014**, *127*, 3893-3901.
34. Brum, A.M.; van der Leije, C.S.; Schreuders-Koedam, M.; Verhoeven, J.; Janssen, M.; Dekkers, D.H.; Demmers, J.A.; Eijken, M.; van de Peppel, J.; PTM van Leeuwen, J., *et al.* Identification of chloride intracellular channel protein 3 as a novel gene affecting human bone formation. *Journal of Bone and Mineral Research* **2017**, *1*, 16-26.
35. Miura, N.; Sato, R.; Tsukamoto, T.; Shimizu, M.; Kabashima, H.; Takeda, M.; Takahashi, S.; Harada, T.; West, J.E.; Drabkin, H., *et al.* A noncoding rna gene on chromosome 10p15.3 may function upstream of htert. *BMC Mol Biol* **2009**, *10*, 5.
36. Tsuno, S.; Wang, X.; Shomori, K.; Hasegawa, J.; Miura, N. Hsa-mir-520d induces hepatoma cells to form normal liver tissues via a stemness-mediated process. *Sci Rep* **2014**, *4*, 3852.
37. Tsukerman, P.; Yamin, R.; Seidel, E.; Khawaled, S.; Schmiedel, D.; Bar-Mag, T.; Mandelboim, O. Mir-520d-5p directly targets twist1 and downregulates the metastamir mir-10b. *Oncotarget* **2014**, *5*, 12141-12150.
38. Manh, T.H.; Young, M.K.; Dae-Yeul, Y.; Dae, H.L.; Moonjae, C.; Changlim, H. Tgf- β secreted from activated hepatic stellate cells may induce the transdifferentiation of hepatocytes into hepatocarcinoma in hbx-expressing livers. *J Korean Soc Appl Biol Chem* **2014**, *57*, 529-538.
39. Block, G.J.; Ohkouchi, S.; Fung, F.; Frenkel, J.; Gregory, C.; Pochampally, R.; DiMattia, G.; Sullivan, D.E.; Prockop, D.J. Multipotent stromal cells are activated to reduce apoptosis in part by upregulation and secretion of stanniocalcin-1. *Stem cells* **2009**, *27*, 670-681.
40. Ono, M.; Ohkouchi, S.; Kanehira, M.; Tode, N.; Kobayashi, M.; Ebina, M.; Nukiwa, T.; Irokawa, T.; Ogawa, H.; Akaike, T., *et al.* Mesenchymal stem cells correct inappropriate epithelial-mesenchyme relation in pulmonary fibrosis using stanniocalcin-1. *Mol Ther* **2015**, *23*, 549-560.
41. Kato, N.; Ji, G.; Wang, Y.; Baba, M.; Hoshida, Y.; Otsuka, M.; Taniguchi, H.; Moriyama, M.; Dharel, N.; Goto, T., *et al.* Large-scale search of single nucleotide polymorphisms for hepatocellular carcinoma susceptibility genes in patients with hepatitis c. *Hepatology* **2005**, *42*, 846-853.
42. Chen, H.; Suzuki, M.; Nakamura, Y.; Ohira, M.; Ando, S.; Iida, T.; Nakajima, T.; Nakagawara, A.; Kimura, H. Aberrant methylation of fbn2 in human non-small cell lung cancer. *Lung Cancer* **2005**, *50*, 43-49.
43. Yeung, H.Y.; Lai, K.P.; Chan, H.Y.; Mak, N.K.; Wagner, G.F.; Wong, C.K. Hypoxia-inducible factor-1-mediated activation of stanniocalcin-1 in human cancer cells. *Endocrinology* **2005**, *146*, 4951-4960.
44. Bernard, O.; Jeny, F.; Uzunhan, Y.; Dondi, E.; Terfous, R.; Label, R.; Sutton, A.; Larghero, J.; Vanneaux, V.; Nunes, H., *et al.* Mesenchymal stem cells reduce hypoxia-induced apoptosis in alveolar epithelial cells by modulating hif and ros hypoxic signaling. *Am J Physiol Lung Cell Mol Physiol* **2018**, *314*, L360-L371.
45. Dupuy, F.; Tabaries, S.; Andrzejewski, S.; Dong, Z.; Blagih, J.; Annis, M.G.; Omeroglu, A.; Gao, D.; Leung, S.; Amir, E., *et al.* Pdk1-dependent metabolic reprogramming dictates metastatic potential in breast cancer. *Cell Metab* **2015**, *22*, 577-589.
46. Lockyer, P.J.; Kupzig, S.; Cullen, P.J. Capri regulates ca(2+)-dependent inactivation of the ras-mapk pathway. *Curr Biol* **2001**, *11*, 981-986.
47. Delire, B.; Starkel, P. The ras/mapk pathway and hepatocarcinoma: Pathogenesis and therapeutic implications. *Eur J Clin Invest* **2015**, *45*, 609-623.