

Article

LC-HRMS Screening and Identification of Novel Peptide Markers of Ricin Based on Multiple Protease Digestion Strategies

Long-Hui Liang², Chang-Cai Liu^{1,2*}, Bo Chen², Long Yan², Hui-Lan Yu^{1,2}, Yang Yang^{1,2}, Ji-Na Wu², Xiao-Sen Li², and Shi-Lei Liu^{1,2*}

¹ State Key Laboratory of NBC Protection for Civilian, Beijing 102205, China;

² The laboratory of Analytical Chemistry, Research Institute of Chemical Defence, Beijing 102205, China;

* Correspondence: liuchangcai_0@126.com; Tel.: +86-177-4446-6227

* Correspondence: liu_shilei@lacricd.com; Tel.: +86-136-6120-8709

Abstract: Both ricin and *R. communis agglutinin* (RCA120), belonging to the type II ribosome-inactivating proteins (RIPs-II), are derived from the seeds of castor bean plant. They share very similar amino acid sequences, but ricin is much more toxic than RCA120. It is urgently necessary to distinguish ricin and RCA120 in response to public safety. Currently, mass spectrometric assays are well established for unambiguous identification of ricin by accurate analysis of differentiated amino acid residues after trypsin digestion. However, diagnostic peptides are relatively limited for unambiguous identification of trace ricin, especially in complex matrices. Here, we demonstrate a digestion strategy of multiple proteinases to produce novel peptide markers for unambiguous identification of ricin. LC-HRMS was used to verify the resulting peptides, among which only the peptides with uniqueness and good MS response were selected as peptide markers. Seven novel peptide markers were obtained from tandem digestion of trypsin and endoproteinase Glu-C in PBS buffer. From the chymotrypsin digestion under reduction and non-reduction conditions, eight and seven novel peptides were selected respectively. Using pepsin under pH 1~2 and proteinase K digestion, 6 and 5 peptides were selected as novel peptide markers. In conclusion, the obtained novel peptides from the established digestion methods can be recommended for the unambiguous identification of ricin during the investigation of illegal use of the toxin.

Keywords: ricin; marker peptides; unambiguous identification; mass spectrometry

Key Contribution: Novel peptide markers of ricin were obtained based on multiple protease digestion strategies for unambiguous identification of ricin.

1. Introduction

Type II ribosome-inactivating proteins (RIPs-II) are a class of heterodimeric protein toxins, which consists of two disulfide-linked polypeptide chains. The A-chain is functional chain of N-glycosidase activity responsible for irreversibly inactivating the large ribosome subunits through depurination of a specific adenine from 28S rRNA. The B-chain has lectin activity on binding to the terminal galactose receptors on cell surface allowing for A-chain internalization by endocytosis [1,2]. Ricin is one of the best known RIP-II toxins, which was extracted from the castor bean plant *Ricinus communis*. Exposure symptoms include nausea, vomiting, fever and even death[3]. Because of its wide availability, high toxicity and ease of preparation, ricin has been used in several threat incidents since the beginning of 20th century[4]. As one of the most potential chemical warfare agents and bioterrorism agents, production and possession of ricin have been prohibited by the Chemical Weapons Convention (CWC) since 1997 [5]. In response to the potential use of ricin for

bioterrorism, criminal, or military purposes, establishing accurate methods is very necessary for identification of ricin and other RIP-II protein toxins.

Immunological and functional assay methods have been developed to determine ricin and other RIP-II toxins [6-10]. However, the presence of *R. communis agglutinin* (RCA120) make it difficult for these methods to unambiguously identify ricin due to its identical functional activity and high amino acid sequence similarity (93% in A-chain and 84% in B-chain) with ricin. Furthermore, because of 2-3 orders of less toxicity of RCA120 than ricin [11,12], it is very essential to distinguish ricin from RCA120 for the unambiguous identification of ricin.

Mass spectrometric (MS) techniques provide an alternative approach for unambiguous identification of ricin by accurate determination of amino acid sequence differences between ricin and RCA120. Before MS detection, it is essential to produce unique peptide markers of ricin with suitable molecular weight and enough MS response using proteinase digestion. Currently, tryptic peptides of T7A, T11A, T6B and T18B unique to ricin have become the common and representative peptide markers [13-16]. The peptide markers were widely used for unambiguous identification of ricin in complex matrices including crude extracts of castor bean, protein and stevia powder, beverages and soil, et al [17]. Some efforts were made to enhance efficiency of trypsin digestion. For example, instead of trypsin digestion following the denaturation, reduction and alkylation, the direct trypsin digestion of intact ricin under the solvent-assisted condition was performed with no missed cleavage and reduced digestion time [13,18]. Besides, two peptide markers (C18A and C18B) generated from chymotrypsin digestion were also applied for unambiguous identification of ricin from crude extracts of castor bean and soil samples [19]. However, the selected peptide markers are still relatively limited. Thus, it is necessary to obtain abundant peptide markers produced from more types of proteinase digestion for unambiguous identification of trace ricin, especially in complex matrices.

Here, novel peptide markers of ricin were screened and identified using LC high-resolution (HR) MS based on tandem and multiple protease digestion strategies. Typical proteinases including trypsin, Glu-C, chymotrypsin, pepsin, and proteinase K were attempted to generate abundant peptides. The digested peptides from different proteinases were screened using *In-Silico* BLASTp search against non-redundant protein database. The unique peptides with good MS response were screened and identified by LC-HRMS and selected as peptide markers.

2. Results and discussion

2.1 Amino acid Sequence difference between Ricin and RCA120

To understand the sequence differences between ricin and RCA120, sequence alignment is performed using Clustal W program of BioEdit 7.0.9.0 (Figure 1). The two ricin isoforms of D and E have identical amino acid sequence in A-chain, only 15% sequence difference in B-chain. Ricin E is considered as one gene recombination product between ricin D and RCA120, because of its identical N-terminus in B-chain with that of ricin D and the identical C-terminus in B-chain with that of RCA120 [20]. Based on their identical sequence of A-chain, ricin D and E share similar toxicity. Meanwhile, because of the extremely low content of ricin E in all *R. communis* ecotypes, "ricin" is specially referred as ricin D throughout this article.

RCA120 shows a high sequence homology with ricin D, with 93% for A-chain and 84% for B-Chain (Figure 1). It has the same N-glycosidase activity of A-chain and lectin activity of B-chain with ricin, but less toxicity. Therefore, it is of great importance to discriminate ricin and RCA120 in complex matrices such as castor bean extracts. Currently, MS-based technique can be used to detect and distinguish between ricin and RCA120 through the small number of different amino acid sequences in peptide markers.




Figure 1. The amino sequence alignment of Ricin D (Uniprot ID: P02879), Ricin E (B chain GI:225419) and RCA120 (Uniprot ID: P06750) using Clustal W program of BioEdit 7.0.9.0. *Gray shading* represents 40% identical residues among the sequences. Amino acids that differ among ricin D, ricin E, and RCA120 are indicated with outlined. The amino acid similarity was also outlined at the end of each chain.

2.2. Ricin peptide markers from trypsin digestion

Peptide markers of ricin from trypsin digestion have been mainly produced by digestion after denaturation, reduction and alkylation [14,21], or direct digestion of intact ricin [13,18]. To fully understand the profile of tryptic peptide markers of ricin, efforts were made to screen them by digestion under both reduction and non-reduction. Tryptic peptides unique to ricin D (P02879) were screened by BLASTp searching tryptic peptides across all protein sequences from the updated UniprotKB/Swiss-Prot database. *In silico* digestion of ricin was carried out using the peptide mass tool (https://web.expasy.org/peptide_mass/) to obtain tryptic peptides. Tryptic peptides unique to ricin across all protein sequences were considered as candidates of its peptide markers. There are nine specific peptides identified from both A-chain and B-chain under the reduction condition (Supplementary material, Table S1); Nine specific peptides in A-chain and eight specific peptides in B-chain were identified under the non-reduction condition (Supplementary material, Table S2). Under both conditions, TA5 (LTTGADVR) and TB19 (ASDPSLK) had been used as peptide markers

for differentiating ricin from RCA120 [22,23], but they were confirmed not to be unique since they are also present in the proteins from apple snail (P83672) and winter moth (A0AL7KVU4). Taking account of this reason, the two peptides of TA5 and TB19 were not considered as candidates of ricin peptide markers.

Subsequently, tryptic digestion of purified ricin D standard was performed using our improved method of tryptic digestion under condition of denaturation and reduction, in which one denaturant resistant trypsin/lys-C mix (Promega) was applied to reduce enzymatic activity inhibition and missed cleavage. To reduce the labor procedure and elevate ionization efficiency, ACN of 20% was used as denaturation agent instead of urea or guanidine chloride. Besides, the traditional alkylation step using iodoacetamide (IAM) modification was removed because of low percentage of cysteine residues. As a result, all ricin specific peptides were detected by LC-Q/TOF-MS except for T12B, probably due to its longest length of 67 amino acid residues. The most sensitive digested peptides in order were T7A, T11A, T9A, T10A, T13A and T12A in A-chain, T20B, T6B, T14B and T18B in B-chain. These unique peptides were considered as the peptide markers suitable for unambiguous identification of ricin. The three tryptic peptides of T10A, T6B and T18B had not been selected as peptide markers of ricin [22], probably due to missed cleavage by non-denaturant resistant trypsin. The use of the trypsin/lys-C mix resistant to denaturant reduced the missed cleavage and increase the yields of these peptides. Direct tryptic digestion of purified ricin D standard was performed using our unpublished 10%ACN-assisted trypsin digestion system. It was found that similar peptide markers were obtained with the top four MS abundant peptides of T9A, T7A, T6B and T20B. The unique intra-disulfide linked peptides of both T3B-ss-T5B and T14B-ss-T16B were identified. In addition, one common inter-disulfide T24A-ss-T1B between intact ricin and RCA120 was also detected, which has been considered as an important peptide marker for the presence of intact ricin or RCA120 [18].

2.3. Novel peptide markers of ricin from tandem digestion with endoprotease Glu-C after trypsin

Followed by the above trypsin digestion, several specific peptides have been selected as peptide markers with good MS response. However, some unique peptides (eg. T8A, T12A and T10B) with MW over 3kDa are not suitable as peptide markers due to their lower MS response. Notably, we found in these larger unique peptides there were common aspartic acid (D) or glutamic acid (E) residues, which of C-terminus can be specially cleaved by endoprotease Glu-C. To confirm this, firstly, the mixed proteinase system with both trypsin and Glu-C was attempted to digest these large peptides in NH_4HCO_3 buffer. However, the result was not satisfied probably due to the low cleavage efficiency of Glu-C in NH_4HCO_3 buffer. Tandem digestion strategy was then introduced to digest the large peptides by Glu-C in phosphate buffer after trypsin digestion in NH_4HCO_3 buffer according to 3.2.3 protocol. *In silico* Glu-C digestion followed by trypsin shows more unique peptides than separate tryptic digestion (Supplementary material, Table S3). It was found that *In silico* digested peptides using Glu-C after trypsin were well consistent with the resulting peptides analyzed by LC-Q/TOF-MS by their extracted ion chromatographs (Figure 2).




Figure 2. LC-high resolution accurate mass analysis of the trypsin combined with Glu-C digest of 0.1 mg/mL of purified ricin under denaturation and reduction. Base peak chromatogram overlaid on extracted ion chromatograms of ricin peptide makers. Total ion chromatography of ricin at top right corner window.

As a result, we obtained three more novel peptide markers of TG13A, TG26A and TG28A in A-chain with tandem digestion strategy than with single trypsin digestion. There is no D and E residue sites in the amino acid sequence of T2A-glyc and T7A, the two peptides after Glu-C digestion were respectively denominated as TG2A-glyc and TG9A. The larger unique peptides such as T8A and T12A, were subsequently digested by Glu-C into shorter peptides suitable for MS analysis. The uniqueness of shorter peptides is re-confirmed by BLASTp search all other proteins. Amongst these unique peptides, TG13A from T8A, as well as TG26A and TG28A from T12A in A-chain were identified as novel peptide markers with good MS response. Because of strongly hydrophobic property, T13A from tryptic digestion has been confirmed to be poor chromatographic behavior and MS response [13]. After subsequent Glu-C digestion, T13A was cleavage into TG29A with weaker hydrophobic property than T13A based on Kyte-Doolittle hydrophobic index analysis. Correspondingly, TG29A exhibits better chromatographic behavior than T13A but the MS response is still relatively low. In addition, the TG20A from T10A has relative low MS response, probably due to the common intermediate resistance at the region of DR by proteinase cleavage [13].

Table 1 The specific ricin peptide markers under direct digestion with Glu-C after trypsin

TG#&chain	Amino acid sequence	(M+H) ⁺	(M+2H) ²⁺	(M+3H) ³⁺	(M+4H) ⁴⁺
TG2A-glyc*	QYPIINFITAGATVQSYTNFIR	3675.6946	1838.3509	1225.9031	919.6791
TG9A	VGLPINQR	896.5312	448.7692	299.5152	-
TG13A	VTNAYVVGYR	1141.6000	571.3036	381.2048	-
TG20A	YTFAFGGNYD	1154.4789	577.7431	385.4978	-
TG26A	LGNGPLE	699.3671	350.1872	233.7939	-
TG28A	AISALYYSTGGTQLPTLAR	2146.1179	1073.5626	716.0441	537.2849
TG29A	SFIICQMISE	1283.6374	642.3223	428.5506	-
TG47A-ss-TG2B ^a	CAPPPSSQF-ss-VCMD	1397.5535	699.2804	466.5226	-
TG6B-ss-TG9B ^a	NGLCVD-ss-FHNGNAIQLWPCK	2145.0003	1073.0038	715.7154	537.0555
TG11B	ANQLWTLK	973.5465	487.2769	325.1870	-
TG15B-ss-TG16B ^a	CLTTYGYSPGVYVMYD-ss-CNTA ATD	2637.1194	1319.0633	879.7113	660.0353
TG20B	SSLVLAATSGNSGTTTLTVQTNIYA VSQGWLPNTNTQPFVTTIVGLYGL CLQANSQVWIED	6284.1859	3142.5966	2095.4002	1571.8024
TG27B-ss-TG30B ^a	NCLTSD-ss-ILSCGPASSQR	1824.7458	912.9094	608.9639	456.9912
TG33B	GTILNLYSGLVLD	1377.7624	689.3848	459.9256	-
TG37B	QIILYPLHGD	1168.6360	584.8217	390.2169	-
TG38B	PNQIWLPLF	1127.6248	564.3160	376.5464	-

a: Disulfide bound peptide

The LC-MS observed ions were indicated in bold.

In B-chain, four more novel peptide markers of TG6B-ss-TG9B, TG11B, TG27B-ss-TG30B, and TG37B were obtained. In details, following Glu-C digestion the two tryptic peptides of T18B and T20B were cleaved into the three peptides of TG33B, TG37B, and TG38B. TG37B was identified as novel peptide markers based on their uniqueness to ricin and exhibit good chromatographic behavior and MS response (Figure 2). The peptide of TG11B with 7 amino acid residues from T6B was also confirmed to be unique and possess good response in MS analysis. We also identified two novel peptides of TG6B-ss-TG9B and TG27B-ss-TG30B with intramolecular disulfide bond in MS analysis. The peptide of T10B should have been transferred into a novel intramolecular disulfide bond peptide of TG15B-ss-TG16B, but it was not observed in MS analysis probably due to its low ionization efficiency. Notably, the A-B chain linker peptide TG47A-ss-TG2B from T24A-ss-T1B was identified with comparably chromatography retention and MS response with TG26A. Similar to T24A-ss-T1B, TG47A-ss-TG2B also can provides an important information for the presence intermolecular disulfide bond in spite of the consistence between ricin and RCA120.

2.4 Novel Peptide markers of ricin from chymotrypsin digestion

Recently, one LC-MS method combined with chymotrypsin digestion was developed for identification of ricin [19]. Two specific peptides of C18A (TDVQNRV) and VL-C18B (VL-AATSGNSGTTL) with one missed cleavage site were selected as peptide markers for accurate identification of ricin in plant extracts and soil samples [19]. However, the selected peptide markers are limited for unambiguous identification of ricin in complicated matrix. In addition, we found that in our digestion system, the MS response for VL-C18B was low because of missed cleavage at the first Leucine residue using the previous method of chymotrypsin digestion. To completely screen chymotryptic peptide markers, chymotrypsin digestion of ricin under both reduction and non-reduction conditions were optimized and performed followed by LC-Q-TOF/MS analysis of the digested peptides. The uniqueness of each chymotryptic peptide was verified by BLASTp searching against NCBI UniprotKB/Swiss-Prot protein database. As a result, we find that there are seven specific peptides of C2A, C4A, C5A, C18A, C27A, C31A and C33A in A-chain, and 14 specific peptides of C2B-C5B, C8B, C14B, C16B, C18B, C26B, C27B, C31B, C40B, C41B and C43B in B-chain under digestion with denaturation and reduction (Table 2). As for non-reduction conditions, besides the above specific peptides, both the A-B chain linker peptide of C49A-ss-C1B (RCAPPPSSQF-ss-ADVCM) and the intermolecular disulfide bond peptide of C3B-ss-C5B (CVDVRDGRF-ss-WPCKSNTDANQL) were different peptides from digestion with denaturation and reduction.

Table 2 The ricin specific chymotryptic peptides under digestion with denaturation and reduction

C#&chain	Amino acid sequence	(M+H) ⁺	(M+2H) ²⁺	(M+3H) ³⁺	(M+4H) ⁴⁺
C2A	TTAGATVQSY	998.4789	499.7431	333.4978	-
C4A	IRAVRGRL	940.6163	470.8118	314.2103	-
C5A	TTGADVREHPVLPNRVGLP INQRF	2799.5376	1400.2724	933.8507	700.6418
C18A	TDVQNRV	895.4268	448.2170	299.1471	-
C27A	EEAISAL	732.3774	366.6923	265.1449	-
C31A	STGGTQLPTL	974.5153	487.7613	325.5099	-
C33A	IICIQM	720.3783	360.6928	240.7976	-
C2B	DPEPIVRIVGRNGL	1534.8699	767.9386	512.2948	384.4734
C3B	CVDVRDGRF	1066.5098	533.7585	356.1748	-
C4B	HNGNAIQL	866.4479	433.7276	289.4875	-
C5B	WPCKSNTDANQL	1376.6263	688.8168	459.5469	344.9125
C8B	KRDNTIRSNGKCL	1504.8012	752.9043	502.2719	376.9562
C14B	DCNTAATDATRW	1324.5586	643.8386	442.1911	331.8956
C16B	DNGTIINPRSSL	1286.6699	643.8386	429.5615	322.4234
C18B	AATSGNSGTTL	979.4691	490.2382	327.1612	-
C26B	QANSGQVW	889.4162	445.2118	297.1436	-
C27B	IEDCSSEKAEQQW	1552.6584	776.8328	518.2243	388.9205
C31B	TSDSNIRETVVKIL	1574.8748	787.9410	525.6298	394.4746
C40B	DVRASDPSL	959.4792	480.2432	320.4979	-
C41B	KQIIL	614.4235	307.7154	205.4794	-
C43B	HGDPNQIW	966.4428	483.7250	322.8191	-

The LC-MS observed ions were indicated in bold.

Under the condition of denaturation and reduction, peptide mapping analyses were performed using our optimized digestion conditions with 20% ACN as denaturation agent to cleave purified ricin. In A-chain, all 7 specific chymotryptic fragments were verified using LC-MS analysis as shown in Figure 3 and listed in Table 2. The three peptides of C18A, C4A and C31A with 7-10 amino acid residues were selected as novel peptide markers based on good MS response and suitable MW. In the B-chain, 11 out of 14 specific chymotryptic peptides were identified in MS analysis (Figure 3) and listed in Table 2. The chymotryptic peptides with the best MS response were C8B, C27B, C31B, C18B, and C4B in order. They were considered as the peptide markers suitable for unambiguous identification of ricin. Notably, we observed no missing cleavage peptides of C18B AATSGNSGTTL, suggesting that the better accuracy of digestion was achieved than that of previous studies. Both C40B and C43B peptides were not identified in MS probably because they were located in the polar region in B-chain by Kyte-Doolittle hydrophobic index analysis. This result is well consistent with previous findings that polar regions may be more resistant to digestion than other regions of protein [13].




Figure 3. LC-high resolution accurate mass analysis of the chymotrypsin digest of 0.1 mg/mL of purified ricin under denaturation and reduction. Base peak chromatogram overlaid on extracted ion chromatograms of ricin maker peptides. Total ion chromatography of ricin at top right corner window.

Under the condition of ACN-assisted direct digestion, similar number of chymotryptic fragments were selected as peptide markers including three peptide markers of C18A, C4A and C31A in A chain, as well as four peptide markers of C18B, C8B, C41B and C31B in B chain (Figure 4). An exception for this condition was that, the unique intra-disulfide linked peptide C3B-ss-C5B was identified in the extraction ion chromatography of the triply charged molecular ion at m/z 814.0431. The common inter-chain disulfide peptide C49A-ss-C1B between ricin and RCA120 was also detected in extracted ion chromatograms (Figure 4). Although both the inter-chain disulfide peptide C49A-ss-C1B and intra-disulfide linked peptide C3B-ss-C5B were identified, the C2B situated between them was not observed probably due to intrinsically low MS response.




Figure 4. LC-high resolution accurate mass analysis of the direct chymotrypsin digestion of 0.1 mg/mL of ricin. Base peak chromatogram overlaid on extracted ion chromatograms of ricin maker peptides. Total ion chromatography of ricin at top right corner window.

2.5. Novel peptide markers of ricin from other proteinase digestion

Peptide markers produced from other proteinase digestion are essential for verification of trace ricin in complex matrices. Here, two other proteinases of pepsin and proteinase K were also attempted to digest ricin into potential peptide markers.

2.5.1 Novel peptide markers of ricin from pepsin digestion

Pepsin is a protease with high digestion efficiency, which preferentially cleaves phenylalanine (F), leucine (L), and tyrosine (Y) at the C-terminus [24]. According to the product instruction, the

specificity for cleavage at F and L is best at pH 1.3. In case the pH is above 2.0, the specificity for F and L cleavage decreases heavily with the increased specificity for Y. When the pH is above 5.0, the pepsin will lose its activity. Pepsin can be used alone or in combination with other proteases for protein analysis by mass spectrometry or other applications[24]. Here, ricin was separately digested with pepsin under the two conditions of pH 1~2 (0.67% FA) and pH 2~4 (0.1% FA) and then evaluated. Following theoretical enzyme digestion and specificity assessment by NCBI BLASTp search, the specific pepsin digest peptides were obtained in pH 1~2 (Table S4) and pH 2~4 (Table S5). During the digestion of ricin at pH 1~2, 10 peptides from A chain and 9 peptides from B chain were unique to ricin. At pH 2~4, there are only 7 unique peptides and all of them are from B chain. This phenomenon should be resulted from the reduced cleavage selectivity of pepsin at pH value above 2. Under the pH value, ricin is preferred to being cleaved into short peptides because of pepsin's more cleavage sites of F, L, Y, E, Q, and A.

Under the optimized conditions at pH 1~2 according to the procedure of 3.2.6, 7 out of 10 ricin A-chain specific peptides were verified by LC-Q-TOF/MS (Figure 5). Three peptides, P24A, P13A, and P7A exhibits the best MS abundance, and they can be considered as novel peptide markers. The A-B chain disulfide peptide P36A-ss-P1B was not observed, probably because of its large molecular weight and disulfide bond. In the B-chain, all ricin specific peptides were identified except P7B that is a long peptide with 42 amino acid residues. We selected the three peptides of P26B, P23B and P2B as novel peptide markers based on their highest MS intensity among specific peptides. At pH 2~4, all 7 ricin unique peptides were identified including disulfide peptide P2B-ss-P4B as showed in Figure 6 and listed in Supplementary material, Table S5. These identified unique peptides form B-chain at pH 2~4 will not be enough for unambiguous identification of ricin with both chains. As a consequence, it is recommended to digest ricin by pepsin under pH 1~2 for its unique identification and quantification.




Figure 5. LC-high resolution accurate mass analysis of the pepsin digestion production of 0.1 mg/mL of purified ricin (pH 1~2). Base peak chromatogram overlaid on extracted ion chromatograms of ricin maker peptides. Total ion chromatography of ricin at top right corner window.




Figure 6. LC-high resolution accurate mass analysis of the pepsin digest of 0.1 mg/mL of purified ricin (pH 2~4). Base peak chromatogram overlaid on extracted ion chromatograms of ricin maker peptides. Total ion chromatography of ricin at top right corner window.

2.5.2 Novel peptide markers of ricin from Proteinase K digestion

We also used Proteinase K isolated from *Tritirachium album* to digest ricin into novel peptide markers. Under the optimized conditions in 3.2.7, one A-Chain and five B-Chain peptides were confirmed as ricin-specific peptides by *in silico* digestion and BLASTp search (Table 3). All these peptides were identified by MS with good intensity (Supplementary material, Figure S1). In spite of the limited number of peptide markers, they may have good sensitivity and cost short digestion time. Thus, these peptide markers can be used as a supplement for unambiguous identification of ricin.

Table 3 The theoretical ricin peptides under proteinase K direct digestion

K#&chain	Amino acid sequence	(M+H) ⁺	(M+2H) ²⁺	(M+3H) ³⁺	(M+4H) ⁴⁺
K71A	STGGTQL	663.3308	332.1690	221.7818	-
K15B	PCKSNTDA	835.3614	418.1843	279.1253	-
K41B	TSGNSGTTL	837.3948	419.2011	279.8031	-
K61B	EDCSSEKA	868.3353	434.6713	290.1166	-
K92B	HGDPNQI	780.3635	390.6854	260.7927	-

The LC-MS observed ions were indicated in bold.

3. Materials and Methods

3.1 Sequence alignment and identification of uniqueness of digested peptides

Amino acid sequence of ricin was downloaded from NCBI UniProt database (P02879). *In silico* digestion of ricin into particular peptides by various proteinases were generated using ExPASy peptide mass tool (https://web.expasy.org/peptide_mass/). The *in silico* enzymatic peptides were selected under the following criteria: no missed cleavages, more than 5 amino acid numbers. Uniqueness of digested peptides were identified by BLASTp searching them against the UniProtKB/SwissProt protein sequence database (~470034 protein sequences).

3.2 Digestion of Ricin using multiple proteinases

3.2.1 Trypsin digestion of ricin with denaturation and reduction

Fifty microliter of 0.2 mg/mL purified ricin was transferred into a 200 μ L PCR tube, containing 42 μ L of the denaturing and reducing solution composed of 20 μ L aliquot of acetonitrile, 1 μ L of 0.5 M dithiothreitol (DTT), 5 μ L of 1.0 M ammonium bicarbonate (pH 8.5) and 16 μ L of H₂O. The mixture was incubated at 60°C for 50 min and then cooled at room temperature for 10 min. 4 μ g of trypsin/lys-C mix (Promega, Mass Spec Grade) was added and then the reaction solution was kept

at 40°C for 4 h. The reaction was terminated by adding the appropriate volume of 10% formic acid (FA) to achieve 0.5% final concentration and transferred to vials for LC-MS analysis.

3.2.2 Solvent-assisted direct digestion of ricin

Digestion was performed according to our unpublished acetonitrile-assisted trypsin digestion method. Briefly, 50 µL of 0.2 mg/mL of ricin were transferred to a 200 µL PCR Tube, which contains 10 µL of acetonitrile and 27 µL of H₂O. Then, 5 µL of 1.0 M ammonium bicarbonate (pH 8.8) and 4 µg of trypsin/lys-C mix (Promega, Mass Spec Grade) were added and incubated at 40°C for 4 h. After digestion, the reaction was terminated by adding appropriate volume of 10% FA to 0.5% final concentration. The resulting samples were transferred to vials for LC-MS analysis.

3.2.3 Tandem digestion with Glu-C after trypsin

After solvent-assisted trypsin digestion of ricin according to the procedure of section 3.2.2, the digestion mixture was concentrated to dryness using vacuum concentrator. Then, the sample was reconstituted with 80 µL 20mM PBS (15 mM NaH₂PO₄ and 5 mM Na₂HPO₄). A total of 8 µL of 0.25 µg/µL aqueous Glu-C (Promega, Mass Spec Grade) solution was added, and the digestion was performed at 45 °C for 4 h. Five µL of 10% FA was added to terminate the digestion and then transferred to vials for LC-MS analysis.

3.2.4 Chymotrypsin digestion with denaturation and reduction

One 50 µL of 0.2 mg/mL of ricin were added to a 200 µL PCR tube, containing the denaturing and reducing solution composed of 20 µL aliquot of acetonitrile, 1 µL of 0.5 M dithiothreitol (DTT), and 10 µL of 1.0 M tris-HCl (pH 8.0), and 11 µL of H₂O. The mixture was then incubated at 60°C for 50 min. Following the sample being cooled to room temperature, 8 µL of 0.25 µg/µL chymotrypsin (Promega, Sequence Grade) was added and incubated at 40°C for 4 h. After digestion, samples were acidified by adding 8 µL of 5 % FA and transferred to vials for LC-MS analysis.

3.2.5 Solvent-assisted chymotrypsin digestion of ricin

Firstly, 50 µL of purified ricin sample was transferred in a 200 µL PCR tube, which contains 10 µL of 1.0 M Tris-HCl (pH 8.0) and 10 µL acetonitrile. Subsequently, aliquot of 8 µL of 0.25 µg/µL chymotrypsin (Promega, Sequencing Grade) was added and then the mixture was incubated for 4 h at 40°C. The digestion was terminated by adding 8 µL of 5 % FA. The resulting sample was transferred to an autosampler vial for LC-MS analysis.

3.2.6 Pepsin digestion

Briefly, 12 µL of 2 mg/mL pepsin (Sigma-Aldrich) was added to 50 µL of 0.2 mg/mL of ricin in H₂O. As for the digestion reaction under pH 1~2, 7 µL of 10% FA was added to the sample. As for that under pH 2~4, 1 µL of 10% FA was added. At last, the mixture was diluted to 100 µL with H₂O. The mixture was incubated at 60°C for 4 h. After digestion, the mixture reaction was terminated by adding appropriate volume of 2% ammonium hydroxide and transferred to vials for LC-MS analysis.

3.2.7 Proteinase K digestion

Firstly, 20 µL of 20 mg/mL proteinase K (Merck Millipore) in 20 mM Tris-HCl (pH 7.8) were added to 50 µL of 0.2 mg/mL of ricin in H₂O. Then, 2 µL of 1.0 M Tris-HCl and 28 µL of H₂O were added. The sample was mixed and incubated for 2 h at 60°C. After digestion, 10% FA was added in the sample to terminate the reaction and the reaction solution was transferred to vials for LC-MS analysis.

3.3 Accurate Mass LC-MS Qualitative Analysis

An Agilent 1200 HPLC system combined with a 6520 high-resolution accurate Q-TOF mass spectrometer (Agilent technology, MA, USA) was used for sample analysis. The digests peptides were verified by extract the accurate molecular weight of the *in silico* digested peptides. Sample aliquots were separated on an Advanced Peptide Map 2.7- μ m, 2.1 \times 150 mm column (Agilent technology, MA, USA). Mobile phase A consisted of 0.1% formic acid in water. Mobile phase B consisted of 0.1% formic acid in acetonitrile. The flow rate for analysis was 250 μ L/min and the injection volume was 5 μ L. The optimized gradient elution program is as follows: 0-3 min 5% B; 3-45 min linear gradient to 70%B; 45-53 min 70%B; 53-54 min linear gradient to 100%B; 54-60 min, 100%B. The column was re-equilibrated with 5 % B for 4 min before the next run. Eluted peptides were injected to the mass spectrometer equipped with electrospray ionization source in positive mode MS scan was acquired over the mass range 150–3000 at 1.1 spectra/s; gas temperature=330 °C, gas flow =8 L/min, nebulizer=30 psi, and capillary voltage=3500 V; Fragmentor is 150 V and skimmer is 65 V. Mass Hunter workstation B.05.00 software was used (Agilent Scientific, USA) for instrument control and to process the data files.

Supplementary Materials: The following are available online at www.mdpi.com/xxx/s1, Figure S1: LC-high resolution accurate mass analysis of the proteinase K digest of 0.2 mg/mL of purified ricin. Base peak chromatogram overlaid on extracted ion chromatograms of ricin maker peptides. Total ion chromatography of ricin at top right corner window, Table S1: The specific ricin tryptic peptides under digestion with denaturation and reduction, Table S2: The specific ricin tryptic peptides under direct digestion, Table S3: Theoretical mass to charge values of ricin digested peptides from Glu-C after trypsin, Table S4: The ricin specific peptides under pepsin direct digestion at pH 1~2, Table S5: The theoretical ricin peptides under pepsin digestion at pH 2~4,

Acknowledgments:

Author Contributions:

Conflicts of Interest:

References

1. Fabbri, M.S.; Katayama, M.; Nakase, I.; Vago, R. Plant Ribosome-Inactivating Proteins: Progresses, Challenges and Biotechnological Applications (and a Few Digressions). *Toxins* **2017**, *9*, doi:10.3390/toxins9100314.
2. Van Damme, E.J.M.; Hao, Q.; Chen, Y.; Barre, A.; Vandenbussche, F.; Desmyter, S.; Rougé, P.; Peumans, W.J. Ribosome-Inactivating Proteins: A Family of Plant Proteins That Do More Than Inactivate Ribosomes. *Critical Reviews in Plant Sciences* **2010**, *20*, 395-465, doi:10.1080/07352689.2001.10131826.
3. Jennifer Audi, M.B., Manish Patel, Joshua Schier, John Osterloh. Ricin Poisoning: A Comprehensive Review. *JAMA* **2005**, *294*, 2342-2451, doi:10.1001/jama.294.18.2342.
4. Worbs, S.; Kohler, K.; Pauly, D.; Avondet, M.A.; Schaer, M.; Dorner, M.B.; Dorner, B.G. Ricinus communis intoxications in human and veterinary medicine—a summary of real cases. *Toxins* **2011**, *3*, 1332-1372, doi:10.3390/toxins3101332.
5. Stirpe, F.; Battelli, M.G. Ribosome-inactivating proteins: progress and problems. *Cellular and molecular life sciences : CMLS* **2006**, *63*, 1850-1866, doi:10.1007/s00018-006-6078-7.
6. Wang, D.; Baudys, J.; Barr, J.R.; Kalb, S.R. Improved Sensitivity for the Qualitative and Quantitative Analysis of Active Ricin by MALDI-TOF Mass Spectrometry. *Analytical chemistry* **2016**, *88*, 6867-6872, doi:10.1021/acs.analchem.6b01486.
7. Tang, J.J.; Sun, J.F.; Lui, R.; Zhang, Z.M.; Liu, J.F.; Xie, J.W. New Surface-Enhanced Raman Sensing Chip Designed for On-Site Detection of Active Ricin in Complex Matrices Based on Specific Depurination. *ACS applied materials & interfaces* **2016**, *8*, 2449-2455, doi:10.1021/acsami.5b12860.
8. Falach, R.; Sapozhnikov, A.; Gal, Y.; Israeli, O.; Leitner, M.; Seliger, N.; Ehrlich, S.; Kronman, C.; Sabo, T. Quantitative profiling of the *in vivo* enzymatic activity of ricin reveals disparate depurination of different pulmonary cell types. *Toxicology letters* **2016**, *258*, 11-19, doi:10.1016/j.toxlet.2016.06.003.
9. Simon, S.; Worbs, S.; Avondet, M.A.; Tracz, D.M.; Dano, J.; Schmidt, L.; Volland, H.; Dorner, B.G.; Corbett, C.R. Recommended Immunological Assays to Screen for Ricin-Containing Samples. *Toxins* **2015**, *7*, 4967-4986, doi:10.3390/toxins7124858.

10. Gaylord, S.T.; Dinh, T.L.; Goldman, E.R.; Anderson, G.P.; Ngan, K.C.; Walt, D.R. Ultrasensitive Detection of Ricin Toxin in Multiple Sample Matrixes Using Single-Domain Antibodies. *Analytical chemistry* **2015**, *87*, 6570-6577, doi:10.1021/acs.analchem.5b00322.
11. Worbs, S.; Skiba, M.; Soderstrom, M.; Rapinoja, M.L.; Zeleny, R.; Russmann, H.; Schimmel, H.; Vanninen, P.; Fredriksson, S.A.; Dorner, B.G. Characterization of Ricin and *R. communis* Agglutinin Reference Materials. *Toxins* **2015**, *7*, 4906-4934, doi:10.3390/toxins7124856.
12. Sylvia, W.; Kernt, K.; Diana, P.; Marc-André, A.; Martin, S.; Dorner, M.B.; Dorner, B.G. Ricinus communis Intoxications in Human and Veterinary Medicine-A Summary of Real Cases. *Toxins* **2011**, *3*, 1332-1372.
13. Ostin, A.; Bergstrom, T.; Fredriksson, S.-A.; Nilsson, C. Solvent-Assisted Trypsin Digestion of Ricin for Forensic Identification by LC-ESI MS/MS. *Analytical chemistry* **2007**.
14. McGrath, S.C.; Schieltz, D.M.; McWilliams, L.G.; Pirkle, J.L.; Barr, J.R. Detection and quantification of ricin in beverages using isotope dilution tandem mass spectrometry. *Analytical chemistry* **2011**, *83*, 2897-2905, doi:10.1021/ac102571f.
15. Ma, X.; Tang, J.; Li, C.; Liu, Q.; Chen, J.; Li, H.; Guo, L.; Xie, J. Identification and quantification of ricin in biomedical samples by magnetic immunocapture enrichment and liquid chromatography electrospray ionization tandem mass spectrometry. *Analytical and bioanalytical chemistry* **2014**, *406*, 5147-5155, doi:10.1007/s00216-014-7710-2.
16. Schieltz, D.M.; McWilliams, L.G.; Kuklenyik, Z.; Prezioso, S.M.; Carter, A.J.; Williamson, Y.M.; McGrath, S.C.; Morse, S.A.; Barr, J.R. Quantification of ricin, RCA and comparison of enzymatic activity in 18 *Ricinus communis* cultivars by isotope dilution mass spectrometry. *Toxicon : official journal of the International Society on Toxinology* **2015**, *95*, 72-83, doi:10.1016/j.toxicon.2015.01.003.
17. Worbs, S.; Skiba, M.; Bender, J.; Zeleny, R.; Schimmel, H.; Luginbühl, W.; Dorner, B. An International Proficiency Test to Detect, Identify and Quantify Ricin in Complex Matrices. *Toxins* **2015**, *7*, 4987-5010, doi:10.3390/toxins7124859.
18. Fredriksson, S.A.; Artursson, E.; Bergstrom, T.; Ostin, A.; Nilsson, C.; Astot, C. Identification of RIP-II toxins by affinity enrichment, enzymatic digestion and LC-MS. *Analytical chemistry* **2015**, *87*, 967-974, doi:10.1021/ac5032918.
19. Braun, A.V.; Taranchenko, V.F.; Tikhomirov, L.A.; Grechukhin, A.P.; Rybal'chenko, I.V. Detection of Ricin in Plant Extracts and Soil Using Liquid Chromatography-High-Resolution Mass Spectrometry. *Journal of Analytical Chemistry* **2018**, *73*, 786-795, doi:10.1134/s1061934818080026.
20. Araki, T.; Funatsu, G. The complete amino acid sequence of the B-chain of ricin E isolated from small-grain castor bean seeds. Ricin E is a gene recombination product of ricin D and *Ricinus communis* agglutinin. *Biochimica et biophysica acta* **1987**, *911*, 191-200.
21. Sten-Ajke Fredriksson, A.G.H., Elisabeth Artursson, Ad. L. de Jong, Calle Nilsson, and Ben L. M. van Baar. Forensic Identification of Neat Ricin and of Ricin from Crude Castor Bean Extracts by Mass Spectrometry. *Anal. Chem.* **2005**, *77*, 1545-1555.
22. Fredriksson, S.-Å.; Hulst, A.G.; Artursson, E.; Jong, A.L.d.; Nilsson, C.; Baar, B.L.M.v.; Chem, A. Forensic identification of neat ricin and of ricin from crude castor bean extracts by mass spectrometry. *Anal. Chem.* **2005**, *77*, 1545.
23. Ostin, A.; Bergstrom, T.; Fredriksson, S.A.; Nilsson, C. Solvent-assisted trypsin digestion of ricin for forensic identification by LC-ESI MS/MS. *Anal. Chem.* **2007**, *79*, 6271-6278, doi:10.1021/ac070174.
24. Sweeney, P.J.; Walker, J.M. Proteolytic Enzymes for Peptide Production. *Methods in Molecular Biology* **1993**, *16*, 277-303.