

Article

Four-Dimensional Almost Einstein Manifolds with Skew-Circulant Structures

Iva Dokuzova ^{1,‡,*}  and Dimitar Razpopov ² 

¹ Department of Algebra and Geometry, Faculty of Mathematics and Informatics, University of Plovdiv Paisii Hilendarski, Plovdiv, Bulgaria, dokuzova@uni-plovdiv.bg

² Department of Mathematics and Informatics, Faculty of Economics, Agricultural University of Plovdiv, Bulgaria, razpopov@au-plovdiv.bg

* Correspondence: dokuzova@uni-plovdiv.bg; Tel.: 359887977638

‡ These authors contributed equally to this work.

Abstract: We consider a four-dimensional Riemannian manifold M equipped with an additional tensor structure S , whose fourth power is minus identity and the second power is an almost complex structure. In a local coordinate system the components of the metric g and the structure S form skew-circulant matrices. Both structures S and g are compatible, such that an isometry is induced in every tangent space of M . By a special identity for the curvature tensor, generated by the Riemannian connection of g , we determine classes of an Einstein manifolds and an almost Einstein manifolds. For such manifolds we obtain propositions for the sectional curvatures of some special 2-planes in a tangent space of M . We consider an almost Hermitian manifold associated with the studied manifold and find conditions for g , under which it is a Kähler manifold. We construct some examples of the considered manifolds on Lie groups.

Keywords: Riemannian manifold; Einstein manifold; sectional curvatures; Ricci curvature; Lie group

MSC: 53B20; 53C15; 53C25; 53C55; 22E60

1. Introduction

The right circulant matrices and the right skew-circulant matrices are Toeplitz matrices, which are thoroughly studied in [1] and [3]. The set of invertible circulant (skew-circulant) matrices form a group with respect to the matrix multiplication. Such matrices have application to geometry, linear codes, graph theory, vibration analysis (for example [2,7,9,11–13]).

A. Gray, L. Hervella and L. Vanhecke used curvature identities to classify and to study the almost Hermitian manifolds (for instance in [4–6,15]). The Hermitian manifolds form a class of manifolds with an integrable complex structure J . The class of the Kähler manifolds is their subclass and such manifolds have a parallel structure J . According to A. Gray, the Kähler manifolds have an especially rich geometric structure, due to the Kähler curvature identity $R(\cdot, \cdot, J\cdot, J\cdot) = R(\cdot, \cdot, \cdot, \cdot)$. Some of the recent investigations on the curvature properties of the almost Hermitian manifolds are made in [8,10,14,16].

In the present work we study a four-dimensional differentiable manifold M with a Riemannian metric g . The manifold M is equipped with an additional tensor structure S of type $(1,1)$, which satisfies $S^4 = -\text{id}$. Moreover, the component matrix of S is a special skew-circulant matrix. The structure S is compatible with g , such that an isometry is induced in every tangent space of M . Such

29 a manifold (M, g, S) is associated with an almost Hermitian manifold (M, g, J) , where $J = S^2$ is an
30 almost complex structure.

31 The paper is organized as follows. In Sect. 2, we introduce the manifold (M, g, S) . In Sect. 3, we
32 find conditions under which an orthogonal basis of the type $\{x, Sx, S^2x, S^3x\}$ exists in every tangent
33 space of (M, g, S) . In Sect. 4, we consider a class of almost Einstein manifolds (M, g, S) . Also, we
34 obtain conditions for (M, g, S) to be an Einstein manifold. In Sect. 5, we find some curvature properties
35 of these manifolds. In Sect. 6, we obtain a necessary and sufficient condition for S to be parallel with
36 respect to the Riemannian connection of g . Also, we get conditions for (M, g, J) to be a Kähler manifold.
37 In Sect. 7, we construct examples of the considered manifolds on Lie groups and find some of their
38 geometric characteristics.

39 **2. Preliminaries**

Let M be a 4-dimensional Riemannian manifold equipped with a tensor structure S in every tangent space $T_p M$ at a point p on M . Let S have a skew-circulant matrix, with respect to some basis $\{e_i\}$, as follows

$$(S_j^k) = \begin{pmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ -1 & 0 & 0 & 0 \end{pmatrix}. \quad (1)$$

Then S has the property

$$S^4 = -\text{id}. \quad (2)$$

Let the metric g and the structure S satisfy

$$g(Sx, Sy) = g(x, y). \quad (3)$$

40 Here and anywhere in this work, x, y, z, u will stand for arbitrary elements of the algebra on smooth
41 vector fields on M or vectors in $T_p M$. The Einstein summation convention is used, the range of the
42 summation indices being always $\{1, 2, 3, 4\}$.

The conditions (1) and (3) imply that the matrix of g has the form

$$(g_{ij}) = \begin{pmatrix} A & B & 0 & -B \\ B & A & B & 0 \\ 0 & B & A & B \\ -B & 0 & B & A \end{pmatrix}, \quad (4)$$

i.e. it is skew-circulant. Here $A = A(p)$ and $B = B(p)$ are smooth functions of an arbitrary point $p(X^1, X^2, X^3, X^4)$ on M . The determinant of g has a value $\det(g_{ij}) = (A^2 - 2B^2)^2$. It is supposed that

$$A(p) > \sqrt{2}B(p) > 0 \quad (5)$$

43 in order g to be positive definite. A manifold M introduced in this way we denote by (M, g, S) .

Now, we consider an associated metric \tilde{g} with g , determined by

$$\tilde{g}(x, y) = g(x, Sy) + g(Sx, y). \quad (6)$$

Using (1), (4) and (6) we get that the matrix of its components is

$$(\tilde{g}_{ij}) = \begin{pmatrix} 2B & A & 0 & -A \\ A & 2B & A & 0 \\ 0 & A & 2B & A \\ -A & 0 & A & 2B \end{pmatrix}. \quad (7)$$

44 Since (5) is valid, it is easy to see that \tilde{g} is an indefinite metric.

The inverse matrices of (g_{ij}) and (\tilde{g}_{ij}) are as follows:

$$(g^{ij}) = \frac{1}{D} \begin{pmatrix} A & -B & 0 & B \\ -B & A & -B & 0 \\ 0 & -B & A & -B \\ B & 0 & -B & A \end{pmatrix}, \quad (8)$$

$$(\tilde{g}^{ij}) = \frac{1}{2D} \begin{pmatrix} -2B & A & 0 & -A \\ A & -2B & A & 0 \\ 0 & A & -2B & A \\ -A & 0 & A & -2B \end{pmatrix}, \quad (9)$$

45 where $D = A^2 - 2B^2$.

46 **3. Orthogonal S -basis of $T_p M$**

47 If x is a nonzero vector on (M, g, S) , then according to (1) we have $Sx \neq \pm x$. Therefore the angle
48 φ between x and Sx belongs to the interval $(0, \pi)$. Evidently, the vectors x, Sx, S^2x and S^3x determine
49 six angles, which belong to $(0, \pi)$. For these angles we establish the next statement.

Theorem 1. Let x be a nonzero vector on (M, g, S) . Then

$$\angle(x, Sx) = \angle(Sx, S^2x) = \angle(S^2x, S^3x) = \varphi, \quad \angle(x, S^3x) = \pi - \varphi, \quad \angle(x, S^2x) = \angle(Sx, S^3x) = \frac{\pi}{2}, \quad (10)$$

50 where $\varphi \in (0, \pi)$.

Proof. Let $x = (x^1, x^2, x^3, x^4)$ be a nonzero vector on (M, g, S) . By using (1), we get

$$Sx = (x^2, x^3, x^4, -x^1), \quad S^2x = (x^3, x^4, -x^1, -x^2), \quad S^3x = (x^4, -x^1, -x^2, -x^3). \quad (11)$$

From (2) and (3) it follows

$$g(x, Sx) = -g(x, S^3x), \quad g(x, S^2x) = 0. \quad (12)$$

Having in mind (4) and (11), we calculate

$$\begin{aligned} g(x, x) &= A((x^1)^2 + (x^2)^2 + (x^3)^2 + (x^4)^2) + 2B(x^1x^2 + x^2x^3 + x^3x^4 - x^1x^4), \\ g(x, Sx) &= A(x^1x^2 + x^2x^3 + x^3x^4 - x^1x^4) + B((x^1)^2 + (x^2)^2 + (x^3)^2 + (x^4)^2). \end{aligned} \quad (13)$$

Now, due to (3) and (5), we can determine the angle between x and Sx and the angle between x and S^2x as follows:

$$\cos \varphi = \frac{g(x, Sx)}{g(x, x)}, \quad \cos \phi = \frac{g(x, S^2x)}{g(x, x)}. \quad (14)$$

We apply (12) and (13) in (14) and find

$$\cos \varphi = \frac{A(x^1x^2 + x^2x^3 + x^3x^4 - x^1x^4) + B((x^1)^2 + (x^2)^2 + (x^3)^2 + (x^4)^2)}{A((x^1)^2 + (x^2)^2 + (x^3)^2 + (x^4)^2) + 2B(x^1x^2 + x^2x^3 + x^3x^4 - x^1x^4)},$$

$$\cos \phi = 0.$$

51 Then, bearing in mind (3) and (12), we get (10). \square

⁵² **Definition 1.** A basis of type $\{x, Sx, S^2x, S^3x\}$ of $T_p M$ is called a *S-basis*. In this case we say that the vector x
⁵³ induces a *S-basis* of $T_p M$.

⁵⁴ The following statements hold.

Theorem 2. Every nonzero vector $x = (x^1, x^2, x^3, x^4)$, which satisfies

$$4x^2x^4((x^1)^2 - (x^3)^2) + 4x^1x^3((x^4)^2 - (x^2)^2) + ((x^1)^2 + (x^3)^2)^2 + ((x^2)^2 + (x^4)^2)^2 \neq 0, \quad (15)$$

⁵⁵ induces a *S-basis* of $T_p M$.

Proof. If a nonzero vector $x \in T_p M$ has coordinates (x^1, x^2, x^3, x^4) , then using (11) we get the determinant formed by the coordinates of the vectors x, Sx, S^2x and S^3x . It is

$$\Delta = 4x^2x^4((x^1)^2 - (x^3)^2) + 4x^1x^3((x^4)^2 - (x^2)^2) + ((x^1)^2 + (x^3)^2)^2 + ((x^2)^2 + (x^4)^2)^2.$$

⁵⁶ In case that (15) is valid, we have $\Delta \neq 0$, i.e. x, Sx, S^2x and S^3x form a basis. \square

Lemma 1. Let a vector x induce a *S-basis* and let φ be the angle between x and Sx . The following inequalities are valid:

$$\frac{\pi}{4} < \varphi < \frac{3\pi}{4}. \quad (16)$$

Proof. We suppose without loss of generality that $g(x, x) = 1$. Then, from (3), (12) and (14), we find

$$g(x, Sx) = g(Sx, S^2x) = g(S^2x, S^3x) = -g(x, S^3x) = \cos \varphi, \quad g(x, S^2x) = g(Sx, S^3x) = 0. \quad (17)$$

We consider a nonzero vector y , such that

$$y = -\cos \varphi x + Sx - \cos \varphi S^2x. \quad (18)$$

Since g is a Riemannian metric we have $g(y, y) > 0$. Substituting (18) into the latter inequality, and using (17), we get

$$1 - 2\cos^2 \varphi > 0.$$

⁵⁷ Then, taking into account $0 < \varphi < \pi$, we obtain (16). \square

⁵⁸ Bearing in mind Theorem 1, Theorem 2 and Lemma 1, we arrive at the following

⁵⁹ **Theorem 3.** For every manifold (M, g, S) an orthogonal *S-basis* of $T_p M$ exists.

⁶⁰ **4. Almost Einstein manifolds**

Let ∇ be the Riemannian connection of g . The curvature tensor R of ∇ is determined by

$$R(x, y)z = \nabla_x \nabla_y z - \nabla_y \nabla_x z - \nabla_{[x, y]} z. \quad (19)$$

The tensor of type $(0, 4)$ associated with R is defined by

$$R(x, y, z, u) = g(R(x, y)z, u). \quad (20)$$

The Ricci tensor ρ with respect to g is given by the well-known formula

$$\rho(y, z) = g^{ij}R(e_i, y, z, e_j). \quad (21)$$

The scalar curvature τ with respect to g and its associated quantity are determined by

$$\tau = g^{ij}\rho(e_i, e_j), \quad \tau^* = \bar{g}^{ij}\rho(e_i, e_j). \quad (22)$$

Now, we consider a manifold (M, g, S) with the condition

$$\nabla S = 0. \quad (23)$$

61 i.e., S is a parallel structure with respect to ∇ .

Proposition 1. *Every manifold (M, g, S) with a parallel structure S satisfies the curvature identity*

$$R(x, y, Sz, Su) = R(x, y, z, u). \quad (24)$$

Proof. The well-known formula $(\nabla_x S)y = \nabla_x S y - S \nabla_x y$, together with (23), yields

$$\nabla_x S y = S \nabla_x y. \quad (25)$$

On the other hand (19) implies $R(x, y, Sz, Su) = g(R(x, y)Sz, Su)$. Because of the latter identity, using (3), (19) and (25), we have successively

$$\begin{aligned} R(x, y, Sz, Su) &= g(\nabla_x \nabla_y Sz - \nabla_y \nabla_x Sz - \nabla_{[x,y]} Sz, Su) \\ &= g(\nabla_x S(\nabla_y z) - \nabla_y S(\nabla_x z) - S(\nabla_{[x,y]} z), Su) \\ &= g(S(\nabla_x \nabla_y z - \nabla_y \nabla_x z - \nabla_{[x,y]} z), Su) = g(R(x, y)z, u), \end{aligned}$$

62 which completes the proof. \square

63 We will note that the identity (24) defines a more general class of manifolds (M, g, S) than the
64 class with the condition (23). Farther in this paper, we investigate the properties of manifolds in these
65 two classes.

Let R_{ijkl} be the components of the curvature tensor R of type $(0, 4)$. The local form of (24) is
 $R_{ijlm}S_k^l S_h^m = R_{ijkl}$. Then, using (1), we find the equalities

$$\begin{aligned} R_{1313} &= R_{2424} = R_{1324}, \\ R_{1212} &= R_{1414} = R_{2323} = R_{3434} = R_{1223} = R_{1214} = R_{1434} = R_{1234} = R_{2334} = R_{2314}, \\ R_{1213} &= R_{1224} = R_{1413} = R_{2414} = R_{2423} = R_{2313} = R_{1334} = R_{2434}. \end{aligned}$$

By applying the Bianchi identity to the above components of R , we obtain

$$\begin{aligned} R_{1313} = R_{2424} = R_{1324} &= 2R_{1212} = 2R_{1414} = 2R_{2323} = 2R_{3434} = 2R_{1223} = 2R_{1214} \\ &= 2R_{1434} = 2R_{1234} = 2R_{2334} = 2R_{2314}, \\ R_{1213} = R_{1224} = R_{1413} &= R_{2414} = R_{2423} = R_{2313} = R_{1334} = R_{2434}. \end{aligned} \quad (26)$$

66 Vice versa, from (1) and (26) it follows (24).

67 Hence we arrive at the following

68 **Proposition 2.** *The property (24) of the curvature tensor R of the manifold (M, g, S) is equivalent to the
69 conditions (26).*

Proposition 3. *If a manifold (M, g, S) has the property (24), then the components of the Ricci tensor ρ satisfy*

$$\rho_{11} = \rho_{22} = \rho_{33} = \rho_{44}, \quad \rho_{12} = \rho_{23} = \rho_{34} = -\rho_{14}, \quad \rho_{13} = \rho_{24} = 0. \quad (27)$$

Proof. Due to Proposition 2 the components of the curvature tensor R satisfy (26). For brevity, we denote

$$R_1 = R_{1313}, \quad R_2 = R_{1213}. \quad (28)$$

Thus, having in mind (8), (21), (26) and (28), we get the components of ρ , as follows:

$$\begin{aligned} \rho_{11} = \rho_{22} = \rho_{33} = \rho_{44} &= \frac{2}{D} (-AR_1 + 2BR_2), \\ \rho_{12} = \rho_{23} = \rho_{34} = -\rho_{14} &= \frac{2}{D} (BR_1 - AR_2), \\ \rho_{13} = \rho_{24} &= 0. \end{aligned} \quad (29)$$

70 i.e. the equalities (27) are valid. \square

A Riemannian manifold is said to be Einstein if its Ricci tensor ρ is a constant multiple of the metric tensor g , i.e.

$$\rho(x, y) = \alpha g(x, y). \quad (30)$$

71 In [17], for locally decomposable Riemannian manifolds is defined a class of almost Einstein
72 manifolds. For the considered in our paper manifolds, we give the following

Definition 2. A Riemannian manifold (M, g, S) is called almost Einstein if the metrics g and \tilde{g} satisfy

$$\rho(x, y) = \alpha g(x, y) + \beta \tilde{g}(x, y), \quad (31)$$

73 where α and β are smooth functions on M .

74 **Theorem 4.** If a manifold (M, g, S) has the property (24), then it is almost Einstein.

Proof. Due to Proposition 3, for (M, g, S) the equalities (27) are valid. Consequently, from (22), using (8), (9) and (27), we get the values of the scalar curvatures τ and τ^* , as follows:

$$\tau = \frac{4}{D} (A\rho_{11} - 2B\rho_{12}), \quad \tau^* = \frac{4}{D} (-B\rho_{11} + A\rho_{12}).$$

Immediately from the latter equalities we have

$$\rho_{11} = \frac{\tau}{4} A + \frac{2\tau^*}{4} B, \quad \rho_{12} = \frac{\tau}{4} B + \frac{\tau^*}{4} A, \quad (32)$$

and bearing in mind (4) and (7) we get

$$\rho_{11} = \frac{\tau}{4} g_{11} + \frac{\tau^*}{4} \tilde{g}_{11}, \quad \rho_{12} = \frac{\tau}{4} g_{12} + \frac{\tau^*}{4} \tilde{g}_{12}.$$

Then, taking into account (4), (7), (27) and (32), we obtain

$$\rho_{ij} = \frac{\tau}{4} g_{ij} + \frac{\tau^*}{4} \tilde{g}_{ij}, \quad (33)$$

i.e.

$$\rho(x, y) = \frac{\tau}{4} g(x, y) + \frac{\tau^*}{4} \tilde{g}(x, y). \quad (34)$$

75 Therefore, comparing (34) with (31), we state that (M, g, S) is an almost Einstein manifold. \square

76 **Corollary 1.** The manifold (M, g, S) with (24) is Einstein if and only if the scalar curvature τ^* vanishes.

Proof. If (M, g, S) has the scalar curvature which satisfies

$$\tau^* = 0, \quad (35)$$

then the equality (34) implies $\rho(x, y) = \frac{\tau}{4}g(x, y)$, i.e. (M, g, S) is an Einstein manifold.

Conversely. Since (M, g, S) is an Einstein manifold its Ricci tensor ρ has the form (30). Thus (34) implies (35). \square

In the next theorem, we explicitly express the curvature tensor R of an almost Einstein manifold (M, g, S) by both structures g and S .

Theorem 5. *Let (M, g, S) have the property (24). Then the curvature tensor R has an expression*

$$R = \frac{\tau}{16}(2\pi_1 + \pi_3) + \frac{\tau^*}{8}\pi_2, \quad (36)$$

where

$$\begin{aligned} \pi_1(x, y, z, u) &= g(y, z)g(x, u) - g(x, z)g(y, u), \\ \pi_2(x, y, z, u) &= g(y, z)\tilde{g}(x, u) + g(x, u)\tilde{g}(y, z) - g(x, z)\tilde{g}(y, u) - g(y, u)\tilde{g}(x, z), \\ \pi_3(x, y, z, u) &= \tilde{g}(y, z)\tilde{g}(x, u) - \tilde{g}(x, z)\tilde{g}(y, u). \end{aligned} \quad (37)$$

Proof. Due to Proposition 3, the components of the Ricci tensor ρ of (M, g, S) are given by (29). Therefore, by straightforward computation, we get

$$R_1 = -\frac{1}{2}(A\rho_{11} + 2B\rho_{12}) \quad R_2 = -\frac{1}{2}(B\rho_{11} + A\rho_{12}).$$

We substitute (32) into the above equalities and obtain

$$R_1 = -\frac{1}{8}((A^2 + 2B^2)\tau + 4AB\tau^*), \quad R_2 = -\frac{1}{8}(2AB\tau + (2B^2 + A^2)\tau^*). \quad (38)$$

From (4), (7), (28) and (38) it follows

$$\begin{aligned} R_{1313} &= \frac{\tau}{16}(2(g_{13}g_{31} - g_{11}g_{33}) + \tilde{g}_{13}\tilde{g}_{31} - \tilde{g}_{11}\tilde{g}_{33}) + \frac{\tau^*}{8}(g_{13}\tilde{g}_{31} + \tilde{g}_{13}g_{31} - \tilde{g}_{11}g_{33} - g_{11}\tilde{g}_{33}), \\ R_{1213} &= \frac{\tau}{16}(2(g_{13}g_{21} - g_{11}g_{23}) + \tilde{g}_{13}\tilde{g}_{21} - \tilde{g}_{11}\tilde{g}_{23}) + \frac{\tau^*}{8}(g_{13}\tilde{g}_{21} + \tilde{g}_{13}g_{21} - \tilde{g}_{11}g_{23} - g_{11}\tilde{g}_{23}), \end{aligned}$$

Consequently, using (4), (7), (26), (28) and (38), we have

$$R_{ijkh} = \frac{\tau}{16}(2(g_{ih}g_{jk} - g_{ik}g_{jh}) + \tilde{g}_{ih}\tilde{g}_{jk} - \tilde{g}_{ik}\tilde{g}_{jh}) + \frac{\tau^*}{8}(g_{ih}\tilde{g}_{jk} + \tilde{g}_{ih}g_{jk} - \tilde{g}_{ik}g_{jh} - g_{ik}\tilde{g}_{jh}),$$

which is equivalent to (36) with (37). \square

5. Curvature properties of (M, g, S)

The sectional curvature of a non-degenerate 2-plane $\{x, y\}$ spanned by the vectors $x, y \in T_p M$ is the value

$$k(x, y) = \frac{R(x, y, x, y)}{g(x, x)g(y, y) - g^2(x, y)}. \quad (39)$$

Let x induce a S -basis of $T_p M$ for (M, g, S) and let $\sigma = \{x, Sx\}$ be a 2-plane. Evidently, if $y \in \sigma$ and $y \neq x$, then $Sy \notin \sigma$. Consequently, σ has only two S -bases: $\{x, Sx\}$ and $\{-x, -Sx\}$. Thus the sectional curvature $k(x, Sx)$ depends only on $\varphi = \angle(x, Sx)$.

Theorem 6. Let (M, g, S) have the property (24) and let a vector x induce a S -basis. Then the sectional curvatures, determined by the S -basis, are

$$\begin{aligned} k(x, Sx) &= k(Sx, S^2x) = k(x, S^3x) = k(S^2x, S^3x) \\ &= \frac{1}{16(\cos^2 \varphi - 1)} (\tau(1 + 2\cos^2 \varphi) + 4\tau^* \cos \varphi), \\ k(x, S^2x) &= k(Sx, S^3x) = -\frac{1}{8} (\tau(1 + 2\cos^2 \varphi) + 4\tau^* \cos \varphi), \end{aligned} \quad (40)$$

87 where $\varphi = \angle(x, Sx)$.

Proof. Let a vector x induce a S -basis. The equalities (3), (12) and (14) imply

$$\begin{aligned} g(x, Sx) &= g(Sx, S^2x) = g(S^2x, S^3x) = -g(x, S^3x) = g(x, x) \cos \varphi, \\ g(x, S^2x) &= g(Sx, S^3x) = 0. \end{aligned} \quad (41)$$

88 Due to Lemma 1, the angle $\varphi = \angle(x, Sx)$ satisfies (16).

Now, from (2), (3), (6) and (41) we find

$$\tilde{g}(x, x) = 2g(x, x) \cos \varphi, \quad \tilde{g}(x, Sx) = g(x, x), \quad \tilde{g}(x, S^2x) = 0, \quad \tilde{g}(x, S^3x) = -g(x, x). \quad (42)$$

89 Applying (36), (37), (41) and (42) in (39), we obtain (40). \square

Corollary 2. Let a vector x induce an orthonormal S -basis. Then

$$\begin{aligned} k(x, Sx) &= k(Sx, S^2x) = k(x, S^3x) = k(S^2x, S^3x) = -\frac{\tau}{16}, \\ k(x, S^2x) &= k(Sx, S^3x) = -\frac{\tau}{8}. \end{aligned}$$

90 **Proof.** The proof follows directly from (40), when $\varphi = \frac{\pi}{2}$. \square

91 Due to Theorem 6 and Corollary 1 we establish the following

Proposition 4. If (M, g, S) with (24) is an Einstein manifold, then the sectional curvatures, determined by an S -basis, are

$$\begin{aligned} k(x, Sx) &= k(Sx, S^2x) = k(x, S^3x) = k(S^2x, S^3x) = \frac{\tau(1 + 2\cos^2 \varphi)}{16(\cos^2 \varphi - 1)}, \\ k(x, S^2x) &= k(Sx, S^3x) = -\frac{\tau}{8}(1 + 2\cos^2 \varphi). \end{aligned}$$

Now, we recall that the Ricci curvature in the direction of a non-zero vector x is the value

$$r(x) = \frac{\rho(x, x)}{g(x, x)}. \quad (43)$$

Theorem 7. Let (M, g, S) have the property (24) and let a vector x induce a S -basis. Then the Ricci curvatures are

$$r(x) = r(Sx) = r(S^2x) = r(S^3x) = \frac{\tau}{4} + \frac{\tau^*}{2} \cos \varphi, \quad (44)$$

92 where $\varphi = \angle(x, Sx)$.

Proof. According to Theorem 4, the Ricci tensor ρ is given by (34). Then, using (3), we find

$$\rho(x, x) = \rho(Sx, Sx) = \rho(S^2x, S^2x) = \rho(S^3x, S^3x) = \frac{\tau}{4}g(x, x) + \frac{\tau^*}{4}\tilde{g}(x, x). \quad (45)$$

93 Let a vector x induce a S -basis. From (3), (42), (43) and (45) it follows (44). \square

94 Further, Theorem 7 and Corollary 1 imply the next statement.

Proposition 5. *Let (M, g, S) with (24) be an Einstein manifold. Then the Ricci curvatures are*

$$r(x) = r(Sx) = r(S^2x) = r(S^3x) = \frac{\tau}{4}.$$

95 **Proof.** These equalities follow directly by substituting $\tau^* = 0$ into (44). \square

96 **6. Manifolds with parallel structures**

In this section we study a manifold (M, g, S) , whose structure S satisfies (23). Also, we consider an associated manifold (M, g, J) with a structure $J = S^2$. Bearing in mind (2) and (3), we get that the manifold (M, g, J) is almost Hermitian and the structure J is almost complex. In case that J is parallel (M, g, J) is a Kähler manifold. The characteristic condition of a Kähler manifold is

$$\nabla J = 0. \quad (46)$$

97 We note that equalities (23) and $J = S^2$ imply (46).

Theorem 8. *Let (M, g, S) have the property (23). Then the scalar curvatures τ and τ^* satisfy*

$$3\tau_1 = \tau_2^* - \tau_4^*, \quad 3\tau_2 = \tau_1^* + \tau_3^*, \quad 3\tau_3 = \tau_2^* + \tau_4^*, \quad 3\tau_4 = -\tau_1^* + \tau_3^*, \quad (47)$$

98 where $\tau_i = \frac{\partial \tau}{\partial X^i}$, $\tau_i^* = \frac{\partial \tau^*}{\partial X^i}$.

Proof. It is known that in a Riemannian manifold for the scalar curvature τ and the Ricci tensor ρ it is valid

$$\nabla_i \rho_k^i = \frac{1}{2} \nabla_k \tau, \quad (48)$$

99 where $\rho_k^i = \rho_{ak} g^{ai}$.

On the other hand, if (M, g, S) satisfies (23), then it satisfies (24). Therefore, the Ricci tensor has the expression (33). Hence, from (1), (4), (7), (8) and (33), we get

$$\rho_k^i = \frac{\tau}{4} \delta_k^i + \frac{\tau^*}{4} (S_k^i - (S_k^i)^3),$$

where δ_k^i are the Kronecker symbols. Using the above equalities, (23) and (48) we obtain

$$\tau_k = \frac{\tau_i}{4} \delta_k^i + \frac{\tau_i^*}{4} (S_k^i - (S_k^i)^3).$$

100 Then, from (1) it follows (47). \square

101 According to Theorem 8 and Corollary 1 we establish the following

102 **Proposition 6.** *If (M, g, S) with (23) is an Einstein manifold, then the scalar curvature τ is a constant.*

103 6.1. Conditions for parallel structures

Theorem 9. The manifold (M, g, S) satisfies (23) if and only if

$$A_1 = B_2 - B_4, \quad A_2 = B_1 + B_3, \quad A_3 = B_2 + B_4, \quad A_4 = B_4 = B_3 - B_1, \quad (49)$$

104 where $A_i = \frac{\partial A}{\partial X^i}$, $B_i = \frac{\partial B}{\partial X^i}$.

Proof. If Γ_{ij}^s are the Christoffel symbols of ∇ , then

$$\nabla_i S_j^t = \partial_i S_j^t + \Gamma_{ik}^t S_j^k - \Gamma_{ij}^k S_k^t. \quad (50)$$

Together with (23), (50) yields

$$\Gamma_{ik}^t S_j^k = \Gamma_{ij}^k S_k^t. \quad (51)$$

From (1) and (51) we get

$$\begin{aligned} \Gamma_{11}^1 &= \Gamma_{12}^2 = \Gamma_{13}^3 = \Gamma_{14}^4 = \Gamma_{22}^3 = \Gamma_{23}^4 = -\Gamma_{24}^1 = -\Gamma_{33}^1 = -\Gamma_{34}^2 = -\Gamma_{44}^3, \\ \Gamma_{11}^2 &= \Gamma_{12}^3 = \Gamma_{13}^4 = -\Gamma_{14}^1 = \Gamma_{22}^4 = -\Gamma_{23}^1 = -\Gamma_{24}^2 = -\Gamma_{33}^2 = -\Gamma_{34}^3 = -\Gamma_{44}^4, \\ \Gamma_{11}^3 &= \Gamma_{12}^4 = -\Gamma_{13}^1 = -\Gamma_{14}^2 = -\Gamma_{22}^1 = -\Gamma_{23}^2 = -\Gamma_{24}^3 = -\Gamma_{33}^3 = -\Gamma_{34}^4 = \Gamma_{44}^1, \\ \Gamma_{11}^4 &= -\Gamma_{12}^1 = -\Gamma_{13}^2 = -\Gamma_{14}^3 = -\Gamma_{22}^2 = -\Gamma_{23}^3 = -\Gamma_{24}^4 = -\Gamma_{33}^4 = \Gamma_{34}^1 = \Gamma_{44}^2. \end{aligned} \quad (52)$$

Now, using (1), (4), (8) and the well known identities

$$2\Gamma_{ij}^s = g^{as}(\partial_i g_{aj} + \partial_j g_{ai} - \partial_a g_{ij}),$$

we calculate

$$\begin{aligned} \Gamma_{11}^1 &= \frac{1}{2D}(A A_1 - B(4B_1 - A_2 + A_4)), & \Gamma_{11}^2 &= \frac{1}{2D}(A(2B_1 - A_2) + B(A_3 - A_1)), \\ \Gamma_{11}^3 &= \frac{1}{2D}(B(A_2 + A_4) - A A_3), & \Gamma_{11}^4 &= \frac{1}{2D}(B(A_1 + A_3) - A(2B_1 + A_4)), \\ \Gamma_{12}^1 &= \frac{1}{2D}(A A_2 - B(A_1 + B_2 + B_4)), & \Gamma_{12}^2 &= \frac{1}{2D}(A A_1 - B(A_2 + B_1 - B_3)), \\ \Gamma_{12}^3 &= \frac{1}{2D}(A(B_1 - B_3) - B(A_1 - B_2 - B_4)), & \Gamma_{12}^4 &= \frac{1}{2D}(B(A_2 + B_3 - B_1) - A(B_2 + B_4)), \\ \Gamma_{13}^1 &= \frac{1}{2D}(A A_3 - 2B B_3), & \Gamma_{13}^2 &= \frac{1}{2D}(A(B_1 + B_3) - B(A_1 + A_3)), \\ \Gamma_{13}^3 &= \frac{1}{2D}(A A_1 - 2B B_1), & \Gamma_{13}^4 &= \frac{1}{2D}(A(B_1 - B_3) + B(A_3 - A_1)), \\ \Gamma_{14}^1 &= \frac{1}{2D}(A A_4 + B(A_1 - B_2 - B_4)), & \Gamma_{14}^2 &= \frac{1}{2D}(A(B_2 + B_4) - B(A_4 + B_1 + B_3)), \\ \Gamma_{14}^3 &= \frac{1}{2D}(A(B_1 + B_3) - B(A_1 + B_2 + B_4)), & \Gamma_{14}^4 &= \frac{1}{2D}(A A_1 + B(A_4 - B_1 - B_3)), \\ \Gamma_{22}^1 &= \frac{1}{2D}(A(2B_2 - A_1) - B(A_2 + A_4)), & \Gamma_{22}^2 &= \frac{1}{2D}(A A_2 - B(4B_2 - A_1 - A_3)), \\ \Gamma_{22}^3 &= \frac{1}{2D}(A(2B_2 - A_3) + B(A_4 - A_2)), & \Gamma_{22}^4 &= \frac{1}{2D}(B(A_3 - A_1) - A A_4), \\ \Gamma_{23}^1 &= \frac{1}{2D}(A(B_3 - B_1) - B(A_3 - B_2 + B_4)), & \Gamma_{23}^2 &= \frac{1}{2D}(A A_3 + B(B_1 - B_3 - A_2)), \\ \Gamma_{23}^3 &= \frac{1}{2D}(A A_2 - B(B_2 - B_4 + A_3)), & \Gamma_{23}^4 &= \frac{1}{2D}(A(B_2 - B_4) - B(A_2 + B_1 - B_3)), \\ \Gamma_{24}^1 &= \frac{1}{2D}(A(B_4 - B_2) - B(A_4 - A_2)), & \Gamma_{24}^2 &= \frac{1}{2D}(A A_4 - 2B B_4), \\ \Gamma_{24}^3 &= \frac{1}{2D}(A(B_2 + B_4) - B(A_2 + A_4)), & \Gamma_{24}^4 &= \frac{1}{2D}(A A_2 - 2B B_2), \\ \Gamma_{33}^1 &= \frac{1}{2D}(B(A_2 - A_4) - A A_1), & \Gamma_{33}^2 &= \frac{1}{2D}(A(2B_3 - A_2) + B(A_1 - A_3)), \\ \Gamma_{33}^3 &= \frac{1}{2D}(A A_3 - B(4B_3 - A_2 - A_4)), & \Gamma_{33}^4 &= \frac{1}{2D}(A(2B_3 - A_4) - B(A_1 + A_3)), \\ \Gamma_{34}^1 &= \frac{1}{2D}(B(A_3 + B_2 - B_4) - A(B_1 + B_3)), & \Gamma_{34}^2 &= \frac{1}{2D}(A(B_4 - B_2) + B(B_3 + B_1 - B_4)), \\ \Gamma_{34}^3 &= \frac{1}{2D}(A B_4 - B(A_3 - B_2 + B_4)), & \Gamma_{34}^4 &= \frac{1}{2D}(A A_3 - B(B_1 + B_3 + B_4)), \\ \Gamma_{44}^1 &= \frac{1}{2D}(B(A_2 + A_4) - A(2B_4 + A_1)), & \Gamma_{44}^2 &= \frac{1}{2D}(B(A_1 + A_3) - A A_2), \\ \Gamma_{44}^3 &= \frac{1}{2D}(A(2B_4 - A_3) + B(A_2 - A_4)), & \Gamma_{44}^4 &= \frac{1}{2D}(A A_4 - B(4B_4 + A_1 - A_3)). \end{aligned} \quad (53)$$

105 We apply (53) in (52) and obtain the conditions (49).

106 Vice versa. Let (49) hold true. We put equalities (49) into (53) and find (52). Hence (1) and (52)
107 imply (51). Consequently, from (1), (50) and (51) we get (23). \square

¹⁰⁸ **Theorem 10.** *The manifold (M, g, J) is Kähler if and only if the equalities (49) are valid.*

Proof. Having in mind (1), we get that the components of the structure $J = S^2$ on (M, g, J) are given by the skew-circulant matrix

$$(J_j^k) = \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ -1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \end{pmatrix}. \quad (54)$$

Therefore, from (46), (54) and

$$\nabla_i J_j^t = \partial_i J_j^t + \Gamma_{ik}^t J_j^k - \Gamma_{ij}^k J_k^t$$

it follows

$$\Gamma_{ik}^t J_j^k = \Gamma_{ij}^k J_k^t. \quad (55)$$

Together with (54), (55) yields

$$\begin{aligned} \Gamma_{11}^1 &= \Gamma_{13}^3 = -\Gamma_{33}^1, & \Gamma_{14}^4 &= \Gamma_{23}^4 = \Gamma_{12}^2 = -\Gamma_{34}^2, & \Gamma_{22}^3 &= -\Gamma_{24}^1 = -\Gamma_{44}^3, \\ \Gamma_{11}^2 &= \Gamma_{13}^4 = -\Gamma_{33}^2, & \Gamma_{14}^1 &= \Gamma_{23}^1 = -\Gamma_{12}^3 = \Gamma_{34}^3, & \Gamma_{22}^4 &= -\Gamma_{24}^2 = -\Gamma_{44}^4, \\ \Gamma_{11}^3 &= -\Gamma_{13}^1 = -\Gamma_{33}^3, & \Gamma_{14}^2 &= \Gamma_{23}^2 = -\Gamma_{12}^4 = \Gamma_{34}^4, & \Gamma_{22}^1 &= \Gamma_{24}^3 = -\Gamma_{44}^1, \\ \Gamma_{11}^4 &= -\Gamma_{13}^2 = -\Gamma_{33}^4, & \Gamma_{14}^3 &= \Gamma_{23}^3 = \Gamma_{12}^1 = -\Gamma_{34}^1, & \Gamma_{22}^2 &= \Gamma_{24}^4 = -\Gamma_{44}^2. \end{aligned} \quad (56)$$

¹⁰⁹ We apply (53) in (56) and obtain conditions (49).

¹¹⁰ Vice versa. From (49) it follows (23). Obviously (23) implies (46). \square

¹¹¹ Bearing in mind Theorem 9 and Theorem 10 we state the following

¹¹² **Corollary 3.** *The structure S of (M, g, S) is parallel with respect to ∇ if and only if the structure J of (M, g, J) is parallel with respect to ∇ .*

¹¹⁴ **7. Lie groups as 4-dimensional Riemannian manifolds with skew-circulant structures**

Let G be a 4-dimensional real connected Lie group and \mathfrak{g} be its Lie algebra with a basis $\{x_1, x_2, x_3, x_4\}$. We introduce a structure S and left invariant metric g as follows

$$Sx_1 = x_2, Sx_2 = x_3, Sx_3 = x_4, Sx_4 = -x_1, \quad (57)$$

$$g(x_i, x_j) = \begin{cases} 0, & i \neq j; \\ 1, & i = j. \end{cases} \quad (58)$$

¹¹⁵ Obviously (2) and (3) are valid. Therefore (G, g, S) is a Riemannian manifold of the considered type.

For the manifold (G, g, S) we suppose that S is an Abelian structure, i.e.

$$[x_i, x_j] = [Sx_i, Sx_j]. \quad (59)$$

According to (57), (59) and the Jacobi identity for the commutators $[x_i, x_j]$ we obtain

$$\begin{aligned} [x_1, x_2] &= [x_1, x_4] = [x_2, x_3] = [x_3, x_4] = \lambda_1 x_1 + \lambda_2 x_2 + \lambda_3 x_3 + \lambda_4 x_4, \\ [x_1, x_3] &= [x_2, x_4] = (\lambda_2 - \lambda_4) x_1 + (\lambda_1 + \lambda_3) x_2 + (\lambda_2 + \lambda_4) x_3 + (\lambda_3 - \lambda_1) x_4, \end{aligned} \quad (60)$$

¹¹⁶ where $\lambda_i \in \mathbb{R}$.

¹¹⁷ It is easy to see that a manifold (G, g, S) with a Lie algebra \mathfrak{g} , determined by (60), has an Abelian structure S .

¹¹⁹ **Theorem 11.** Let (G, g, S) be a manifold with a Lie algebra \mathbf{g} determined by (60). Then (G, g, S) satisfies the
¹²⁰ identity (23).

Proof. The well-known Koszul formula implies

$$2g(\nabla_{x_i}x_j, x_k) = g([x_i, x_j], x_k) + g([x_k, x_i], x_j) + g([x_k, x_j], x_i),$$

and having in mind (58) and (60), we get

$$\begin{aligned} \nabla_{x_1}x_1 &= -\lambda_1(x_2 + x_4) + (\lambda_4 - \lambda_2)x_3, & \nabla_{x_1}x_2 &= \lambda_1(x_1 - x_3) + (\lambda_4 - \lambda_2)x_4, \\ \nabla_{x_1}x_3 &= \lambda_1(x_2 - x_4) + (\lambda_2 - \lambda_4)x_1, & \nabla_{x_1}x_4 &= \lambda_1(x_1 + x_3) + (\lambda_2 - \lambda_4)x_2, \\ \nabla_{x_2}x_1 &= -\lambda_2(x_2 + x_4) - (\lambda_1 + \lambda_3)x_3, & \nabla_{x_2}x_2 &= \lambda_2(x_1 - x_3) - (\lambda_1 + \lambda_3)x_4, \\ \nabla_{x_2}x_3 &= \lambda_2(x_2 - x_4) + (\lambda_1 + \lambda_3)x_1, & \nabla_{x_2}x_4 &= \lambda_2(x_1 + x_3) + (\lambda_1 + \lambda_3)x_2, \\ \nabla_{x_3}x_1 &= -\lambda_3(x_2 + x_4) - (\lambda_2 + \lambda_4)x_3, & \nabla_{x_3}x_2 &= \lambda_3(x_1 - x_3) - (\lambda_2 + \lambda_4)x_4, \\ \nabla_{x_3}x_3 &= \lambda_3(x_2 - x_4) + (\lambda_2 + \lambda_4)x_1, & \nabla_{x_3}x_4 &= \lambda_3(x_1 + x_3) + (\lambda_2 + \lambda_4)x_2, \\ \nabla_{x_4}x_1 &= -\lambda_4(x_2 + x_4) + (\lambda_1 - \lambda_3)x_3, & \nabla_{x_4}x_2 &= \lambda_4(x_1 - x_3) + (\lambda_1 - \lambda_3)x_4, \\ \nabla_{x_4}x_3 &= \lambda_4(x_2 - x_4) + (\lambda_3 - \lambda_1)x_1, & \nabla_{x_4}x_4 &= \lambda_4(x_1 + x_3) + (\lambda_3 - \lambda_1)x_2. \end{aligned} \tag{61}$$

¹²¹ From (57), (61) and the formula $(\nabla_{x_i}S)x_j = \nabla_{x_i}Sx_j - S\nabla_{x_i}x_j$ we get $(\nabla_{x_i}S)x_j = 0$, i.e. (23) is valid. \square

Further, using (19), (20), (58), (60) and (61) we calculate the following components of the curvature tensor R :

$$\begin{aligned} R_{1313} = R_{2424} = R_{1324} &= 2R_{1212} = 2R_{1414} = 2R_{2323} = 2R_{3434} = 2R_{1223} = 2R_{1214} \\ &= 2R_{1434} = 2R_{1234} = 2R_{2334} = 2R_{2314} = 2(\lambda_1^2 + \lambda_2^2 + \lambda_3^2 + \lambda_4^2), \\ R_{1213} = R_{1224} = R_{1413} &= R_{2414} = R_{2423} = R_{2313} = R_{1334} = R_{2434} \\ &= 2(\lambda_1\lambda_2 + \lambda_2\lambda_3 + \lambda_3\lambda_4 - \lambda_1\lambda_4). \end{aligned} \tag{62}$$

The rest of the nonzero components are obtained from the properties

$$R_{ijks} = R_{ksij}, R_{ijks} = -R_{jiks} = -R_{isks}.$$

From (58), (62) and the formula (21) we get the components of the Ricci tensor ρ :

$$\begin{aligned} \rho_{11} = \rho_{22} = \rho_{33} = \rho_{44} &= -4(\lambda_1^2 + \lambda_2^2 + \lambda_3^2 + \lambda_4^2), \\ \rho_{12} = \rho_{23} = \rho_{34} &= -4(\lambda_1\lambda_2 + \lambda_2\lambda_3 + \lambda_3\lambda_4 - \lambda_1\lambda_4), \\ \rho_{13} = \rho_{24} &= 0, \quad \rho_{14} = -\rho_{12}. \end{aligned} \tag{63}$$

Now, using (6) and (58), we find the components of \tilde{g} and the components of its inverse. They are as follows:

$$\begin{aligned} \tilde{g}_{11} = \tilde{g}_{22} = \tilde{g}_{33} = \tilde{g}_{44} &= 0, \quad \tilde{g}_{12} = \tilde{g}_{23} = \tilde{g}_{34} = -\tilde{g}_{14} = 1, \quad \tilde{g}_{13} = \tilde{g}_{24} = 0, \\ \tilde{g}^{11} = \tilde{g}^{22} = \tilde{g}^{33} = \tilde{g}^{44} &= 0, \quad \tilde{g}^{12} = \tilde{g}^{23} = \tilde{g}^{34} = -\tilde{g}^{14} = \frac{1}{2}, \quad \tilde{g}^{13} = \tilde{g}^{24} = 0. \end{aligned}$$

Then, applying (58), (63) in (22), we get the values of the scalar curvatures τ and τ^* as follows:

$$\tau = -16(\lambda_1^2 + \lambda_2^2 + \lambda_3^2 + \lambda_4^2), \quad \tau^* = -16(\lambda_1\lambda_2 + \lambda_2\lambda_3 + \lambda_3\lambda_4 - \lambda_1\lambda_4). \tag{64}$$

¹²² Consequently, the equalities (58), (63) and (64) imply (33), i.e. (G, g, S) is an almost Einstein manifold. Further, using (39), (58) and (62), for the sectional curvatures of the basic 2-planes we find

$$\begin{aligned} k(x_2, x_4) = k(x_1, x_3) &= 2(\lambda_1^2 + \lambda_2^2 + \lambda_3^2 + \lambda_4^2), \\ k(x_1, x_2) = k(x_1, x_4) &= k(x_2, x_3) = k(x_3, x_4) = \lambda_1^2 + \lambda_2^2 + \lambda_3^2 + \lambda_4^2. \end{aligned} \tag{65}$$

¹²³ Therefore, we arrive at the following

¹²⁴ **Theorem 12.** *Let (G, g, S) be a manifold with a Lie algebra \mathbf{g} determined by (60). Then*

- ¹²⁵ (i) *The nonzero components of the curvature tensor R are (62);*
- ¹²⁶ (ii) *The components of the Ricci tensor ρ are (63);*
- ¹²⁷ (iii) *The scalar curvatures τ and τ^* are (64). The manifold is almost Einstein;*
- ¹²⁸ (iv) *The sectional curvatures of the basic 2-planes are (65).*

¹²⁹ Bearing in mind Corollary 1 and the second equality of (64) we construct two examples of Einstein
¹³⁰ manifolds (G, g, S) .

Proposition 7. *Let (G, g, S) be a manifold with a Lie algebra \mathbf{g} determined by*

$$\begin{aligned}[x_1, x_2] &= [x_1, x_4] = [x_2, x_3] = [x_3, x_4] = \lambda_1 x_1 + \lambda_1 x_3 + \lambda_4 x_4, \\ [x_1, x_3] &= [x_2, x_4] = -\lambda_4 x_1 + 2\lambda_1 x_2 + \lambda_4 x_3.\end{aligned}$$

¹³¹ Then

- (i) *The nonzero components of the Ricci tensor ρ are*

$$\rho_{11} = \rho_{22} = \rho_{33} = \rho_{44} = -4(2\lambda_1^2 + \lambda_4^2);$$

- (ii) *The manifold is Einstein and the scalar curvatures τ and τ^* are*

$$\tau = -16(2\lambda_1^2 + \lambda_4^2), \quad \tau^* = 0;$$

- (iii) *The sectional curvatures of the basic 2-planes are*

$$k(x_2, x_4) = k(x_1, x_3) = 2(2\lambda_1^2 + \lambda_4^2), \quad k(x_1, x_2) = k(x_1, x_4) = k(x_2, x_3) = k(x_3, x_4) = 2\lambda_1^2 + \lambda_4^2.$$

¹³² **Proof.** This example we get by substituting $\lambda_3 = \lambda_1$ and $\lambda_2 = 0$ into each of the equalities (60), (63),
¹³³ (64) and (65). \square

Proposition 8. *Let (G, g, S) be a manifold with a Lie algebra \mathbf{g} determined by*

$$\begin{aligned}[x_1, x_2] &= [x_1, x_4] = [x_2, x_3] = [x_3, x_4] = (\lambda_2 + \lambda_4)x_1 + \lambda_2 x_2 + (\lambda_4 - \lambda_2)x_3 + \lambda_4 x_4, \\ [x_1, x_3] &= [x_2, x_4] = (\lambda_2 - \lambda_4)x_1 + 2\lambda_4 x_2 + (\lambda_2 + \lambda_4)x_3 - 2\lambda_2 x_4.\end{aligned}$$

¹³⁴ Then

- (i) *The nonzero components of the Ricci tensor ρ are*

$$\rho_{11} = \rho_{22} = \rho_{33} = \rho_{44} = -12(\lambda_2^2 + \lambda_4^2);$$

- (ii) *The manifold is Einstein and the scalar curvatures τ and τ^* are*

$$\tau = -48(\lambda_2^2 + \lambda_4^2), \quad \tau^* = 0;$$

- (iii) *The sectional curvatures of the basic 2-planes are*

$$k(x_2, x_4) = k(x_1, x_3) = 6(\lambda_2^2 + \lambda_4^2), \quad k(x_1, x_2) = k(x_1, x_4) = k(x_2, x_3) = k(x_3, x_4) = 3(\lambda_2^2 + \lambda_4^2).$$

¹³⁵ **Proof.** We put $\lambda_1 = \lambda_2 + \lambda_4$ and $\lambda_3 = \lambda_4 - \lambda_2$ into each of the equalities (60), (63), (64) and (65). \square

¹³⁶ **Author Contributions:** All authors contributed equally to this work.

¹³⁷ **Funding:** This work is partially supported by project MU19-FMI-020 of the Scientific Research Fund, Paisii
¹³⁸ Hilendarski University of Plovdiv, Bulgaria.

¹³⁹ **Acknowledgments:** The authors are grateful to Professor Dr. G. Dzhelepov for his valuable comments on this
¹⁴⁰ paper.

¹⁴¹ **Conflicts of Interest:** The authors declare no conflict of interest.

¹⁴² **References**

- ¹⁴³ 1. Davis, P. J. *Circulant matrices, A Wiley-Interscience Publication. Pure and Applied Mathematics*. John Wiley and Sons, New York-Chichester-Brisbane, New York, USA, 1979; pp. 250.
- ¹⁴⁴ 2. Dzhelepov G.; Dokuzova I.; Razpopov D. On a three-dimensional Riemannian manifold with an additional structure, *Plovdiv Univ. Paisi Khilendarski Nauchn. Trud. Mat.*, **2011**, 38 (3), 17–27.
- ¹⁴⁵ 3. Gray, R. M. Toeplitz and circulant matrices: A review, *Found. Trends Commun. Inf. Theory* **2006**, 2 (3), 155–239.
- ¹⁴⁶ 4. Gray, A. Curvature identities for Hermitian and almost Hermitian manifolds, *Tôhoku Math. J.* (2) **1976**, 28 (4), 601–612.
- ¹⁴⁷ 5. Gray, A.; Hervella, L.M. The sixteen classes of almost Hermitian manifolds and their linear invariants, *Ann. Mat. Pura Appl.* (4) **1980**, 123, 35–58.
- ¹⁴⁸ 6. Gray, A.; Vanhecke, L. Almost Hermitian manifolds with constant holomorphic sectional curvatures, *Časopis Pěst. Mat.* **1979**, 104 (2), 170–179.
- ¹⁴⁹ 7. Jiang, XY.; Hong, K. Explicit determinants of the k-Fibonacci and k-Lucas RSFPLR circulant matrix in codes. In: Yang Y., Ma M., Liu B. (eds) *Information Computing and Applications. ICICA 2013. Communications in Computer and Information Science*, 391. Springer, Berlin, Heidelberg, Germany 2013; 625–637.
- ¹⁵⁰ 8. Liu, K.; Yang, X. Ricci curvatures on Hermitian manifolds, *Trans. American Math. Soc.* **2017**, 369 (7), 5157–5196.
- ¹⁵¹ 9. Muzychuk, M. A Solution of the isomorphism problem for circulant graphs, In: *Proc. London Math. Soc.* (3) **2004**, 88 (1), 1–41.
- ¹⁵² 10. Matsuo, K. Pseudo-Bochner curvature tensor on Hermitian manifolds, *Colloq. Math.* **1999**, 80 (2), 201–209.
- ¹⁵³ 11. Olson, B.; Shaw, S.; Shi, C.; Pierre C.; Parker, R. G. Circulant matrices and their application to vibration analysis, *Appl. Mech. Rev.*, **2014**, 66 (4), 1–41.
- ¹⁵⁴ 12. Razpopov, D. Four-dimensional Riemannian manifolds with two circulant structures, *Math. Educ. Math.* **2015**, 44, In: *Proc. of 44-th Spring Conf. of UBM*, SOK Kamchia, Bulgaria, 179–185.
- ¹⁵⁵ 13. Roth, R. M.; Lempel, A. Application of circulant matrices to the construction and decoding of linear codes, *IEEE Trans. Inform. Theory*, **1990**, 36 (5), 1157–1163.
- ¹⁵⁶ 14. Prvanović, M. Conformally invariant tensors of an almost Hermitian manifold associated with the holomorphic curvature tensor, *J. Geom.* **2012**, 103 (1), 89–101.
- ¹⁵⁷ 15. Vanhecke, L. Some almost Hermitian manifolds with constant holomorphic sectional curvature, *J. Differential Geom.* **1977**, 12(4): 461–471.
- ¹⁵⁸ 16. Yang, B.; Zheng, F. On curvature tensor of Hermitian manifolds, *Comm. Anal. Geom.* **2018**, 26 (5), 1195–1222.
- ¹⁵⁹ 17. Yano K. Differential geometry on complex and almost complex spaces, *International Series of Monographs in Pure and Applied Mathematics* 49, A Pergamon Press Book The Macmillan and Co., New York, USA, 1965; pp. 326.