

Article

# Plasma Spectroscopy of Various Types of Gypsum: An ideal Terrestrial Analogue

Abhishek K. Rai <sup>1</sup>, Jayanta K. Pati <sup>1,2</sup>, Christian G. Parigger <sup>3,\*</sup>, Awadhesh K. Rai <sup>4</sup>

<sup>1</sup> Department of Earth and Planetary Sciences, Nehru Science Centre, University of Allahabad,

Prayagraj-211 002, India; [abhishekraigeology@gmail.com](mailto:abhishekraigeology@gmail.com)

<sup>2</sup> National Center of Experimental Mineralogy and Petrology, 14, Chatham Lines, University of Allahabad,

Prayagraj-211 002, India; [jkpati@gmail.com](mailto:jkpati@gmail.com)

<sup>3</sup> Department of Physics and Astronomy, University of Tennessee/University of Tennessee Space Institute, 411

B. H. Goethert Parkway, Tullahoma, TN – 37388, USA; [cparigge@tennessee.edu](mailto:cparigge@tennessee.edu)

<sup>4</sup> Department of Physics, University of Allahabad, Prayagraj-211002, India; [awadheshkrai@gmail.com](mailto:awadheshkrai@gmail.com)

\* Correspondence: [cparigge@tennessee.edu](mailto:cparigge@tennessee.edu); Tel.: +1-931-841-5690

**Abstract:** The first detection of gypsum ( $\text{CaSO}_4 \cdot 2\text{H}_2\text{O}$ ) by the Mars Science Laboratory (MSL) rover Curiosity in the Gale Crater, Mars created a profound impact on planetary science and exploration. The unique capability of plasma spectroscopy involving in situ elemental analysis in extraterrestrial environments, suggesting the presence of water in the red planet based on phase characterization and providing a clue to Martian paleoclimate. The key to gypsum as an ideal paleoclimate proxy lies in its textural variants, and in this study terrestrial gypsum samples from varied locations and textural types have been analyzed with laser-induced breakdown spectroscopy (LIBS). Petrographic, sub-microscopic and powder X-ray diffraction characterizations confirm the presence of gypsum (hydrated calcium sulphate;  $\text{CaSO}_4 \cdot 2\text{H}_2\text{O}$ ), bassanite (semi-hydrated calcium sulphate;  $\text{CaSO}_4 \cdot 1/2\text{H}_2\text{O}$ ) and anhydrite (anhydrous calcium sulphate;  $\text{CaSO}_4$ ) along with accessory phases (quartz and jarosite). The principal component analysis of LIBS spectra from texturally varied gypsums can be differentiated from one another because of the chemical variability in their elemental concentrations. The concentration of gypsum is determined from the partial least-square regressions model. Rapid characterization of gypsum samples with LIBS is expected to work well in extraterrestrial environments.

**Keywords:** laser-induced plasma; atomic spectroscopy; laser-induced breakdown spectroscopy; atomic spectroscopy; principal component analysis; partial least-square regression; gypsum; Mars

## 1. Introduction

Laser Induced Breakdown Spectroscopy (LIBS) is an atomic emission spectroscopic technique in which high power laser pulse is focused on the surface of the material and plasma is created. Light from the plasma is fed to the spectrometer to get spectral lines of the corresponding elements present in the material. The LIBS technique is applied in mineral and rock analyses in field and laboratory environments [1–2] for rapid measurements without involving sample preparation. It has a unique in-situ point detection capability. LIBS technique was introduced in planetary exploration for the first time in 2012 when the Mars Science Laboratory (MSL) rover Curiosity studied the Gale Crater [3–4]. The ChemCam identified gypsum, a water-bearing non-metallic calcium sulphate mineral ( $\text{CaSO}_4 \cdot 2\text{H}_2\text{O}$ ) as veins on the rim of Endeavour Crater, Mars as veins [5–6]. Consequently, it was suggested that gypsum on Mars formed due to evaporation of a large lake [7–9] under a suitable

physico-chemical environment. Sulphate salts are useful markers to decode paleoenvironments as they precipitate from aqueous fluids suggesting the presence of liquid water. Further the aqueous geochemistry promotes the proliferation of microbial life and its consequent preservation in planetary environments [10].

Gypsum provides ideal habitat for endolithic microbial communities, protection from harmful ultra-violet radiations and supports their proliferation even under High Arctic conditions [11] to hot springs [12] environments with temperatures larger than 50°C. Microbes thrive within gypsum crystals in terrestrial impact craters, such as Haughton structure [13-15] and the extraterrestrial landscapes are endowed with such structures [16]. Based on terrestrial evidences, the possible presence of microbial life is predicted in gypsum veins and in other sulfate minerals on Mars [17-18], other planets of the solar system, their moons, asteroids and comets.

The MSL Curiosity rover characterized many calcium sulfate veins at Gale crater, Mars including gypsum at Martian surface with nominal change in elemental composition [19]. Dunes of gypsum are also detected at the Martian North pole [20-21]. Presence of sulphates associated with clays at the base of the Mount Sharp by Compact Reconnaissance Imaging Spectrometer (CRISM) is also pointed out by Milliken *et al.* 2010 [22] and also by Opportunity rover at Meridiani Planum [23].

Various hypotheses explaining the origin of the massive Martian sulphate deposits exist advocating hydrothermal processes, oxidation of sulphides, evaporative deposition, and bedrock leaching by volcanic vapors [23-29]. The report of gypsum veins from Mars, its formation in various terrestrial environments and suitability for the proliferation of life have made this hydrated sulfate mineral an ideal terrestrial analogue [30]. In addition, the texturally different types of gypsum, which in turn control the physical and chemical conditions of their formation, have attracted significant attention [31]. In the present work, gypsum samples from varied terrestrial environments and textural types have been studied for their inter-sample comparison using LIBS data combined with multivariate statistical analysis methods. The data generated can be compared with the available ChemCam data on gypsum from Mars and upcoming Chandrayaan-2 experiments from Moon [32].

## 2. Mineralogy and geochemistry of gypsum

Gypsum crystallizes in monoclinic system and its structure contains parallel layers of  $(\text{SO}_4)^{2-}$  group bonded with  $(\text{Ca})^{+2}$ . The  $\text{CaSO}_4$  molecules are separated by sheets of  $\text{H}_2\text{O}$ -molecules having very weak bonds and breakdown on heating [33-34]. Calcium atoms are enclosed by two water molecules and six sulphate oxygen atoms, though the sulphur atoms are at the middle of sulphate oxygen tetrahedra. The water molecules have two oxygen atoms and calcium as their first neighbours. The arrangement consists of  $\text{SO}_4$  tetrahedra,  $\text{CaO}_8$  polyhedra and weak hydrogen bonds made among the O atoms of nearest-neighbour  $\text{SO}_4$  tetrahedra and intercalated  $\text{H}_2\text{O}$  molecules [35]. The diaphaneity of gypsum varies from transparent to opaque and is usually light in color. The habit of gypsum is platy, columnar, fibrous, needle-like, lenticular, forms massive aggregates and twins-swallowtail. It is a relatively soft mineral (1.5-2: Mohs' Scale of Hardness) and gets easily weathered. Gypsum constitutes 79.1% of calcium sulphate and 20.9 % water by weight. Since gypsum is known to form by several processes (diagenesis [36]; chemical weathering [37]; evaporation [38]; hydrothermal activities etc. [39]) and under varied physical and chemical environments and this has led to useful reconstruction of the paleoenvironment [40-41]. Sometimes

lamination occurring in gypsum (medium to coarse grains) from the evaporate basin can lead us to know about the changes in the basin i.e. water composition and water level. Gypsum laminas (like biolamina) crystallizes in the shallow part, approximately 200m can be either deformed by cyclic droughts like mudcracks or ruptured by crystallizing sulphates. Gypsums crystallizing in the deeper zones constitute elongated crystals (nearly 20-30 cm long), distorted in one direction due to the low floor current activity, and represent as proper paleocurrent indicators. Selenite is one the varieties of gypsum, which forms under stable condition at few to several meters depth and they can grow up to 10 meters. Laminated selenites forming under deeper zones show ripple marks, turbidities and slump structure with fragments of older and more lithified gypsum.

Calcium sulfate can crystallize in three phases when it comes with contact with water gypsum ( $\text{CaSO}_4 \cdot 2\text{H}_2\text{O}$ ), hemihydrate or bassinite ( $\text{CaSO}_4 \cdot \frac{1}{2}\text{H}_2\text{O}$ ) and anhydrite ( $\text{CaSO}_4$ ). Gypsum may lose crystallization water molecules, becoming bassanite (hemihydrate:  $\text{CaSO}_4 \cdot \frac{1}{2}\text{H}_2\text{O}$ ) or anhydrite ( $\text{CaSO}_4$ ) under suitable geological and temperature conditions, pressure and dissolved electrolyte [42-43]. Formation of gypsum takes place in through a complex multi-step processes [43]. It can form in evaporite beds during evaporation of water near lake and sea water. Gypsum can form diagenetically by oxidation. It can form in association with sedimentary rocks as oxidation of existing sediments leading to gypsum formation and replaces the other minerals in the sediments [44]. It can also occur in hot spring from volcanic vapors as well as in sulfate solutions in veins [19]. Hydrothermal anhydrite in veins is commonly hydrated to gypsum by groundwater in near-surface exposures. Gypsum can also occur in a flower-like form with embedded sand grains called desert rose in arid areas [45].

### 3. Materials and methods


The five gypsum samples are collected by one of us (JKP) in person from different geographic locations and environments of formation. They are shown in Figure 1 and include following varieties: 1: "Desert Rose" Gypsum, Sahara Desert, Morocco (MDRG). 2: Acicular Gypsum, Morocco (MAG); 3: Platy Gypsum, Kachchh, Gujarat (KG); 4 Flaky Gypsum, Rajpura-Dariba Cu-Pb-Zn Mines, Rajasthan (RDG1); 5: Bladed Gypsum, Rajpura-Dariba Cu-Pb-Zn Mines, Rajasthan (RDG2). The SEM images of all five samples are shown in Figure 2.

#### 3.1 Sample description

Desert Rose sample (MDRG; Sahara, Morocco) comprise 5 rose-like formations of platy crystals. They form an array of circular nearly flat plates, where plates are having sharper edges and relatively thick center. The diameter of biggest plate is about 5 cm and smallest is around 1.5 cm. The color is pinkish brown similar to sand (5YR 4/6; Munsell Color Chart). The sand encrustation on the top surface is about a few mm thick [45].



**Figure 1.** Megascopic specimens used in the present study and they include (a) Desert Rose Gypsum, Sahara Desert, Morocco (MDRG). (b) Acicular Gypsum, Morocco (MAG)., (c) Gypsum, Kachchh, Gujarat, India (KG). (d) Gypsum, Rajpura-Dariba Cu-Pb-Zn Mine, Rajasthan, India (RDG1). (e) Gypsum, Rajpura-Dariba Cu-Pb-Zn Mine, Rajasthan, India (RDG2).



**Figure 2.** Scanning electron microscope (SEM) images of the samples used in the present study which include: 1: Desert Rose Gypsum, Sahara Desert, Morocco (MDRG), Gypsum, Gy, and quartz, Qtz, are identified. 2: Acicular Gypsum, Morocco (MAG). 3: Gypsum, Kachchh, Gujarat, India (KG). 4: Gypsum, Rajpura-Dariba Cu-Pb-Zn Mine, Rajasthan, India (RDG1). 5: Gypsum, Rajpura-Dariba Cu-Pb-Zn Mine, Rajasthan, India (RDG2).



Acicular gypsum (MAG; Morocco) constitutes column-like aggregate of needle-shaped gypsum crystals. The length of the gypsum is around 24 cm and width around 3 cm. The color of the gypsum is white (5GY 9/2; Munsell Color Chart). Visibly the sample does not contain any impurity [46].

Kachchh gypsum (KG; Gujarat, India) is light brown color (7.5 YR 7/8; Munsell Color Chart), contains randomly arranged sub-millimeter-thin flakes/plates. The effect of surficial weathering in the sample is evident. The length of the sample is about 10 cm, and its width is 2 cm [48-51].

Rajpura- Dariba gypsum 1 (RDG1; Dariba Mines, Rajasthan) is transparent, colorless. The sample is having sub-millimeter platy layers. The length of the sample is 11.7 cm, with width of 6 cm [52].

Rajpura-Dariba gypsum 2 (RDG2; Dariba Mines, Rajasthan) is off-white in color (5 Y 9/2; Munsell Color Chart). The sample is having blade like petal having no fixed orientation and the blades are very thin (around 0.5 cm). The sample length and width are 12.3 and 2.5 cm, respectively [52].

### 3.2 Powder X-ray diffraction (XRD)

Powdered gypsum samples were characterized by X-ray Diffractometer (Philips PanAnalytical) using CuK $\alpha$  radiation and Ni filter. The scans were made between 10 to 75° 2 $\theta$  at a speed of 2° per minute, divergence slit = 1°, anti-scatter slit = 1°, receiving slit = 0.30 mm, generator voltage of 40 kV and tube current of 30 mA.

### 3.3 Experimental Set-up

The experimental setup consists of a frequency-doubled (532 nm) Nd:YAG laser (Continuum Surelite III-10). The plasma is created using a 15 mJ laser pulse having a pulse width of 4 ns with 10 pulses per second of repetition rate. A convex lens of focal length 15 cm is used to focus the laser on the sample to create hot plasma. The light emitted by the plasma at the sample surface was collected by a set of collection optics (CC52 collimator, Andor Technology, USA) and focused into the optical fibre bundle. The optical fiber bundle delivered the light to the Mechelle spectrograph (ME5000, Andor Technology, USA) equipped with an ICCD camera (iStar 334, Andor Technology, USA). The spectral signatures of atoms and ions from the plasma are collected after a short interval of time to avoid the background continuum. Therefore, the gate delay and gate width are 0.7  $\mu$ s and 4  $\mu$ s, respectively. A detailed analysis of spectral lines uses Andor SOLIS software and NIST atomic spectroscopic database [53].

### 3.4 Multivariate analysis

Multivariate analysis is a tool that can be used to maximize the information extracted from a huge number of datasets, here in the present case spectral data of LIBS spectra of the samples. Single spectra of LIBS can contain more than 8000 data points thus it is very difficult to distinguish the spectral variables (wavelength, intensity) of the samples with chemical heterogeneity [54-55].


Multivariate analyses techniques such as PCA and PSLR are used in the present study. PCA is an unsupervised method that reduces the dimensionality (number of features) within a dataset while still retaining as much information as possible. It is an orthogonal transformation that reduces the spectral data by projecting them into lower dimensions called principal components [56]. For the computation of the first principal component it is the linear combination of variables on which the most data variation can be projected. And for succeeding component it is perpendicular to the

previous principal component so that it contains maximum variation. Partial Least Squares Regression (PLSR) is one of the multivariate methods used in the interpretation of spectra (independent variables) to predict concentrations (dependent variables) of element in the sample. For making the regression model the concentration of Calcium and sulfur is taken from the analysis of Electron Microprobe Analyzer [57].

## 4. Results and discussion

### 4.1 X-ray diffraction analysis


X-ray diffraction studies on powdered samples reveal the presence of gypsum and anhydrite in all samples, illustrated in Figure 3, having a prominent d-spacing of 7.66 Å, 4.27 Å, 3.82 Å, 3.08 Å and 2.69 Å 1.82 Å, respectively. Presence of Jarosite (d-values: 5.59 Å and 3.72 Å) is noticed in gypsum from Matanumadh, Kachchh whereas in Desert Rose gypsum (Morocco) sample, quartz is observed with characteristic d-values of 3.36 Å (I/I<sub>0</sub>: 59.42) and 1.38 Å (I/I<sub>0</sub>: 5.59).



**Figure 3.** Powder XRD patterns of different gypsum samples showing the presence of gypsum (Gy; d=7.66, 4.27, 3.82 and 3.08Å), anhydrite (Anh; d=2.69, 2.46 and 2.29Å), quartz (Qtz; d=3.36Å) and jarosite (Jt; 5.59 and 3.72Å).


### 4.2 LIBS analysis

172 The LIBS spectra of different gypsum samples are recorded in the wavelength range between 200  
173 and 800 nm. Figure 4 illustrates the results.




**Figure 4.** Typical LIBS spectra (250–600 nm) of different gypsum samples analyzed during the present study stacked to show the relative intensity variations amongst the samples.


174 Differentiation of LIBS spectra of similar types of samples is difficult by visual inspection, but a  
175 careful analysis of the spectra reveals differences in the intensity of the gypsum samples because of  
176 their variable elemental concentrations. The wavelengths of different spectral lines present in the  
177 LIBS spectra of all five gypsum samples have been identified using National Institute of Standards  
178 and Technology (NIST) atomic spectroscopic database [53]. Figures 5 to 8 display details of the  
179 spectra together with the identified elements.




**Figure 5.** Spectral regions of recorded Al, Mg, Ca, and Fe data for the gypsum samples.



**Figure 6.** Spectral regions of recorded Mn, Sr, Ca, and Ba atomic and CaOH molecular data for the gypsum samples.



**Figure 7.** Spectral regions of recorded H, K, and O atomic data for the gypsum samples.



**Figure 8.** Spectral regions of recorded H, K, and O atomic data for the gypsum samples.



Figures 5 to 8 indicate that the intensities of spectral lines of the elements are different in the LIBS spectra of gypsums formed in different conditions/environment. Thus, with the information of elemental analysis or composition of gypsum one can predict the environment and the process of formation which cannot be achieved by their physical appearance. Persistent spectral lines along with the other weak spectral lines of calcium are observed in all five samples. Similarly, in the observation of Forni *et al.* [58, 59], a molecular band near 606 nm probably due to presence of CaOH is also observed in the LIBS spectra of all five gypsum samples.

If the concentration of any element in the sample is very high, its molecular bands must be present in the laser induced plasma of the sample [60]. The presence of sulfur lines (542.8 nm, 543.2 nm, 545.3 nm) in the LIBS spectra of gypsum also confirms its presence, as indicated in Figure 8. Similarly, the presence of carbon line at 247.8 nm and oxygen at 777.1 nm indicates the presence of carbonates in the present study. Strong spectral lines of magnesium at wavelengths 279.5 nm, 285.2 nm, 383.8 nm etc. are present in four gypsum samples except in acicular sample (MAG). Magnesium occurs in the defect sites of the gypsum lattice since they form large soluble complexes [61]. The intensity of spectral line of magnesium varies depending upon its concentration in the sample. Presence of prominent line of Si in MDRG is due to the presence of quartz grains from the sand which bonds with moisture on the top of gypsum plates. Ca lines are prominent in all gypsum samples except MDRG as the analyzed spots are dominantly masked by quartz grains. Presence of spectral lines of Fe in KG and MDRG are due to jarosite and quartz, respectively. Gypsum comprises abundant fluid phase, which is a mixture of various elements K, Ti, Mn and Sr in minor to trace amounts. Their concentration in gypsum is used in stratigraphic correlation, paleosalinity estimation and to decipher the origin of brine and diagenesis [62, 63]. The presence of spectral lines of Mn and Fe in KG indicates the terrigenous deposit mainly siliciclastic sediments or clay minerals into the sedimentary basin. Sr lines are found in RDG2 as Sr can replace Ca in the gypsum lattice [64]. Presence of strontium in gypsum can take place in many ways like diagenetic process, bacterial sulphate reduction or by dissolution or recrystallization. Abundance and scarcity of Sr content in gypsum indicate the episodic salinity fluctuation of the basin and this also reveals the paleohydrology of the basin [65]. Similarly, Presence of spectral line of Ba (455.4 nm and 614.1 nm) may be correlated as a possible tracer of fluid circulation in hydrothermal water [66]. The wavelength of the spectral lines observed in the LIBS spectra of all five gypsums are summarized in Tables 1 and 2. The same calcium lines are identified in all samples and the wavelengths are listed below Table 2.

Hydration level in gypsum depends upon the hydrogen and oxygen content, calcium sulfate either occurs as non-hydrated (anhydrite  $\text{CaSO}_4$ ) or hydrated phase (bassinite:  $\text{CaSO}_4 \cdot 1/2\text{H}_2\text{O}$  and gypsum:  $\text{CaSO}_4 \cdot 2\text{H}_2\text{O}$ ). H alpha emission line at 656.5 nm is sensitive to hydrogen content which can differentiate hydration levels of calcium sulfates in the samples. Similarly, the emission line of oxygen at 777.1 nm is sensitive to correlate the hydration level in the sample [67]. Spectral lines of both elements are detected but respective intensities of the spectral lines of hydrogen and oxygen in samples RDG-1 are very low. Thus, their concentration in RDG1 is also low which establishes the calcium sulfates observed may be bassanite.

**Table 1.** Spectral wavelengths of atoms observed in laser induced breakdown spectroscopic spectra of different gypsum samples: Hydrogen, nitrogen, oxygen, sodium, magnesium, aluminium, and silicon.

Element	Acicular Gypsum	Desert Rose	Kachchh Gypsum	RajpuraDariba Gypsum 1	RajpuraDariba Gypsum 2
H (1)	656.3 (I)	656.3 (I)	656.3 (I)	656.3 (I)	656.3 (I)
N (7)	744.2(I), 746.8(I), 868.3(I)	744.2(I), 746.8(I), 868.3(I)	744.2(I), 746.8(I), 868.3(I)	744.2(I), 746.8(I), 868.3(I)	744.2(I), 746.8(I), 868.3(I)
O (8)	777.4(I), 844.6(I), 926.6(I)	777.4(I), 844.6(I), 926.6(I)	777.4(I), 844.6(I), 926.6(I)	777.4(I), 844.6(I), 926.6(I)	777.4(I), 844.6(I), 926.6(I)
Na (11)	588.9(I), 589.5(I)	588.9(I), 589.5(I)	588.9(I), 589.5(I)	588.9(I), 589.5(I)	588.9(I), 589.5(I)
Mg (12)	-	279.0(II), 279.5(II), 280.2(II), 285.2(I),	279.0(II), 279.5(II), 280.2(II), 285.2(I),	279.0(II), 279.5(II), 280.2(II), 285.2(I),	279.0(II), 279.5(II), 280.2(II), 285.2(I),
Al (13)	-	308.2(I), 309.2(I), 394.3(I), 396.1(I)	308.2(I), 309.2(I), 394.3(I), 396.1(I)	-	-
Si (14)	-	220.7(I),221.0(I), 221.6(I),250.6(I), 251.4(I),251.6(I), 251.9(I), 252.8(I),288.1(I), 390.5(I), 413.1(II),557.6(II), 634.7(II), 716.5(I),741.5(I)	220.7(I),221.0(I), 221.6(I),250.6(I), 251.4(I),251.6(I), 251.9(I), 252.8(I),288.1(I), ,390.5(I) 413.1(II),557.6(II), 634.7(II)	252.8(I),288.1(I)	-

222

223

224

225

226


**Table 2:** Spectral wavelengths of atoms observed in laser induced breakdown spectroscopic spectra of different gypsum samples: Sulfur, potassium, titanium, manganese, iron, strontium, barium.

Element	Acicular Gypsum	Desert Rose	Kachchh Gypsum	RajpuraDariba Gypsum 1	RajpuraDariba Gypsum 2
S (16)	542.8(I), 543.2(I), 545.3(I)	542.8(I), 543.2(I), 545.3(I)	542.8(I), 543.2(I), 545.3(I)	542.8(I), 543.2(I), 545.3(I)	542.8(I), 543.2(I), 545.3(I)
K (19)	766.4(I), 769.8(I)	766.4(I), 769.8(I)	766.4(I), 769.8(I)	766.4(I), 769.8(I)	766.4(I), 769.8(I)
Ti (22)	-	-	334.9(II), 336.1(II)	-	-
Mn (25)	-	-	257.6(II), 403.0(I)	-	-
Fe (26)	-	238.2(II),239.5(II), 240.4(II),249.3(II), 252.3(I), 259.9(II),258.5(II), 259.8(II),261.1(II) 263.1(II),271.9(I), 273.9(II) 274.9(II),275.5(II), 344.0(I), 358.1(I), 373.7(I)	238.2(II),239.5(II), 240.4(II),249.3(II), 252.3(I), 259.9(II),258.5(II), 259.8(II),261.1(II) 263.1(II),271.9(I), 273.9(II), 274.9(II),275.5(II), 344.0(I)	-	-
Sr (38)	-	-	-	-	407 (I)
Ba (56)	-	-	-	-	455.4(II), 493.3(II)


The observed calcium atomic lines, Ca (20), are the same for all samples. The wavelengths of Ca I and Ca II are: 315.8(II), 317.9(II), 370.6(II), 373.6(II), 393.2(II), 396.7(II), 370.5(II), 422.6(I), 430.2(I), 442.5(I), 443.9(I), 445.4(I), 501.9(II), 518.8(I), 558.7(I), 610.2(I), 612.2(I), 616.2(I), 645.6(II), 646.2(I), 649.5(I), 647.2(I), 720.1(I) nm.

4.3 Principal component analysis (PCA)

In the present study PCA is used to analyze the LIBS spectra of gypsum samples. The spectra obtained from different gypsum samples are arranged in the form of matrix which is made up with variables having 8848 features. The Unscrambler-X software (CAMO Software India Pvt. Ltd.) is used to solve the data matrix to generate PCA plot. Figure 9 illustrates that the combination of PC1 and PC2 is 92% which is very close to 100% that explain most of the variance present in the dataset. Five replicate spectra of each gypsum sample are classified in five different groups using PCA. These groups indicate variation in elemental concentrations in gypsums. Replicates of each gypsum sample are very close to each other forming a clustering in the score plot. Clearly, the samples of each class tend to cluster and in almost all cases they are well-separated from the other classes. The PC1 classifies gypsum samples into two groups (see Fig. 9). The first group contains MAG, RDG1 and RDG2 having positive correlation. The second groups KG and MRD show negative correlation. MRD and KG are different from MAG, RDG1 and RDG2 as they contain Si, Fe, Al, Mg and Ti. The above observation is also consistent with the loading plot, as illustrated in Figure 10. A positive correlation observed in the loading plot corresponds to the emission lines of Ca which is greater in RDG1, RDG2 and MAG while negative correlation is noted in Si, Al, Ti, Mg and Fe. In the second group (MRD and KG), Fe and Mn are present in only KG. The Si content is very high in MRD in comparison to KG. This is the main reason leading to the classification of these samples into two different clusters based on PC2. Further the cluster of KG is comparatively spread out suggesting sample heterogeneity. Similarly, in the first group PC2 is classified into two different categories (RDG1 and MAG) due to the less hydration in RDG1. The clustering of RDG2 and MAG are very close to each other as they contain meager impurities. But RDG2 and MAG also form two groups because of Sr and Ba present in RDG2.



**Figure 9.** PCA plot of the samples. The labels indicate the gypsum K= Kachchch; M= Acicular; D1=Rajpura-Dariba 1; D2= Rajpura Dariba 2; R= Desert Rose.



**Figure 10.** Loading plot for the principal component analysis of the LIBS spectral data.

#### 4.4 Partial least-square regression (PLSR)

PLSR is used to construct a model for prediction of the elements. In this experiment it is used to predict the concentration of main constituent of gypsum i.e. calcium and sulfur. The wavelength region having maximum number of spectral lines of corresponding elements has been included for building the calibration model. PLSR model is employed to construct the calibration model for the gypsum samples whose concentration was obtained from EPMA analysis. The graphical results are called Predicated vs. Reference plots, see Figure 11. Similarly, PLSR plots are generated for sulfur.

The performance of the model depends upon the two parameters (i) coefficient of determination ( $R^2$ ) and (ii) root mean square error (RMSE). The model is said to be strongly correlated when the RMSE and  $R^2$  values tend to zero and one, respectively in the plots. In this study model  $R^2$  values are nearly 0.96 and RMSE value is 0.02 and the calibration (blue line) and validation (red line) deviate very little from each other. Thus, the model adopted in the present study is strongly correlated. Concentration of all calcium and sulfur is determined using above PLSR model and the values are tabulated in Table 3.





273

274

274

$$^* \text{Accuracy} = \frac{\text{PLS predicted concentration}}{\text{Reference Concentration}} \times 100$$

## 5. Conclusions

In this work, terrestrial gypsum samples from varied locations and textural types have been analyzed by laser-induced breakdown spectroscopy (LIBS). Petrographic, sub-microscopic and X-ray diffraction (XRD) investigations confirm the gypsums. The spectral lines and their respective intensities for various light (H, C, N and O) and trace elements (Ti, Mn, Fe, Ba and Sr) present in the gypsum samples are in close agreement with the results reported by conventional analytical techniques (INAA, XRF and LA-ICP-MS), moreover, minor variations could also be gauged. The texturally varied gypsum samples from different locations bearing information on specific geological environments can be compared using LIBS data. The study highlights the LIBS sensitivity to the water content for hydration levels of calcium sulfates. One can distinguish rapidly between anhydrite ( $\text{CaSO}_4$ ), bassanite,  $\text{CaSO}_4 \cdot 1/2\text{H}_2\text{O}$ , and gypsum using a multi-elemental analysis technique. The data collected with LIBS in this experimental study can be used as mineral references for detection of similar formation of gypsum wherever it occurs. KG gypsum contains Mn and Fe in appreciable amount indicating the availability of terrigenous sediments in the source basin.

PCA coupled LIBS shows the similarity and differences between various gypsum samples in terms of their elemental concentrations because the samples are very well grouped into distinct clusters. The PSLR method is utilized for the prediction of the concentrations of calcium and sulfur, subsequently, this information can be used to predict the concentration of same elements in any other gypsum samples. The texturally different gypsum samples that originate from different locations can be compared, and the gypsum samples contain information on specific geological environments.

After the discovery of gypsum by the ChemCam rover, study of Martian sediments (either *in situ* or in returned samples) is expected to become possible in the future. The present study demonstrated the potential application of LIBS for rapid characterization of gypsum samples which can be extended to the study of gypsum samples in extraterrestrial environments.

**Acknowledgement:** One of the authors, Abhishek K. Rai, expresses thanks to the UGC (CRET) of India for financial assistance of his doctoral research at the University of Allahabad, UP, India.

**Author Contributions:** The samples are provided by Jayanta K. Pati, Abhishek K. Rai conducted the experimental investigations, and all authors contributed to the writing of this article.

**Conflicts of Interest:** The authors declare no conflict of interest

## References

- [1] Gaft M; Sapir-Sofer I; Modiano H; Stana R. Laser induced breakdown spectroscopy for bulk minerals online analyses. *Spectrochim. Acta Part B: At. Spectrosc.* **2007**, *62*, 1496–1503; doi.org/10.1016/j.sab.2007.10.041.
- [2] Díaz Pace D.M.; Gabriele N.A.; Garcimuño M.; D'Angelo C.A.; Bertuccelli G.; Bertuccelli D. Analysis of Minerals and Rocks by Laser-Induced Breakdown Spectroscopy. *Spectrosc. Lett.* **2011**, *44*, 399–411; dx.doi.org/10.1080/00387010.2011.573048.

- [3] Wiens, R.C.; Maurice, S.; Barraclough, B.; Saccoccio, M.; Barkley, W.C.; Bell III, J.F.; Bender, S.; Bernardin, J.; Blaney, D.; Blank, J.; Bouyé, M.; Bridges, N.; Bultman, N.; Caïs, P.; Clanton, R.C.; Clark, B.; Clegg, S.; Cousin, A.; Cremers, D.; Cros, A.; DeFlores, L.; Delapp, D.; Dingler, R.; D'Uston, C.; Dyar, M.D.; Elliott, T.; Enemark, D.; Fabre, C.; Flores, M.; Forni, O.; Gasnault, O.; Hale, T.; Hays, C.; Herkenhoff, K.; Kan, E.; Kirkland, L.; Kouach, D.; Landis, D.; Langevin, Y.; Lanza, N.; LaRocca, F.; Lasue, J.; Latino, J.; Limonadi, D.; Lindensmith, C.; Little, C.; Mangold, N.; Manhes, G.; Mauchien, P.; McKay, C.; Miller, E.; Mooney, J.; Morris, R.V.; Morrison, L.; Nelson, T.; Newsom, H.; Ollila, A.; Ott, M.; Pares, L.; Perez, R.; Poitrasson, F.; Provost, C.; Reiter, J.W.; Roberts, T.; Romero, F.; Sautter, V.; Salazar, S.; Simmonds, J.J.; Stiglich, R.; Storms, S.; Striebig, N.; Thocaven, J.J.; Trujillo, T.; Ulibarri, M.; Vaniman, D.; Warner, N.; Waterbury, R.; Whitaker, R.; Witt, J.; Wong-Swanson, B. The ChemCam Instrument Suite on the Mars Science Laboratory (MSL) Rover: Body Unit and Combined System Tests. *Space Sci. Rev.* **2012**, *170*, 67–227; doi.org/10.1007/s11214-012-9902-4
- [4] Grotzinger, J.P.; Arvidson, R.E.; Bell, J.; Calvin, W.; Clark, B.C.; Fike, D.A.; Golombek, M.; Greeley, R.; Haldemann, A.; Herkenhoff, K.E.; Jolliff, B.L.; Knoll, A.H.; Malin, M.; McLennan, S.M.; Parker, T.; Soderblom, L.; Sohl-Dickstein, J.N.; Squyres, S.W.; Tosca, N.J.; Watters, W.A. Stratigraphy and sedimentology of a dry to wet eolian depositional system, Burns formation, Meridiani Planum, Mars. *Earth Planet. Sci. Lett.* **2005**, *240*, 11–72; doi.org/10.1016/j.epsl.2005.09.039.
- [5] Knoll, A.H.; Squyres, S. W.; Arvidson, R. E.; Bell, J. F., III; Calef, F., III; Clark, B. C.; Cohen, B. A.; Crumpler, L. A.; de Souza, P. A. Jr.; Farrand, W. H.; Gellert, R.; Grant, J.; Hurowitz, J. A.; Herkenhoff, K. E.; Johnson, J. R.; Jolliff, B. L.; Li, R.; McLennan, S. M.; Ming, D. W.; Mittlefehldt, D. W.; Parker, T. J.; Paulsen, G.; Rice, M. S.; Ruff, S. W.; Schröder, C.; Yen, A. S.; Zacny, K. Ancient Impact and Aqueous Processes at Endeavour Crater, Mars. *Sci.* **2012**, *336*, 570–576; doi.org/10.1126/science.1220476.
- [6] Squyres, S.W.; Knoll, A.H. Sedimentary rocks at Meridiani Planum: Origin, diagenesis, and implications for life on Mars. *Earth Planet. Sci. Lett.* **2005**, *240*, 1–10; doi.org/10.1016/j.epsl.2005.09.038
- [7] Elwood Madden, M.E.; Bodnar, R.J.; Rimstidt, J.D. Jarosite as an indicator of water-limited chemical weathering on Mars. *Nat.* **2004**, *431*, 821–823; doi.org/10.1038/nature02971.
- [8] M. Nachon, M.; Clegg, S.M.; Mangold, N.; Schröder, S.; Kah, L.C.; Dromart, G.; Ollila, A.; Johnson, J.R.; Oehler, D.C.; Bridges, J.C.; Le Mouélic, S.; Forni, O.; Wiens, R.C.; Anderson, R.B.; Blaney, D.L.; Bell III, J.F.; Clark, B.; Cousin, A.; Dyar, M.D.; Ehlmann, B.; Fabre, C.; Gasnault, O.; Grotzinger, J.; Lasue, J.; Lewin, E.; Léveillé, R.; McLennan, S.; Maurice, S.; Meslin, P.-Y.; Rapin, W.; Rice, M.; Squyres, S.W.; Stack, K.; Sumner, D.Y.; Vaniman, D.; Wellington, D. Calcium sulfate vein observations at Yellowknife Bay using ChemCam on the Curiosity Rover. American Geophysics Union, Fall Meeting, **2013** Abstract P23C-1797.
- [9] Crumpler, L.S.; Arvidson, R.E.; Bell, J.; Clark, B.C.; Cohen, B.A.; Farrand, W.H.; Gellert, R.; Golombek, M.; Grant, J.A.; Guinness, E.; Herkenhoff, K.E.; Johnson, J.R.; Jolliff, B.; Ming, D.W.; Mittlefehldt, D.W.; Parker, T.; Rice, J.W. Jr.; Squyres, S.W.; Sullivan, R.; Yen, A.S. Context of ancient aqueous environments on Mars from in situ geologic mapping at Endeavour Crater. *J. Geophys. Res.* **2015**, *120*, 538–569; doi.org/10.1002/2014JE004699.

- [10] Anderson, D.E.; Ehlmann, B.L.; Forni, O.; Clegg, S.M.; Cousin, A.; Thomas, N.H.; Lasue, J.; Delapp, D.M.; McInroy, R.E.; Gasnault, O.; Dyar, M.D.; Schröder, S.; Maurice, S.; Wiens, R. C. Characterization of LIBS emission lines for the identification of chlorides, carbonates, and sulfates in salt/basalt mixtures for the application to MSL ChemCam data. *J. Geophys. Res.* **2017**, *122*, 744–770; doi.org/10.1002/2016JE005164.
- [11] Ziolkowski, L.A.; Mykytczuk, N.C.S.; Omelon, C.R.; Johnson, H.; Whyte, L.G.; Slater, G.F. Arctic Gypsum Endoliths: a biogeochemical characterization of a viable and active microbial community. *Biogeosciences Discuss.* **2013**, *10*, 2269 – 2304; doi:10.5194/bgd-10-2269-2013.
- [12] Bonny, S.; Jones, B. Microbes and mineral precipitation, Miette Hot Springs, Jasper National Park, Alberta, Canada. *Can. J. Earth Sci.* **2003**, *40*, 1483–1500; doi.org/10.1139/e03-060.
- [13] Parnell, J.; Lee, P.; Cockell, C.S.; Osinski, G.R. Microbial colonization in impact-generated hydrothermal sulphate deposits, Haughton impact structure, and implications for sulphates on Mars. *Int. J. Astrobiol.* **2004**, *3*, 247-256; doi:10.1017/s1473550404001995.
- [14] Tang, M.; Ehreiser, A.; Li, Y.-L. Gypsum in modern Kamchatka volcanic hot springs and the Lower Cambrian black shale: Applied to the microbial-mediated precipitation of sulfates on Mars. *Am. Mineral.* **2014**, *99*, 2126-2137; doi:10.2138/am-2014-4754.
- [15] Rasuk, M.C.; Kurth, D.; Flores, M.R.; Contreras, M.; Novoa, F.; Poiré, D.; Farias, M.E. Microbial Characterization of Microbial Ecosystems Associated to Evaporites Domes of Gypsum in Salar de Llamara in Atacama Desert. *Microb. Ecol.* **2014**, *68*, 483-494. doi:10.1007/s00248-014-0431-4.
- [16] Pati, J. K.; Reimold, W. U. Impact cratering -fundamental process in geoscience and planetary science. *J. Earth Syst. Sci.* **2007**, *116*, 81-98; doi:10.1007/s12040-007-0009-3.
- [17] Aubrey, A.; Cleaves, H.J.; Chalmers, J.H.; Skelley, A.M.; Mathies, R.A.; Grunthaner, F.J.; Ehrenfreund, P.; Bada, J. L. Sulfate minerals and organic compounds on Mars. *Geol.* **2006**, *34*, 357; doi:10.1130/g22316.1.
- [18] Benison, K.C.; Karmanocky, F.J. Could microorganisms be preserved in Mars gypsum? Insights from terrestrial examples. *Geol.* **2014**, *42*, 615-618; doi:10.1130/g35542.1.
- [19] Ritsema, C.J.; Groenenberg, J.E., Pyrite Oxidation, Carbonate Weathering, and Gypsum Formation in a Drained Potential Acid Sulfate Soil. *Soil Sci Soc Am J.* **1993**, *57*, 968; doi:10.2136/sssaj1993.03615995005700040015x.
- [20] Langevin, Y.; Poulet, F.; Bibring, J.-P.; Gondet, B. Sulfates in the North Polar Region of Mars Detected by OMEGA/Mars Express. *Sci.* **2005**, *307*, 1584-1586; doi.org/10.1126/science.1109091.
- [21] Massé, M.; Le Mouélic, S.; Bourgeois, O.; Combe, J.-P.; Le Deit, L.; Sotin, C.; Bibring, J.-P.; Gondet, B.; Langevin, Y. Mineralogical composition, structure, morphology, and geological history of Aram Chaos crater fill on Mars derived from OMEGA Mars Express data. *J. Geophys. Res.* **2008**, *113*, E12; doi:10.1029/2008je003131.
- [22] Milliken, R. E.; Grotzinger, J. P.; Thomson, B. J. Paleoclimate of Mars as captured by the stratigraphic record in Gale Crater. *Geophys. Res. Lett.* **2010**, *37*(4); doi:10.1029/2009gl041870. 1
- [23] Bibring, J.-P. Mars Surface Diversity as Revealed by the OMEGA/Mars Express Observations. *Sci.* **2005**, *307*, 1576-1581; doi:10.1126/science.1108806.
- [24] Tanaka, K.L. Mars' north polar gypsum: Possible origin related to early Amazonian magmatism at Alba Patera and aeolian mining. In Fourth International Conference on Mars Polar Science and Exploration, no. 8024, Davos, **2006** Switzerland.

- [25] Fishbaugh, K. E.; Poulet, F.; Chevrier, V.; Langevin, Y.; Bibring, J.-P. On the origin of gypsum in the Mars north polar region. *J. Geophys. Res.* **2007**, *112*, E7; doi:10.1029/2006je002862.
- [26] Squyres, S.W.; Aharonson O.; Clark B.C.; Cohen B.A.; Crumpler L.; de Souza P.A.; Farrand W.H.; Gellert R.; Grant J.; Grotzinger J.P.; Haldemann A.F.; Johnson J.R.; Klingelhöfer G.; Lewis K.W.; Li R.; McCoy T.; McEwen A.S.; McSween H.Y.; Ming D.W.; Moore J.M.; Morris R.V.; Parker T.J.; Rice J.W. Jr.; Ruff S.; Schmidt M.; Schröder C.; Soderblom L.A.; Yen A. Pyroclastic Activity at Home Plate in Gusev Crater, Mars, *Sci.* **2007**, *316*, 738–742; doi:10.1126/science.1139045
- [27] Szyrkiewicz, A.; Ewing, R.C.; Moore, C.H.; Glamoclija, M.; Bustos, D.; Pratt, L. M. (2010). Origin of terrestrial gypsum dunes-Implications for Martian gypsum-rich dunes of Olympia Undae. *Geomorph.* **2010**, *121*, 69–83; doi:10.1016/j.geomorph.2009.02.017
- [28] Dehouck, E.; Chevrier, V.; Gaudin, A.; Mangold, N.; Mathé, P.-E.; Rochette, P. Evaluating the role of sulfide-weathering in the formation of sulfates or carbonates on Mars. *Geochim. Cosmochim. Acta.* **2012**, *90*, 47–63; doi:10.1016/j.gca.2012.04.057.
- [29] Squyres, S.W.; Arvidson, R.E.; Bell J.F. III, Brückner J.; Cabrol N.A.; Calvin W.; Carr M.H.; Christensen P.R.; Clark B.C.; Crumpler L.; Marais D.J.; d'Uston C.; Economou T.; Farmer J.; Farrand W.; Folkner W.; Golombek M.; Gorevan S.; Grant J.A.; Greeley R.; Grotzinger J.; Haskin L.; Herkenhoff K.E.; Hviid S.; Johnson J.; Klingelhöfer G.; Knoll A.H.; Landis G.; Lemmon M.; Li R.; Madsen M.B.; Malin M.C.; McLennan S.M.; McSween H.Y.; Ming D.W.; Moersch J.; Morris R.V.; Parker T.; Rice J.W. Jr.; Richter L.; Rieder R.; Sims M.; Smith M.; Smith P.; Soderblom L.A.; Sullivan R.; Wänke H.; Wdowiak T.; Wolff M.; Yen A. The Opportunity Rover's Athena Science Investigation at Meridiani Planum, Mars, *Sci.* **2004**, *306*, 1698–1703; doi:10.1126/science.1106171.
- [30] Young, B.W.; Chan, M.A. Gypsum veins in Triassic Moenkopi mudrocks of southern Utah: Analogs to calcium sulfate veins on Mars, *J. Geophys. Res.* **2017**, *122*, 150–171; doi:10.1002/2016je005118.
- [31] Kasprzyk, A. Sedimentological and diagenetic patterns of anhydrite deposits in the Badenian evaporite basin of the Carpathian Foredeep, southern Poland. *Sed. Geol.* **2003**, *158*, 167–194; doi:10.1016/s0037-0738(02)00265-8.
- [32] Laxmiprasad, A.S.; Sridhar Raja, V.L.N.; Menon, S.; Goswami, A.; Rao, M.V.H.; Lohar, K. A. An in situ laser induced breakdown spectroscopy (LIBS) for Chandrayaan-2 rover: Ablation kinetics and emissivity estimations. *Adv. Space Res.* **2013**, *52*, 332–341; doi:10.1016/j.asr.2013.03.021.
- [33] Deer, W.A.; Howie, R.A.; Zussman, J. An Introduction to the Rock Forming Minerals, London: Longman. 469, **1966**.
- [34] Mandal, P.K.; Mandal, T.K. Anion water in gypsum ( $\text{CaSO}_4 \cdot 2\text{H}_2\text{O}$ ) and hemihydrate ( $\text{CaSO}_4 \cdot 1/2\text{H}_2\text{O}$ ), *Cem. Concr. Res.* **2002**, *32*, 313–316; doi:10.1016/s0008-8846(01)00675-5.
- [35] Heijnen, W.M.M.; Hartman, P. Structural morphology of gypsum ( $\text{CaSO}_4 \cdot 2\text{H}_2\text{O}$ ), brushite ( $\text{CaHPO}_4 \cdot 2\text{H}_2\text{O}$ ) and pharmacolite ( $\text{CaHAsO}_4 \cdot 2\text{H}_2\text{O}$ ). *J Cryst Growth.* **1991**, *108*, 290–300; doi:10.1016/0022-0248(91)90376-g.
- [36] Yen, A.S.; A.S.; Ming, D.W.; Vaniman, D.T.; Gellert, R.; Blake, D.F.; Morris, R.V.; Morrison, S.M.; Bristow, T.F.; Chipera, S.J.; Edgett, K.S.; Treiman, A.H.; Clark, B.C.; Downs, R.T.; Farmer, J.D.; Grotzinger, J.P.; Rampe, E.B.; Schmidt, M.E.; Sutter, B.; Thompson, L.M.; MSL Science



- Team. Multiple stages of aqueous alteration along fractures in mudstone and sandstone strata in Gale Crater, Mars. *Earth Planet Sci. Lett.* **2017**, *471*, 186–198; doi:10.1016/j.epsl.2017.04.033.
- [37] Liao, X.; Zhang, J.; Li, J.; Chigira, M.; Wu, X. Study on the chemical weathering of black shale in northern Guangxi, China. 10 Asian Regional Conference of IAEG, 2015.
- [38] Al-Youssef, M. Gypsum Crystals Formation and Habits, Dukhan Sabkha, Qatar. *Journal of Earth Science & Climatic Change*, **2015**, *6*, 321; doi:10.4172/2157-7617.1000321.
- [39] Kuhn, T.; Herzig, P.; Hannington, M.; Garbe-Schönberg, D.; Stoffers, P. Origin of fluids and anhydrite precipitation in the sediment-hosted Grimsey hydrothermal field north of Iceland. *Chem. Geol.* **2003**, *202*, 5–21; doi:10.1016/s0009-2541(03)00207-9.
- [40] Stawski, T. M.; van Driessche, A.E.S.; Ossorio, M.; Diego Rodriguez-Blanco, J.; Besselink, R.; Benning, L.G. Formation of calcium sulfate through the aggregation of sub-3 nanometre primary species. *Nat. Commun.* **2016**, *7*; doi:10.1038/ncomms11177.
- [41] Palacio, S.; Azorín, J.; Montserrat-Martí, G.; Ferrio, J. P. The crystallization water of gypsum rocks is a relevant water source for plants. *Nat. Commun.* **2014**, *5*; doi:10.1038/ncomms5660.
- [42] Joanna, J. Crystallization, Alternation and Recrystallization of Sulphates. *Advances in Crystallization Processes*, **2012**.
- [43] Freyer, D.; Voigt, W. Crystallization and Phase Stability of CaSO<sub>4</sub> and CaSO<sub>4</sub> - Based Salts. *Monatshefte für Chemie*. **2003**, *134*, 693–719; doi:10.1007/s00706-003-0590-3.
- [44] Pirlet, H.; Wehrmann, L.M.; Brunner, B.; Frank, N.; DeWanckele, J.; Van Rooij, D.; Foubert, A.; Swennen, R.; Naudts, L.; Boone, M.; Cnudde, V.; Henriët, J.-P. Diagenetic formation of gypsum and dolomite in a cold-water coral mound in the Porcupine Seabight, off Ireland. *Sed.* **2009**, *57*, 786–805; doi:10.1111/j.1365-3091.2009.01119.x.
- [45] Whitlock, H.P. Desert Roses. *Natural History (New York)*, **1930**, 421–425.
- [46] Macfadyen, W.A. Sandy Gypsum Crystals from Berbera, British Somaliland. *Geol. Mag.* **1950**, *87*(6), 409–420.
- [47] Eswaran, H.; Zi-Tong, G. Properties, genesis, classification, and distribution of soils with gypsum. In Nettleton, W.D. (ed.), *Occurrence, Characteristics, and Genesis of Carbonate, Gypsum, and Silica Accumulations in Soils*. Soil Science Society of America Special Publication **1991**, *26*, SSSA, Madison, 89–119.
- [48] Watson, A. Structure, chemistry and origins of gypsum crusts in southern Tunisia and the central Namib Desert. *Sed.* **1985**, *32*, 855–875; doi:10.1111/j.1365-3091.1985.tb00737.x.
- [49] Chitale, D.V. Natroalunite in a Laterite Profile over Deccan Trap Basalts at Matanumad, Kutch, India. *Clays Clay Miner.* **1987**, *35*, 196–202; doi:10.1346/ccmn.1987.0350305.
- [50] Greenberger, R.N.; Mustard, J.F.; Kumar, P.S.; Dyar, M.D.; Breves, E.A.; Sklute, E.C. Sklute, Low temperature aqueous alteration of basalt: Mineral assemblages of Deccan basalts and implications for Mars, *J. Geophys. Res. Planets.* **2012**, *117*, E00J12; doi:10.1029/2012je004127.
- [51] Bhattacharya, S.; Mitra, S.; Gupta, S.; Jain, N.; Chauhan, P.; Parthasarathy, G.; Ajai. Jarosite occurrence in the Deccan Volcanic Province of Kachchh, western India: Spectroscopic studies on a Martian analog locality. *J. Geophys. Res. Planets.* **2016**, *121*-3, 402–431; doi:10.1002/2015je004949.
- [52] Halder, S.K.; M. Deb, M. Geology and mineralization of Rajpura-Dariba Lead-Zinc belt, Rajasthan. International workshop proceedings, Delhi-Udaipur, India, **2001**, 177–187.

- [53] National Institute of Standards and Technology, Electronic database; <http://physics.nist.gov/PhysRefData/ASD/lines.form.html> (accessed June 11, 2019).
- [54] Rai, A.K.; Maurya, G.S.; Kumar, R.; Pathak, A.K.; Pati, J.K.; Rai, A.K. Analysis and Discrimination of Sedimentary, Metamorphic, and Igneous Rocks Using Laser-Induced Breakdown Spectroscopy. *J. Appl. Spectrosc.* **2017**, *83*, 1089-1095; doi:10.1007/s10812-017-0412-8.
- [55] Gottfried, J. L.; Harmon, R.S.; De Lucia, F.C.; Miziolek, A.W. Multivariate analysis of laser-induced breakdown spectroscopy chemical signatures for geomaterial classification. *Spectrochim. Acta Part B: At. Spectrosc.* **2009**, *64*, 1009-1019; doi:10.1016/j.sab.2009.07.005.
- [56] Rai, A.K.; Pati, J.K.; Kumar, R. Spectro-chemical study of moldavites from Ries impact structure (Germany) using LIBS. *Opt Laser Technol.* **2019**, *117*, 146-157; doi:10.1016/j.optlastec.2019.01.028.
- [57] Rai, A.K.; Singh, A.K.; Pati, J.K.; Gupta, S.; Chakarvorty, M.; Niyogi, A.; Pandey, A.; Dwivedi, M.M.; Kuldeep, Pandey, K.; Prakash, K.; Assessment of topsoil contamination in an urbanized interfluvial region of Indo-Gangetic Plains (IGP) using magnetic measurements and spectroscopic techniques. *Environ. Monit. Assess.* **2019**, *191*, 403; doi.org/10.1007/s10661-019-7525-x.
- [58] Forni, O.; Maurice, S.; Gasnault, O.; Wiens, R.C.; Cousin, A.; Clegg, S.M.; Sirven, J.-B.; Lasue, J. Independent component analysis classification of laser induced breakdown spectroscopy spectra. *Spectrochim. Acta Part B: At. Spectrosc.* **2013**, *86*, 31-41; doi:10.1016/j.sab.2013.05.003.
- [59] Forni, O.; Gaft, M.; Toplis, M.; Clegg, S.M.; Ollila, A.; Sautter, V.; Nachon, M.; Gasnault, O.; Mangold, N.; Maurice, S.; Meslin, P.-Y.; Schröder, S.; Wiens, R.C.; Blaney, D.; Cousin, A.; DeFlores, L.P.; Vaniman, D.T. First fluorine detection on Mars with ChemCam on-board MSL, 456<sup>th</sup> Lunar and Planetary Science Conference **2014**, paper 1328.pdf.
- [60] Cremers, D.A.; Radziemski, L.J. Handbook of Laser Induced Breakdown Spectroscopy, JohnWiley, Chichester, England, **2006**.
- [61] Rinaudo, C.; Franchini-Angela, M.; Boistelle, R. Curvature of gypsum crystals induced by growth in the presence of impurities. *Mineral. Mag.* **1989**, *53*, 479-482; doi:10.1180/minmag.1989.053.372.08.
- [62] Moore, G.W. Origin and chemical composition of evaporite deposits. U.S. Geological Survey Open-File Rep.174, **1960**.
- [63] Dean, W.E.; Tung, A.L. Trace and minor elements in anhydrite and halite, Supai Formation (Permian), East central Arizona. 4th International Symposium on Salt, Cleveland, Northern Ohio. *Geol. Soc.* **1974**, *1*, 75-285.
- [64] Kushnir, J. The coprecipitation of strontium, magnesium sodium, potassium and chloride ions with gypsum: A1n experiment study. *Geochim. et Cosmochim. Acta* **1980**, *44*, 1471-1482.
- [65] Kushnir, J. The composition and origin of brines during the Messinian desiccation event in the Mediterranean Basin as deduced from concentrations of ions coprecipitated with gypsum and anhydrite, *Chem. Geol.* **1982**, *35*, 333-350; doi:10.1016/0009-2541(82)90010-9
- [66] Warren, J. Evaporites: Their Evolution and Economics. Blackwell Science, Oxford, U. K. **1999**.
- [67] Rapin, W.; Meslin, P.-Y.; Maurice, S.; Vaniman, C.; Nachon, M.; Mangold, N.; Schröder, S.; Gasnault, O.; Forni, O.; Wiens, R.C.; Martínez, G.M.; Cousin, A.; Sautter, V.; Lasue, J.; Rampe, E.B.; Archeri, D. Hydration state of calcium sulfates in Gale crater, Mars: Identification of bassanite veins. *Earth Planet Sci Lett.* **2016**, *452*, 197-205.