

Overexpression of MiR482c in Tomato Induces Enhanced Susceptibility to Late Blight

Yu-Hui Hong¹, Jun Meng², Xiao-Li He¹, Yuan-Yuan Zhang¹, Yu-Shi Luan^{1*}

¹School of Bioengineering, Dalian University of Technology, Dalian 116024, China.

²School of Computer Science and Technology, Dalian University of Technology, Dalian 116024,

China.

*Corresponding author

Yu-Shi Luan: ysluan@dlut.edu.cn;

Abstract

Tomato is the highest-value fruit/vegetable crop worldwide. However, the quality and yield of tomatoes are severely affected by late blight. MicroRNA482s (miR482s) are involved in plant immune system. In this study, miR482c was transiently and stably overexpressed in tomatoes in transgenic plants to explore its mechanism in tomato resistance against late blight. Tomato in transgenic plants transiently overexpressed miR482c displayed larger lesion area than the control plants upon infection. Furthermore, compared with the WT tomato plants, the transgenic tomato plants stably overexpressing miR482c displayed decreased expression of target genes accompanied by lower POD, SOD, and PAL activity activities and higher MDA content, thereby leading to a decline in the ROS scavenging ability and aggravating the damage of lipid peroxidation product accumulation on the cell membrane, eventually enhancing plant susceptibility. This finding indicates that miR482c may act as a negative regulator in tomato resistance by regulating NBS-LRR expression levels and ROS levels.

Keywords: tomato; late blight; miRNA; overexpression; resistance

INTRODUCTION

Tomato (*Solanum lycopersicum*) is the highest-value fruit/vegetable crop worldwide with considerable significance for human nutrition and health [1, 2]. In the last five years, the average annual production of tomatoes was nearly 96 million tons according to data from the Food and Agriculture Organization of the United Nations (FAOSTAT) [3]. The yield and quality of tomatoes are seriously affected by the late blight in both pre- and postharvest stages [4]. It has been shown that an unprotected tomato crop (greenhouse, field) can be devastated within seven to ten days once infected with late blight [5]. The management of late blight is relatively difficult, as this disease destroys foliage and fruits throughout the whole growth and development period of tomato [5, 6]. Traditional strategies for plant resistance rely on the frequent use of antimicrobial, leading to the serious pollution of the environment. To date, genetic engineering resistant cultivars has become the most efficient and economical long-term approach to strengthen plant resistance to various forms of stress. Due to a long breeding cycle and rapid pathogen variation, it is difficult for traditional disease-resistant breeding approaches to meet the needs of agricultural production. Comprehending the molecular mechanism of plant-pathogen interactions and exploring disease resistance genes are key to enhancing plant resistance.

MicroRNAs (miRNAs), as a class of 22-24 nucleotide (nt) noncoding RNAs, play significant roles in most eukaryotes by regulating the posttranscription of target genes [7, 8]. MiRNAs have been shown to bind target mRNA molecules at complementary sites and thereby trigger target degradation or translational inhibition [9]. In recent years, mounting studies have reported that miRNA cascades influence the plant response to biotic stress [10]. For instance, using small RNA sequencing, Luan et al [11]. identified 70 significantly changed miRNAs in tomato in response to late blight. MiR393 in *Arabidopsis* was shown to enhance plant immune systems and contribute to plant defense against

bacteria [12]. MiR164a in rice was down-regulated, and its target transcription factor OsNAC60 was derepressed correspondingly to enhance plant defenses against rice blast [13]. MiR1916 in tomato was demonstrated to negatively regulate tomato resistance to late blight and gray mold [14]. In addition to acting as regulators under biotic stress, miRNAs also participate in the response to abiotic stress and plant development. For example, miR169c contributes to tomato tolerance to drought by regulating targeted three nuclear factor Y subunit genes and one resistance-associated gene [15]. MiR171b participates in organ development by downregulating several GRAS genes [16]. The direct binding of ripening inhibitor (RIN) to the promoter of miR172a in tomato suggested in fruit development and ripening there was a close correlation between RIN and miRNA expression [17].

Conserved miRNAs mostly exist as gene families in the genome, and more than 90% of the conserved miRNA gene families are multigene families [18]. The conservation of different member sequences in the plant miRNA family suggests a similar or synergistic function. For instance, activating miR166k-166h expression in rice enhanced plant resistance to rice blast and bakanae disease [19]. Tomato plants overexpressing miR172a and miR172b exhibited enhanced resistance to late blight by suppressing an AP2/ERF transcription factor [20]. Silencing miR2118b (corresponding to sly-miR482d in the miRBase) and miR482e conferred resistance to late blight to tomato [21]. In addition, several studies have shown that members of the same gene family may play different roles in the immune system of plants. For example, Csukasi et al [22]. documented that during the course of strawberry receptacle development, treatment with gibberellin down-regulated miR159a expression but that miR159b expression remained unchanged. Ma et al [23]. found via small RNA sequencing that miR156b and miR156e were significantly induced in two peanut eighth-generation recombinant inbred lines (RIL8) at 35 days after flowering, whereas the expression of miR156o was significantly reduced.

Moreover, in rice, overexpression of miR164c led to reduced germination rates [24], while miR164b negatively regulated drought resistance by cleaving NAC genes [25]. These studies suggested that members of miRNA families may act differently in response to diverse stresses in various plant species.

MiR482 is an ancient and extensive family present in all land plants and performs a regulatory function by cleaving and inhibiting the expression of target nucleotide binding site and leucine-rich repeat (NBS-LRR) mRNAs [26]. NBS-LRR, as a class of Resistance (R) genes, recognize effectors in a direct or indirect way and can redirect the defense signaling and trigger R-gene-mediated immunity [27]. The miR482 family is more labile in sequence than other miRNA families [26-28]. To some extent, its diversity is a result of the amino acid variation in its target sequence [29]. Additionally, the expression of miR482 in the Solanaceae is higher than that in other species [26]. The miR482 family in *Solanum lycopersicum* possesses five members, namely, sly-miR482a/b/c/d/e [30-32]. The length of all five members is 22 nt rather than the typical 21 nt [26]. As members of the miR482 family in tomato display low sequence conservation with sequence variation at 5/6 variable sites [21], we hypothesize that the functions of members may differ. A recent study showed that sly-miR482e and sly-miR2118b (corresponding to sly-miR482d in the miRBase) differed in structure and function, which resulted in differences in their targets and mechanisms [21]. In our previous study, silencing sly-miR482b with a short tandem target mimic (STTM) was able to confer enhanced resistance to late blight in transgenic tomato [33]. The sequence alignment of sly-miR482b and sly-miR482c showed that their sequences were identical at 19 sites and varied at three sites. This difference indicated the possibility that miR482b and miR482c target distinct genes and function in different ways. Here, we overexpressed miR482c in tomato to study its mechanism. Two NBS-LRR genes with coiled-coil domains (CNLs),

Solyc07g049700 and *Solyc11g006530*, were predicted to be targeted by miR482c according to previous studies [26, 34]. The co-regulation of miR482c and these two genes was identified in tomato (*S. lycopersicum* cv. Heinz 1706) resistant to *Phytophthora infestans* IPO-C [34]. The genes *Solyc07g049700* and *Solyc11g006530* were selected and named *SICNL1* and *SICNL2*, respectively. Here, our study complemented the previous work and further explained the relationship and difference between miR482c and miR482b, laying a solid foundation for continued work.

MATERIALS AND METHODS

Plant Strains, *Phytophthora infestans* Infections

Tomato Zaofen No. 2 (susceptible cultivar) plants were raised in a growth room with a 16 hour light (at 22°C) and 8 hour dark (at 18°C) period. The agent of late blight, *Phytophthora infestans* (*P. infestans*) “P12103”, was cultured in oat medium at 20°C without light for 20 days, and the plates covered with mycelium were rinsed with distilled water until the concentration of spores reached 10^6 zoospores \cdot mL⁻¹. Afterwards, the spore suspensions were placed at 4°C for 1-2 hours to release the zoospores. Tomato plants with six fully expanded true leaves were sprayed with the prepared spore suspension. The control group consisted of tomato plants sprayed with sterile water. Leaf samples were collected at 0, 6, 24, 48, 72 and 96 hours postinoculation (hpi) according to a prior study [34]. All samples were quickly frozen in liquid nitrogen and subsequently stored at -80°C until DNA and RNA extraction.

Construction of the MiR482c-Overexpression Plasmids

The precursor sequence of miR482c (miRBase MI0020250) was amplified from tomato genomic DNA. The gene-specific primer pairs (listed in Table S1) were designed by Oligo7 software. The PCR products were cloned into the pBI121 vector driven by the 35S promoter. The sequence was identified

by DNA sequencing.

Transient Overexpression of MiR482c in Tomato

The plasmids pBI121-miR482c were introduced into the *Agrobacterium tumefaciens* (*A. tumefaciens*) strain “GV3101” by the freeze-thaw method [35]. The culture of *A. tumefaciens* was centrifuged for 10 min at $4000 \times g$, and the pellet was resuspended in infiltration medium (10 mM MES, 10 mM MgCl₂, and 20 μ M acetosyringone; OD₆₀₀ = 1.0). Next, the *A. tumefaciens* cultures containing pBI121-miR482c were agroinoculated into the leaves of the tomato plants. The infiltration with *A. tumefaciens* cultures containing empty vectors was used as the control. Three days after infiltration, each infiltrated leaf was inoculated with 20 μ L of *P. infestans* spore suspension (10^6 zoospores \cdot mL⁻¹) at the same position. The necrotic areas were observed after 3 days. Statistical differences in the diameters of necrotic lesions (DOL) between the control and infections were estimated using Duncan multiple range tests.

Generation of Transgenic Tomato Overexpressing MiR482c

The transformation of tomato was performed via the *Agrobacterium*-mediated leaf disk method [36]. Explants were obtained from cotyledons excised from one-week-old seedlings. The putative resistant buds were selected on 1/2-strength Murashige and Skoog medium containing 50 mg \cdot L⁻¹ kanamycin and 200 mg \cdot L⁻¹ carbenicillin with a 16 hour light (at 22°C) and 8 hour dark (at 18°C) period. After nearly two months, the putative positive transgenic lines were further confirmed by PCR amplification of the kanamycin-resistant gene neomycin phosphoryl transferase II (*nptII*) (listed in Table S1). The expression levels of miR482c in the selected transgenic lines were determined using quantitative real-time PCR (qRT-PCR).

Disease Resistance of Transgenic Plants Overexpressing MiR482c

The transgenic plants and wild-type (WT) plants with consistent growth were selected for the evaluation of resistance. In the detached-leaf infection assay, three leaves per plant and three plants per line were detached from one-month-old plants and placed on filter paper in a glass garden. A pipette tip (0.5-10 μL) was used to gently stab a wound to the right of the midvein on the abaxial side of the leaf. Next, 20 μL spore suspensions of *P. infestans* were injected into the wound. The plates were incubated in the dark at 20°C. The disease symptoms on inoculated leaves were observed and measured at 5 days postinfection. Subsequently, the leaves were dyed in trypan blue solution (2.5 mg · mL⁻¹) at 23 ± 1°C for 12 hours and discolored by being soaked in lactophenol/ethanol solution [14].

In the whole-plant infection assay, the WT plants and transgenic plants were sprayed with the same zoospore suspension and maintained in the greenhouse without light at 20 ± 1°C. The humidity was kept at 95%. After one day, a 16 hour light and 8 hour dark period was provided. Five days after infection, the disease severity rating (DSR) [37]. was counted. Additionally, the abundance of miR482c and its target genes in the WT and transgenic plants were detected using qRT-PCR.

DAB/NBT Staining and Measurements of Hydrogen Peroxide/MDA Content

At 5 days postinfection, the transgenic plants and WT plants with consistent growth were selected for analysis of H₂O₂/O₂⁻ with diamino benzidine (DAB)/nitro blue tetrazolium (NBT) staining as reported by previous methods [38]. Moreover, the activities of peroxidase (POD) and superoxide dismutase (SOD) were measured according to a previous study [35]. Furthermore, the content of malondialdehyde (MDA) and the activity of phenylalanine ammonia-lyase (PAL) were determined according to methods described by Hong et al [37].

qRT-PCR Analysis

The relative expression levels of all genes in this work were determined by qRT-PCR. The tomato

housekeeping gene *actin* was selected as the reference gene. All reactions were repeated three times.

Together, three independent biological assays were performed. The relative expression levels of each gene were calculated using the $2^{-\Delta\Delta CT}$ method.

Statistical Analysis

Statistical differences in data were estimated by Duncan multiple range tests using IBM SPSS Statistics19 software. Significant differences ($P < 0.05$) are presented with different lowercase letters.

All data are presented as the means \pm standard deviations (SDs).

RESULTS

Expression Patterns of MiR482c and Its Target Genes upon Infection

The relative expression levels of miR482c and *SICNLs* upon infection are shown in Figure 1. The relative expression of miR482c was significantly up-regulated at 6 hpi, subsequently down-regulated at 48 hpi, and eventually up-regulated at 96 hpi (Figure 1A). Moreover, the expression patterns of *SLCNL1* (Figure 1B) and *SLCNL2* (Figure 1C) was significantly down-regulated at 6 hpi, then up-regulated at 24 hpi, showing contrasting trends to that of miR482c. After 48 hpi, the expression pattern of *SLCNL1* and *SLCNL2* were similar to that of miR482c. It is speculated that miR482c and *SLCNLs* were induced early in infection, and there is a targeted relationship between miR482c and *SLCNLs*.




Figure 1. MiR482c and SICNL expression patterns upon infection. (A, B, C) The abundance of

miR482c (A), SLCNL1 (B) and SICNL2 (C) in WT plants upon infection. hpi: hours postinfection. The tomato housekeeping gene actin was selected as the reference gene. SDs of three replicates are indicated with error bars. The letters above the column indicate a significant difference ($P < 0.05$) between the samples.

Transient Overexpression of MiR482c Compromised Tomato Resistance

The framework of the pBI121-miR482c overexpression plasmids is shown in Figure 2A. The control transiently overexpressed empty vector (EV) in tomato leaves. The abundance of miR482c in tomato transiently overexpressing miR482c (TO482c) was nearly 20-fold higher than that in the control (Figure 2B). Compared with the control plants, the TO482c plants displayed more necrotic areas at 3 days after leaf infiltration (Figure 2C). In addition, the DOL on the TO482c leaves were larger than those on the control leaves (Figure 2D). The above results indicated that transiently overexpressing miR482c might compromise tomato resistance.




Figure 2. Transient overexpression of miR482c in tomato leaves. (A) The framework of the miR482c-overexpression vectors. (B) The expression level of miR482c in tomato that transiently overexpressed miR482c. EV: empty vector; TO482c: transiently overexpressing pBI121-miR482c. (C) The necrotic lesions on the leaves of EV and TO482c plants on the 3rd day after inoculation. (D) The

DOL in EV and TO482c plants. SDs of three replicates are indicated with error bars. The letters above the column indicate a significant difference ($P < 0.05$) between the samples.

MiR482c Had a Negative Effect on Tomato Resistance

The pBI121-miR482c plasmids were introduced into tomato to generate transgenic plants to further determine the regulation of miR482c in the resistance of tomato to late blight. Three positive transgenic lines (L1, L2, and L3) were selected for further examination. The expression patterns of miR482c and *SICNLs* in the transgenic lines were determined by qRT-PCR. As shown in Figure 3A, miR482c was more abundant in the three transgenic lines than in the WT plants, with the expression levels of the three transgenic lines L1, L2, and L3 ~2-fold, 2.5-fold, and 2.8-fold higher, respectively, than those of WT. Moreover, the transcript levels of *SICNLs* in the miR482c-overexpressing plants were lower than those in the WT plants (Figure 3B, C). The decline in *SICNL* mRNA levels in the transgenic plants further demonstrated that *SICNL* mRNAs were targeted by miR482c.




Figure 3. Disease resistance of transgenic plants stably overexpressing miR482c. (A, B, C) The relative expression levels of miR482c (A), *SICNL1* (B) and *SICNL2* (C) in transgenic tomato lines. WT: wide type; L1: line 1; L2: line 2; L3: line 3. (D) The disease symptoms of WT and L1-L3 in the whole-plant infection assay on the 5th day after infection. (E) The DSR of WT and L1-L3 on the 5th day

after infection. (F) Necrotic areas on detached leaves of the WT and L1-L3 plants on the 5th day after inoculation. Top: necrotic areas; bottom: trypan blue staining of the detached leaves. (G) The ratio of lesion area to leaf area in the detached-leaf infection assay. SDs of three replicates were displayed by error bars. The letters above the column indicate a significant difference ($P < 0.05$) between the samples.

In the whole-plant infection assay, more severe disease symptoms were observed on the leaves of the L1-L3 plants than on the leaves of the WT plants (Figure 3D). The DSR of the L1-L3 plants was almost 4-fold higher than that of the WT plants (Figure 3E). In the detached-leaf inoculation assay, dead cells were detected in the leaves stained with trypan blue after infection. The L1-L3 leaves had darker blue marks than did the WT leaves, indicating increased cell death in the L1-L3 plants (Figure 3F). Additionally, the ratio of lesion area to leaf area in the leaves of the WT plants was much less than that of the leaves of the transgenic tomato plants (Figure 3G).

Overexpression of MiR482c Contributed to Changes in Physiological Indicators

In plant-pathogen interactions, oxidative bursts frequently occur and produce large amounts of reactive oxygen species (ROS), mainly including H_2O_2 and O_2^- . In this study, DAB staining and NBT staining were conducted to determine the accumulation of H_2O_2 and O_2^- in the transgenic plants and WT plants after treatment. As shown in Figure 4A, the leaves of the transgenic plants exhibited more intensely stained spots than those of WT. Additionally, the enzyme activities of POD and SOD were significantly lower in the L1-L3 plants than in the WT plants upon infection (Figure 4B, C). This was consistent with the notably decreased expression of *SISOD* and *SIPOD* (genes encoding the SOD and POD enzymes in tomato) [20, 35]. in the L1-L3 plants compared with the WT plants upon infection (Figure 4D, E). These results indicated that the L1-L3 plants accumulated more ROS than the WT

plants.




Figure 4. Detection of physiological and biochemical indicators of the miR482c-overexpression

plants. (A) DAB staining and NBT staining of tomato leaves in the whole-plant infection assay on the 5th day after infection. (B, C) SOD (B) and POD (C) activity of WT and L1-L3 after infection for 5 days. (D, E) The expression levels of *SISOD* (D) and *SIPOD* (E) in WT and transgenic lines on the 5th day after infection. (F) The MDA content of WT and L1-L3 after infection for 5 days. (G) The PAL activity of WT and L1-L3 after infection for 5 days. SDs of three replicates are indicated with error bars. The letters above the column indicate a significant difference ($P < 0.05$) between the samples.

In addition, a higher MDA content was found in the L1-L3 plants, indicating that the L1-L3 plants suffered more severe membrane damage than the WT plants (Figure 4F). PAL is a key enzyme in the phenylpropanoid metabolic pathway. The level of PAL activity can also be used as an important physiological indicator for measuring plant disease resistance. The activity of the PAL enzyme in the L1-L3 plants was higher than that in the WT plants (Figure 4G). These results suggested that overexpressing miR482c in tomato aggravated membrane damage.

DISCUSSION

MiRNAs, as key regulators, have been widely suggested to play crucial roles in plant resistance. A large number of studies have documented that miRNAs function in the plant response to biotic stress. For instance, *osa-miR167d* has been reported to act as a negative regulator in rice resistance to rice blast disease [39]. *Osa-miR396* has been demonstrated to negatively regulate the resistance of rice to brown planthopper (BPH) [40]. In Arabidopsis, *miR825* and *miR825** were shown to compromise plant resistance to *Pseudomonas syringae* pv. *tomato* (*Pst*) DC3000 [41]. Tomato plants are generally subjected to various pathogens leading to severe diseases and substantial losses. However, the mechanism of *miR482c* in tomato disease resistance remains elusive. Our study indicated that overexpression of *miR482c* in tomato led to enhanced susceptibility to late blight disease.

We found that the expression pattern of *miR482c* was induced at 6 hpi and 24 hpi, subsequently declined at 48 hpi and 72 hpi, and finally increased at 96 hpi (Figure 1A). The relative expression levels of both *SLCNL1* and *SLCNL2* decreased at 6 hpi and subsequently were up-regulated at 24 hpi in contrast. Afterwards, at 48 hpi, *SLCNL1* increased, whereas *SLCNL2* decreased. Both *SLCNLs* declined at 72 hpi and were eventually up-regulated at 96 hpi (Figure 1B, C). A previous study demonstrated that the co-regulations between *miR482*-*NBS-LRRs* were not exactly the same [34]. Such differences in co-regulation suggest that despite active targeting by *miR482c* in *S. lycopersicum*, *SLCNL1* and *SLCNL2* are likely to be regulated by other mechanisms in addition to the regulation by *miR482c*. Furthermore, upregulation or downregulation is not static between *miR482* and *NBS-LRR* mRNA but can shift between time points. Switches between translational and posttranscriptional may account for such shifted regulation [34]. Additionally, Jiang et al. demonstrated that the abundance of *miR482b* in L3708 tomato (resistant cultivar) was lowest at 24 hpi but increased to a moderate level at 96 hpi [33]. The different expression patterns of *mi482b* and *miR482c* may result from divergent cultivars.

To further confirm the participation of miR482c in tomato resistance, three transgenic lines overexpressing miR482c (L1-L3) were generated. After five days postinfection, the transgenic lines displayed more severe symptoms than WT (Figure 3D) with greater DSRs (Figure 3E). Moreover, in the detached-leaf inoculation assay, necrotic areas on the leaves of transgenic lines were larger than those on the leaves of WT (Figure 3F). This indicated that the resistance was impaired after overexpression of miR482c. Previous work reported similar results; the mature sequences of *stu*-miR482e in potato and *sly*-miR482c in tomato are identical, and there are only two site variations between the precursor sequences of *stu*-miR482e and *sly*-miR482c. Overexpression of *stu*-miR482e in potato was demonstrated to enhance plant sensitivity to *Verticillium dahliae* infection [42].

In the early stages of plant-pathogen interactions, oxidative bursts frequently occur and produce large amounts of ROS, mainly including H₂O₂ and O₂⁻ [43]. Excessive ROS may result in lipid peroxidation and oxidative damage to the membrane or even cell death. In rice, Wu et al. showed that miR528 acts as a negative regulator in viral resistance and is accompanied by an increased accumulation of ROS [44]. In our study, the DAB and NBT staining assay showed that there were more dark spots on the leaves of the transgenic tomato plants than the leaves of the WT tomato plants, indicating increased cell death and accumulation of H₂O₂ and O₂⁻ in the transgenic tomato plants (Figure 4A). Plants have evolved a complete antioxidant defense system that removes excess ROS and maintains homeostasis. The activities of SOD and POD related to ROS clearance in the transgenic tomato plants were considerably lower upon infection than those in the WT plants (Figure 4B, C). Together, the relative expression levels of the antioxidant enzyme genes *SIPOD* and *SISOD* were also down-regulated in the transgenic plants (Figure 4D, E). Similar results have been reported in tomato, for example, Zhang et al. reported that *slmapk3* mutants and WT showed increased accumulation of

ROS upon *Botrytis cinerea* infection, and *slmapk3* mutants showed reduced resistance with an increased level of ROS content [45].

In summary, we proposed a working model to describe the mechanism of miR482c in tomato resistance to late blight (Figure 5). Overexpression of miR482c could reduce the expression of its targeted *SLCNL* genes and antioxidant enzyme genes, thereby leading to a decline in the ROS scavenging ability and aggravating the damage of lipid peroxidation product accumulation on the cell membrane, eventually enhancing plant susceptibility. Compared with the function of miR482b which was reported earlier [33], miR482c showed analogous negative regulation with miR482b in tomato resistance to late blight. Thus, despite having individual characteristics, miR482c and miR482b may play analogous roles in the plant immune system. Results have been reported that coincide with this finding. For instance, Li et al. reported that miR169a and miR169c in Arabidopsis functioned as negative regulators under drought stress by cleaving NFYA5 [46]. Furthermore, Zhao et al. demonstrated that miR169g and miR169n both targeted an NF-YA gene exhibiting distinct and overlapping responses to salt and drought stresses [47].




Figure 5. The regulation network of miR482c-NBS-LRR involved in tomato resistance.

Overexpressing miR482c in tomato enhanced plant susceptibility to late blight by reducing the targeted *SICNL* genes, accompanied by lower POD, SOD, and PAL activities and higher MDA content. Red lines represent negative regulated gene or function, while green arrows represent positive regulated

gene or function. Red dotted arrow indicates the effectors produced by pathogens invading the host plant acts on the production of ROS by direct or indirect way.

Our findings provide insight into the method by which miR482c–NBS-LRR participated in the response of tomato to late blight. Future work should therefore include follow-up work designed to study whether silencing miR482c enhances disease resistance and whether the miR482c cascade plays a role in resistance to other forms of stress.

Author Contributions

Methodology and investigation, Y.-S.L., Y.-H.H., X.-L.H., and Y.-Y.Z.; writing-original draft preparation, Y.-S.L., Y.-H.H.; supervision, Y.-S.L., J.M.; writing-review and editing, Y.-S.L., Y.-H.H., J.M..

Funding

This research was funded by the National Natural Science Foundation of China (Nos. 31872116 and 61872055).

Notes

The authors declare no competing financial interest.

Abbreviations Used

H₂O₂, hydrogen peroxide; MDA, malondialdehyde; NBT, nitro blue tetrazolium; O₂⁻, superoxide; PAL, phenylalanine ammonia-lyase; POD, peroxidase; qRT-PCR, quantitative real-time PCR; ROS, reactive oxygen species; SD, standard deviation; SOD, superoxide dismutase; WT, wild type

References

1. Tieman, D.; Zhu, G.; Resende, M. F., Jr.; Lin, T.; Nguyen, C.; Bies, D.; Rambla, J. L.; Beltran, K. S.; Taylor, M.; Zhang, B.; Ikeda, H.; Liu, Z.; Fisher, J.; Zemach, I.; Monforte, A.; Zamir, D.; Granell, A.; Kirst, M.; Huang, S.; Klee, H., A chemical genetic roadmap to improved tomato flavor. *Science* **2017**, *355*, 391-394; DOI: 10.1126/science.aal1556.
2. Fei, Z. J.; Joung, J. G.; Tang, X. M.; Zheng, Y.; Huang, M. Y.; Lee, J. M.; McQuinn, R.; Tieman, D. M.; Alba, R.; Klee, H. J.; Giovannoni, J. J., Tomato Functional Genomics Database: a comprehensive resource and analysis package for tomato functional genomics. *Nucleic Acids Res.* **2011**, *39*, 1156-1163; DOI: 10.1093/nar/gkq991.
3. Food and Agricultural Organization of the United Nations. <http://www.fao.org/faostat/>.
4. Fry, W. E.; Birch, P. R.; Judelson, H. S.; Grunwald, N. J.; Danies, G.; Everts, K. L.; Gevens, A. J.; Gugino, B. K.; Johnson, D. A.; Johnson, S. B.; McGrath, M. T.; Myers, K. L.; Ristaino, J. B.; Roberts, P. D.; Secor, G.; Smart, C. D., Five reasons to consider *phytophthora infestans* a reemerging pathogen. *Phytopathology* **2015**, *105*, 966-981; DOI: 10.1094/PHYTO-01-15-0005-FI.
5. Nowicki, M.; Foolad, M. R.; Nowakowska, M.; Kozik, E. U., Potato and tomato late blight caused by *Phytophthora infestans*: an overview of pathology and resistance breeding. *Plant Dis.* **2012**, *96*, 4-17; DOI: 10.1094/PDIS-05-11-0458.
6. Kelley, B. S.; Lee, S. J.; Damasceno, C. M.; Chakravarthy, S.; Kim, B. D.; Martin, G. B.; Rose, J. K., A secreted effector protein (SNE1) from *Phytophthora infestans* is a broadly acting suppressor of programmed cell death. *Plant J.* **2010**, *62*, 357-366; DOI: 10.4161/psb.5.6.11778.

7. Bartel, D. P., MicroRNAs: target recognition and regulatory functions. *Cell* **2009**, *136*, 215-233; DOI: 10.1016/j.cell.2009.01.002.
8. Zeng, J.; Gupta, V. K.; Jiang, Y. M.; Yang, B.; Gong, L.; Zhu, H., Cross-Kingdom Small RNAs among Animals, Plants and Microbes. *Cells* **2019**, *8*, 371; DOI: 10.3390/cells8040371.
9. D'Ario, M.; Griffiths-Jones, S.; Kim, M., Small RNAs: big impact on plant development. *Trends Plant. Sci.* **2017**, *22*, 1056-1068; DOI: 10.1016/j.tplants.2017.09.009.
10. Sun, T.; Li, M. Y.; Li, P. F.; Cao, J. M., MicroRNAs in Cardiac Autophagy: Small Molecules and Big Role. *Cells* **2018**, *7*, 15; DOI: 10.3390/cells7080104.
11. Luan, Y.; Cui, J.; Zhai, J.; Li, J.; Han, L.; Meng, J., High-throughput sequencing reveals differential expression of miRNAs in tomato inoculated with *Phytophthora infestans*. *Planta* **2015**, *241*, 1405-1416; DOI: 10.1007/s00425-015-2267-7.
12. Navarro, L.; Dunoyer, P.; Jay, F.; Arnold, B.; Dharmasiri, N.; Estelle, M.; Voinnet, O.; Jones, J. D., A plant miRNA contributes to antibacterial resistance by repressing auxin signaling. *Science* **2006**, *312*, 436-439; DOI: 10.1126/science.1126088.
13. Wang, Z.; Xia, Y.; Lin, S.; Wang, Y.; Guo, B.; Song, X.; Ding, S.; Zheng, L.; Feng, R.; Chen, S.; Bao, Y.; Sheng, C.; Zhang, X.; Wu, J.; Niu, D.; Jin, H.; Zhao, H., Osa-miR164a targets *OsNAC60* and negatively regulates rice immunity against the blast fungus *Magnaporthe oryzae*. *Plant J.* **2018**, *95*, 584-597; DOI: 10.1111/tpj.13972.
14. Chen, L.; Meng, J.; He, X. L.; Zhang, M.; Luan, Y. S., Sly-miR1916 targets multiple target genes and negatively regulates the immune response in tomato. *Plant, Cell Environ.* **2019**, *42*, 1393-1407; DOI: 10.1111/pce.13468.

- 15.Zhang, X. H.; Zou, Z.; Gong, P. J.; Zhang, J. H.; Ziaf, K.; Li, H. X.; Xiao, F. M.; Ye, Z. B.,
Over-expression of microRNA169 confers enhanced drought tolerance to tomato. *Biotechnol. Lett.*
2011, *33*, 403-409; DOI: 10.1007/s10529-010-0436-0.
- 16.Li, H.; Zhang, Q.; Li, L.; Yuan, J.; Wang, Y.; Wu, M.; Han, Z.; Liu, M.; Chen, C.; Song, W.; Wang,
C., Ectopic overexpression of bol-miR171b increases chlorophyll content and results in sterility in
broccoli (*Brassica oleracea* L var. *italica*). *J. Agri. Food Chem.* **2018**, *66*, 9588-9597; DOI:
10.1021/acs.jafc.8b01531.
- 17.Gao, C.; Ju, Z.; Cao, D. Y.; Zhai, B. Q.; Qin, G. Z.; Zhu, H. L.; Fu, D. Q.; Luo, Y. B.; Zhu, B. Z.,
MicroRNA profiling analysis throughout tomato fruit development and ripening reveals potential
regulatory role of RIN on microRNAs accumulation. *Plant Biotechnol. J.* **2015**, *13*, 370-382; DOI:
10.1111/pbi.12297.
- 18.Nozawa, M.; Miura, S.; Nei, M., Origins and evolution of microRNA genes in plant species.
Genome Biol. Evol. **2012**, *4*, 230-239; DOI: 10.1093/gbe/evs002.
- 19.Salvador-Guirao, R.; Hsing, Y. I.; San Segundo, B., The polycistronic miR166k-166h positively
regulates rice immunity via post-transcriptional control of *EIN2*. *Front. Plant Sci.* **2018**, *9*, 337;
DOI: 10.3389/fpls.2018.00337.
- 20.Luan, Y.; Cui, J.; Li, J.; Jiang, N.; Liu, P.; Meng, J., Effective enhancement of resistance to
Phytophthora infestans by overexpression of miR172a and b in *Solanum lycopersicum*. *Planta*
2018, *247*, 127-138; DOI:10.1007/s00425-017-2773-x.
- 21.Canto-Pastor, A.; Santos, B.; Valli, A. A.; Summers, W.; Schornack, S.; Baulcombe, D. C.,
Enhanced resistance to bacterial and oomycete pathogens by short tandem target mimic RNAs in
tomato. *Proc. Natl. Acad. Sci. U. S. A.* **2019**, *116*, 2755-2760; DOI: 10.1073/pnas.1814380116.

- 22.Csukasi, F.; Donaire, L.; Casanal, A.; Martinez-Priego, L.; Botella, M. A.; Medina-Escobar, N.; Llave, C.; Valpuesta, V., Two strawberry miR159 family members display developmental-specific expression patterns in the fruit receptacle and cooperatively regulate Fa-GAMYB. *New Phytol.* **2012**, *195*, 47-57; DOI: 10.1111/j.1469-8137.2012.04134.x.
- 23.Ma, X.; Zhang, X.; Zhao, K.; Li, F.; Li, K.; Ning, L.; He, J.; Xin, Z.; Yin, D., Small RNA and degradome deep sequencing reveals the roles of microRNAs in seed expansion in peanut (*Arachis hypogaea* L.). *Front. Plant Sci* **2018**, *9*, 349; DOI: 10.3389/fpls.2018.00349.
- 24.Zhou, Y.; Zhou, S.; Wang, L.; Wu, D.; Cheng, H.; Du, X.; Mao, D.; Zhang, C.; Jiang, X., *miR164c* and *miR168a* regulate seed vigor in rice. *J. Integr. Plant Biol.* **2019**; DOI: 10.1111/jipb.12792.
- 25.Fang, Y.; Xie, K.; Xiong, L., Conserved miR164-targeted NAC genes negatively regulate drought resistance in rice. *J. Exp. Bot.* **2014**, *65*, 2119-2135; DOI: 10.1093/jxb/eru072.
- 26.Shivaprasad, P. V.; Chen, H. M.; Patel, K.; Bond, D. M.; Santos, B. A.; Baulcombe, D. C., A microRNA superfamily regulates nucleotide binding site-leucine-rich repeats and other mRNAs. *Plant Cell.* **2012**, *24*, 859-874; DOI: 10.1105/tpc.111.095380.
- 27.de Vries, S.; Kloesges, T.; Rose, L. E., Evolutionarily dynamic, but robust, targeting of resistance genes by the miR482/2118 gene family in the Solanaceae. *Genome Biol. Evol.* **2015**, *7*, 3307-3321; DOI: 10.1093/gbe/evv225.
- 28.Zhai, J.; Jeong, D. H.; De Paoli, E.; Park, S.; Rosen, B. D.; Li, Y.; Gonzalez, A. J.; Yan, Z.; Kitto, S. L.; Grusak, M. A.; Jackson, S. A.; Stacey, G.; Cook, D. R.; Green, P. J.; Sherrier, D. J.; Meyers, B. C., MicroRNAs as master regulators of the plant *NB-LRR* defense gene family via the production of phased, *trans*-acting siRNAs. *Genes Dev.* **2011**, *25*, 2540-2553; DOI: 10.1101/gad.177527.111.

- 29.Zhang, Y.; Xia, R.; Kuang, H.; Meyers, B. C., The diversification of plant *NBS-LRR* defense genes directs the evolution of microRNAs that target them. *Mol. Biol. Evol.* **2016**, *33*, 2692-2705; DOI: 10.1093/molbev/msw154.
- 30.Kravchik, M.; Sunkar, R.; Damodharan, S.; Stav, R.; Zohar, M.; Isaacson, T.; Arazi, T., Global and local perturbation of the tomato microRNA pathway by a *trans*-activated *DICER-LIKE 1* mutant. *J. Exp. Bot.* **2014**, *65*, 725-739; DOI: 10.1093/jxb/ert428.
- 31.Mohorianu, I.; Schwach, F.; Jing, R.; Lopez-Gomollon, S.; Moxon, S.; Szittyá, G.; Sorefan, K.; Moulton, V.; Dalmay, T., Profiling of short RNAs during fleshy fruit development reveals stage-specific sRNAome expression patterns. *Plant J.* **2011**, *67*, 232-246; DOI: 10.1111/j.1365-313x.2011.04586.x.
- 32.Li, F.; Pignatta, D.; Bendix, C.; Brunkard, J. O.; Cohn, M. M.; Tung, J.; Sun, H.; Kumar, P.; Baker, B., MicroRNA regulation of plant innate immune receptors. *Proc. Natl. Acad. Sci. U. S. A.* **2012**, *109*, 1790-1795; DOI: 10.1073/pnas.1118282109.
- 33.Jiang, N.; Meng, J.; Cui, J.; Sun, G.; Luan, Y., Function identification of miR482b, a negative regulator during tomato resistance to *Phytophthora infestans*. *Hortic. Res.* **2018**, *5*, 9; DOI: 10.1038/s41438-018-0017-2.
- 34.de Vries, S.; Kukuk, A.; von Dahlen, J. K.; Schnake, A.; Kloesges, T.; Rose, L. E., Expression profiling across wild and cultivated tomatoes supports the relevance of early miR482/2118 suppression for *Phytophthora* resistance. *P. Roy. Soc. B-Biol. Sci.* **2018**, *285*, 20172560; DOI: 10.1098/rspb.2017.2560.

- 35.Cui, J.; Jiang, N.; Zhou, X.; Hou, X.; Yang, G.; Meng, J.; Luan, Y., Tomato MYB49 enhances resistance to *Phytophthora infestans* and tolerance to water deficit and salt stress. *Planta* **2018**, *248*, 1487-1503; DOI: 10.1007/s00425-018-2987-6.
- 36.Khare, N.; Goyary, D.; Singh, N. K.; Shah, P.; Rathore, M.; Anandhan, S.; Sharma, D.; Arif, M.; Ahmed, Z., Transgenic tomato cv. Pusa Uphar expressing a bacterial mannitol-1-phosphate dehydrogenase gene confers abiotic stress tolerance. *Plant Cell, Tissue Organ Cult.* **2010**, *103*, 267-277; DOI: 10.1007/s11240-010-9776-7.
- 37.Hong, Y.; Cui, J.; Liu, Z.; Luan, Y., *SpWRKY6* acts as a positive regulator during tomato resistance to *Phytophthora infestans* infection. *Biochem. Biophys. Res. Commun.* **2018**, *506*, 787-792; DOI: 10.1016/j.bbrc.2018.10.155.
- 38.Cui, J.; Luan, Y.; Jiang, N.; Bao, H.; Meng, J., Comparative transcriptome analysis between resistant and susceptible tomato allows the identification of lncRNA16397 conferring resistance to *Phytophthora infestans* by co-expressing glutaredoxin. *Plant J.* **2017**, *89*, 577-589; DOI: 10.1111/tpj.13408.
- 39.Zhao, Z. X.; Feng, Q.; Cao, X. L.; Zhu, Y.; Wang, H.; Chandran, V.; Fan, J.; Zhao, J. Q.; Pu, M.; Li, Y.; Wang, W. M., *Osa-miR167d* facilitates infection of *Magnaporthe oryzae* in rice. *J. Integr. Plant Biol.* **2019**; DOI: 10.1111/jipb.12816.
- 40.Dai, Z.; Tan, J.; Zhou, C.; Yang, X.; Yang, F.; Zhang, S.; Sun, S.; Miao, X.; Shi, Z., The OsmiR396-OsGRF8-OsF3H-flavonoid pathway mediates resistance to the brown planthopper in rice (*Oryza sativa*). *Plant Biotechnol. J.* **2019**; DOI: 10.1111/pbi.13091.
- 41.Niu, D.; Xia, J.; Jiang, C.; Qi, B.; Ling, X.; Lin, S.; Zhang, W.; Guo, J.; Jin, H.; Zhao, H., *Bacillus cereus* AR156 primes induced systemic resistance by suppressing miR825/825* and activating

- defense-related genes in Arabidopsis. *J. Integr. Plant Biol.* **2016**, *58*, 426-439; DOI: 10.1111/jipb.12446.
42. Yang, L.; Mu, X.; Liu, C.; Cai, J.; Shi, K.; Zhu, W.; Yang, Q., Overexpression of potato miR482e enhanced plant sensitivity to *Verticillium dahliae* infection. *J. Integr. Plant Biol.* **2015**, *57*, 1078-1088; DOI: 10.1111/jipb.12348.
43. Apel, K.; Hirt, H., Reactive oxygen species: Metabolism, oxidative stress, and signal transduction. *Annu. Rev. Plant Biol.* **2004**, *55*, 373-399; DOI: 10.1146/annurev.arplant.55.031903.141701.
44. Wu, J.; Yang, R.; Yang, Z.; Yao, S.; Zhao, S.; Wang, Y.; Li, P.; Song, X.; Jin, L.; Zhou, T.; Lan, Y.; Xie, L.; Zhou, X.; Chu, C.; Qi, Y.; Cao, X.; Li, Y., ROS accumulation and antiviral defence control by microRNA528 in rice. *Nat. Plants* **2017**, *3*, 16203; DOI: 10.1038/nplants.2016.203.
45. Zhang, S.; Wang, L.; Zhao, R.; Yu, W.; Li, R.; Li, Y.; Sheng, J.; Shen, L., Knockout of *SIMAPK3* Reduced Disease Resistance to *Botrytis cinerea* in Tomato Plants. *J. Integr. Plant Biol.* **2018**, *66*, 8949-8956; DOI: 10.1021/acs.jafc.8b02191.
46. Li, W. X.; Oono, Y.; Zhu, J.; He, X. J.; Wu, J. M.; Iida, K.; Lu, X. Y.; Cui, X.; Jin, H.; Zhu, J. K., The *Arabidopsis* NFYA5 transcription factor is regulated transcriptionally and posttranscriptionally to promote drought resistance. *Plant Cell.* **2008**, *20*, 2238-2251; DOI: 10.1105/tpc.108.059444.
47. Zhao, B.; Ge, L.; Liang, R.; Li, W.; Ruan, K.; Lin, H.; Jin, Y., Members of miR-169 family are induced by high salinity and transiently inhibit the NF-YA transcription factor. *BMC Mol. Biol.* **2009**, *10*, 29; DOI: 10.1186/1471-2199-10-29.

Table S1 Primer sequence used in this study.

Purpose	Gene Name	Primer sequence (5' - 3')
clone of pre-miR482c	c-miR482c	FP:CGAGCTCCCCAGCTTTCATCATGTTTTTCA
		RP:CGGGATCCGAAGCCATTGGAGTTAAGTTGACA
identification of transgenic lines	<i>nptII</i>	FP: CTCACCTTGCTCCTGCCGAGA RP: CGCCTTGAGCCTGGCGAACAG
	miR482c	FP:TCTTGCCAATACCGCCCATTCC
qRT-PCR	tomato <i>actin</i>	FP:ACCTTCAACGTTCCAGCTATG RP:TCACCAGAGTCCAACACAATAC
	<i>SLCNL1</i>	FP:TGGATGCACATGAAATCGGC RP:TCCAAATCTCCGTTGGGAGC
	<i>SLCNL2</i>	FP:TGTTGGGATGGAGGATGACTT RP:TCTTACCTATAACCGCCCATTCC
	<i>SIPOD</i>	FP: GAATGTCCGGGAGTTGTTTCT RP: TCTTCTACCAGTTGGCACATTC
	<i>SISOD</i>	FP: AGAAAGCTGTTGCTGTCCTTA RP: CCAGGAGCAAGTCCAGTTATAC