

Article

Benthic Habitat Morphodynamics- Using Remote Sensing to Quantify Storm-Induced Changes in Nearshore Bathymetry and Surface Sediment Texture at Assateague National Seashore

Arthur Trembanis ^{1,*}, Alimjan Abila², Ken Haulsee ¹, and Carter DuVal¹

¹ School of Marine Science and Policy
University of Delaware
Lewes, DE 19958 USA
art@udel.edu

² Middle East Technical University
Institute of Marine Science, Turkey
Visiting PhD Fellow at UD 2016-2017
alimjan_abila@yahoo.com

* Correspondence: art@udel.edu; Tel.: (+1 302-831-2498)

Abstract: This study utilizes repeated geoacoustic mapping to quantify the morphodynamic response of the nearshore to storm-induced changes. The aim of this study was to quantitatively map the nearshore zone of Assateague Island National Seashore (ASIS) to determine what changes in bottom sediments, benthic fauna and fish habitat are attributable to storm events including hurricane Sandy and the passage of hurricane Joaquin. Specifically, (1) the entire domain of the National Parks Service offshore area was mapped with side-scan sonar and multibeam bathymetry at a resolution comparable to that of the existing pre-storm survey, (2) a subset of the benthic stations were resampled that represented all sediment strata previously identified, and (3) newly obtained data were compared to that from the pre-storm survey to determine changes that could be attributed to specific storms such as Sandy and Joaquin. Capturing event specific dynamics requires rapid response surveys in close temporal association of the before and after period. The time-lapse between the pre-storm surveys for Sandy and our study meant that only a time and storm integrated signature for that storm could be obtained whereas with hurricane Joaquin we could identify impacts to the habitat type and geomorphology more directly related to that particular storm. This storm impacts study provides for the National Park Service direct documentation of storm-related changes in sediments and marine habitats on multiple scales: from large scale, side-scan sonar maps and interpretation of acoustic bottom types, to characterize as fully as possible habitats from 1 to 10 m up to many kilometer scales, as well as from point benthic samples within each sediment stratum and these results can help guide management of the island resources.

Keywords: side-scan sonar, swath bathymetry, habitat monitoring, hurricane Sandy, hurricane Joaquin, climate change, shoreline detection, remote sensing

1. Introduction

The overall goal of this study was to map the morphodynamic changes to the nearshore zone of Assateague Island National Seashore (ASIS) to determine what changes in bottom sediments, benthic fauna and fish habitat occurred are attributable to storm impacts such as Superstorm Sandy and hurricane Joaquin. Specifically, we (1) mapped the full survey area with side-scan sonar and multibeam bathymetry at a resolution comparable to that of the pre-storm survey, (2) resampled a

subset of the benthic stations that represented all sediment strata previously identified, and (3) compared newly obtained data to that from the pre-storm survey to examine storm impacts. This survey design aimed to document storm-related changes in sediments and marine habitats on multiple scales: from large scale, side-scan sonar maps and interpretation of acoustic bottom types, to characterize as fully as possible habitats from 1 to 10 m up to many kilometer scales, as well as from point benthic samples within each sediment stratum.

1.1 Background on Acoustic Mapping

Human reliance upon, and interference with benthic ecosystems necessitates an understanding of the spatial extent, structure, and function of these unique ecosystems (Balthis et al., 2006; Engle et al., 2009). This project work utilized advanced geoacoustic techniques for remote benthic habitat mapping and employed both traditional technologies for wide area coverage combined with point sampling and robotic systems including an autonomous underwater vehicle (AUV) and a remotely operated vehicle (ROV) for gathering data in an unprecedented level of resolution over targeted regions of interest. Knowledge gained from this study thus allows coastal and estuary environmental stakeholders responsible for the Assateague Island National Seashore (ASIS) to better manage resources to protect these environments by providing information on the location of habitats, species' distributions and associations to varying sedimentary and morphological regimes. The new benthic habitat and geomorphologic datasets will serve as a useful reference for an array of mapping and management efforts particularly providing insights into temporal dynamics from before and after storms.

Estuarine and coastal regions of the US face multiple marine spatial planning issues and challenges, including the effects of episodic storms, climate change, and other stressors (Douvere and Ehler, 2005). In particular, there has been a strong interest in and multiple efforts to examine the effects of hurricane Sandy on the seabed within the Mid-Atlantic Bight (Goff et al., 2013; Hapke et al., 2013; Trembanis et al., 2013). The anticipated impacts of offshore development on the seabed and to the ecology and, more generally, NOAA's initiatives in marine spatial planning (Douvere and Ehler, 2009) point to an ever-increasing need for both base-line mapping and monitoring efforts directed at biological habitats and geological features of the littoral zone.

In this study, we realized the need to fuse disparate data streams and recognized the most effective means of analyzing and understanding them is through data visualization. By repeated benthic habitat and morphology mapping, this field effort and the subsequent data analysis provides a critical extension to the previous study of the ASIS site (2011-2012). We recognize that together these data sets collected over multiple years represent a knowledge base much greater than the sum of its parts, and it is this fact that motivated the project. The ecological services provided by these benthic communities (Diaz et al., 2004), e.g., physical structure, prey species refuge and recreational fisheries, justify the renewed interest in these communities and habitats.

Recent benthic mapping on similar prior projects to this project (Trembanis et al., 2012; Miller et al., 2010; Raineault et al., 2012) have demonstrated that, in general, shallow coastal ecosystems are far more complex and heterogeneous in bottom type than typically anticipated. Furthermore, conventional sampling by direct methods (i.e., bottom grabs or dredges) is also generally recognized as severely limited by several factors including gear bias, spatial averaging and limited sample density. It is only from the careful fusion of multiple technologies (i.e., ship-based, ROV, AUV, grab samples, and dredge) that one is able to compile the complete picture of the nature and composition of the bottom (Diaz et al., 2004, Miller et al., 2010; Raineault et al., 2012; Raineault et al., 2013).

Accurate bathymetric and backscatter data are essential for hydrographic charts, dredging, and habitat mapping. Previously, the primary method for seafloor mapping was the use of single-

beam bathymetry. It is an accurate technique for collecting seafloor data, yet there are a number of limitations. In single-beam surveys, a discrete footprint of the seabed is ensconced leaving large gaps between adjacent survey tracks that are not surveyed. Single-beam surveys typically cover only 5-10% of the total seafloor area leaving large portions unsampled. Therefore, the maps generated from single-beam sonar samples may lack high-enough resolution to detect small and ecologically important habitat features (Parnum et al., 2009). In a recent study, Parnum et al. (2009) found that a full coverage swath sonar system resulted in a 90% classification accuracy compared to 70-80% accuracy from a single beam system approach. Using conventional methods like RoxAnn (e.g., Hauser 2002; Wilson and Madsen, 2006), found that the sampling is limited to a single sonar point footprint directly below the vessel, and maps necessitate heavy interpolation between track lines. In shallow settings, single-beam systems therefore require progressively more and more tightly spaced lines as the local depth decreases. In contrast, multi-beam echosounders, interferometric sonar, and side-scan sonar systems characterize a wide swath on either side of the survey vessel. Surfaced towed acoustic sensors suffer from positioning errors associated with the ambiguity of the towfish location relative to the ship GPS. Surface vessel mounted sensors also suffer from reduced resolution (larger footprint) as water depth increases. Wide swath acoustic methods afford the potential of effectively mapping large areas of the seafloor at a resolution appropriate for habitat mapping and correlation with biological communities. Seafloor echoes can be interpreted in terms of hardness and roughness and correlated with the sediment and faunal characteristics.

Several reviews and studies of coastal sedimentary systems (e.g., Holland et al., 2003; Trembanis et al., 2004; Green et al., 2004) have repeatedly demonstrated that dramatic spatiotemporal heterogeneity dominates seabed settings with respect to composition and morphology and that this heterogeneity is more the norm than the exception. Further exacerbating the problem, seabed heterogeneity often exists at spatial scales on the order of 1-10's m, well below typical single-beam survey spacing and gridding schemes. Understanding and being able to map this spatial patchiness at sufficient coverage and resolution scales is a critical component of habitat mapping efforts and an important motivation behind this project.

1.2 Study Design and Objectives

This study was designed to characterize the physical and biological components of the nearshore benthic zone of Assateague Island National Seashore (ASIS), including delineations of habitat and assemblage types using the CMECS system. Here "assemblage" refers to the list of resident species in a particular sample or site, and "habitat" refers to the spatial context of those assemblages, as defined by sedimentary characteristics like grain size or organic content. We characterized benthic invertebrate assemblages and their associated sedimentary habitats by analyzing Young grab samples for grain size distribution, and total organic carbon (TOC). The study methods were modeled off of the 2011 ASIS benthic habitat and assemblage assessment projects undertaken by Versar, Inc. (Columbia, MD) and the Maryland Geological Survey (MGS), originally contracted by the National Park Service (NPS). A Natural Resource Condition Assessment undertaken by ASIS and the University of Maryland (Carruthers *et al.* 2011) ranked the Atlantic subtidal habitats as reaching 99% of their reference condition (as determined by historical observations and data), albeit with very limited confidence or knowledge of indicators for the reference condition. Thus, data from the assessments by Versar, Inc. and the Maryland Geological Survey serve as a crucial pre-Sandy (2011) point of comparison for the possible effects of the physical stressors of hurricane Sandy on sediment distribution and benthic invertebrate assemblage structure. Here we make a detailed comparison of benthic assemblage patterns between the pre-Sandy (2011) and post-storm surveys a central part of the study, while considering inherent interpretive constraints due to the temporal scope of the project and the timing of sampling events around storms. While this project was one of four simultaneous benthic mapping regimes funded by the NPS in the aftermath of

Sandy, this study covered the largest habitat area, focused on the Atlantic subtidal zone, and was unique in its availability of pre-Sandy (2011) reference data.

The updated knowledge of benthic invertebrate assemblages at Assateague Island is intended to aid NPS staff in preserving park natural resources, as well as the historical and economic value of park activities. Benthic invertebrates provide ecosystem services—principally as crucial food sources for shelf fisheries (Molina *et al.* 2014)—that are necessary for the continued success of recreational fishing practices at ASIS. Benthos may also act as indicators of habitat health, since certain species are known to thrive in organically enriched or polluted conditions, while others perish (Pearson and Rosenberg, 1978; Pocklington and Wells, 1992 Dean, 2008).

In this paper we set out to: (1) provide an updated subtidal resource inventory to the National Park Service by determining benthic assemblage “ensemble” metrics including abundance, diversity, species richness, assemblage similarity, characteristic species, and associations with environmental variables; (2) analyze pre-Sandy (2011) and post-Sandy (2014–2015) assemblages with a uniform statistical approach and compared them to determine what changes occurred through the study period; (3) examine inter-annual variability in bottom type and macrobenthos over a four-year study to determine if any of the observed changes can likely be attributed to hurricane Sandy; (4) conduct targeted acoustic mapping of areas of particular interest before and after storms to provide an unprecedented higher resolution of these target locations.

1.3 Study Site

Assateague Island is a 58-km long barrier island stretching from the Ocean City inlet in Maryland, down past Chincoteague Island in northern Virginia. This dynamic island is characterized by barrier beach and back-bay environments that are subject to the constant eroding and re-shaping forces of the Atlantic Ocean. The pre-storm and post-storm surveys of Assateague Island subtidal resources discussed here include the majority of the length of the island (Figure 1). We sampled benthic sediments up to 1 km offshore in accordance with the pre-storm survey and NPS jurisdictions (Figure 2), whereas the USGS surveyed areas further offshore of Assateague Island. Historically the nearshore subtidal habitats at ASIS are comprised mostly of fine sand, with patches and swaths of mud and coarser sediments. Much of the observed heterogeneity in sediments is due to the relict sediments left behind by the characteristic landward ‘rolling-over’ of barrier islands.




Figure 1. Pre-Sandy (2011) survey sites (n=125) sampled in October 2011 on the beach (n=15) and subtidally up to 1 km offshore (n=110) of Assateague Island National Seashore, Maryland. Image taken from Google Earth Pro Landsat, 2016.



Figure 2. Post-storm (n=120 grabs) sampled in October 2014 (n=60 grabs, white) and 2015 (n=60 grabs, yellow), up to 1 km offshore of Assateague Island National Seashore survey sites, Maryland. Image taken from Google Earth Pro Landsat, 2016.

2. Materials and Methods

2.1. Acoustic Mapping: Side-scan Sonar Workflow

The side-scan sonar data collected was processed using Chesapeake Technology, Inc., SonarWiz sonar processing software. Sonar data processing followed industry standard procedures for side-scan data, focusing on gain corrections to remove the variation in across track brightness inherent in side-scan sonar performance in order to achieve the most representative mosaic of the seafloor. Standardized gain settings were chosen to make the sonar data as internally consistent as possible between different daily mission datasets, while also attempting to replicate the products produced from the Versar sonar data collected off Assateague in 2011. Raw Edgetech sonar files (.jsf) were imported using a time-varying gain, to compensate for signal loss on the outer swath of the sonar files. Once the files were imported, bottom-tracking corrections were applied if automatic bottom tracking had failed. Once verified or corrected, an empirical gain normalization correction was applied. This gain correction makes the data set consistent from a file-to-file basis, removing gain biases caused by variations in backscatter intensity from differing sediments between each file. This

produces an interiorly consistent mosaic. From this work flow (Figure 3), several mapping products were created. A standard georeferenced image file (geotiff) and Google Earth georeferenced file (.kmz) are generated. Vessel navigation or track-lines files were exported as shapefiles (.shp). Each gain-corrected file was exported as a waterfall image (.jpg). Target reports were generated (in .pdf format), highlighting unique features or objects on the seafloor. Finally, coverage reports (.xls) list the metadata for each file (start/end latitude & longitude, length, time, etc.), as well as the total area and line-line overlap achieved for the complete mission.




Figure 3. Side-scan sonar post processing workflow used by the University of Delaware, Assateague Island National Seashore mapping project: from EdgeTech raw data to SonarWiz to shapefiles and Google Earth data products.

2.2. Bathymetry Workflow

The bathymetric data collected was processed using Chesapeake Technology, Inc., SonarWiz sonar processing software. Sonar data processing followed industry standard procedures for bathymetric data, focusing on injecting tidal offsets, sound velocity offsets, and configuring the vessel files in order to achieve the most representative mosaic of the seafloor. The bathymetric processing workflow (Figure 4) was standardized to keep the bathymetric representations as internally consistent as possible. The first step was to configure a vessel offset file for the vessel that was recording the sonar data. All of the research vessels measurements were captured including the inertial measurement unit (I.M.U.) to the echosounder and the I.M.U to the antennas were recorded and saved in a Sonarwiz6 vessel file. After the vessel file is created, the raw sonar data (.jsf) was imported into a new Sonarwiz6 project that was set-up with an internally consistent naming convention based off the date of the survey. Before the data was merged, the sound velocity profiles, tide file, and the patch test values were imported into the project. After the echosounder was installed and before surveying operations began, a patch test was completed to ensure that each installation of the echosounder provided internally consistent data. The process accounts for offsets in roll, pitch, time latency, and heading. These patch test values were then entered into the vessel offsets file. In the field, sound velocity profiles were collected every hour on a Castaway CTD. The sound velocity files

(.csv) were then imported into the Sonarwiz6 project. A tide file from the days that we surveyed was downloaded from NOAA Tides and Current website in the form of .csv and then imported into the Sonarwiz6 project. After the patch test values, sound velocity




Figure 4. Schematic of workflow used by the University of Delaware mapping team for multibeam bathymetry data from EdgeTech 6205 acquisition to gridded data products. Data collected as part of a submerged mapping study for Assateague Island National Seashore in 2014-2015.

profiles, and the tide file were imported into the project, the data was merged. This merging process injects all of the above information onto the XYZ point cloud that was recorded during the survey to constrain the positional error, while providing the most replicable data products that are internally consistent. Gridded mosaics, at a resolution of .75 meters, were then exported as a .grd file using the CUBE algorithm, which is now the industry standard for providing the most accurate gridded solutions. The backscatter from the bathy was also processed in Sonarwiz6 and was exported as a .grd file. In all, a geotiff, .kmz, .xyz, and .grd of both the bathymetry data and the backscatter data were exported as bathymetric data products.

2.3. Acoustic Seabed Classification Map Preparation

Our three, 2014-2015 acoustic surveys allowed us to make both inter and intra-annual comparisons for storm-related changes in bottom sediments. The first survey resulted in side-scan sonar mosaics and sediment acoustic class shapefiles from June 2014. These data include 5 side-scan sonar mosaics and sediment class shapefiles that were generated from the side-scan sonar data collected during June 20 – 25, 2014 (Figure 6 upper plot). The second survey was the most broadly defined and resulted in side-scan sonar mosaics and sediment class shapefiles for May 2015 (Figure 8). These data include 12 side-scan sonar mosaics and sediment class shapefiles that were generated from the side-scan sonar data collected during May 12 – 21, 2015.

Finally, we conducted a set of spot selected surveys in October 2015 producing side-scan sonar mosaics and sediment class shapefiles. This survey was spatially focused on areas of change following hurricane Joaquin. Data were collected during October 13 – 16, 2015.

2.3.2. Acoustic Seabed Classification

First, all the available datasets were imported into ArcGIS platform under coordinate system - WGS_1984_UTM_Zone_18N. Then, manually digitized acoustic classes were matched against side-scan sonar mosaics and grain size data of grab samples to make sure the side-scan sonar mosaics were segmented correctly. After that, sediment class boundaries were matched between neighboring sediment classes to correct any inconsistencies in the interpretation of side-scan sonar mosaics, (i.e. to keep the class assignments consistent from one survey day to the next). Minor discrepancies with the side-scan sonar data interpretation and inconsistencies in sediment class boundaries were corrected manually using ArcGIS Edit tool.

After fixing all the discrepancies related with side-scan sonar data interpretation, all the same sediment classes in different shapefiles were merged into a single shapefile using the “Merge” tool in Arc Toolbox (Figure 5). Overlapping areas of the same sediment classes between different shapefiles were dissolved into one using the “Dissolve” tool in Arc Toolbox. At the end, 4 single shapefiles representing 4 different sediment classes namely Coarse Sand, Medium Sand, Fine Sand, and Mud, were merged into one single shapefile named Sediment Class_201406_201505.shp. A new field named “Area” was added to the table of contents of the newly generated shapefile and areas in m² were calculated for the sediment classes using “Calculate Geometry” method in ArcGIS.




Figure 5. Flow-chart of data processing steps for manually segmented side-scan sonar - sediment classification into a habitat map for Assateague Island National Seashore, developed by the University of Delaware 2014-2015.

The acoustic backscatter data collected in the field surveys was used to generate mosaic sonar images as outlined in the section above and then digitized using ArcGIS to manually outline and build categorical maps of the Assateague Island survey domain. The same acoustic class types as used in the 2011 precursor study by Versar/MGS were adopted here for consistency and the previous acoustic class maps were used along with the newly acquired and created side-scan sonar mosaics to guide the human operator in the segmentation process. All segmentation effort was conducted by the same team member for consistency and then reviewed by the principle investigator. The backscatter segments were then assigned to the acoustic class and were assigned similar colors again adopting the same nomenclature and color scheme as used in the precursor study to aid in comparison. Note that up to this point the classes are simply acoustic classes and require some external ground-truthing to further relate the classes to actual seabed types. The next step of relating the acoustic classes to seabed types comes through the correlation of class maps to the benthic grab samples and grain size analysis samples.

Figures 3 and 4 illustrate the workflows used in data processing both the side-scan sonar and phase measuring bathymetric sonar.

2.3.3. Ground Truthing

Essential for any geoacoustic marine habitat mapping initiative is the need to gather ground truthing observations to compare to the acoustically derived maps. Throughout our previous habitat mapping campaigns we have utilized benthic grab samples for ground truthing (Raineault et al., 2012; Trembanis et al., 2013). Grab samples were analyzed for grain size distribution and benthic faunal composition using standard approaches (see below and references therein for details). We used a modified Young grab sampler for recovering surface sediment ground truthing samples. The available grain size data included grain size analysis results from 98 grab samples taken during June of 2011 and 146 grab samples taken during October of 2014 and October of 2015 for benthic samples plus additional ones for ground truthing only. However, only the ones taken during 2014 and 2015 were used to check the side-scan sonar data interpretation in this study. Additionally, a small portable YSI Castaway CTD (<http://www.ysi.com/castaway>) was used to gather vertical profiles of

salinity, temperature, and sound speed for use in environmental characterization and swath bathymetry data processing.

3. Results

3.1. Acoustic mapping products and findings?

In total, over 65 km² of geoacoustic mapping was accomplished along the entire length of Assateague Island from as near to the shore as was possible out to at least 1km offshore. Additional sonar lines were acquired in order to connect the nearshore ASIS park boundary to the regional USGS mapping effort (Figure 6). The mapping area covered depths from 2 m-12 m extending from the beach to ~2 nm offshore, with final map resolution 50 cm/pixel for side-scan sonar and 1m²/pixel for bathymetry products. The field survey design as developed and agreed to by all project partners conducting mapping at this and other post-Sandy NPS sites was to achieve 100% side-scan sonar coverage and as much bathymetric coverage as possible given the constraints imposed by such shallow water survey operations. In this study we were able to achieve approximately 50% swath bathy coverage (Figure 7) and 100% side-scan sonar coverage (Figure 8).




Figure 6. Overview of study site area showing park boundary (green) and recent USGS coverage along with UD mapping team gap filling coverage (red).



319
320 Figure A7. Bathymetry mosaic (1m/pixel) from 2014-2015 acoustic surveys (550 kHz) for the entire
321 study area. Notable features captured in the bathymetry included pronounced shore-attached
322 oblique bars.

323 Resulting swath bathymetry data for the study side was gridded to a resolution of 1m/pixel.
324 Reliable bathymetry data was obtained for approximately half the total sonar swath width resulting
325 in 50% seafloor coverage in depths ranging from 2 to 12 m. Early sampling in 2014 around Fishing
326 Point was conducted using spacing resulting in 100% bathymetry coverage (200% side-scan sonar)
327 but the survey rate was unsustainable for the entire study site and it was agreed upon by all project
328 partners to expand the line spacing to achieve 100% side-scan and 50% bathymetry. This survey
329 strategy is also in keeping with the same approach used by the USGS in their larger regional
330 coverage provided for a meaningful connection of this study to that regional effort. The resulting
331 swath bathymetric grids extend what was only single-beam echosounder data from the precursor
332 2011 project and thus provide an enhanced baseline for AINS resource managers. Notable
333 features captured in the bathymetry included pronounced shore-attached oblique bars.




Figure 8. Side-scan mosaic from 2014-2015 acoustic surveys for the entire study area.

A complete set of side-scan sonar mosaics with a resolution of 50 cm/pixel was generated for the entire length of the island and included extensions to fill gaps between the ASIS park boundary and the USGS regional survey effort. Gain settings and post-processing were set to generate an internally consistent sonar backscatter map (Figure 8) that allowed for consistent segmentation and interpretation with respect to sediment class. The convention used in the backscatter mosaics is for bright yellow color for high-amplitude returns (i.e. coarse sand and gravel) and dark brown/black for low amplitude returns (i.e. mud). Bright high-amplitude returns in the backscatter mosaic were clearly associated with the raised features of shore-attached oblique sand ridges noted in the bathymetry (Figure 7). Mud patches were most notable in the adjoining troughs associated with the sand ridges or from relict marsh outcrops exposed by scour and barrier island migration.




Figure 9. Sediment classification from 2014–2015 for the entire study area. Four sediment classes (Coarse and medium sand, Mud and Fine Sand) were recognized and shaded with different colors. Mapping survey conducted 2014–2015.

As is shown in Figure 9, the sediment classification map includes four different sediment classes were recognized and shaded with different colors.

Sediment Type	Classification Map Coverage %
Coarse Sand	10%
Medium Sand	1%
Fine Sand	82%
Mud	7%

Table 1. Surficial sediment distribution by class type based on the 2014–2015 habitat mapping expedition.

Segmentation and acoustic classification of the side-scan sonar backscatter mosaic was performed via manual heads-up segmentation utilizing the same class types from the 2011 precursor study but informed by the sediment grab samples collected in this study. The GIS shapefile vectors for each acoustic class type are shown in Figure 9 using the same color and naming convention from 2011 for ease of comparison. The distribution of class type (Table 1) followed similar overall trends as documented in 2011 with medium and coarse sands associated with positive relief features like the large shore-attached oblique ridges and mud associated with the adjoining troughs. Notably absent in the 2014–2015 data were any mappable patches of gravel. The percentage area distribution of each class type are summarized in table 1.

The following three figures illustrate an example of an area with pronounced spatiotemporal changes in acoustic class texture between the 2011 precursor map (Figure 10) and repeated surveys with the EdgeTech 6205 in 2014 (Figure 11) and 2015 (Figure 12). The same backscatter amplitude convention is used but the gain settings and sonar frequencies are slightly different between the 2011 and 2014/15 surveys. Nevertheless, the general trend shows a migration of a set of coarse sand patches and the burial of a previously exposed mud patch.




Figure 10. Side-scan mosaic and classification shapefiles based on 2011 survey data.




Figure 11. May 2015 side-scan mosaic and sediment classification shapefile compared to 2011 from Figure 9.





Figure 12. October 2015 side-scan mosaic and sediment classification shapefile compared to May 2015 (Figure 11) and 2011 (Figure 10).

3.2. Temporal changes in seafloor sediment type

Select portions of the survey domain were mapped both in May 2015 and October 2015 capturing before and after the passage of hurricane Joaquin. Sediment class maps were determined in the same manner and using the same grain size informed class types in order to examine changes in the before and after storm surveys. Sediment types were coded from 1 to 4 from smallest to largest grain size with 1 as mud, 2 as fine sand, 3 as medium sand and 4 as coarse sand and these codes were manually inserted into the table of contents of sediment class shapefiles. Sediment class shapefiles were then converted into raster files with sediment class type code being the value field. In order to understand if the seafloor sediments are coarsening or fining from May 2015 to October 2015, a sediment property change map was generated by subtracting sediment class raster data of May 2015 from sediment class raster data of October 2015 (Figure 13). This map shows the pattern and intensity of change in sediment properties from May 2015 to October 2015 as a magnitude of sediment property change index (Figure 13). For example, positive sediment property change index means the seabed became coarser, negative sediment property change index means the seabed became finer and zero value means sediment property was unchanged. Higher absolute value means higher intensity of change, for example +3 area represents a change from mud to coarse sand -3 area represents a change from coarse sand to mud.



396

397 Figure 13. Sediment property change map for submerged resources of Assateague Island National
398 Seashore between May 2015 and October 2015. Positive sediment property change index means
399 seafloor became coarser, negative sediment property change index means seafloor became finer and
400 zero value means sediment texture did not change.

401 Comparison of sediment class boundaries between May 2015 and October 2015 suggest a major
402 shift (mostly south to eastward) in different sediment class boundaries. The distance of the shift in
403 the fine-coarse sediment class boundary is about 20 - 40 m in the northern portion of the study area
404 (Figure 14 and 15) and up to 80 m in the southern portion (Figures 16-19).The shift in mud-coarse
405 sediment class boundary however, was measured to be up to 180 m in the southern part (Figure 17).




Figure 14: Side-scan sonar mosaic interpretation, Assateague Island National Seashore, Maryland
(A) Side-scan sonar mosaics collected during May 13, 2015; (B) Interpretation of side-scan sonar
data collected in May 13, 2015; (C) Side-scan sonar mosaics collected in October 2015; (D)
Interpretation of side-scan sonar mosaics collected in October 2015. Note that red polygons
represent the same areas mapped during both May 13, 2015 and October 2015.




Figure 15: Temporal changes in sediment class boundaries, Assateague Island National Seashore, Maryland. The underlying image on the bottom is the side-scan sonar mosaic collected during May 2015. Shaded polygons on the top represent sediment classes inferred from the side-scan sonar mosaics collected during October 2015. Note the southeastward shift (18 - 50 m) in the boundary of fine-coarse sand sediment classes.




Figure 16: Side-scan sonar mosaic interpretation, Assateague Island National Seashore, Maryland. (A) Side-scan sonar mosaics collected during May 13, 2015; (B) Interpretation of side-scan sonar data collected in May 13, 2015; (C) side-scan sonar mosaics collected in October 2015; (D) Interpretation of side-scan sonar mosaics collected in October 2015. Note that red polygons represent the same areas mapped during both May 13, 2015 and October 2015.



427

428

429

430

431

432

433

Figure 17: Temporal changes in sediment class boundaries, Assateague Island National Seashore, Maryland. The profile in the bottom is the side-scan sonar mosaics collected during May 2015. Shaded polygons on the top represent sediment classes inferred from the side-scan sonar mosaics collected during October 2015. Please note the northward shift (180 m) in the boundary of mud-coarse sand sediment class and Southeastward shift (37- 40m) in the boundary of fine - coarse sand sediment class.




Figure 18: Side-scan sonar mosaic interpretation, Assateague Island National Seashore, Maryland. (A) Side-scan sonar mosaics collected during May 14, 2015; (B) Interpretation of side-scan sonar data collected in May 14, 2015; (C) Side-scan sonar mosaics collected in October 2015; (D) Interpretation of side-scan sonar mosaics collected in October 2015. Note that red polygons represent the same areas mapped during both May 14, 2015 and October 2015.



Figure 19: Temporal changes in sediment class boundaries, Assateague Island National Seashore, Maryland. The profile in the bottom is the side-scan sonar mosaics collected during May 2015. Shaded polygons on the top represent sediment classes inferred from the side-scan sonar mosaics collected during October 2015. Please note the west-south-eastward shift (38 - 57 m) in the boundaries of fine - coarse sand sediment classes.

3.3. Temporal changes in bathymetry

Storms and extreme weathering events (Morton, 2002 and Morton and Sallenger, 2003) or human induced disturbances such as dredging (Cooper et al., 2007) have the potential to change the nearshore bathymetry. These changes can have serious impacts to the marine flora and fauna that most marine environment protection efforts are focused on.

The purpose of the bathymetric change comparison is to see if it is possible to determine significant changes in the bathymetry after the passage of hurricane Joaquin, which passed through the area in late September and early October of 2015. A total of 18 bathymetric cross sections were constructed

using Interpolate Line tool in ArcScene across areas with overlapping survey coverage from before (May 2015) and after (October 2015) the passage of hurricane Joaquin. Nine of the cross sections (A1, B1 to J1) were made from May 2015 bathymetry, and the other nine cross section (A2, B2 to J2) were constructed from October 2015 bathymetry data along the same trackline as their pairs in May 2015. These bathymetry profiles were then exported as Excel spreadsheets. Sediment class type was added to each of the points along the bathymetry profiles based on the attributes from the acoustic sediment class maps discussed previously. Next, the Excel spreadsheets of the bathymetry profiles were imported into MatLab to generate comparison charts between pairs of bathymetry profiles. Sediment types were coded from 1 to 4: 1 as mud, 2 as fine sand, 3 as medium sand and 4 as coarse sand. Bathymetry and sediment class comparison charts were generated by subtracting sediment type values of May 2015 profiles from October 2015 profiles, in order to understand if the seafloor sediments are coarsening or fining with the lateral shift in bathymetry (Figures 20 and 21).

In the areas where cross sections were constructed more or less parallel to the direction of sediment class boundary shift, southwestward lateral shift of sand bars (up to 60 meters) were recognized in bathymetry cross section profiles (Figures 22 and 23). In the areas where cross sections were made more or less normal to the direction of sediment class boundary shift, vertical shifts of up to 65 centimeters are recognized (Figure 24). The largest vertical change was recognized in October 2015 profile J2 (Figure 24), which suggests axial incision of about 65 centimeters. This is the closest profile to the shoreline. Other profiles parallel to profile J2 suggests that vertical change in bathymetry seems to decrease with increasing distance from the shoreline.

Apart from the axial incision recognized in October 2015 profile J2 (Figure 24), comparison of May and October profiles suggests that seabed changed from fairly smooth in May 2015 to incised by shore oblique/perpendicular bars in October 2015(Figures 22 and 23). These are clear signs of hurricane effect on the seafloor. Decrease in vertical offset towards the sea may suggest an inverse relationship between hurricane effect on the seafloor and the distance from the shoreline. A Comparison between bathymetry change and sediment class boundary shift in this area suggests that sediments in the top of sand bars went coarser after the hurricane, while the ones in the trough of sand bars went finer (Figures 20 and 21).




Figure 20: Bathymetry profile change versus sediment class boundary shift after Hurricane Joaquin. Curved bathymetry profiles A1 (May 2015) and A2 (October 2015) are color coded according to the sediment types in the chart located in the top. Horizontal scale are meter in both panels, and vertical are meters in the top panel and size class index in the bottom panel. Sediment classes were indexed to 4, 1 representing the finest and 4 representing the coarsest. The chart in the bottom was generated by subtracting sediment type values of A1 (May 2015) from A2 (October 2015) to understand if the seafloor sediments are coarsening or fining with the lateral shift in bathymetry. Please note the coarsening of sediments in the top of the sand bars and fining of sediments in the trough. Inset scale is in meters on both axes.




Figure 21: Bathymetry profile change versus sediment class boundary shift after Hurricane Joaquin, Assateague Island National Seashore. Curved bathymetry profiles C1 (May 2015) and C2 (October 2015) in Fig. 25 are color coded according to the sediment types in the chart located in the top. Horizontal scale are meters in both panels, and vertical are meters in the top panel and size class index in the bottom panel. Sediment classes were indexed from 1 to 4, 1 representing the finest and 4 representing the coarsest. The chart in the bottom was generated by subtracting sediment type values of C1 (May 2015) from C2 (October 2015) to understand if the seafloor sediments are coarsening or fining with the lateral shift in bathymetry. Please note the coarsening of sediments in the top of the sand bars and fining of sediments in the trough. Inset scale is in meters on both axes.




Figure 22. Bathymetry profile change after Hurricane Joaquin, Assateague Island National Seashore, Maryland. Red curved line represents the bathymetry profile C1 in May 2015 and blue curved line represents the bathymetry profile C2 in October 2015. Please note the Southwestward lateral shift of sand bars in the bathymetry profile. Inset scale is in meters on both axes.




Figure 23: bathymetry profile change after Hurricane Joaquin. Red curved line represents the bathymetry profile G1 in May 2015 and blue curved line represents the bathymetry profile G2 in October 2015. Please note the change in surface roughness and the vertical offset in the bathymetry.




Figure 24: bathymetry profile change after Hurricane Joaquin. Red curved line represents the bathymetry profile J1 in May 2015 and blue curved line represents the bathymetry profile J2 in October 2015. Please note the axial incision in J2 profile.




Figure 25. Timeline of Storm events and seafloor mapping studies and resulting benthic habitat and morphology products.




Figure 26. Significant wave height (H_s) in meters (m) at offshore NDBC Buoy 44009 (38.461 N 74.703 W) between 2011-2016 including the passage of hurricanes Sandy and Joaquin near the study site.

4. Discussion

This nearshore benthic morphodynamic study at Assateague Island addresses a large gap in knowledge about the spatiotemporal dynamics of subtidal habitats and geomorphology in response to storm events (Figure 25). This knowledge will inform subsequent general management plans to best safeguard subtidal resources from anthropogenic and climate change stressors. This study also provides relevant data and recommendations for future assessments at ASIS.

Although storms have varied morphologic impacts on different benthic substrates (i.e. coarse sand versus mud), the ASIS nearshore zone experienced large lateral shifts on the order of 30 -70 m of sediment type contacts resulting in changes from fine to coarse sediment type with additional vertical depth changes from erosion and migration of between 50 cm to as much as 2 m associated with migration of nearshore oblique sand bars as a result the passage of storms.

We are not able to directly link the assemblage and habitat changes outlined here to Hurricane Sandy, excluding the possibility of impacts from other storms throughout the overall study period. This is largely due to the study timeline (Figure 26) and constraints of sampling logistics around unexpected storm events. It is more realistic to characterize the sum of assemblage differences between the two surveys as integrated changes over a multi-year scale. Despite the inherent uncertainty in the study, this benthic morphodynamic study is a valuable resource to inform management of publicly-valued and historically under-studied habitats.

Author Contributions: For research articles with several authors, a short paragraph specifying their individual contributions must be provided. The following statements should be used “conceptualization, X.X. and Y.Y.; methodology, X.X.; software, X.X.; validation, X.X., Y.Y. and Z.Z.; formal analysis, X.X.; investigation, X.X.; resources, X.X.; data curation, X.X.; writing—original draft preparation, X.X.; writing—review and editing, X.X.; visualization, X.X.; supervision, X.X.; project administration, X.X.; funding acquisition, Y.Y.”,

551 **Funding:** This study was funded by the National Park Service under CESU Task P14AC00380 and modification
552 14-0001.

553 **Acknowledgments:** Pre-storm data was provided by Versar, Inc. We thank the Assateague Island National
554 Seashore staff and project partners for their input, including Monique Lafrance-Bartley, Bill Hulslander, Charles
555 Roman, Mark Finkbeiner, Sara Stevens and Robin Baranowski. Field and lab work was made possible by
556 Captains Kevin Beam and Evan Falgowski, Ellie Rothermel, Hannah Rusch, Danielle Ferraro, Stephanie Dohner,
557 Tim Pileguard, Trevor Metz, Anna Schutschkow, Annie Daw, Jason Button, Leah Morgan, Drew Friedrichs,
558 Rachel Dalafave, Alec Halbruner, Meghan Owings, Rachel Oidtman, and Alex Itin. We thank the
559 Environmental Protection Agency Region III’s Ocean and Dredge Disposal Program Team, especially Renee
560 Searfoss and Sherilyn Morgan Lau, for the loan of a Young grab sampler, as well as the University of Delaware’s
561 Chris Sommerfield and Adam Marsh for the loans of additional laboratory and field equipment.

562 **Conflicts of Interest:** The authors declare no conflict of interest. The funders had no role in the design of the
563 study; in the collection, analyses, or interpretation of data; in the writing of the manuscript, or in the decision to
564 publish the results.
565

566

567 **References**

568 1. Abe, Hirokazu, Genki Kobayashi, and Waka Sato-Okoshi. 2015. Impacts of the 2011 tsunami on the subtidal
569 polychaete assemblage and the following recolonization in Onagawa Bay, northeastern Japan. *Marine*
570 *environmental research* 112: 86–95. doi:10.1016/j.marenvres.2015.09.011.

571 2. Aller, Josephine Y., and Ian Stupakoff. 1996. The distribution and seasonal characteristics of benthic
572 communities on the Amazon shelf as indicators of physical processes. *Continental Shelf Research* 16: 717–751.
573 doi:10.1016/0278-4343(96)88778-4.

574 3. Anderson, J.R., K. J. Storey, and R. Carolane. 1981. *Macrobenthic fauna and sediment data for eight estuaries on*
575 *the south coast of New South Wales, Technical Memorandum 81/22*. Canberra.

576 4. Anderson, M.J., R.N. Gorley, and K.R. Clarke. 2008. PERMANOVA+ for PRIMER: Guide to software
577 and statistical methods. Plymouth, U.K.: PRIMER-E.

578 5. Avila, L., and J. Cangialosi. 2011. *Tropical Cyclone Report: Hurricane Irene (AL092011)*.

579 6. Avila, Lixion A. 1998. *Preliminary Report: Hurricane Bonnie (AL021998)*. National Hurricane Center.
580 https://www.nhc.noaa.gov/data/tcr/AL021998_Bonnie.pdf

581 7. Balthis, W.L., J.L. Hyland, and D.W. Bearden. 2006. Ecosystem responses to extreme natural events: impacts
582 of three sequential hurricanes in fall 1999 on sediment quality and condition of benthic fauna in the Neuse
583 River Estuary, North Carolina. *Environmental monitoring and assessment* 119: 367–89. doi:10.1007/s10661-005-
584 9031-6.

585 8. Bartholomew, A. 2001. Polychaete Key for Chesapeake Bay and Coastal Virginia. Virginia Institute of
586 Marine Science. http://www.vims.edu/research/units/legacy/benthic_ecology/polychaete_key/index.php

587 9. Bender, Morris A, Thomas R Knutson, Robert E Tuleya, Joseph J Sirutis, Gabriel A Vecchi, Stephen T Garner,
588 and Isaac M Held. 2010. Modeled impact of anthropogenic warming on the frequency of intense Atlantic
589 hurricanes. *Science (New York, N.Y.)* 327: 454–8. doi:10.1126/science.1180568.

590 10. Berg, R. 2015. *Tropical Cyclone Report: Hurricane Arthur (AL012014)*.

591 11. Biernbaum, Charles K. 1979. Influence of sedimentary factors on the distribution of benthic amphipods of
592 Fishers Island Sound, Connecticut. *Journal of Experimental Marine Biology and Ecology* 38: 201–223.
593 doi:10.1016/0022-0981(79)90068-6.

594 12. Blake, E, T Kimberlain, R Berg, J Cangialosi, and J Beven. 2013. *Tropical Cyclone Report: Hurricane Sandy.*
595 *NOAA National Hurricane Center*.

596 13. Blott, S.J., and K. Pye. 2001. Gradistat: a grain size distribution and statistics package for the analysis of
597 unconsolidated sediments. *Earth Surface Processes and Landforms* 26: 1237–1248.

598 14. Buttolph, A. M., W. G. Grosskopf, G. P. Bass, and N. C. Kraus. 2006. Natural sand bypassing and response
599 of ebb shoal to jetty rehabilitation, Ocean City Inlet, MD, USA. 30th Coastal Engineering Conference. San
600 Diego, CA:3,344-3,356. World Scientific Press, Singapore.

601 15. Case, R. A. 1986. Annual Summary: Atlantic Hurricane Season of 1985. *Monthly Weather Review* 114: 1390–
602 1405.

603 16. Carruthers, T., K. Beckert, B. Dennison, J. Thomas, T. Saxby, M. Williams, T. Fisher, J. Kumer, C. Schupp,
604 B. Sturgis, and C. Zimmerman. 2011. Assateague Island National Seashore Natural Resource Condition
605 Assessment: Maryland, Virginia. Natural Resource Report NPS/ASIS/NRR—2011/405. National Park
606 Service, Fort Collins Colorado.

607 17. Clarke, K.R. and R.N. Gorley. 2015. PRIMER v7 User Manual. Plymouth, U.K.: PRIMER-E.

608 18. Clarke, K.R. and R.N. Gorley. 2006. PRIMER v6: User manual/tutorial. PRIMER-E: Plymouth.

609 19. Clarke, K.R. and R.M. Warwick. 2001. Change in marine communities: an approach to statistical analysis
610 and interpretation. 2nd. Edition. PRIMER-E: Plymouth.

611 20. Colden, A, and R Lipcius. 2015. Lethal and Sublethal effects of sediment burial on the eastern oyster
612 *Crassostrea virginica*. *Mar Ecol Progress Series* 527: 105–117.

613 21. Dauer, Daniel M., Catherine A. Maybury, and R.Michael Ewing. 1981. Feeding behavior and general
614 ecology of several spionid polychaetes from the chesapeake bay. *Journal of Experimental Marine Biology and*
615 *Ecology* 54: 21–38. doi:10.1016/0022-0981(81)90100-3.

616 22. Diaz, Robert J, Martin Solan, and Raymond M Valente. 2004. A review of approaches for classifying benthic
617 habitats and evaluating habitat quality. *Journal of environmental management* 73: 165–81.
618 doi:10.1016/j.jenvman.2004.06.004.

619 23. Dittmann, Sabine, C Günther, and U Schleier. 1999. Recolonization of Tidal Flats After Disturbance. In *The*
620 *Wadden Sea Ecosystem*, ed. Sabine Dittmann, 175–192. Berlin, Heidelberg: Springer Berlin Heidelberg.
621 doi:10.1007/978-3-642-60097-5.

24. Dean, H. K. 2008. The use of polychaetes (Annelida) as indicator species of marine pollution: a review. *Journal of Tropical Biology* 56: 11-38. <http://www.redalyc.org/articulo.oa?id=44919934004>
25. Dernie, K.M, M.J Kaiser, E.A Richardson, and R.M Warwick. 2003. Recovery of soft sediment communities and habitats following physical disturbance. *Journal of Experimental Marine Biology and Ecology* 285-286: 415–434. doi:10.1016/S0022-0981(02)00541-5.
26. Douvere F., and Ehler C. 2009. Ecosystem-based Marine Spatial Management: An Evolving Paradigm for the Management of Coastal and Marine Places. *Ocean Yearbook*, Volume 23:-26
27. Downes, B.J., L.A. Barmuta, P.G. Fairweather, D.P. Faith, M.J. Keough, P.S. Lake, B.D. Mapstone and G.P. Quinn. 2002. Monitoring Ecological Impacts. Concepts and Practice in Flowing Waters. Cambridge.
28. Dreyer, J, J H Bailey-Brock, and S A McCarthy. 2005. The immediate effects of Hurricane Iniki on intertidal fauna on the south shore of O'ahu. *Marine Environmental Research* 59: 367–80. doi:10.1016/j.marenvres.2004.04.006.
29. Engle, Virginia D, Jeffrey L Hyland, and Cynthia Cooksey. 2009. Effects of Hurricane Katrina on benthic macroinvertebrate communities along the northern Gulf of Mexico coast. *Environmental monitoring and assessment* 150: 193–209. doi:10.1007/s10661-008-0677-8.
30. Etter, Ron J., and J. Frederick Grassle. 1992. Patterns of species diversity in the deep sea as a function of sediment particle size diversity. *Nature* 360: 576–578. doi:10.1038/360576a0.
31. Federal Geographic Data Committee. 2012. Coastal and Marine Ecological Classification Standard, June 2012. FGDC-STD-018-2012. Reston, VA: Federal Geographic Data Committee
32. Flemer, David A., Barbara F. Ruth, and Charles M. Bundrick. 2016. Effects of sediment type on macrobenthic infaunal colonization of laboratory microcosms. *Hydrobiologia* 485. Kluwer Academic Publishers: 83–96. doi:10.1023/A:1021385901210.
33. Fonseca, L., and Mayer, L.A., 2007. Remote estimation of surficial seafloor properties through the application of Angular Range Analysis to multibeam sonar data, *Marine Geophysical Researches*, 28(2):119-126, DOI 10.1007/s11001-007-9019-4
34. Forrest, A.L., Trembanis, A.C., and Todd, W.L. 2012. Ocean floor mapping as a precursor for space exploration. *J. Ocean Tech.* 7(2):71–87.
35. Frangipane, Gretel, Mario Pistolato, Emanuela Molinaroli, Stefano Guerzoni, and Davide Tagliapietra. 2009. Comparison of loss on ignition and thermal analysis stepwise methods for determination of sedimentary organic matter. *Aquatic Conservation: Marine and Freshwater Ecosystems* 19: 24–33. doi:10.1002/aqc.970.
36. Goff, John A., James A. Austin, Roger D. Flood, Beth Christensen, Cassandra M. Browne, and Steffen Saustrup. 2013. Rapid Response Survey Gauges Sandy's Impact on Seafloor. *Eos, Transactions American Geophysical Union* 94: 337–338. doi:10.1002/2013EO390001.
37. Goodsell, P.J., and S.D Connell. 2005. Historical configuration of habitat influences the effects of disturbance on mobile invertebrates. *Mar Eco Progress Series* 299: 79-87.
38. Gosner, Kenneth L. 1978. *A Field Guide to the Atlantic Seashore: From the Bay of Fundy to Cape Hatteras*. Edited by R. T. Peterson. Houghton Mifflin Harcourt.
39. Green, M.O., Vincent, C.E., and Trembanis, A.C. 2004. Suspension of coarse and fine sand on a wave-dominated shoreface, with implications for the development of rippled scour depressions. *Continental Shelf Research* vol. 24 pp. 317-335.
40. Hapke, Cheryl J., Hilary F. Stockdon, William C. Schwab, and Mary K. Foley. 2013. Changing the Paradigm of Response to Coastal Storms. *Eos, Transactions American Geophysical Union* 94: 189–190. doi:10.1002/2013EO210001.
41. Hauser, O. 2002. Tubeworm nodule reefs of Delaware Bay: An evaluation of two acoustic survey methods and lab experiments of nodule formation. University of Delaware M.S. Thesis.
42. Hedges, J.I., and R.G. Keil. 1995. Sedimentary organic matter preservation: an assessment and speculative synthesis. *Marine Chemistry* 49: 81–115.
43. Hendrick, Vicki J, Zoë L Hutchison, and Kim S Last. 2016. Sediment Burial Intolerance of Marine Macroinvertebrates. *PloS one* 11. Public library science, 1160 Battery Street, ste 100, San Francisco, CA 94111 USA: e0149114. doi:10.1371/journal.pone.0149114.
44. Hernández-Arana, Hector A., Ashley A. Rowden, Martin J. Attrill, Richard M. Warwick, and Gerardo Gold-Bouchot. 2003. Large-scale environmental influences on the benthic macroinfauna of the southern Gulf of Mexico. *Estuarine, Coastal and Shelf Science* 58: 825–841. doi:10.1016/S0272-7714(03)00188-4.
45. Holland, K.T., Keen, T., and Kaihatu, J.M. 2003. Understanding coastal dynamics in heterogeneous sedimentary environments. *Coastal Sediments '03*, Clearwater Beach, FL.
46. Hoogsteen, M. J. J., E. A. Lantinga, E. J. Bakker, J. C. J. Groot, and P. A. Tittone. 2015. Estimating soil organic carbon through loss on ignition: effects of ignition conditions and structural water loss. *European Journal of Soil Science* 66: 320–328. doi:10.1111/ejss.12224.

47. Hughes, C., C.A. Richardson, M. Luckenbach, and R. Seed. 2009. Difficulties in separating hurricane induced effects from natural benthic succession: Hurricane Isabel, a case study from Eastern Virginia, USA. *Estuarine, Coastal and Shelf Science* 85: 377–386. doi:10.1016/j.ecss.2009.08.023.
48. Hurrell, J., and National Center for Atmospheric Research staff (Eds). 2016. The Climate Data Guide: Hurrell North Atlantic Oscillation (NAO) Index (station-based).
49. Huston, M. 1979. A General Hypothesis of Species Diversity. *American Naturalist* 113: 81–101.
50. Johnson, Ralph Gordon. 1971. Animal-sediment relations in shallow water benthic communities. *Marine Geology* 11: 93–104. doi:10.1016/0025-3227(71)90069-7.
51. Kirby-Smith, William W., and Joseph Ustach. 1986. Resistance to hurricane disturbance of an epifaunal community on the continental shelf off North Carolina. *Estuarine, Coastal and Shelf Science* 23: 433–442. doi:10.1016/0272-7714(86)90001-6.
52. Knutson, Thomas R., Joseph J. Sirutis, Stephen T. Garner, Gabriel A. Vecchi, and Isaac M. Held. 2008. Simulated reduction in Atlantic hurricane frequency under twenty-first-century warming conditions. *Nature Geoscience* 1. Nature Publishing Group: 359–364. doi:10.1038/ngeo202.
53. Kröncke, I, JW Dippner, H Heyen, and B Zeiss. 1998. Long-term changes in macrofaunal communities off Norderney (East Frisia, Germany) in relation to climate variability. *Marine Ecology Progress Series* 167: 25–36. doi:10.3354/meps167025.
54. Kröncke, Ingrid, Bernd Zeiss, and Christa Rensing. 2001. Long-term variability in macrofauna species composition off the island of Norderney (East Frisia, Germany) in relation to changes in climatic and environmental conditions. *Senckenbergiana maritima* 31: 65–82. doi:10.1007/BF03042837.
55. Lawrence, M.B. 1987. Annual Summary: Atlantic Hurricane Season of 1986. *Monthly Weather Review* 115: 2155–2160.
56. Little, Colin. 2000. *The Biology of Soft Shores and Estuaries. Biology of Habitats*. 1st ed. New York: Oxford University Press.
57. Lomovasky, Betina J., Fausto N. Firstater, Alex Gamarra Salazar, Jaime Mendo, and Oscar O. Iribarne. 2011. Macro benthic community assemblage before and after the 2007 tsunami and earthquake at Paracas Bay, Peru. *Journal of Sea Research* 65: 205–212. doi:10.1016/j.seares.2010.10.002.
58. Magurran, A.E and B.J. McGill, eds. 2011. *Biological Diversity: Frontiers in Measurement and Assessment*. Oxford.
59. Mallin, M, and C Corbett. 2006. How Hurricane Attributes Determine the Extent of Environmental Effects: Multiple Hurricanes and Different Coastal Systems. *Estuaries and Coasts* 29: 1046–1061.
60. Mancinelli, G., S. Fazi, and L. Rossi. 2016. Sediment structural properties mediating dominant feeding types patterns in soft-bottom macrobenthos of the Northern Adriatic Sea. *Hydrobiologia* 367. Kluwer Academic Publishers: 211–222. doi:10.1023/A:1003292519784.
61. Miller, D., N. Raineault, and A. Trembanis, 2010. Mapping Patchy Hard-Bottom Habitats with Multiple Technologies from Ship, ROV and Autonomous Underwater Vehicles in Delaware Bay. AGU Ocean Sciences meeting.
62. Miselis, Jennifer L., Brian D. Andrews, Robert S. Nicholson, Zafer Defne, Neil K. Ganju, and Anthony Navoy. 2016. Evolution of Mid-Atlantic Coastal and Back-Barrier Estuary Environments in Response to a Hurricane: Implications for Barrier-Estuary Connectivity. *Estuaries and Coasts*. doi:10.1007/s12237-015-0057-x.
63. Molina, Pilar, F. Patricio Ojeda, Marcela Aldana, V.M. Pulgar, M. Roberto García-Huidobro, and José Pulgar. 2014. Spatial and temporal variability in subtidal macroinvertebrates diversity patterns in a management and exploitation area for benthic resources (MEABRs). *Ocean & Coastal Management* 93: 121–128. doi:10.1016/j.ocecoaman.2014.03.005.
64. Munroe, D., A. Tabatabai, I. Burt, D. Bushek, E.N. Powell, and J. Wilkin. 2013. Oyster mortality in Delaware Bay: Impacts and recovery from Hurricane Irene and Tropical Storm Lee. *Estuarine, Coastal and Shelf Science* 135: 209–219. doi:10.1016/j.ecss.2013.10.011.
65. National Hurricane Center, National Oceanic and Atmospheric Association. 2011. 2011 Atlantic Hurricane Season. <http://www.nhc.noaa.gov/data/tcr/index.php?season=2011&basin=atl>
66. National Hurricane Center, National Oceanic and Atmospheric Association. 2012. 2012 Atlantic Hurricane Season. <http://www.nhc.noaa.gov/data/tcr/index.php?season=2012&basin=atl>
67. National Hurricane Center, National Oceanic and Atmospheric Association. 2013. 2013 Atlantic Hurricane Season. <http://www.nhc.noaa.gov/data/tcr/index.php?season=2013&basin=atl>
68. National Hurricane Center, National Oceanic and Atmospheric Association. 2014. 2014 Atlantic Hurricane Season. <http://www.nhc.noaa.gov/data/tcr/index.php?season=2014&basin=atl>
69. National Hurricane Center, National Oceanic and Atmospheric Association. 2015. 2015 Atlantic Hurricane Season. <http://www.nhc.noaa.gov/data/tcr/index.php?season=2015&basin=atl>

70. Natural History Museum, London. 1999. Polychaete Identification Keys. <http://www.nhm.ac.uk/research-curation/research/projects/taxinfo/index2.html>
71. Negrello Filho, Orlei A., and Paulo C. Lana. 2013. Short-term stability of estuarine benthic assemblages: are storms pattern-defining events? *Zoologia (Curitiba)* 30. Sociedade Brasileira de Zoologia: 266–272. doi:10.1590/S1984-46702013000300003.
72. Pansu, M., and J. Gauthierou. 2006. *Handbook of Soil Analysis: Mineralogical, Organic and Inorganic Methods*. Springer-Verlag Berlin Heidelberg.
73. Parnum, I., J. Siwabessy, A. Gavrilov, and M. Parsons. 2009. A comparison of single beam and multibeam sonar systems in seafloor habitat mapping. *Underwater Acoustic Measurements*.
74. Pasch, R.J., and L. A. Avila. 1992. Annual Summary: Atlantic Hurricane Season of 1991. *Monthly Weather Review* 120: 2671–2687.
75. Pearson, T. H., and R. Rosenberg. 1977. Pearson TH, Rosenberg R. Macrobenthic succession in relation to organic enrichment and pollution of the marine environment. *Oceanogr Mar Biol Ann Rev* 16: 229–311. *Oceanography and marine biology* 16.
76. Pedrosa-Pàmies, R., A. Sanchez-Vidal, A. Calafat, M. Canals, and R. Durán. 2013. Impact of storm-induced remobilization on grain size distribution and organic carbon content in sediments from the Blanes Canyon area, NW Mediterranean Sea. *Progress in Oceanography* 118: 122–136. doi:10.1016/j.pocean.2013.07.023.
77. Pocklington, Patricia, and P.G. Wells. 1992. Polychaetes: Key Taxa for Marine Environmental Quality Monitoring. *Marine Pollution Bulletin* 24: 593–598.
78. Pollock, Leland W. 1998. *A Practical Guide to the Marine Animals of Northeastern North America*. Rutgers University Press.
79. Posey, M., W. Lindberg, T. Alphin, and F. Vose. 2016. Influence of storm disturbance on an offshore benthic community. *Bulletin of Marine Science* 59. Allen: 523–529. <http://cat.inist.fr/?aModele=afficheN&cpsidt=2489234>. Accessed April 20.
80. Probert, P. K. 1984. Disturbance, sediment stability, and trophic structure of soft-bottom communities. *Journal of Marine Research* 42. Sears Foundation for Marine Research: 893–921. doi:10.1357/002224084788520837.
81. Quester Tangent. 2010. QTC SWATHVIEW 1.0 User Manual and Reference: DMN-SWVW-0000-R00. Saanichton, British Columbia, Canada.
82. Quinn, G.P and M.J. Keough. 2002. *Experimental Design and Data Analysis for Biologists*. Cambridge.
83. Ramey, Patricia A., and E. Bodnar. 2008. Selection by a deposit-feeding polychaete, *Polygordius jouinae*, for sands with relatively high organic content. *Limnology and Oceanography* 53: 1512–1520. doi:10.4319/lo.2008.53.4.1512.
84. Raineault, Nicole A., Arthur C. Trembanis, and Douglas C. Miller. 2011. Mapping Benthic Habitats in Delaware Bay and the Coastal Atlantic: Acoustic Techniques Provide Greater Coverage and High Resolution in Complex, Shallow-Water Environments. *Estuaries and Coasts* 35: 682–699. doi:10.1007/s12237-011-9457-8.
85. Raineault, Nicole A., Arthur C. Trembanis, Douglas C. Miller, and Vince Capone. 2013. Interannual changes in seafloor surficial geology at an artificial reef site on the inner continental shelf. *Continental Shelf Research* 58: 67–78. doi:10.1016/j.csr.2013.03.008.
86. Rees, E.I.S., A. Nicholaidou, and P. Laskaridou. 1977. The effects of storms on the dynamics of shallow water benthic associations. In *Biology of Benthic Organisms: 11th European Symposium on Marine Biology*, ed. B. F. Keegan, P. J. S. Boaden, and P. O. Ceidigh, 465–474. Galway, Ireland: Elsevier. doi:10.1016/B978-0-08-021378-1.50052-X.
87. Salen-Picard, C., A. Darnaude, D. Arlhac, and M. Harmelin-Vivien. 2002. Fluctuations of macrobenthic populations: a link between climate-driven river run-off and sole fishery yields in the Gulf of Lions. *Oecologia* 133: 380–388. doi:10.1007/s00442-002-1032-3.
88. Sánchez, María A, María L Jaubet, Griselda V Garaffo, and Rodolfo Elías. 2013. Spatial and long-term analyses of reference and sewage-impacted sites in the SW Atlantic (38°S, 57°W) for the assessment of sensitive and tolerant polychaetes. *Marine Pollution Bulletin* 74: 325–33. doi:10.1016/j.marpolbul.2013.06.033.
89. Santos, M.F.L.d., and A.M.S. Pires-Vanin. 2004. Structure and dynamics of the macrobenthic communities of ubatuba bay, southeastern brazilian coast. *Brazilian Journal of Oceanography* 52: 59–73.
90. Seiderer, L.J., and R.C. Newell. 1999. Analysis of the relationship between sediment composition and benthic community structure in coastal deposits: Implications for marine aggregate dredging. *ICES Journal of Marine Science*, 56: 757–765.
91. Slacum, H.W. JR., W.H. Burton, E. T. Methratta, E. D. Weber, R. J. Llansó, and J. Dew-Baxter. 2010. Assemblage structure in shoal and flat-bottom habitats on the inner Continental Shelf of the Middle

- Atlantic Bight, USA. *Marine and Coastal Fisheries: Dynamics, Management, and Ecosystem Science* 2:277–298.
92. Slacum, H.W. Jr., J.H. Vølstad, E. Weber, W. Richkus, R. Diaz, and C. Tallent. 2008. The value of applying commercial fishers' experience with designed surveys to identify characteristics of essential fish habitat for adult summer flounder (*Paralichthys dentatus*). *North American Journal of Fisheries Management*. Volume 28(3):710-721.
93. Slacum H.W., Jr., R. Llansó, J. Dew, and L. Methratta. 2009. An Assessment and Characterization of Benthic Habitats Delineated by Acoustic Mapping in Chesapeake Bay. Prepared for NOAA Chesapeake Bay Office, Annapolis, Maryland, by Versar, Inc., Columbia, Maryland. Contract DG 133F-05-NC-3216
94. Snyder, Noah P. 2004. *Sediment grain-size and loss-on-ignition analyses from 2002 Englebright Lake coring and sampling campaigns*. USGS Open-File Report 2004-1080.
95. Sopkin, K.L., H.F. Stockdon, K.S. Doran, N.G. Plant, K.L.M. Morgan, K.K. Guy, and K.E.L. Smith. 2014. *Hurricane Sandy—Observations and Analysis of Coastal Change: USGS Open-File Report 2014–1088*. Reston, VA. doi:http://dx.doi.org/10.3133/ofr20141088.
96. Sousa, W.P. 2001. Natural disturbance and the dynamics of marine benthic communities. In *Marine Community Ecology*, ed M.D. Bertness, S.D. Gaines, and M.E. Hay, 85-130. Sunderland, MA: Sinauer Associates.
97. Subramaniam, S., H. Barad, A.B. Martinez, and B. Bourgeois, Seafloor Characterization using Texture. 1993. Southeastcon '93, Proceedings, IEEE, pp. 8.
98. Taghon, Gary L., Patricia A. Ramsey, Charlotte M. Fuller, Rosemarie F. Pectrecca, and Judith P. Grassle. 2016. Benthic Community Structure and Sediment Properties in Barnegat Bay, New Jersey, Before and After Hurricane Sandy. *Estuaries and Coasts*. doi:10.1007/s12237-016-0133-x.
99. Thrush, Simon F., John S. Gray, Judi E. Hewitt, and Karl I. Ugland. 2006. Predicting the effects of habitat homogenization on marine biodiversity. *Ecological Applications* 16: 1636–1642. doi:10.1890/10510761(2006)016[1636:pteohh]2.0.CO;2.
100. Trembanis, Arthur, Carter DuVal, Jonathan Beaudoin, Val Schmidt, Doug Miller, and Larry Mayer. 2013. A detailed seabed signature from Hurricane Sandy revealed in bedforms and scour. *Geochemistry, Geophysics, Geosystems* 14: 4334–4340. doi:10.1002/ggge.20260.
101. Trembanis AC, Forrest AL, Miller DC, Lim DSS, Gernhardt ML, Todd WL (2012) Multiplatform ocean exploration: insights from the NEEMO space analog mission. *J Mar Tech Soc* 46(4):7–19.
102. Trembanis, A.C., Gutsche, J., and Nebel, S. 2010. Precursor exploration missions in Kelly Lake, British Columbia-MARSLIFE project: San Francisco, American Geophysical Union, Fall 2010 Meeting, abstract P13B-1399.
103. Trembanis, A.C., J. Walker, D. Miller, B. Phoel, 2012. A High-Density Image Analysis of Sea Scallop Habitat in the Mid-Atlantic Bight. GeoHab 2012 Conference, May 1-4, 2012. Orcas Island, WA.
104. Trembanis, A.C., J.H. Walker, A. Ochse, K. Ochse. 2012. A demonstration
105. sea scallop survey of the Federal Inshore Areas of the New York Bight using a
106. camera mounted autonomous underwater vehicle. 181 pages. Final report, submitted to NMFS August, 2012.
107. Trembanis, A.C., Wright, L.D., Friedrichs, C.T., Green, M.O., and Hume, T. 2004. The effects of spatially complex inner shelf roughness on boundary layer turbulence and current and wave friction: Tairua embayment, New Zealand. *Cont. Shelf Res.* 24:1549-1571.
108. Van Hoey, Gert, Steven Degraer, and Magda Vincx. 2004. Macrobenthic community structure of soft-bottom sediments at the Belgian Continental Shelf. *Estuarine, Coastal and Shelf Science* 59: 599–613. doi:10.1016/j.ecss.2003.11.005.
109. Vaughan, Nancy D, Thomas C Johnson, David L Mearns, Albert C Hine, William W Kirby-Smith, Joseph F Ustach, and Stanley R Riggs. 1987. The impact of Hurricane Diana on the north Carolina continental shelf. *Marine Geology* 76: 169–176. doi:10.1016/0025-3227(87)90026-0.
110. Vecchi, Gabriel A., and Thomas R. Knutson. 2011. Estimating Annual Numbers of Atlantic Hurricanes Missing from the HURDAT Database (1878–1965) Using Ship Track Density. *Journal of Climate* 24: 1736–1746.
111. Vecchi, Gabriel A., and Thomas R. Knutson. 2008. On Estimates of Historical North Atlantic Tropical Cyclone Activity. *Journal of Climate* 21: 3580–3600. doi:10.1175/2008jcli2178.1.
112. Versar C-MAP. 2013. Assateague Island National Seashore Atlantic Subtidal and Beach Benthic Biological Resource Inventory and Access database, unpublished final reporting to the National Park Service. Versar, Inc. Columbia, MD.
113. Warwick, R.M., and K.R. Clarke. 1993. Increased variability as a symptom of stress in marine communities. *Journal of Experimental Marine Biology and Ecology* 172: 215–226. doi:10.1016/0022-0981(93)90098-9.

853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871

114. Whitlatch, Robert. 1977. Seasonal changes in the community structure of the macrobenthos inhabiting the intertidal sand and mud flats of Barnstable Harbor, Massachusetts. *The Biological Bulletin* 152.
115. Whitlatch, Robert B. 1981. Animal-sediment relationships in intertidal marine benthic habitats: Some determinants of deposit-feeding species diversity. *Journal of Experimental Marine Biology and Ecology* 53: 31–45. doi:10.1016/0022-0981(81)90082-4.
116. Wigley, R.L. and R.B. Theroux. 1981. Atlantic continental shelf and slope of the United States—macrobenthic invertebrate fauna of the Middle Atlantic Bight region—faunal composition and quantitative distribution. Geologic Survey Professional Paper 529-N. USGPO, Washington, DC.
117. Wilber, D.H., G.L. Ray, D.G. Clarke, and R.J. Diaz. 2008. Responses of Benthic Infauna to Large-Scale Sediment Disturbance in Corpus Christi Bay, Texas. *Journal of Experimental Marine Biology and Ecology* 365: 13–22. doi:10.1016/j.jembe.2008.07.029.
118. Williams, K.L. 2001. An effective density fluid model for acoustic propagation in sediments derived from Biot theory. *J. Acoust. Soc. Am.* 110:2276-2281.
119. Wilson, B.D. and J.A. Madsen. 2006. Acoustical methods for bottom and sub-bottom imaging in estuaries: benthic mapping project to identify and map the bottom habitat and sub-bottom sediments of Delaware Bay. *Sea Technologies* 47(6):43-46.
120. Zajac, Roman N., Robert B. Whitlatch, and Simon F. Thrush. 1998. Recolonization and succession in soft-sediment infaunal communities: the spatial scale of controlling factors. *Hydrobiologia* 375. Kluwer Academic Publishers: 227–240. doi:10.1023/A:1017032200173.s