

1 Article

# 2 Performance of winter wheat cultivars grown 3 organically and conventionally with focus on 4 Fusarium head blight and *Fusarium* trichothecene 5 toxins

6 Tomasz Góral <sup>1\*</sup>, Aleksander Łukanowski <sup>2</sup>, Elżbieta Małuszyńska <sup>3</sup>, Kinga Stuper-Szablewska <sup>4</sup>,  
7 Maciej Buśko <sup>4</sup> and Juliusz Perkowski <sup>4</sup>

8 <sup>1</sup> Department of Plant Pathology, Plant Breeding and Acclimatization Institute – National Research Institute,  
9 Radzików, 05-870 Błonie, Poland

10 <sup>2</sup> Department of Phytopathology and Molecular Mycology, Faculty of Agriculture and Biotechnology, UTP  
11 University of Sciences and Technology, Al. prof. S. Kaliskiego 7, bldg. I, 85-796 Bydgoszcz, Poland

12 <sup>3</sup> Department of Seed Science and Technology, Plant Breeding and Acclimatization Institute – National  
13 Research Institute, Radzików, 05-870 Błonie, Poland

14 <sup>4</sup> Department of Chemistry, Poznań University of Life Sciences, ul. Wojska Polskiego 75, 60-625 Poznań,  
15 Poland

16 \*Correspondence: t.goral@ihar.edu.pl

17 **Abstract:** Growing acreage and changing consumer preferences cause increasing interest in the  
18 cereal products originating from organic farming. Lack of results of objective test, however, does  
19 not allow drawing conclusions about the effects of cultivation in the organic system and comparison  
20 to currently preferred conventional system. Field experiment was conducted in organic and  
21 conventional fields. Thirty modern cultivars of winter wheat were sown. They were characterized  
22 for disease infection including Fusarium head blight, seed sowing value, the amount of DNA of the  
23 six species of *Fusarium* fungi as well as concentration of ergosterol and trichothecenes in grain. The  
24 intensity Fusarium head blight was at a similar level in both systems. However, *Fusarium*  
25 colonization of kernels expressed as ergosterol level or DNA concentration was higher for the  
26 organic system. It did not reflect in an increased accumulation of trichothecenes in grain, which was  
27 similar in both systems, but sowing value of organically produced seeds was lower. Significant  
28 differences between analyzed cropping systems and experimental variants were found. The  
29 selection of the individual cultivars for organic growing in terms of resistance to diseases and  
30 contamination of grain with *Fusarium* toxins was possible. Effects of organic growing differ  
31 significantly from the conventional and grain obtained such way can be recommended to  
32 consumers. There are indications for use of particular cultivars bred for conventional agriculture in  
33 the case of organic farming, and the growing organic decreases plant stress resulting from intense  
34 fertilization and chemical plant protection.

35 **Keywords:** Fusarium head blight; *Fusarium* species; soil minerals; mycotoxins; organic farming;  
36 sowing value; winter wheat

## 38 1. Introduction

39 A way of growing crops is changing because of the geopolitical situation and consumer  
40 preferences. In recent years, high interest in organic farming has been observed in Europe  
41 (<https://ec.europa.eu/eurostat/statistics-explained/pdfscache/5461.pdf>). In 2002, organic farming took  
42 up 5.0 million hectares, while in 2017 it was 12.6 million hectares. Austria, Estonia, Sweden, Italy,  
43 Czech Republic and Latvia were the countries with the highest share of organic farmland, while the  
44 largest areas of organic farmland were in Spain, Italy, France and Germany. In Poland in 2016 it was  
45 536.6 thousand ha – 3.7% of all agricultural land [1].

46 This is due to the awareness that in organic farming practices the use of artificial fertilizers as  
47 well as pesticides is not allowed. There is limited list of substances, which can be used as natural  
48 fungicides to protect crops against fungal diseases. Lack of fungicide protection can result in higher  
49 severity of fungal diseases. Chemical seed treatment is not applied which leads to increased incidence  
50 of seed borne diseases [2,3]. Thus, seed transmitted diseases are considered the most harmful in  
51 organic farming. Leaf diseases (not seed transmitted) and foot rots are less important. Severity of  
52 these diseases correlates with high nitrogen doses and high crop density, so under organic farming  
53 conditions they are less damaging [4]. Diseases caused by fungi surviving on crop debris (including  
54 *Fusarium* head blight) can be controlled by cultural practices, so they are less damaging than seed  
55 borne ones [3]. However, *Fusarium* fungi causing *Fusarium* head blight are able to produce toxic  
56 secondary metabolites – mycotoxins contaminating grain. The main *Fusarium* species causing  
57 *Fusarium* head blight are *F. culmorum*, *F. graminearum* and *F. avenaceum* [5,6]. Cereal heads are  
58 infected mainly during the flowering period [7]. This is the stage where cereals are the most  
59 susceptible to infection with *Fusarium* fungi spores. After infection, the fungus develops in infected  
60 flower spreading then to other flowers in the spikelet. Then through rachis, the fungus spreads to  
61 another spikelets causing necrosis and bleaching individual spikelets [8,9]. The invaded cereal grain,  
62 even visually healthy looking, is contaminated with fungal mycotoxins, which are phyto- and  
63 zootoxic. *Fusarium* spp. affecting cereals are known as potent producers of type A trichothecenes (T-  
64 2 and HT-2 toxins, diacetoxyscirpenol et al.) and of type B (deoxynivalenol, nivalenol et al.) as well  
65 as moniliformin, zearalenone, enniatins, beauvericin and the other toxins [5,6,10].

66 Avoiding the presence of *Fusarium* mycotoxins in food is very important, thus organic food is  
67 perceived as “food without chemistry” of higher quality than conventional [11,12]. In the literature,  
68 you can find analyses on this issue comparing wheat from organic and conventional cropping  
69 systems [13–17]. Mäder et al. [18] analyzed *Fusarium* metabolites, deoxynivalenol (DON) and  
70 nivalenol (NIV), content in wheat grain produced in a 21-year conventional and organic agrosystems.  
71 It was found higher concentration of DON in samples from conventional fields in both years of  
72 mycotoxin analysis; however, differences were not significant. NIV concentration was similar in both  
73 cropping systems. Magkos et al. [19] in their review summarized results of 12 papers on  
74 contamination of organic and conventional cereals with *Fusarium* mycotoxins. Organically grown  
75 cereals has been reported to be either more, less, or equally contaminated compared with  
76 conventional cereals. Authors concluded that this variability resulted from different cultivars,  
77 geographical locations of fields and time of harvest in different studies. It makes data not directly  
78 comparable.

79 In the literature, it can be found a number of analyses of effects of organic cultivation of wheat.  
80 However, experimental data that can verify the views presented there are still not very numerous.  
81 Considering this, we decided to carry out a field experiment on sowing 30 cultivars of winter wheat  
82 in the same location, at the same time on conventional and organic plots. The aim of the experiment  
83 was a comprehensive comparison of the results obtained for both cropping systems through the  
84 analysis in a series of elements that describe the structure of the yield, fungal diseases, presence of  
85 *Fusarium* fungi through analysis of the DNA content, production of mycotoxins in grain. The results  
86 were subject of the widest possible statistical analysis with the aim of finding relevant or irrelevant  
87 differences in both cultivation systems.

## 88 2. Material and methods

### 89 2.1. Field experiments

90 Thirty cultivars of winter wheat (*Triticum aestivum* L.) were evaluated (Table 1). The cultivars  
91 were listed in the Polish National List of the Research Centre for Cultivar Testing (COBORU) and  
92 were added to the list between 1998 (‘Mewa’) and 2009 (‘Belenus’). The cultivars were described in  
93 details in the paper of Góral et al. [20]. They differed in the pedigree, morphological characters, and  
94 resistance to *Fusarium* head blight (FHB). Cultivars were grouped in four classes of FHB resistance:  
95 susceptible (S), medium susceptible (MS), medium resistant (MR) and resistant (R).

96

**Table 1.** List of winter wheat cultivars used in this study.

No.	Cultivar	No.	Cultivar	No.	Cultivar
1	Akteur MS*	11	Jenga MS	21	Naridana MS
2	Alcazar S	12	Kampana S	22	Nateja R
3	Anthus MS	13	Kohelia MR	23	Ostka Strzelecka MS
4	Batuta MS	14	Legenda MR	24	Ostroga MR
5	Belenus MS	15	Ludwig MS	25	Slade MS
6	Bogatka MR	16	Markiza MS	26	Smuga S
7	Boomer MR	17	Meteor MS	27	Sukces MR
8	Dorota MR	18	Mewa MS	28	Tonacja MR
9	Figura MS	19	Mulan MS	29	Türkis MS
10	Garantus MS	20	Muszelka S	30	Zyta MR

97 \* - group of resistance to Fusarium head blight [20]; S=susceptible, MS=medium susceptible,  
 98 MR=medium resistant, R= resistant.

99 Field experiments were established in 2014 in the experimental fields of state-owned research  
 100 institute - Plant Breeding and Acclimatization Institute (IHAR-PIB) in Radzików, Central Poland.  
 101 First experiment was sown in the conventional field (GPS coordinates: 52.212517, 20.634765). Pre-  
 102 crop was oilseed rape. Artificial fertilizers were applied according to standard agricultural practices  
 103 in IHAR-PIB in particular. In the autumn 3 dt ha<sup>-1</sup> of 'Polifoska 6' fertilizer was applied (N -18 kg ha<sup>-1</sup>,  
 104 P - 45 kg ha<sup>-1</sup>, K -72 kg ha<sup>-1</sup>). In the spring, after the start of vegetation ammonium nitrate fertilizer  
 105 was applied in an amount providing 68 kg N ha<sup>-1</sup>. Weeds and pests were controlled with herbicides  
 106 and insecticides. Immediately after sowing weeds were controlled with herbicide 'Maraton 375SC' in  
 107 a dose of 4 L ha<sup>-1</sup>. In spring weeds were controlled using the herbicide 'Attribut 70GS' in a dose of 60  
 108 mg ha<sup>-1</sup>. Cereal leaf beetle and aphids were controlled with 'Fastac Active 050ME' in a dose of 250 ml  
 109 ha<sup>-1</sup>. No fungicides were applied.

110 Simultaneously the same wheat cultivars were sown in the experimental organic field of IHAR-  
 111 PIB (GPS coordinates: 52.216319, 20.638653). Wheat was grown according to organic farming  
 112 practices with no chemical disease control and application of fertilizers. Pre-crop was pea. Weeds  
 113 were controlled mechanically. No fertilizers or other components allowed in organic farming were  
 114 applied. Distance between two experimental fields was about 500 m. Single plot size in both  
 115 experiments was 5 m<sup>2</sup>. In both fields, cultivars were sown in three randomized blocks (replications)  
 116 distant from each other by 2 meters.

117 Heading and full flowering dates for individual plots were recorded. Plant height was measured  
 118 after the end of heading stage. Fusarium head blight was scored based on the mean percentage of  
 119 blighted spikelets per infected head (disease severity) and the percentage of infected heads per plot  
 120 (disease incidence). Fusarium head blight index (FHB<sub>i</sub>) was calculated as the combination of disease  
 121 severity and disease incidence.

$$\text{FHB}_i = (\text{FHB}_{\text{severity}} \times \text{FHB}_{\text{incidence}}) / 100 \quad (1)$$

122 Presence of other fungal diseases were also recorded. They were as follows: yellow rust (*Puccinia*  
 123 *striiformis*), leaf rust (*P. triticina*), Septoria tritici blotch (*Zymoseptoria tritici*), Stagonospora nodorum  
 124 blotch (*Parastagonospora nodorum*) and tan spot (*Pyrenophora tritici-repentis*). These diseases were  
 125 scored according to percentage of leaf area per plot with symptoms of disease – necrosis and/or  
 126 sporulation.

## 127 2.2. Analysis of mineral elements in soil

128 In spring, soil samples were collected from conventional and organic fields. Twenty soil cores  
 129 were taken from experimental plots in both fields using soil sampler. Soil cores from plots were mixed  
 130 thoroughly.

131 The material was mineralized with a CEM Mars 5 Xpress (CEM, Matthews, NC, USA)  
 132 microwave mineralization system (55 ml vessels) using 8 ml HNO<sub>3</sub> (65%) and 2 ml H<sub>2</sub>O<sub>2</sub>, according

133 to the program comprising three stages: first stage – power 800 W, time 10 min, temperature 120 °C;  
134 second stage – power 1600 W, time 10 min, temperature 160 °C; third stage – power 1600 W, time 10  
135 min, temperature 200 °C [21]. Materials after digestion were filtered through 45 mm filters  
136 (Qualitative Filter Papers Whatman, Grade 595: 4 – 7 µm; GE Healthcare, Buckinghamshire, UK), and  
137 filtrate completed with deionized water from Milli-Q Academic System (non-TOC (Total Organic  
138 Carbon); Millipores. A.S., Molsheim, France) to a final volume of 50 ml. Concentration of particular  
139 trace elements was analyzed by the flame atomic absorption spectrometry (Cd, Cu, Mn, Cr, Co, Si,  
140 Ni and Zn), atomic emission spectrometry (Mg, Ca, Na, K, B) using an AA Duo – AA280FS/AA280Z  
141 spectrometer (Agilent Technologies, Mulgrave, Victoria, Australia), equipped with a Varian hollow-  
142 cathode lamp (HCL; Varian, Mulgrave, Victoria, Australia). Calibration curves were prepared in four  
143 replicates per each trace element concentration. Detection limit for the analysed metals was, ng kg<sup>-1</sup>:  
144 Ca 0.015, Na 0.10, K 0.09, Mg 0.003, B 0.06, Cu 0.18, Zn 0.06, Cr 0.005, Mn 0.005, Co 0.011, Si 0.12, Ni  
145 0.005, Cd 0.01.

### 146 2.3. Seed quality

147 For the evaluation of germination ability, 3 x 50 seeds from each experimental plot (180 samples)  
148 were sown in plastic boxes between two layers of moistened (to 60% WR) filter paper. After sowing,  
149 the samples were prechilled at 7 °C for 3 days and placed in Sanyo growth chamber (Sanyo Electric  
150 Co., Ltd., Japan) at constant temperature 20 °C. After four days, first count (germination energy) was  
151 made. The normal seedlings were counted and share in percent was evaluated. According to present  
152 International Seed Testing Association Rules [22] after eight days, the final count (germination  
153 capacity) and evaluation of normal seedlings, abnormal seedlings (AS), dead seeds (DS) and fresh  
154 ungerminated (FUS) seeds were made.

### 155 2.4. *Fusarium* DNA quantification with Real-Time PCR

#### 156 2.4.1. Isolation of total DNA from grain

157 DNA was extracted according to Doyle and Doyle [23] protocol with modifications of  
158 Department of Phytopathology and Molecular Mycology UTP.

159 Ten grams of grain was homogenized to fine powder and 100 mg of such prepared sample was  
160 taken for DNA isolation. Samples were transferred into 2.0 ml tubes and poured with 600 µl of the  
161 extraction buffer containing CTAB 5.0 %, EDTA 0.5 M, NaCl 5.0 M, Tris-HCl (pH 8.0) 1.0 M, β-  
162 mercaptoethanol, PVP (2.0%) and water. DNA was purified by addition of phenol:chloroform:  
163 isoamyl alcohol mixture (25:24:1) followed by centrifugation and taking the upper phase  
164 (supernatant) to the new tube, where equal volume of chloroform:isoamyl alcohol mixture (24:1) was  
165 added, mixed by inverting and centrifuged. Supernatant was taken and DNA was precipitated with  
166 cold ethanol. DNA pellet was washed with 70% cold ethanol, left to dry for 25-30 minutes and poured  
167 with TE buffer or sterile water to dissolve. Samples were stored at -20°C for further analyses.

#### 168 2.4.2. Preparation of standard curve

169 Material for preparation of standard curve was a series of 10-fold dilutions of DNA isolated from  
170 pure culture of researched six *Fusarium* species (*F. avenaceum*, *F. culmorum*, *F. graminearum*, *F.*  
171 *langsethiae*, *F. poae* and *F. sporotrichioides*). Pure fungal cultures were grown on PDA medium on Petri  
172 dish and DNA was isolated from scraped and lyophilized mycelium using the same protocol as for  
173 grain.

#### 174 2.4.3. Preparation of DNA samples for Real-Time PCR

175 Concentrations of DNA obtained from kernels were measured with Quantus fluorometer  
176 (Promega, USA). All samples were diluted in sterile deionized water to 10 ng·µl<sup>-1</sup>. Final concentration  
177 of *Fusarium* DNA in a sample was expressed in picograms per 100 ng of total DNA.

#### 178 2.4.4. Real-Time PCR reaction conditions

179 Amplification was performed with LightCycler 480II (Roche) using LightCycler 480 SYBR Green  
180 I Master (Roche) in a volume of 10 ml per sample (5.5  $\mu$ l premix + 4.5  $\mu$ l DNA) in 45 cycles according  
181 to thermal profiles specific to each *Fusarium* species.

182 The primers used for each researched *Fusarium* species were as follows:

183 *Fusarium avenaceum*: JIAF / JIAR

184 (GCTAATTCTTAACCTACTAGGGGCC / CTGTAATAGGTTATTTACATGGGCG) [24];

185 *Fusarium culmorum*: Fc01F / Fc01R

186 (ATGGTGAACCTCGTCGTGGC / CCCTTCTTACGCCAATCTCG) [25];

187 *Fusarium graminearum*: Fg16F / Fg16R

188 (CTCCGGATATGTTGCGTCAA / GGTAGGTATCCGACATGGCAA) [25];

189 *Fusarium langsethiae*: FlangF3 / LanspoR1

190 (CAAAGTTCAGGGCGAAAACCT / TACAAGAAGACGTGGCGATAT) [26];

191 *Fusarium poae*: Fp82F / Fp82R

192 (CAAGCAAACAGGCTCTTACC / TGTTCCACCTCAGTGACAGGTT) [27]

193 *Fusarium sporotrichioides*: FsporF1 / LanspoR1

194 (CGCACAACGCAAACCTCATC / TACAAGAAGACGTGGCGATAT) [26]

## 195 2.5. Analysis of trichothecenes

196 Grain samples (60) were analyzed for the presence of trichothecenes according to Perkowski et  
197 al. [28]. Subsamples (10 g) were extracted with acetonitrile:water (82:18) and purified on a charcoal  
198 column (Celite 545/charcoal Draco G/60/activated alumina neutral 4:3:4 (w/w/w)).

199 Type A trichothecenes (HT-2 toxin [HT-2], T-2 toxin [T-2], T-2 tetraol, T-2 triol,  
200 diacetoxyscirpenol [DAS], scirpentriol [STO]) were analysed as TFAA (trifluoroacetic anhydride)  
201 derivatives. To the dried sample 100  $\mu$ l of trifluoroacetic acid anhydride were added. After 20 min.,  
202 the reacting substance was evaporated to dryness under nitrogen. The residue was dissolved in 500  
203  $\mu$ l of isooctane and 1  $\mu$ l was injected onto a gas chromatograph-mass spectrometer (GC/MS). Type B  
204 trichothecenes (DON, NIV, 3-acetyldeoxynivalenol [3-AcDON], 15-acetyldeoxynivalenol [15-  
205 AcDON], fusarenon X [FUS-X]) were analyzed as TMS (trimethylsilylsilyl ethers) derivatives. To the  
206 dried extract, the amount of 100  $\mu$ l of TMSI/TMCS (trimethylsilyl imidazole/trimethylchlorosilane;  
207 v/v 100/1) mixture was added. After 10 min. 500  $\mu$ l of isooctane were added and the reaction was  
208 quenched with 1ml of water. The isooctane layer was used for the analysis and 1  $\mu$ l of the sample  
209 was injected on a GC/MS system.

210 The analyses were run on a gas chromatograph (Hewlett Packard GC 6890, Waldbronn,  
211 Germany) hyphenated to a mass spectrometer (Hewlett Packard 5972 A, Waldbronn, Germany),  
212 using an HP-5MS, 0.25 mm x 30 m capillary column. The injection port temperature was 280°C, the  
213 transfer line temperature was 280°C and the analyses were performed with programmed  
214 temperature, separately for type A and type B trichothecenes. The type A trichothecenes were  
215 analysed using the following programmed temperatures: initial 80°C held for 1 min., from 80°C to  
216 280°C at 10°C min<sup>-1</sup>, the final temperature being maintained for 4 min. For the type B trichothecenes  
217 initial temperature of 80°C was held for 1 min., from 80°C to 200°C at 15°C min<sup>-1</sup> held for 6 min and  
218 from 200°C to 280°C at 10°C/min, with the final temperature being maintained for 3 min. The helium  
219 flow rate was held constant at 0.7 ml min<sup>-1</sup>.

220 Quantitative analysis was performed in the single ion monitored mode (SIM) using the  
221 following ions for the detection of STO: 456 and 555; T-2 tetraol 455 and 568; T-2 triol 455 and 569 and  
222 374; HT-2 455 and 327; T-2 327 and 401. DON: 103 and 512; 3-AcDON: 117 and 482; 15-AcDON: 193  
223 and 482; NIV: 191 and 600. Qualitative analysis was performed in the SCAN mode (100 – 700 amu).  
224 Recovery rates for the analyzed toxins were as follows: STO 82±5.3%; T-2 triol 79±5.1%; T-2 86±3.8%;  
225 T-2 tetraol 88±4.0%; HT-2 91±3.3%; DON 84±3.8%; 3AcDON 78±4.8%; 15 AcDON 74±2.2%; and NIV  
226 81±3.8%. The limit of detection was 0.01  $\mu$ g kg<sup>-1</sup>.

## 227 2.6. Chemical analysis of ergosterol

228 Ergosterol (ERG) in 60 grain samples was determined by HPLC as described by Young [29] with  
229 modifications [30,31]. A detailed evaluation of the method was given in a study by Perkowski et al.  
230 [31]. Samples containing 100 mg of ground grains were placed into 17-ml culture tubes, suspended  
231 in 2ml of methanol, treated with 0.5 ml of 2M aqueous sodium hydroxide and tightly sealed. The  
232 culture tubes were then placed within 250-ml plastic bottles, tightly sealed and placed inside a  
233 microwave oven (Model AVM 401/1WH, Whirlpool, Sweden) operating at 2450 MHz and 900 W  
234 maximum output. Samples were irradiated (370 W) for 20 s and after about 5 min for an additional  
235 20 s. After 15 min. the contents of culture tubes were neutralized with 1M aqueous hydrochloric acid,  
236 2 ml MeOH were added and extraction with pentane (3 x 4 ml) was carried out within the culture  
237 tubes. The combined pentane extracts were evaporated to dryness in a nitrogen stream. Before  
238 analysis samples were dissolved in 4 ml of MeOH, filtered through 13-mm syringe filters with a 0.5  
239 mm pore diameter (Fluoropore Membrane Filters, Millipore, Ireland) and evaporated to dryness in a  
240 N<sub>2</sub> stream. The sample extract was dissolved in 1ml of MeOH and 50 µl were analyzed by HPLC.  
241 Separation was performed on a 150 x 3.9 mm Nova Pak C-18, 4 mm column and eluted with  
242 methanol/acetonitrile (90:10) at a flow rate of 0.6 ml min<sup>-1</sup>. Ergosterol was detected with a Waters 486  
243 Tunable Absorbance Detector (Milford, MA, USA) set at 282 nm. The presence of ERG was confirmed  
244 by a comparison of retention times and by co-injection of every tenth sample with an ergosterol  
245 standard.

## 246 2.7. Statistics

247 The statistical analysis was performed using Microsoft® Excel 2016/XLSTAT© Ecology (Version  
248 18.07.38413, Addinsoft, Inc., Brooklyn, NY, USA). Differences between variable means for the two  
249 experimental variants were compared using parametric Student's t-test (XLSTAT procedure: *Two-*  
250 *sample t and z tests*). Variables distribution in samples from the two experimental variants were  
251 compared using the Kruskal-Wallis one-way analysis of variance (XLSTAT procedure: *Comparison of*  
252 *k samples - Kruskal-Wallis, Friedman*). The Kruskal-Wallis test was selected because some of the  
253 variables did not follow normal distribution.

254 The relationships between FHBi, seed quality and concentration of ergosterol, mycotoxins and  
255 *Fusarium* DNA were investigated by Pearson correlation tests (XLSTAT procedure: *Correlation tests*).  
256 Prior to analysis, data that did not follow normal distribution was log<sub>10</sub> transformed to normalize  
257 residual distributions. Multivariate data analysis method was applied to the data on FHB (FHBi, DS,  
258 mycotoxin concentrations, *Fusarium* DNA concentrations) resistance. Principal component analysis  
259 (XLSTAT procedure: *Principal Component Analysis PCA*) was used to show how wheat cultivars  
260 within two experimental variants (60 observations) are distributed with respect to the main variation  
261 described in the first two components and how variables (FHBi, DS, ERG, sum of type A  
262 trichothecenes, sum of type B trichothecenes, *Fusarium* DNA) influence the construction of the two  
263 components. PCA results also revealed associations among variables measured by the angle between  
264 variable vectors.

265 Differences between two variants for all variables were analyzed using multidimensional tests  
266 (XLSTAT procedure: *Multidimensional tests (Mahalanobis)*) and multivariate analysis of variance  
267 (XLSTAT procedure: *MANOVA*).

268 Cultivars in the organic field were grouped according to their resistance to infection of heads  
269 with *Fusarium* fungi measured by FHB index, dead seeds proportion, ERG, sum of type A  
270 trichothecenes, sum of type B trichothecenes, *Fusarium* DNA. 'K-means clustering' procedure of  
271 XLSTAT was applied. Results were visualized using 'Discriminant analysis' procedure of XLSTAT.  
272 Classes obtained from K-means analysis were applied as a qualitative depended variable in DA  
273 analysis.

## 274 3. Results

### 275 3.1. Concentration of mineral elements in soil

276 In order to determine soil conditions in both experimental fields analysis of a number of  
 277 elements that occur in these environments was made (Table S1). For the most of 13 analyzed  
 278 compounds significant differences between organic and conventional fields were found. Only for Co,  
 279 concentration difference was not significant. In soil of conventional field, concentration of K, Mg, Cd,  
 280 Cr, Cu, Ni, and Zn was higher than in soil of organic field. The highest differences were found for Zn  
 281 (7-fold) and Cd (3-fold). On the other hand, in soil of organic field, concentration of Ca, Na, Si, B, and  
 282 Mn was higher than in soil of conventional field.

### 283 3.2. Phenotypic data and fungal diseases

284 Wheat cultivars differed in heading and flowering time. In the conventional field, heading time  
 285 was 29.7 days from 1st May, at a range 24.0 ('Smuga') – 35.0 days ('Sukces') (Table S2). Flowering  
 286 time was on average 31.6 days from 1st May, at a range 26.0 ('Smuga') – 35.0 days ('Sukces', 'Boomer').  
 287 In the organic field, heading time was 28.0 days from 1st May, at a range 24.0 ('Smuga') – 33.0 days  
 288 ('Sukces'). Flowering time was on average 29.8 days from 1st May, at a range 26.0 ('Smuga', 'Ludwig')  
 289 – 35.0 days ('Sukces'). Heading and flowering time were significantly earlier in organic field than in  
 290 conventional one (Table 2).

291 **Table 2.** Phenotypic characters, grain yield and FHB infection of 30 wheat cultivars grown in  
 292 conventional and organic field.

Variant	Heading (days from 1st May)	Flowering (days from 1st May)	Plant height (cm)	Grain yield per plot (kg)	FHBi (%)
Conventional					
Mean	29.7 b	31.6 b	97.8 a	5.0 a	0.74 a
Std. deviation	2.53	2.39	12.22	0.89	1.00
Organic					
Mean	28.0 a	29.8 a	99.0 a	5.1 a	0.66 a
Std. deviation	2.39	2.55	10.39	0.76	0.77

293 Values within the same column followed by the different letters are significantly different at the level  
 294 of probability < 0.01.

295 On average, plant height of wheat cultivars did not differ between organic and conventional  
 296 fields (Table 2). In organic field plant height ranged between 73.7 cm ('Muszelka') and 114.3 cm  
 297 ('Akteur') (Table S2). In conventional field, this parameter ranged between 76.3 cm ('Alcazar') and  
 298 118.7 cm ('Ludwig').

299 Fusarium head blight severity was low and average values for conventional and organic fields  
 300 did not differ significantly (Table 2, Figure 1). In conventional field, FHB index range was from 0 to  
 301 4.4%. Cultivars 'Nateja' and 'Legenda' showed no symptoms of FHB and cultivars 'Kampana',  
 302 'Muszelka' and 'Slade' were the most infected (FHBi = 4.4%, 3.5%, and 2.1%, respectively) (Table S2).  
 303 In organic field, FHB index range was from 0 to 3.2%. Cultivars 'Nateja' and 'Mewa' showed no  
 304 symptoms of FHB and cultivars 'Slade', 'Kampana', 'Turkis' and 'Belenus' were the most infected  
 305 (FHBi = 3.2%, 2.3%, 1.8% and 1.8%, respectively). FHB indexes for conventional and organic fields  
 306 correlated significantly ( $r = 0.776$  at  $P < 0.001$ ).

307 Heading and flowering time did not correlated with FHB severity. In both fields plant height  
 308 significantly negatively correlated with FHB indexes ( $r = -0.519$  for organic field and  $r = -0.589$  for  
 309 conventional field; at  $P < 0.001$ ).

310  
 311



312

313 **Figure 1.** FHB symptoms on heads of wheat grown in organic field. Clockwise from top left: 'Figura',  
 314 'Muszelka', 'Kampna', 'Slade'. Wheat at early milk growth stage (BBCH 73).

315 Symptoms of leaf diseases in both experimental plots were observed starting from half of May,  
 316 when yellow rust was detected. Seventeen cultivars were fully resistant to yellow rust and showed  
 317 no symptoms of disease (Table S3). On average yellow rust severity was slightly higher in  
 318 conventional field; however, difference with organic field was not significant. Leaf rust severity was  
 319 low. On average, it was 0.7% in organic field and 1.0% in conventional one. (Table S3). Symptoms of  
 320 *Septoria tritici* blotch were observed on most (28) cultivars in conventional field and on 17 cultivars  
 321 in organic field. Symptoms of *Stagonospora nodorum* blotch were observed on 12 cultivars in  
 322 conventional field and on 23 cultivars in organic field. Tan spot was observed only in organic field  
 323 with average severity 2.0%. This disease affected fourteen cultivars.

324 Grain yield per plot was similar for both field and do not differ statistically significantly (Table  
 325 2). In organic field grain yield ranged between 3.1 kg ('Nateja') and 6.7 kg ('Jenga') (table S1). In  
 326 conventional field, this parameter ranged between 2.9 kg ('Nateja') and 6.6 kg ('Boomer'). In both  
 327 field grain yield was significantly negatively correlated with yellow rust severity ( $r = -0.517$  for  
 328 organic field and  $r = 0.647$  for conventional field; at  $P < 0.001$ ) and not correlated with FHB indexes.  
 329

### 330 3.3. Characteristic of seed germination

331 Sowing quality of seeds from conventional field was significantly higher than those from organic  
 332 one were (Table 3). The mean value for the germination energy for the conventional seeds was much  
 333 higher (87%) than for the organic seeds (63.2%). Similar was found for the germination capacity. In

334 organic seed material lower percent's share of normal seedlings, but higher number of abnormal  
335 seedlings, dead seeds and fresh ungerminated seeds was observed.

336 **Table 3.** Germination characteristic of 30 wheat cultivars grown in conventional and organic field.

Variant	Germination energy (%)	Germination capacity (%)	Abnormal seedlings (%)	Dead seeds (%)	Fresh, ungerminated seeds (%)
Conventional					
Mean	87.0 b	93.4 b	3.6 a **	2.5 a **	0.6 a
Std. deviation	10.86	3.81	2.26	1.79	0.63
Organic					
Mean	63.2 a	89.3 a	5.3 b **	4.2 b **	1.2 a
Std. deviation	24.00	4.81	2.10	3.01	1.91

337 Values within the same column followed by the different letters are significantly different at the level  
338 of probability < 0.001 or \* < 0.01.

339 Values of the germination energy ranged from 54.0% ('Mewa') to 98.0% ('Nateja') in  
340 conventional samples and from 11.5% ('Belenus') to 93.0% ('Mewa') in organic samples (Table S4).  
341 The germination capacity ranged from 83.5% ('Slade') to 98.5% ('Nateja') in conventional material  
342 and from 75.0% ('Belenus') to 96.0% ('Batuta') in organic material. The percent shares of abnormal  
343 seedlings as well as dead seeds were significantly higher in organic samples (Table 3). These variables  
344 ranged from 0.5% ('Markiza') to 8.5% ('Alcazar', 'Garantus') and from 0.5% ('Batuta') to 15.0%  
345 ('Belenus') in organic field and 0 ('Figura', 'Belenus') to 7.5% ('Mewa', 'Ostroga', 'Slade') and from 0  
346 ('Batuta', 'Nateja') to 7.5% ('Jenga') in conventional field. Additionally, percentage of fresh,  
347 ungerminated seeds was twice higher in organic material than in conventional, however difference  
348 was not significant. It was the highest in organic seed material of cultivars 'Belenus' (7.5%), 'Akteur'  
349 (6.0%), and 'Ostroga' (5.0%).

#### 350 3.4. Concentration of ergosterol and trichothecenes

351 Concentration of ergosterol in grain was significantly higher in samples from organic field than  
352 from conventional one (Table 4). Level of ERG varied from 0.26 ('Ostka Strzelecka') to 1.85 mg kg<sup>-1</sup>  
353 ('Mulan') in conventional field and from 0.26 ('Boomer') to 3.46 mg kg<sup>-1</sup> ('Akteur') in organic field  
354 (Table S4).

355 Amount of type B trichothecenes was low and varied from 8.9 to 460.2 µg kg<sup>-1</sup> in conventional  
356 field and from 10.1 to 384.5 µg kg<sup>-1</sup> in organic field. On average, more trichothecenes were present in  
357 grain from conventional field; however, difference was statistically insignificant. Regarding specific  
358 toxins, only concentration of 3-AcDON was significantly higher in conventional samples.  
359 Concentration of NIV was higher in samples from organic field; however, difference was not  
360 significant. Distributions for FUS-X and 3-AcDON in organic and conventional samples were  
361 significantly different. In conventional samples, these toxins were detected in higher amounts in  
362 single samples whereas they were more evenly distributed in organic samples.

363 **Table 4.** Concentrations of ergosterol (mg kg<sup>-1</sup>) and type B trichothecenes (µg kg<sup>-1</sup>) in grain of 30 wheat  
364 cultivars grown in conventional and organic fields.

Variant	ERG	DON	FUS-X	3-AcDON	15-AcDON	NIV	Total TCT B
Conventional							
Mean	0.74 a *	84.8 a	0.9 a	7.3 b	1.5 a	5.6 a	100.0 a
Range	0.26–1.85	5.8–444.4	0–11.6	2.3–30.3	0–14.3	0–19.0	8.9–460.2
Std. deviation	0.39	97.3	2.4	5.9	2.6	5.0	101.4
Organic							

Mean	1.42 b *	63.7 a	0.9 a	3.1 a	1.1 a	7.4 a	76.2 a
Range	0.26–3.46	2.2–348.4	0–2.9	0–6.2	0–3.3	0–29.5	10.1–384.5
Std. deviation	0.87	86.2	1.0	1.5	1.2	7.3	93.6

365 Values within the same column followed by the different letters are significantly different at the level  
366 of probability < 0.001 or \* < 0.01.

367 The highest concentrations of type B trichothecenes were found in grain of cultivars 'Anthus',  
368 'Ostroga', and 'Garantus' in conventional field (460.2  $\mu\text{g kg}^{-1}$ , 321.1  $\mu\text{g kg}^{-1}$ , 308.9  $\mu\text{g kg}^{-1}$ , respectively)  
369 and in grain of 'Alcazar', 'Kampana', 'Muszelka' and 'Anthus' (384.5  $\mu\text{g kg}^{-1}$ , 278.  $\mu\text{g kg}^{-1}$ , 257.6  $\mu\text{g kg}^{-1}$ ,  
370 244.6  $\mu\text{g kg}^{-1}$ , respectively) in organic field (Table S5).

371 Amount of type A trichothecenes was very low and similar in conventional and organic samples  
372 (5.5 and 5.1  $\mu\text{g kg}^{-1}$ , respectively) (Table S6). It varied from 0.7 ('Boomer') to 30.5  $\mu\text{g kg}^{-1}$  ('Garantus')  
373 in conventional samples and from 1.0 ('Figura') to 14.5  $\mu\text{g kg}^{-1}$  ('Zyta') in organic samples. Average  
374 concentrations of type A trichothecenes were similar in both groups, and they not differ significantly.  
375 Only average concentration of DAS was significantly higher in conventional samples.

### 376 3.5. *Fusarium* species

377 Presence of biomass of six *Fusarium* species was detected in wheat grain. *Fusarium langsethiae*  
378 was detected only in six samples in trace amounts. On average, the highest amount of DNA was  
379 found as follows for *F. poae*, *F. graminearum*, *F. sporotrichioides*, *F. culmorum* and *F. avenaceum* (Table  
380 5). It was true in organic field. In conventional field concentration of *F. culmorum* DNA was higher  
381 than *F. graminearum* and *F. sporotrichioides* DNA.

382 **Table 5.** Concentration of DNA (pg 100ng<sup>-1</sup> of wheat DNA) of five *Fusarium* species in grain of 30  
383 wheat cultivars grown in conventional and organic fields.

Variant	<i>F. a.</i> DNA	<i>F. c.</i> DNA	<i>F. g.</i> DNA	<i>F. p.</i> DNA	<i>F. sp.</i> DNA	<i>Fusarium</i> DNA
Conventional						
Mean	10.8 a	23.3 a	22.7 a	34.7 a	15,1 a*	106.6 a
Range	0–106.7	0–346.2	0.7–79.5	9.4–132.4	0–113.3	15.4–405.2
Std. deviation	19.61	63.14	21.58	24.82	29.10	90.91
Organic						
Mean	30.2 b	41.1 a	67.0 b	98.2 b	50.5 b*	285.7 b
Range	0.2–184.6	0–415.5	0.5–280.0	14.1–222.0	0–350.0	15.3–1205.8
Std. deviation	44.88	85.73	62.92	56.44	94.3	244.83

384 Values within the same column followed by the different letters are significantly different at the level  
385 of probability < 0.001 or \* < 0.05. *F. a.* – *F. avenaceum*, *F. c.* – *F. culmorum*, *F. g.* – *F. graminearum*, *F. p.* –  
386 *F. poae*, *F. sp.* – *F. sporotrichioides*, *Fusarium* DNA – total DNA of five species.

387 Total *Fusarium* DNA concentration in organic samples was more than twice higher than in  
388 conventional samples. It ranged from 15.4 ('Batuta') to 405.2 pg 100ng<sup>-1</sup> ('Figura') in conventional  
389 samples and from 15.3 ('Batuta') to 1205.8 pg 100ng<sup>-1</sup> ('Slade') in organic samples. The difference was  
390 statistically significant. Similarly, significantly higher (about three times) were concentrations of *F.*  
391 *poae*, *F. graminearum*, *F. avenaceum* and *F. sporotrichioides* in organic samples. Amount of *F. culmorum*  
392 was about twice higher in organic samples; however, difference was not statistically significant. The  
393 most *Fusarium* colonized was grain of 'Figura', 'Kampana' and 'Alcazar' cultivars in conventional  
394 field and 'Muszelka', 'Kampana', 'Turkis', 'Ostroga', 'Meteor', 'Bogatka', 'Alcazar', and 'Slade' in  
395 organic field (Table S7).

### 396 3.6 Correlation between experimental components

397 Result's correlations of the evaluation of conventional wheat material showed that the  
398 proportion of the dead seed was poorly correlated with the germination energy, but highly negatively  
399 correlated with germination capacity (Table 6). In the case of organic seed, the dependence was the  
400 same, except that the negative values of the correlation coefficients were higher (Table 7). The  
401 proportion of abnormal seedlings was significantly negatively correlated with the energy and  
402 germination capacity, as well as the number of dead seeds in conventional material. In contrast to the  
403 results of organic material, where the relevant dependencies for these traits were not found.  
404 However, organic material has demonstrated a highly significant negative relationship between the  
405 proportion of fresh ungerminated seeds and the energy and germination capacity and a positive  
406 correlation between fresh ungerminated and dead seed.

407 In conventional and ecological material, the same negative relationship between FHBi and  
408 germination was found, and the positive relationship between FHBi and the share of dead seeds  
409 occurred. These two parameters in conventional samples correlated also significantly with  
410 concentration of type B trichothecenes in grain.

411 We observed significant effect of colonization of kernels with *Fusarium* species on seed quality  
412 (Tabs 6 and 7). The proportion of dead seed in conventional material was highly correlated with the  
413 quantity of the DNA of *F. graminearum*, while in organic material with the amount of DNA of *F. poae*,  
414 *F. sporotrichioides* and total *Fusarium* DNA.

415 There was lack of correlation between *Fusarium* head blight index and amount of ERG and  
416 type A and B trichothecenes in grain in both variants (Tabs 6 and 7). However, in conventional  
417 samples positive tendency FHBi *versus* type B trichothecenes was observed and the same was found  
418 for FHBi *versus* type A trichothecenes in organic samples. *Fusarium* head blight index correlated  
419 significantly with concentration of DNA of three *Fusarium* species – *F. avenaceum*, *F. graminearum* and  
420 *F. poae* in both variants. In organic variant FHBi correlated significantly also with *F. sporotrichioides*  
421 DNA concentration. No correlation was found with *F. culmorum* DNA. Ergosterol content in grain  
422 did not correlated with type A or B trichothecenes as well as with DNA concentration of *Fusarium*  
423 species.

424 In samples from conventional field amount of type B trichothecenes correlated highly  
425 significantly with *F. graminearum* DNA but not with *F. culmorum* DNA. Contradictory, in organic  
426 samples *F. graminearum* did not correlate with type B trichothecenes and for *F. culmorum* there was  
427 found positive tendency however not statistically significant. Regarding specific toxins in  
428 conventional samples *F. graminearum* correlated significantly with DON amount ( $r = 0.531$ ) and for *F.*  
429 *culmorum* some positive tendency was observed for FUS-X and 15-AcDON. In organic samples, only  
430 correlation of *F. culmorum* with 3-AcDON ( $r = 0.421$ ) was found. There was no significant correlation  
431 between amount of type A trichothecenes and potentially producing species *F. sporotrichioides* and *F.*  
432 *poae*.

433 Amounts of DNA of three *Fusarium* species (*F. avenaceum*, *F. graminearum* and *F. poae*) in grain  
434 form both variants correlated statistically significantly (Table 8 and 9). *Fusarium sporotrichioides* DNA  
435 concentration correlated with *F. avenaceum* and *F. poae* DNA. *Fusarium culmorum* DNA concentration  
436 did not correlated with the other species

437 Table 6. Coefficients of correlations between seed quality parameters (germination energy - GE, germination capacity - GC, dead seeds, abnormal seedling - AS, fresh  
 438 ungerminated seeds - FUS), *Fusarium* head blight index (FHBi), ergosterol (ERG) and type A and B trichothecenes concentrations (Total TCT A, Total TCT B) as well as amount  
 439 of DNA of five *Fusarium* species in grain samples from conventional field

Variables	GE	GC	Dead seeds	AS	FUS	FHBi	ERG	Total TCT B	Total TCT A	<i>F. a.</i> DNA	<i>F. c.</i> DNA	<i>F. g.</i> DNA	<i>F. p.</i> DNA	<i>F. sp.</i> DNA
Final count	0.584***													
Dead seeds	-0.419*	-0.803***												
AS	-0.553**	-0.777***	0.436*											
FUS	-0.112	-0.320	0.048	0.012										
FHBi	-0.125	-0.476**	0.545**	0.219	0.058									
ERG	-0.022	0.222	-0.111	-0.190	-0.162	-0.064								
Total TCT B	-0.019	-0.431*	0.369*	0.312	0.160	0.290	-0.163							
Total TCT A	-0.216	-0.252	0.185	0.373*	0.044	0.096	0.011	0.573***						
<i>F. a.</i> DNA	0.258	-0.221	0.412*	0.095	-0.170	0.430*	0.016	-						
<i>F. c.</i> DNA	-0.052	-0.215	0.168	0.002	0.289	0.011	-0.325	0.174	-	-0.013				
<i>F. g.</i> DNA	-0.204	-0.502**	0.661***	0.288	0.010	0.586***	-0.104	0.501**	-	0.583***	0.347			
<i>F. p.</i> DNA	0.259	-0.169	0.393*	-0.020	0.010	0.388*	-0.285	0.202	0.108	0.531**	0.053	0.497**		
<i>F. sp.</i> DNA	0.181	0.016	0.197	0.125	-0.564***	0.193	-0.270	-	-0.162	0.451*	-0.241	0.215	0.447*	
Total DNA	0.124	-0.297	0.565	0.040	-0.107	0.481**	-0.267	0.201	0.063	0.657***	0.343	0.712***	0.762***	0.515**

440 *F.a.* - *F. avenaceum*, *F. c.* - *F. culmorum*, *F. g.* - *F. graminearum*, *F. p.* - *F. poae*, *F. sp.* - *F. sporotrichioides*; coefficients significant at  $P \leq 0.001$  -\*\*\*; 0.01 - \*\*; 0.05\*.

441

442

443

444

445

446 Table 7. Coefficients of correlations between seed quality parameters (germination energy - GE, germination capacity - GC, dead seeds, abnormal seedling - AS, fresh  
 447 ungerminated seeds - FUS) and *Fusarium* head blight index (FHBi), ergosterol (ERG) and type A and B trichothecenes concentrations (Total TCT A, Total TCT B) as well  
 448 as amount of DNA of five *Fusarium* species in grain samples from organic field

Variables	GE	GC	Dead seeds	AS	FUS	FHBi	ERG	Total TCT B	Total TCT B	<i>F. a.</i> DNA	<i>F. c.</i> DNA	<i>F. g.</i> DNA	<i>F. p.</i> DNA	<i>F. sp.</i> DNA
Final count	0.639***													
Dead seeds	-0.511**	-0.874***												
AS	-0.059	-0.327	0.203											
FUS	-0.538**	-0.579***	0.470**	-0.346										
FHBi	-0.274	-0.533**	0.456*	0.360	0.002									
ERG	-0.083	-0.056	0.080	0.257	-0.152	0.036								
Total TCT B	-0.104	-0.219	0.207	0.309	-0.158	0.173	0.182							
Total TCT A	0.097	0.027	0.158	0.056	-0.225	0.315	-0.072	0.162						
<i>F. a.</i> DNA	-0.023	-0.479**	0.557**	0.357	0.117	0.511**	0.192	-	-					
<i>F. c.</i> DNA	-0.052	0.068	0.006	-0.202	-0.111	0.077	0.165	0.235	-	-0.210				
<i>F. g.</i> DNA	-0.127	-0.379*	0.459*	0.370*	-0.003	0.461**	-0.031	-0.006	-	0.506**	-0.039			
<i>F. p.</i> DNA	-0.382*	-0.577***	0.653***	0.220	0.282	0.508**	-0.055	0.208	0.163	0.550**	0.006	0.546**		
<i>F. sp.</i> DNA	-0.338	-0.590***	0.593***	0.302	0.264	0.584***	0.098	-	0.106	0.607***	-0.085	0.323	0.530**	
Total DNA	-0.201	-0.475**	0.557***	0.323	0.089	0.636***	0.004	0.216	0.243	0.640***	0.214	0.760***	0.838***	0.623***


449 *F. a.* - *F. avenaceum*, *F. c.* - *F. culmorum*, *F. g.* - *F. graminearum*, *F. p.* - *F. poae*, *F. sp.* - *F. sporotrichioides*; coefficients significant at  $P \leq 0.001$  -\*\*\*; 0.01 - \*\*; 0.05\*.

### 3.7. Multivariate principal component analysis

Multivariate principal component analysis showed significant difference between two studied populations (wheat cultivars in two environments) in terms of FHB infection (Figure 1). However, this difference was caused by only some cultivars, which showed higher *Fusarium* infection (measured with different parameters) in organic or conventional field. Cultivars from organic field had higher FHB index, proportion of dead seeds and *Fusarium* DNA content. In conventional field, the most infected cultivars had higher toxin content in the grain but moderate FHB index, dead seeds proportion and *Fusarium* biomass amount in kernels. The exception was cultivar 'Kampana' (C\_12) (Figure 2).


There were also carried out other tests - Multidimensional Wilks' Lambda test and Fisher distances test. They pointed to the significance of the separation between the analyzed growing systems at the significance level of  $P < 0.0001$ .

There was also compared which source of variation had higher effect on the obtained results (i.e. FHBi, DS, Total TCT B, Total TCT A and *Fusarium* DNA concentration) using multivariate analysis of variance (MANOVA). Both sources statistically significantly affected the results; however, experimental variant (conventional vs organic field) had much higher significance ( $P < 0.0001$ ). It means that *Fusarium* head blight infection and its effect on grain quality, toxins concentration and *Fusarium* biomass in kernels depended mainly on wheat growing environment. Resistance of cultivars to FHB was less important ( $P < 0.025$ ).



**Figure 2.** Biplot of the principal component analysis for 30 winter wheat cultivars grown in conventional (C) and organic field (O). Two first components explained 55.90% of variability of *Fusarium* head blight index (FHBi), dead seeds proportion (DS), ergosterol (ERG) and type A (Total TCT A) and type B (Total TCT B) trichothecenes content, and concentration of DNA (*Fusarium* DNA) of five *Fusarium* species in grain. Samples from conventional field marked with circles and from organic field with triangles.

Cultivars grown in organic field were compared for their overall performance under such conditions with respect to resistance to *Fusarium* infection. Multivariable analysis (K-means, discriminant analysis) made it possible to divide cultivars into three groups depending on their resistance to head infection, number of dead seeds, accumulation of ergosterol and *Fusarium* toxins in the grain as well as contamination of grain with *Fusarium* fungi (Figure 3, Table 8).



**Figure 3.** Discriminant analysis of 30 cultivars grown in organic field for *Fusarium* head blight index (FHBi), dead seeds proportion (DS), ergosterol (ERG) and type A (Total TCT A) and type B (Total TCT B) trichothecenes content, and concentration of DNA of five *Fusarium* species in grain (*Fusarium* DNA): (a) Observations on the factor axes with marked groups 1-3; (b) Correlation circle.

The most infected five cultivars were in the second group (Figure 3, Table 8). They could be described by the highest FHB index, high number of dead seeds, high accumulation of *Fusarium* toxins and the highest concentration of *Fusarium* biomass in kernels. Only amount of ERG was medium in grains of the cultivars of the group 2. The other 25 cultivars were in two close groups 1 and 3. They mainly differed in amount of ERG in grain, which was the highest in the group 1 while the lowest in the group 3.

The lowest overall infection showed cultivars from the group 3: 'Nateja', 'Mewa' and 'Markiza'. Regarding only accumulation of trichothecenes, it was the lowest in grain of 'Nateja' and 'Ostroga' from the group 3. It was low also in grain of cultivars 'Figura', 'Dorota', 'Mulan', 'Batuta', 'Tonacja' from the group 1, and surprisingly in grain of cultivar 'Belenus' from group of the most infected cultivars. Among the low-toxin accumulating cultivars, 'Ostroga', 'Batuta', 'Mulan', and 'Figura' were in the group of cultivars showing the highest grain yield per plot in organic field (Table S1). The lowest infected cultivar 'Nateja' had low grain yield caused by high yellow rust infection.

**Table 8.** Average values for groups shown in Figure 3A.

Group	Number of cultivars	FHBi (%)	DS (%)	ERG (mg kg <sup>-1</sup> )	Total TCT B (µg kg <sup>-1</sup> )	Total TCT A (µg kg <sup>-1</sup> )	<i>Fusarium</i> DNA (pg 100ng <sup>-1</sup> )
1	13	0.4	3.8	2.10	63	4	206
2	5	2.0	8.7	1.48	191	7	610
3	12	0.4	2.8	0.67	43	6	237

#### 4. Discussion

Until recently, the issue of organic farming was considered marginal. However, the constantly increasing acreage crops grown in this system and fast increasing percentage of consumers interested in obtaining the organic food encouraged a detailed address of this issue, which, so far, was recognized only partially. Thus, it was decided to provide field experiment for a representative sample of winter wheat cultivars for both systems of crops growing - conventional and organic under the same environmental conditions (location, time and weather). Of course, not all growing conditions were the same. This was mainly related to soil conditions. This was reflected in the studies presented by the analysis of the mineral elements. In most cases, we found significant differences in the concentrations occurring in the soil. A higher concentration in organic soil was found for Ca, Na, Si, B and Mn. Lower for K, Mg, Cd, Co, Cr, Cu, Ni and Zn. This is certainly due to organic cultivation resulting from the lack of use of mineral fertilizers. This leads to the relative impoverishment of the soil. However, we found higher concentration of Ca, Na, B and Si in soil from organic field. In review paper by Romero et al. [32] the authors found that silicon shows the beneficial effects on growth, development and health of crops. It activates the defense mechanisms of plants and increases tolerance to fungal diseases [33]. Concentrations of all analyzed metals (despite Mn) was lower in organic soil. Differences were significant for Cr, Cu, Ni and Zn. Depletion of the soil in organic cultivation system has an impact on the amount of soil microorganisms [34,35]. They are responsible for the biochemical processes, and thus, consequently, the resistance mechanism of plants. Studies on the reduction of the content of alkaline metals showed it leads to its acidification, which promotes the growth of micro-organisms for which the acidic (pH 4-5) is beneficial, including microscopic fungi and among them plant pathogenic species [36,37]. With this widespread phenomenon, we have to deal with in our research. These confirms our results indicating a higher concentration of K, Mg, Zn or Cd detected on fertilized, conventional plots. Much higher concentration of Zn and Cd was especially interesting.

The presented experiments concerned 30 cultivars of winter wheat, which were examined comprehensively for several years under conventional conditions to determine their susceptibility to *Fusarium* head blight and were divided into 4 groups as shown in Table 1 [20,38]. They constitute a complete cross-section of widely grown wheat cultivars in Poland, which gives the basis for determining them as model cultivars. In conditions of experimental year *Fusarium* head blight severity was low and average values for conventional and organic field did not differ significantly. Heading and flowering time were significantly earlier in organic field than in conventional. On average, plant height of wheat cultivars did not differ between two analyzed cropping systems. Effect of plant height of FHB severity was cultivar-dependent and similar in both systems. Taller cultivars were less FHB infected.

Sowing quality measured as germination energy and germination capacity of conventional material was significantly higher than for the seeds from organic cultivation system. The percentage's share of abnormal seedlings as well as share of dead seeds was significantly higher in organic seed material. It confirms results obtained for sowing value of conventional and organic oats [39]. Additionally, percentage's share of fresh ungerminated seeds was twice higher in organic seed material than in conventional one. However, difference was not statistically significant.

It is evident that there were significant differences in the seed quality obtained from both cultivation systems. For all parameters, the differences were important to the detriment of organic farming. Particularly large was the difference in germination energy, which for material from the ecological field was almost 25% lower than for the conventional. The above differences may also result

from higher colonization of kernels by mycobiota obtained from the ecological field. This was indicated by twice-higher ergosterol content (the total fungal quantity meter) and a three-fold higher *Fusarium* biomass content in organic grain. The content of ergosterol in grain depends on the type of grain (hulled, hull-less), cereal species and on the level of contamination of the grain with microscopic fungi, both pathogenic strains and native mycobiota [31,40,41]. Its content is affected also by the method of cultivation resulting from the use of fertilizers and treatments related to the use of plant protection chemicals [30].

Plant protection causes a disturbance of the natural homeostasis of the microbiota of kernel's surface, resulting in development of more expansive microbes that can dominate the environment. Thus, unfortunately, probiotic microorganisms, which are a natural barrier to pathogens, are completely removed [42,43]. The logical consequence is that, because of conventional agro-technical management, such crop is more susceptible to colonization by mycobiota. On the other hand, this increases competitiveness with pathogenic fungi producing specific fungal metabolites. Consequently, this has to do with the detection of a twice-higher concentration of ergosterol in organic material. Among the detected types of microscopic fungi, pathogenic ones represent a small percentage. This is related to the presence of large amounts of nonpathogenic mycobiota, which is a competition for pathogens [44,45].

Presence of DNA of six *Fusarium* species was detected in wheat grain. *F. langsethiae* was detected in six samples only in trace amounts. The highest amount of DNA was found as follows for *F. poae*, *F. graminearum*, *F. sporotrichioides*, *F. culmorum* and *F. avenaceum*. It was true in organic field. In conventional field, concentration of *F. culmorum* DNA was higher than *F. graminearum* and *F. sporotrichioides* DNA. The composition of *Fusarium* species was similar to that observed in last years in Europe [5,46–48].

Total *Fusarium* DNA concentration in organic samples was more than twice higher than in conventional samples, what can be explained by the cultivation system. In the case of organic cultivation, an environmental niche with a stabilized microorganism population is formed, which is enriched with probiotic organisms. Living in a symbiosis of microorganisms contribute to the improvement of soil condition, and thus naturally strengthen the resistance mechanisms of plants through i.a. mycorrhiza. In the conventional case, there are stress related to fertilization or the use of pesticides. Some fungi are eliminated, others often having a strong pathogenic remain. An analysis of the DNA content in the grain also gives a lot of interesting information. In the grain of organic farming, almost three times more DNA was found and stronger links between the contents of single species were identified. This was not true for *F. culmorum*, which could be related to the presence of *F. graminearum* in grain species being a competitor in the biosynthesis of type B trichothecenes.

Another legitimate conclusion here is that the testing of the DNA content of the grain is a much more accurate test method than the determination of the fungal concentration by an ERG analysis [49–51]. At the same time, it is emphasized that the amount of ERG gives a full image of the level of contamination with microscopic fungi. This is confirmed by the correlation factors for the ERG. They are, in all cases insignificant what confirms the above argument. Pathogenic fungi in the grain produce various metabolites and among them mycotoxins. This also occurs in the case of fungi of the genus *Fusarium*, which synthesize trichothecene toxins. This was also the case in the analyzed samples.

Based on the above results and conclusions is imposed another one. In grain of organic farming theoretically, the concentration of trichothecenes should be significantly higher. However, it was found that type A and B trichothecenes concentrations in both cases were similar and the differences were not significant. Vanova et al. [52] found higher concentration of DON in grain of wheat grown in three conventional systems. It was significantly higher in two systems where no chemical protection against FHB was applied. Similar tendency was found in barley and oats from organic and traditional farming [53–56]. In their review, Brodal et al. [56] concluded that contamination with *Fusarium* toxins of organically produced cereal grains was similar and sometimes lower than conventionally produced ones.

The established correlation coefficients for both groups were significant for the conventional system only. This is probably due to the fact, that for this system, more fungal biomass of *F. graminearum*

was found in the grain. That resulted in a higher correlation with the sum of type B trichothecenes which *F. graminearum* is an important producer.

The concentration of detected toxins was relatively small. Concentration of DON and T-2/HT-2 toxins was below the European limit and recommendation (Commission Regulation No. 1126/2007 of 28 September 2007; Commission Recommendation No. 2013/165/EU of 27 March 2013). Comparing the two cultivation systems, however, it is evident that in grain the average concentration of type B trichothecenes was lower in the case of organic trials. Differences were not statistically significant, but at the same time, concentration of *Fusarium* DNA was almost 3 times higher in organic grain. Although it can be also found similar data in other papers [52,57–60] it is a positive result. This proves once again that in the organic system determines the community of co-existing microorganisms is established. The most pathogenic and toxigenic are not predominant and environmental stress is not as harmful as the stress associated with significant doses of artificial fertilizers and pesticides as well as simplified rotations [61,62].

When considering the concentration of specific toxins, only concentration of 3-AcDON was significantly higher in conventional samples. Concentration of NIV was higher in samples from organic field; however, difference was not significant. Distributions for FUS-X and 3-AcDON in organic and conventional samples were significantly different. In conventional samples, these toxins were detected in higher amounts in single samples whereas they were more evenly distributed in organic samples. Amount of type A trichothecenes was very low and similar in conventional and organic samples, and they not differ significantly. Only average concentration of DAS was significantly higher in conventional samples.

Comparison of the sum of trichothecene toxins of groups A and B indicate environmental effects. Important correlations were obtained for the conventional system, by the fact that a strong pathogenic species *F. graminearum* stood out, being the most important producer of such toxins as DON, its derivatives and to a lesser extent (depending on the chemotype) NIV. For organic farming, the established coexistence of species was confirmed and no dominance of *F. graminearum* was found. When analyzing occurring mycobiota using ERG as a measure, no significant correlation was found for both environments with the other characteristics. In the case of *Fusarium* DNA testing, such correlations were found with the stronger link found for organic farming both between species (except *F. culmorum*) and other studied traits (mainly FHBi). The data presented is a significant contribution to understanding the philosophy of cultivation system and its effects. A similar method of reasoning and application may be found in paper of Lazzaro et al. [63].

By summarizing this aspect of the research, it is possible to identify clearly the relationship between the analyzed factors in the case of organic cultivation as stronger (Table 7). The above statements was confirmed by a comprehensive statistical analysis. It included a number of tests comparing analyzed populations based on factors such as FHBi, DS, ERG, Total TCT A and B and *Fusarium* DNA concentration. The designated *P*-value for the multidimensional Wilk's test had the value  $< 0.0001$ . *P*-value was similar for Fisher distances. It gives clear grounds for supporting the above conclusions indicating the different mechanism of reaction of plants on environmental stress of both cultivation systems. The MANOVA test was conducted to further validate these conclusions. It clearly showed that its effects depended on the type of cropping system in a very important way ( $P$ -value  $< 0.0001$ ), and to a much lesser extent on the cultivars used in the experiment ( $P$ -value  $< 0.025$ ). Similar observations can be found in the work of e.g. Newton et al. [64].

The issue of wheat cultivars applied in cultivation is often raised [3,4,65,66]. The most important question is whether the same cultivar can be used in both systems. During the study we wanted also, deepen this issue using 30 different cultivars with varying resistance to FHB. The possibility of successfully applying the same cultivar in both systems is becoming increasingly important, also for breeding reasons. Biplot of the principal component analysis shown in Figure 2 indicates the effects of cultivar on the results of the experiment.

It can be concluded that the results indicate a diversified behavior of cultivars, which was characterized by varying distances between cropping effects in two systems. Determined by

multidimensional scaling (MDS) method the average distances between pairs in the conventional and organic systems were for resistant cultivars (R) 0.576; medium resistant (MR) 2.335; medium susceptible (MS) 2.819 and susceptible (S) 3.547. This result is unambiguous and indicates that it is possible to use the cultivars used in conventional crops for organic farming [64,67].

The final stage of the study was comparison of the overall performance of cultivars grown under organic field conditions with respect to resistance to *Fusarium* infection. Using multivariable analysis (K-means, discriminant analysis), it was possible to divide cultivars into 3 groups depending on traits tested as indicated in Table 8 and Figure 3. The division on the three groups finds its justification both in the values shown in the table and separation because of their FHB susceptibility. For five cultivars ('Alcazar' (S), 'Muszelka' (S), 'Kampana' (S), 'Belenus' (MS), 'Slade' (MS)), significantly higher values (excluding ERG) have been obtained for all experimental traits. Discriminant analysis confirmed the condition of these cultivars, which already in other experiments showed low resistance after artificial inoculation of heads when they had high head infection and DON accumulation [20,38].

All the results presented indicate the usefulness of the above studies for the recommendation of individual cultivars to a particular growing method. Such studies requires evidently multiyear or multi location experiments to be fully reliable. Results show differences in effects of the conventional and organic system. The interesting preliminary results obtained, in the meaning of authors, will contribute to a better understanding of the processes of growth and development and effect of cereal farming in certain environmental conditions. They also allow for an objective look at organic farming and perhaps contribute to its rapid growth, as the idea of sustainable cultivation and avoidance of plant stress should gain new supporters.

**Supplementary Materials:** The following are available online. Table S1: Mean concentration (mg kg<sup>-1</sup>) of mineral compounds in soil samples of conventional and organic experimental fields. Table S2: Phenotypic characters of 30 winter wheat cultivars grown in conventional and organic fields. Table S3; Disease infection of 30 winter wheat cultivars grown in conventional and organic fields. Table S4: Sowing quality parameters for of 30 winter wheat cultivars grown in conventional and organic fields. Table S5: Concentrations of ergosterol (mg kg<sup>-1</sup>) and type B trichothecenes (µg kg<sup>-1</sup>) in grain of 30 winter wheat cultivars grown in conventional and organic fields. Table S6: Concentrations of type A trichothecenes (µg kg<sup>-1</sup>) in grain of 30 winter wheat cultivars grown in conventional and organic fields. Table S7: Concentration of DNA (pg 100ng<sup>-1</sup> of wheat DNA) of five *Fusarium* species in grain of 30 winter wheat cultivars grown in conventional and organic fields.

**Author Contributions:** Conceptualization, J.P. and T.G.; methodology, T.G., A.Ł., E.M. K.S.-Sz., M.B. and J.P.; formal analysis, T.G.; investigation, T.G., A.Ł., E.M., K.S.-Sz. and M.B.; writing—original draft preparation, T.G. and J.P.; writing—review and editing, T.G., A.Ł., E.M. and J.P.; supervision, J.P.; project administration, J.P.

**Funding:** This research was financed by National Science Centre (NCN), Poland. Project OPUS4 No. 2012/07/B/NZ9/02385.

**Conflicts of Interest:** The authors declare no conflict of interest

## References

1. Anonym Raport o stanie rolnictwa ekologicznego w Polsce w latach 2013–2014 [Condition of organic farming in Poland. The report 2013–2014]; Agricultural and Food Quality Inspection (GIJHARS), Warszawa 2015, Poland, 2015;
2. Borgen, A. Strategies for regulation of seed borne diseases in organic farming. *Seed Test. Int. - ISTA News Bull.* **2004**, *127*, 19–21.
3. Wolfe, M.S.; Baresel, J.P.; Desclaux, D.; Goldringer, I.; Hoad, S.; Kovacs, G.; Löschenberger, F.; Miedaner, T.; Østergård, H.; Lammerts Van Bueren, E.T. Developments in breeding cereals for organic agriculture. *Euphytica* **2008**, *163*, 323–346.
4. Letourneau, D.; van Bruggen, A.H.C. Crop protection in organic agriculture. In *Organic Agriculture: a Global Perspective*; Kristiansen, P., Acram, T., Reganold, J., Eds.; CABI Publishing, 2006; pp. 93–121.
5. Bottalico, A.; Perrone, G. Toxigenic *Fusarium* species and mycotoxins associated with head blight in small-grain cereals in Europe. *Eur. J. Plant Pathol.* **2002**, *108*, 611–624.

6. Bottalico, A. Fusarium diseases of cereals: Species complex and related mycotoxin profiles, in Europe. *J. Plant Pathol.* **1998**, *80*, 85–103.
7. Wagacha, J.M.; Muthomi, J.W. Fusarium culmorum: Infection process, mechanisms of mycotoxin production and their role in pathogenesis in wheat. *Crop Prot.* **2007**, *26*, 877–885.
8. Parry, D.W.; Jenkinson, P.; McLeod, L. Fusarium ear blight (scab) in small grain cereals—a review. *Plant Pathol.* **1995**, *44*, 207–238.
9. Dweba, C.C.; Figlan, S.; Shimelis, H.A.; Motaung, T.E.; Sydenham, S.; Mwadzingeni, L.; Tsilo, T.J. Fusarium head blight of wheat: Pathogenesis and control strategies. *Crop Prot.* **2017**, *91*, 114–122.
10. Bennett, J.W.; Klich, M. Mycotoxins. *Clin. Microbiol. Rev.* **2003**, *16*, 497–516.
11. Juan, C.; Ritieni, A.; Mañes, J. Occurrence of *Fusarium* mycotoxins in Italian cereal and cereal products from organic farming. *Food Chem.* **2013**, *141*, 1747–1755.
12. Vrček, I.V.; Čepo, D.V.; Rašić, D.; Peraica, M.; Žuntar, I.; Bojić, M.; Mendaš, G.; Medić-Šarić, M. A comparison of the nutritional value and food safety of organically and conventionally produced wheat flours. *Food Chem.* **2014**, *143*, 522–529.
13. Malmauret, L.; Parent-Massin, D.; Hardy, J.-L.; Vergey, P. Contaminants in organic and conventional foodstuffs in France. *Food Addit. Contam.* **2002**, *19*, 524–532.
14. Marx, H.; Gedek, B.; Kollarczik, B. Vergleichende Untersuchungen zum mykotoxikologischen Status von ökologisch und konventionell angebautem Getreide. *Z. Lebensm. Unters. Forsch.* **1995**, *201*, 83–86.
15. McKenzie, A.J.; Whittingham, M.J. Birds select conventional over organic wheat when given free choice. *J. Sci. Food Agric.* **2010**, *90*, 1861–1869.
16. Champeil, A.; Fourbet, J.F.; Doré, T.; Rossignol, L. Influence of cropping system on Fusarium head blight and mycotoxin levels in winter wheat. *Crop Prot.* **2004**, *23*, 531–537.
17. Harcz, P.; De Temmerman, L.; De Voghel, S.; Waegeneers, N.; Wilmart, O.; Vromman, V.; Schmit, J.F.; Moons, E.; Van Peteghem, C.; De Saeger, S.; et al. Contaminants in organically and conventionally produced winter wheat (*Triticum aestivum*) in Belgium. *Food Addit. Contam.* **2007**, *24*, 713–720.
18. Mäder, P.; Hahn, D.; Dubois, D.; Gunst, L.; Alföldi, T.; Bergmann, H.; Oehme, M.; Amadò, R.; Schneider, H.; Graf, U.; et al. Wheat quality in organic and conventional farming: Results of a 21 year field experiment. *J. Sci. Food Agric.* **2007**, *87*, 1826–1835.
19. Magkos, F.; Arvaniti, F.; Zampelas, A. Organic Food: Buying More Safety or Just Peace of Mind? A Critical Review of the Literature. *Crit. Rev. Food Sci. Nutr.* **2006**, *46*, 23–56.
20. Góral, T.; Stuper-Szablewska, K.; Buško, M.; Boczkowska, M.; Walentyn-Góral, D.; Wiśniewska, H.; Perkowski, J. Relationships between genetic diversity and *Fusarium* toxin profiles of winter wheat cultivars. *Plant Pathol. J.* **2015**, *31*, 226–244.
21. Ostrowska-Kolodziejczak, A.; Stuper-Szablewska, K.; Kulik, T.; Busko, M.; Rissmann, I.; Wiwart, M.; Perkowski, J. Concentration of fungal metabolites, phenolic acids and metals in mixtures of cereals grown in organic and conventional farms. *J. Anim. Feed Sci.* **2016**, *25*, 74–81.
22. ISTA International Rules for Seed Testing. **2016**, 2016, i-284(284).
23. Doyle, J.J.; Doyle, J.L. Isolation of plant DNA from fresh tissue. *Focus (Madison)*. **1990**, *12*, 13–15.
24. Turner, A.S.; Lees, A.K.; Rezanoor, H.N.; Nicholson, P. Refinement of PCR-detection of *Fusarium avenaceum* and evidence from DNA marker studies for phenetic relatedness to *Fusarium tricinctum*. *Plant Pathol.* **1998**, *47*, 278–288.
25. Nicholson, P.; Simpson, D.R.; Weston, G.; Rezanoor, H.N.; Lees, A.K.; Parry, D.W.; Joyce, D.; Centre, J.I.; Lane, C. Detection and quantification of *Fusarium culmorum* and *Fusarium graminearum* in cereals using PCR assays. *Physiol. Mol. Plant Pathol.* **1998**, *53*, 17–37.
26. Wilson, A.; Simpson, D.; Chandler, E.; Jennings, P.; Nicholson, P. Development of PCR assays for the detection and differentiation of *Fusarium sporotrichoides* and *Fusarium langsethiae*. *FEMS Microbiol. Lett.* **2004**, *233*, 69–76.
27. Parry, D.W.; Nicholson, P. Development of a PCR assay to detect *Fusarium poae* in wheat. *Plant Pathol.* **1996**, *45*, 383–391.
28. Perkowski, J.; Kiecana, I.; Kaczmarek, Z. Natural occurrence and distribution of *Fusarium* toxins in contaminated barley cultivars. *Eur. J. Plant Pathol.* **2003**, *109*, 331–339.
29. Young, J.C. Microwave-assisted extraction of the fungal metabolite ergosterol and total fatty acids. *J. Agric. Food Chem.* **1995**, *43*, 2904–2910.

30. Perkowski, J.; Wiwart, M.; Busko, M.; Laskowska, M.; Berthiller, F.; Kandler, W.; Krska, R. *Fusarium* toxins and total fungal biomass indicators in naturally contaminated wheat samples from north-eastern Poland in 2003. *Food Addit. Contam.* **2007**, *24*, 1292–1298.
31. Perkowski, J.; Buśko, M.; Stuper, K.; Kostecki, M.; Matysiak, A.; Sz wajkowska-Michałek, L. Concentration of ergosterol in small-grained naturally contaminated and inoculated cereals. *Biologia* **2008**, *63*, 542–547.
32. Romero, A.; Munévar, F.; Cayón, G. Silicon and plant diseases. A review. *Agron. Colomb.* **2011**, *29*, 473–480.
33. Dorneles, K.R.; Dallagnol, L.J.; Pazdiora, P.C.; Rodrigues, F.A.; Deuner, S. Silicon potentiates biochemical defense responses of wheat against tan spot. *Physiol. Mol. Plant Pathol.* **2017**, *97*, 69–78.
34. Wasilkowski, D.; Mrozik, A.; Piotrowska-Seget, Z.; Krzyzak, J.; Pogrzeba, M.; Plaza, G. Changes in enzyme activities and microbial community structure in heavy metal-contaminated soil under in situ aided phytostabilization. *Clean - Soil, Air, Water* **2014**, *42*, 1618–1625.
35. Ahmad, I.; Hayat, S.; Ahmad, A.; Inam, A.; Samiullah, I. Effect of heavy metal on survival of certain groups of indigenous soil microbial population. *J. Appl. Sci. Environ. Manag.* **2005**, *9*, 115–121.
36. Martyniuk, S.; Martyniuk, M. Occurrence of *Azotobacter* spp. in some Polish soils. *Polish J. Environ. Stud.* **2003**, *12*, 371–374.
37. Malik, A.A.; Puissant, J.; Buckeridge, K.M.; Goodall, T.; Jehmlich, N.; Chowdhury, S.; Gweon, H.S.; Peyton, J.M.; Mason, K.E.; van Agtmaal, M.; et al. Land use driven change in soil pH affects microbial carbon cycling processes. *Nat. Commun.* **2018**, *9*, 1–10.
38. Góral, T.; Walentyn-Góral, D. Variation for resistance to *Fusarium* head blight in winter and spring wheat cultivars studied in 2009–2016. Short communication. *Biul. IHAR* **2018**, *284*, 3–12.
39. Małuszyńska, E.; Mańkowski, D.R. Seed sowing value and response to drought stress of organic and conventional oat (*Avena sativa* L.) seeds during 5 years of storage. *Acta Sci. Pol. Agric.* **2016**, *15*, 27–36.
40. Wiwart, M.; Perkowski, J.; Budzyński, M.; Suchowilska, E.; Buśko, M.; Matysiak, A. Concentrations of ergosterol and trichothecenes in the grains of three *Triticum* species. *Czech J Food Sci* **2011**, *29*, 430–440.
41. Buśko, M.; Stuper, K.; Jeleń, H.; Góral, T.; Chmielewski, J.; Tyrakowska, B.; Perkowski, J. Comparison of volatiles profile and contents of trichothecenes group B, ergosterol, and ATP of bread wheat, durum wheat, and triticale grain naturally contaminated by mycobiota. *Front. Plant Sci.* **2016**, *7*, 1243.
42. Karlsson, I.; Friberg, H.; Kolseth, A.K.; Steinberg, C.; Persson, P. Organic farming increases richness of fungal taxa in the wheat phyllosphere. *Mol. Ecol.* **2017**, *26*, 3424–3436.
43. Lori, M.; Symnaczik, S.; Mäder, P.; De Deyn, G.; Gattinger, A. Organic farming enhances soil microbial abundance and activity—A meta-analysis and meta-Regression. *PLoS One* **2017**, *12*, 1–25.
44. Sapkota, R.; Knorr, K.; Jørgensen, L.N.; O’Hanlon, K.A.; Nicolaisen, M. Host genotype is an important determinant of the cereal phyllosphere mycobiome. *New Phytol.* **2015**, *207*, 1134–1144.
45. Hartman, K.; van der Heijden, M.G.A.; Wittwer, R.A.; Banerjee, S.; Walser, J.C.; Schlaeppi, K. Cropping practices manipulate abundance patterns of root and soil microbiome members paving the way to smart farming. *Microbiome* **2018**, *6*, 1–14.
46. Lukanowski, A.; Sadowski, C. *Fusarium langsethiae* on kernels of winter wheat in Poland — occurrence and mycotoxigenic abilities. *Cereal Res. Commun.* **2008**, *36*, 453–457.
47. Stenglein, S.A. *Fusarium poae*: A pathogen that needs more attention. *J. Plant Pathol.* **2009**, *91*, 25–36.
48. Xu, X.M.; Parry, D.W.; Nicholson, P.; Thomsett, M. A.; Simpson, D.; Edwards, S.G.; Cooke, B.M.; Doohan, F.M.; Brennan, J.M.; Moretti, A.; et al. Predominance and association of pathogenic fungi causing *Fusarium* ear blight in wheat in four European countries. *Eur. J. Plant Pathol.* **2005**, *112*, 143–154.
49. Reischer, G.H.; Lemmens, M.; Farnleitner, A.; Adler, A.; Mach, R.L. Quantification of *Fusarium graminearum* in infected wheat by species specific real-time PCR applying a TaqMan probe. *J. Microbiol. Methods* **2004**, *59*, 141–146.
50. Nicholson, P.; Turner, A.S.; Edwards, S.G.; Bateman, G.L.; Morgan, L.W.; Parry, D.W.; Marshall, J.; Nuttall, M. Development of stem-base pathogens on different cultivars of winter wheat determined by quantitative PCR. *Eur. J. Plant Pathol.* **2002**, *108*, 163–177.
51. Justesen, A.F.; Hansen, H.J.; Pinnschmidt, H.O. Quantification of *Pyrenophora graminea* in barley seed using real-time PCR. *Eur. J. Plant Pathol.* **2008**, *122*, 253–263.
52. Váňová, M.; Klem, K.; Míša, P.; Matušinsky, P.; Hajšlová, J.; Lancová, K. The content of *Fusarium* mycotoxins, grain yield and quality of winter wheat cultivars under organic and conventional cropping systems. *Plant, Soil Environ.* **2008**, *54*, 395–402.

53. Ibáñez-Vea, M.; González-Peñas, E.; Lizarraga, E.; López De Cerain, A. Co-occurrence of aflatoxins, ochratoxin A and zearalenone in barley from a northern region of Spain. *Food Chem.* **2012**, *132*, 35–42.
54. Edwards, S.G. *Fusarium* mycotoxin content of UK organic and conventional barley. *Food Addit. Contam. - Part A Chem. Anal. Control. Expo. Risk Assess.* **2009**, *26*, 1185–1190.
55. Edwards, S.G. *Fusarium* mycotoxin content of UK organic and conventional oats. *Food Addit. Contam. - Part A Chem. Anal. Control. Expo. Risk Assess.* **2009**, *26*, 1063–1069.
56. Remža, J.; Lacko-Bartošová, M.; Kosík, T. *Fusarium* mycotoxin content of Slovakian organic and conventional cereals. *J. Cent. Eur. Agric.* **2016**, *17*, 164–175.
57. Brodal, G.; Hofgaard, I.S.; Eriksen, G.S.; Bernhoft, A.; Sundheim, L. Mycotoxins in organically versus conventionally produced cereal grains and some other crops in temperate regions. *World Mycotoxin J.* **2016**, *9*, 755–770.
58. Hoogenboom, L.A.P.; Bokhorst, J.G.; Northolt, M.D.; van de Vijver, L.P.L.; Broex, N.J.G.; Mevius, D.J.; Meijjs, J.A.C.; Van der Roest, J. Contaminants and microorganisms in Dutch organic food products: A comparison with conventional products. *Food Addit. Contam. - Part A Chem. Anal. Control. Expo. Risk Assess.* **2008**, *25*, 1195–1207.
59. Bernhoft, A.; Clasen, P.-E.; Kristoffersen, A.B.; Torp, M. Less *Fusarium* infestation and mycotoxin contamination in organic than in conventional cereals. *Food Addit. Contam. Part A. Chem. Anal. Control. Expo. Risk Assess.* **2010**, *27*, 842–852.
60. Meister, U. *Fusarium* toxins in cereals of integrated and organic cultivation from the Federal State of Brandenburg (Germany) harvested in the years 2000–2007. *Mycotoxin Res.* **2009**, *25*, 133–9.
61. Schmid, F.; Moser, G.; Müller, H.; Berg, G. Functional and structural microbial diversity in organic and conventional viticulture: organic farming benefits natural biocontrol agents. *Appl. Environ. Microbiol.* **2011**, *77*, 2188–2191.
62. Bernhoft, A.; Torp, M.; Clasen, P.-E.-E.; Løes, A.-K.; Kristoffersen, A.B. Influence of agronomic and climatic factors on *Fusarium* infestation and mycotoxin contamination of cereals in Norway. *Food Addit. Contam. Part A. Chem. Anal. Control. Expo. Risk Assess.* **2012**, *29*, 1129–1140.
63. Lazzaro, I.; Moretti, A.; Giorni, P.; Brera, C.; Battilani, P. Organic vs conventional farming: differences in infection by mycotoxin-producing fungi on maize and wheat in Northern and Central Italy. *Crop Prot.* **2015**, *72*, 22–30.
64. Newton, A.C.; Guy, D.C.; Preedy, K. Wheat cultivar yield response to some organic and conventional farming conditions and the yield potential of mixtures. *J. Agric. Sci.* **2017**, *155*, 1045–1060.
65. Mason, H.E.; Spaner, D. Competitive ability of wheat in conventional and organic management systems: a review of the literature. *Can. J. Plant Sci.* **2006**, *86*, 333–343.
66. Osman, A.M.; Almekinders, C.J.M.; Struik, P.C.; Lammerts van Bueren, E.T. Adapting spring wheat breeding to the needs of the organic sector. *NJAS - Wageningen J. Life Sci.* **2016**, *76*, 55–63.
67. Scholten, O.E.; Steenhuis-Broers, G.; Timmermans, B.; Osman, A. Screening for resistance to *Fusarium* head blight in organic wheat production. In Proceedings of the Proceedings of the COST SUSVAR *Fusarium* workshop: *Fusarium* diseases in cereals – potential impact from sustainable cropping systems 01 - 02 June 2007 Velence, Hungary; Vogelgsang, S., Jalli, M., Kovacs, G., Vida, G., Eds.; Velence, Hungary, 2007; pp. 20–23.