

Article

# Associations of Various Nighttime Noise Exposure Indicators with Objective Sleep Efficiency and Subjective Sleep Quality: A Field Study

Martin Rösli<sup>1,2,\*</sup>, Mark Brink<sup>3</sup>, Franziska Rudzik<sup>4,5</sup>, Christian Cajochen<sup>4,5</sup>, Martina S. Ragettli<sup>1,2</sup>, Benjamin Flückiger<sup>1,2</sup>, Reto Pieren<sup>6</sup>, Danielle Vienneau<sup>1,2</sup>, Jean-Marc Wunderli<sup>6</sup>

- <sup>1</sup> Swiss Tropical and Public Health Institute, 4051 Basel, Switzerland
  - <sup>2</sup> University of Basel, 4003 Basel, Switzerland
  - <sup>3</sup> Federal Office for the Environment, 3003 Bern, Switzerland
  - <sup>4</sup> Centre for Chronobiology, Psychiatric Hospital of the University of Basel, 4002 Basel, Switzerland
  - <sup>5</sup> Transfaculty Research Platform Molecular and Cognitive Neurosciences, University of Basel, Switzerland
  - <sup>6</sup> Empa, Laboratory for Acoustics/Noise Control, Swiss Federal Laboratories for Materials Science and Technology, 8600 Dübendorf, Switzerland
- \* Correspondence: martin.roosli@swisstph.ch

**Abstract:** It is unclear which noise exposure time window and noise characteristics during nighttime are most detrimental for sleep quality in real life settings. We have conducted a field study with 105 volunteers wearing a wrist actimeter to record their sleep during seven days, together with concurrent outdoor noise measurements at their bedroom window. Actimetry recorded sleep latency increased by 5.6 minutes (95% confidence interval: 1.6 to 9.6 minutes) per 10 dB(A) increase in noise exposure during the first hour after bedtime. Actimetry assessed sleep efficiency was significantly reduced by 2–3 percent per 10 dB(A) increase in measured outdoor noise ( $L_{eq, 1h}$ ) for the last three hours of sleep. For subjectively reported sleepiness, noise exposure during the last hour prior to wake up was most crucial with an increase in the sleepiness score of 0.31 units (95% CI: 0.08 to 0.54) per 10 dB(A)  $L_{eq, 1h}$ . Associations for estimated indoor noise were not more pronounced than for outdoor noise. Considering noise events in addition to equivalent sound pressure levels ( $L_{eq}$ ) only marginally improved the statistical models. Our study provides evidence that matching the nighttime noise exposure time window to the individual’s diurnal sleep–wake pattern results in a better estimate of detrimental nighttime noise effects on sleep. We found that noise exposure at the beginning and the end of the sleep is most crucial for sleep quality.

**Keywords:** sleep quality; road traffic noise; actimetry; indoor noise; noise measurements; noise annoyance; noise sensitivity; time of day

## 1. Introduction

There is increasing epidemiological research demonstrating negative effects of transportation noise exposure on various chronic diseases such as cardiovascular disease [1–4], metabolic syndrome [5–9], depression [10–12], and cognitive functions [13–16]. Several mechanisms are implicated in these negative noise effects like the activation of the hypothalamic–pituitary–adrenal axis (HPA), which leads to increased cortisol and glucose levels as well as increased blood pressure, with consequences for blood viscosity and blood coagulation [17]. Also, chronic sleep deprivation is a stressor which contributes to allostatic load and has been found to be connected to all these outcomes above. High allostatic load, characterized by repeated stress responses, affects the brain regions involved in memory consolidation and affective processing with potential long term effects on cognitive and mental health [18]. In large population studies, impaired sleep quantity and

quality has been associated with increased risk of developing coronary heart diseases [19]. There is also solid evidence that short sleep duration, lack of slow wave sleep, and circadian desynchronisation of sleep increases the sensitivity to food stimuli contributing to adverse metabolic traits, in particular obesity and type 2 diabetes [20]. Further, sleep deprivation reduces the motivation for physical activity and thus reduces the energy expenditure [21], which further contributes to the risk for cardiometabolic syndromes.

Noise induced sleep disturbance as demonstrated in experimental human laboratory studies [22–24], field trials [25] and observational epidemiological studies [26,27] are thus likely to be on the pathway for detrimental effects on the cardiometabolic system and mental health. Strikingly, evidence for noise effects on various sleep outcomes is only considered “very low” to “moderate” in the recent systematic review of the World Health Organization Environmental Noise Guidelines for the European Region [28]. Thus, many questions remain open.

Sound pressure level at the ear of the sleeper is the relevant entity for quantifying noise effects on sleep. It is possible to quantify reactions of sleepers to noise in field experiments using contrived exposure settings (i.e. reproducing noise in a controlled fashion with loudspeakers). This approach, however, does not adequately consider long-term habituation to a noise source, which may be relevant in a real life situation at home. Observational studies would be appropriate to account for potential habituation effects, but are usually based on noise exposure modeling for the outdoor façade thus rather imprecise regarding noise exposure at the ear.

Further, the average sound pressure level ( $L_{eq}$  and similar indicators) may not be the only relevant factor; other noise exposure characteristics not captured in energy-based exposure indicators may also be important. For instance, in experimental sleep studies on noise effects on sleep, different effects have been observed for road, rail, and aircraft noise [22,23]. Thus, the effects of noise on sleep might either be better predicted by the number of noise events [29], the maximum sound pressure level [30], the sound pressure level slope [24,31] or by the order of events [31]. Confronted with the challenge of how to sum up and weight noise events we have developed an acoustical metric, the intermittency ratio (IR), to characterize short-term temporal variations of transportation noise exposure [32]. In a large cohort study, we found some evidence that IR may have a modifying effect on the cardiovascular mortality risk [33].

Timing of noise exposure is also considered to be relevant for sleep effects. For instance, an experimental sleep study observed that noise curfews at the end of the night are most beneficial for sleep, because noise induced sleep disturbances at the beginning of the night were at least partly compensated during the rest of the night [34].

In order to quantify relevant factors that affect sleep quality through noise exposure in a real life situation, we conducted an observational field study with volunteers wearing wrist actimeters to record their sleep-wake behavior during seven days with concurrent in- and outdoor noise exposure measurements. The study explored i) the relevance of indoor noise compared to outdoor noise, ii) the predictive contribution of IR to sleep effects in addition to equivalent continuous sound pressure levels ( $L_{eq,night}$ ), and iii) the effect of noise exposure at different times during the night. We also tested potential effect modification by noise annoyance, noise sensitivity, and sex.

## 2. Materials and Methods

### 2.1. Study population and procedures

Study participants were recruited from a previous nationwide survey on transportation noise annoyance and sleep disturbances among participants who expressed their willingness to be contacted for further research [35]. Only adults with German language skills were included. People with severe chronic disease or shift work were excluded. Calculated  $L_{den}$  for road traffic at the most exposed façade had to be at least 50 dB to ensure sufficient outdoor noise to be detected inside. In order to explore effects of road traffic noise exposure minimally contaminated by other noise sources, modeled aircraft and railway noise had to be at least 10 dB lower than road traffic noise.

During recruitment we ensured no other relevant noise sources such as construction work or tramways could disturb the sound level measurements.

After recruitment, study participants were visited at home between May and November 2016 and instructed about the procedures. All participants filled in a baseline questionnaire about sociodemographic and other relevant characteristics such as noise annoyance (using the 11-point ICBEN scale for road traffic noise in general [36]), noise sensitivity (using a 6-point Likert scale to rate the statement “I am noise sensitive” from NoiSeQ) [37], and window opening habits during night.

## 2.2. Sleep outcomes

Objective sleep behavior was evaluated using movement data collected by an actimeter device (Daqtometer v2.4 by Daqtix GbR, Oetzen, Germany) continuously worn on the non-dominant wrist during the seven day study period. Data were collected in 10-s to 60-s bins. Participants were provided with diaries in which they logged sleep-relevant information two times during the day. Prior to sleep, they gave information on consumption of coffee and alcohol, daytime naps, screen time use, and medication intake during the preceding day. In the mornings, participants evaluated the preceding sleep episode and noted bed and rise times, and whether or not they used an alarm clock. Additionally, subjective sleep quality was rated on a verbally anchored visual analogue scale from 0 (the worst sleep) to 100 (the best sleep). Subjective sleepiness was rated on a verbally anchored Likert-type scale that ranged from 1 (extremely alert) to 9 (very sleepy–fighting sleep) [38].

Actimetry data were analysed using the “Actiwatch Activity and Sleep Analysis” software (Actiwatch software version 7, Cambridge Neurotechnology Ltd., Cambridge, United Kingdom). Bed and rise times were entered manually based on actimetry records and sleep diary entries, while sleep on- and offset were determined automatically by the algorithm to derive sleep efficiency (proportion of actual sleep time per time in bed), sleep duration, sleep latency (time between bedtime and sleep onset), and moving time (minutes moving per assumed sleep period as time between sleep on- and offset).

## 2.3. Noise measurements and modelling

Noise exposure assessment included a seven-day outdoor measurement, a controlled short-term measurement to derive the sound insulation of each bedroom for different window positions and noise exposure modeling to identify eligible households. We modelled road traffic noise exposure calculation at the most exposed façade of each study participant using the sonROAD emission model [39] and the sound propagation model StL-86 as described in detail in Karipidis et al, 2011 [40].

Outdoor noise at the study participant’s bedroom window was measured with sound level meters type Noise-Sentry RT, a class II measurement device with a measurement uncertainty of about 1 dB(A) [41]. The sound level meters were flush mounted to the outer face of the closed window and logged A-weighted 1-second- $L_{eq}$ ’s. To obtain free field estimates, a correction of –6 dB was applied to the raw data. After a measurement period of seven days, the participants removed the sound level meters and sent them back. From the measurement files  $L_{eq,night}$  and  $IR_{night}$  (23:00–7:00) were computed for each night and participant. Further,  $L_{eq}$  and IR for *a priori* specified time windows during the night were derived: 19:00–23:00, 23:00–1:00, 1:00–5:00, 5:00–6:00, and 6:00–7:00. We also calculated, for each individual,  $L_{eq,1h}$  exposure for the first four hours of sleep as well as the last four hours prior to wake up taking into account their individual sleep pattern on the corresponding night.

Sound transmission of outdoor noise into the bedroom was measured during the home visit. Outdoor and indoor (at the position of the pillow) levels were measured in parallel for three window positions (open, tilted, closed) during three minutes in one-third octave bands from 50 Hz to 10 kHz with a class I sound level meter (type NTI XL2, NTi Audio AG, Schaan, Liechtenstein) and a free field microphone. Temporal resolution was set to one second. The indoor measurements

were compared with the concurrently conducted outdoor measurements and an algorithm was developed to derive the A-weighted attenuation factor in dB based on the correlation and the offset of these parallel measurements [42]. Finally, estimated indoor levels for analysis were obtained for each study participant by subtracting the individual attenuation factor from the outdoor measurements taking into account the preferred window position for the season when the data were collected, as indicated in the baseline questionnaire.

2.4. Data analysis

Primary endpoints were actimeter-derived sleep efficiency, sleep latency, sleep duration, and moving time. Secondary endpoints were subjective sleep quality and sleepiness as rated each morning in the sleep diary.

Association between sleep outcomes and noise exposure was analyzed using mixed regression models with auto-correlated residuals lag=1 and robust standard errors. To account for repeated nights from the same individual and multiple study participation per household a three level random intercept model was applied. Models were adjusted for confounding factors as indicated in the footnote of corresponding result tables (Table 3 and 4). Effect modification by noise annoyance and noise sensitivity was investigated by interaction analysis on dichotomized variables.

3. Results

For the study, we included 107 individuals from 96 households resulting in 720 nights with complete noise exposure data and recorded data on at least one sleep outcome (actimeter-derived or self-reported). Data from two individuals (14 nights) were excluded because their sleep was affected by their children. An additional 10 nights were excluded from the dataset because of sleeping out of home, two nights due to acute respiratory infection, and nine nights due to a recorded sleep duration of less than four hours. This left 694 nights from 105 individuals from 94 households for further analyses, although number of observations were somewhat smaller for specific outcomes because for two individuals only actimetry data was available, and for two individuals only the subjective sleep quality data.

Mean age of the study participants was 52.1 years (SD=14.4 years, age range: 23 to 78 years). Fifty-three participants (51%) were female and 52 male. Twenty-nine individuals (28%) had a University degree, 34 (32%) a higher education, 39 (37%) an apprenticeship, and 3 (3%) compulsory education. Median noise annoyance was 6 with 23% being classified as highly annoyed (score ≥ 8). Forty-one subjects (39%) tended to agree (score 4-6) to the statement “I am noise sensitive”.

Average sleep efficiency as recorded by actimetry was 88% and average sleep duration 7.0 h (Table 1). Subjective sleep quality scores ranged from 4 to 100 with a mean of 65. Average subjective sleepiness score was 4.1. Sleep efficiency was strongly negatively correlated with sleep latency (-0.79) and subjective sleep quality was negatively correlated with subjective sleepiness (-0.49). Correlations of all other outcomes were low (see Supplementary Table 1).

Table 1. Overview of the sleep outcomes (ACT=actimetry; SR=self-reported).

Outcome	N	Mean	SD	Min	Max
ACT Sleep efficiency [%]	634	88.4	7.9	46.3	98.2
ACT Sleep latency [min]	634	29.4	33	2	303
ACT Sleep duration [h]	634	7.0	1.2	4.1	11.7
ACT Moving time [%]	634	7.2	4.7	0.8	35.9
SR Sleep quality [0-100]	639	65	20	4	100
SR Sleepiness [1-9]	633	4.1	1.8	1	9

Table 2 shows the summary estimates for the various noise exposure metrics. Mean measured nighttime noise level ( $L_{eq,night}$ ) outside the window of the study participants' bedroom was 47.0 dB(A) with a maximum of 62.7 dB(A). Estimated mean indoor nighttime noise level at the pillow of the study participants was 30.2 dB(A) (maximum: 55.3 dB(A)). Measured exposure was about 10 dB(A) lower during 01:00 to 05:00 compared to the beginning and end of the night. Correlation between estimated indoor and measured outdoor nighttime exposure was 0.46 reflecting variation in noise attenuation between study participants. Correlations were  $\geq 0.63$  for measured  $L_{eq}$  between different time periods of the night (see Supplementary Table S3).

**Table 2.** Overview of the noise exposure data in dB(A) (for IR, see Supplementary Table S2).

Outcome	N	Mean	SD	Min	Max
Indoor $L_{eq,night}$	685	30.2	7.6	20.0 <sup>1</sup>	55.3
Outdoor $L_{eq,night}$	685	47.0	6.9	29.6	62.7
Outdoor $L_{eq,19-23}$	685	51.2	6.6	33.2	68.6
Outdoor $L_{eq,23-01}$	685	46.1	7.4	29.4	62.9
Outdoor $L_{eq,01-05}$	685	41.8	7.5	27.8	62.7
Outdoor $L_{eq,05-06}$	685	46.3	8.1	28.8	64.7
Outdoor $L_{eq,06-07}$	685	50.6	8.1	29.5	70.3

<sup>1</sup> all estimated indoor values <20 dB(A) have been replaced by 20 dB(A)

Measured nighttime noise ( $L_{eq,night}$ ) tended to be negatively associated with sleep efficiency and positively associated with sleep latency, although not statistically significant (Table 3). For instance sleep efficiency decreased by 1.11% (95%-CI: -2.44% to 0.21%) and sleep latency increased by 5.67 min (95% -CI: -1.00 to 12.34) per 10 dB(A) increase in  $L_{eq,night}$ . No indications of an association ( $p \geq 0.27$ ) were seen for the other sleep outcomes including subjective sleep quality and sleepiness.

**Table 3.** Adjusted association between sleep outcomes and measured outdoor nighttime noise ( $L_{eq,night}$ ) per 10 dB(A) increase; ACT=Actimetry; SR=Self reported,

Outcome	N	Coefficient	Confidence interval	p-value
ACT Sleep efficiency [%]	634	-1.11	-2.44 to 0.21	0.10
ACT Sleep latency [min]	634	5.67	-1.00 to 12.34	0.10
ACT Sleep duration [h]	634	0.01	-0.17 to 0.19	0.94
ACT Moving time [%]	634	-0.41	-1.13 to 0.31	0.27
SR Sleep quality [0-100]	639	-1.09	-4.96 to 2.78	0.58
SR Sleepiness [1-9]	633	0.02	-0.29 to 0.32	0.91

<sup>1</sup> adjusted for age, sex, education, evening caffeine intake, evening alcohol consumption, evening screen time, day of the week, season and whether woken up by an alarm clock

Using estimated indoor noise instead of measured outdoor noise yielded similar results in terms of regression coefficients, but none of the associations was even close to significance (Table 4). Similarly, we did not obtain any indications that IR was associated with any of the objectively recorded or subjectively reported sleep outcomes when also considering  $L_{eq,night}$  in the same model (see Supplementary Table S4). A model with quartiles of  $IR_{night}$  instead of a continuous variable suggests a non-linear association for sleep efficiency and latency. For the second quartile of  $IR_{night}$  (50-63%), sleep efficiency was reduced by 1.24% (95% confidence interval: -2.73 to 0.25) and sleep latency was increased by 3.48 minutes (95% CI: -2.54 to 9.50) compared to the first quartile (4-50%).




**Table 4.** Adjusted association between sleep outcomes and estimated *indoor* nighttime noise ( $L_{eq,night}$ ) per 10 dB(A) increase; ACT=Actimetry; SR=Self reported,

Outcome	N	Coefficient	Confidence interval	p-value
ACT Sleep efficiency [%]	634	-1.06	-2.86 to 0.74	0.25
ACT Sleep latency [min]	634	4.39	-5.54 to 14.32	0.39
ACT Sleep duration [h]	634	-0.06	-0.21 to 0.10	0.48
ACT Moving time [%]	634	-0.24	-0.90 to 0.42	0.47
SR Sleep quality [0-100]	639	0.21	-3.46 to 3.88	0.91
SR sleepiness [1-9]	633	-0.01	-0.28 to 0.26	0.95

<sup>1</sup> adjusted for age, sex, education, evening caffeine intake, evening alcohol consumption, evening screen time, day of the week, season and whether woken up by an alarm clock

Since most indications for noise effects were found for sleep efficiency and sleep latency in relation to outdoor noise, we investigated the effect of noise exposure in potential critical time windows for these two outcomes in more detail (Figure 1). Noise exposure in the evening (19:00-23:00) and in the early morning hours (05:00-06:00) was significantly associated with sleep efficiency, whereas the other time windows reached only borderline significance. For sleep latency, noise exposure until 01:00 was most relevant.




**Figure 1.** Association between measured outdoor noise exposure ( $L_{eq}$ ) at different time windows during night and changes in all-night sleep efficiency and sleep latency per 10 dB(A) increase in noise exposure. Significant associations are highlighted in red. Analyses adjusted for age, sex, education, evening caffeine intake, evening alcohol consumption, evening screen time, day of the week, season, and whether woken up by an alarm clock.


Exposure within fixed time periods (as shown above) may not match the individual time period a person is asleep or in bed. For instance, for somebody rising before 06:00, noise exposure between 06:00 and 07:00 is not relevant regarding sleep disturbance. We have thus calculated hourly noise exposure levels that matched the individual sleeping pattern of each night for the first four and the last four hours of sleep. The corresponding analyses with all actimetry-derived sleep outcomes are shown in Figure 2. This individualized noise exposure provides stronger associations between measured noise exposure and sleep efficiency. Sleep latency increased by five to seven minutes per 10 dB(A) increase in outdoor noise exposure during the first four hours after bedtime. For sleep efficiency, noise exposure during the last three hours prior to wake up was most critical with reductions of 2-3% per 10 dB(A) increase in measured outdoor noise exposure ( $L_{eq,1h}$ ). For sleep duration and moving time, no significant associations were found; although for the latter, noise exposure two to four hours after bedtime tended to increase moving time during sleep and noise exposure two hours before wake-up tended to be negatively associated with moving time.

Subjective sleepiness in the morning was significantly associated with noise exposure in the last hour of sleep, whereas no associations were observed for noise exposure at the beginning of sleep (Figure 3). For subjective sleep quality, none of the noise exposure time windows were significantly associated, but noise exposure in the middle of the sleeping period ( $\pm 4$  hours from bedtime and wake up) and at the end of the sleeping period showed the strongest trends for a relation. For IR matched to individualized sleep patterns, no significant effects on actimetry-derived and subjective sleep outcomes were observed.

Noise annoyance, noise sensitivity and sex were not significant effect modifiers for any of the outcomes and noise exposure time windows. A slight non-significant trend was seen for a stronger association of nighttime noise ( $L_{eq,night}$ ) with sleep efficiency for males ( $p=0.11$ ), people with a high ( $\geq$ median) annoyance score ( $p=0.23$ ), and people not reporting to be noise sensitive ( $p=0.25$ ).



**Figure 2.** Association between actimetry-derived outcomes and outdoor noise exposure for each hour after bedtime (+) or noise exposure for each hour prior to wake up (-). Changes refer to a 10 dB(A) increase in noise exposure in the respective hour. Same adjustments as indicated in Figure 1.



**Figure 3.** Association between self-reported outcomes recorded each morning in the sleep diary and outdoor noise exposure for each hour after bedtime (+) or noise exposure for each hour prior to wake up (-). Changes refer to a 10 dB(A) increase in noise exposure in the respective hour. Same adjustments as indicated in Figure 1.

4. Discussion

Our study suggests that the timing of noise exposure within the night is a relevant factor for the deterioration of objective and subjective sleep quality. Using individual noise exposure time windows, matched to the individual bed- and rise times of each night, provided stronger associations compared to fixed time intervals (such as e.g Leq between 19:00 and 23:00). Sleep latency, as expected, was most consistently associated with noise exposure at the beginning of the night, while for sleep efficiency and self-rated sleepiness noise exposure prior to wake up was most relevant. This suggests that noise exposure in the middle of the night may be less relevant for sleep quantity and quality, whereas noise exposure towards the end of the night, when sleep pressure is reduced and noise levels tend to be higher, is most disturbing for sleep.

This is in line with a field study using contrived aircraft noise exposure observing that aircraft noise events in the early morning elicited stronger reactions, as measured with high-resolution actigraphy, than events at the beginning of the sleep period [31]. Our findings are also in line with a large population based survey conducted 2009-2010 in Oslo with 13,019 participants. Road traffic nighttime noise was mostly associated with waking up too early and with difficulties falling asleep, but barely with awakenings during night [27]. In a Finnish study of 7,019 public sector employees, people exposed to >55 dB road traffic nighttime noise were more likely to report non-restorative sleep (OR=1.29, 95% CI: 1.01-1.65) and waking up too early (OR=1.24, 95% CI: 0.96-1.61) compared to people exposed to 45 dB or less. No associations were observed for frequently waking up during the night, short sleep duration, and difficulties falling asleep, the latter not in line with our results [43]. In a smaller Swiss survey of 1,375 adults, various self-reported sleep quality indices were not significantly related to road traffic nighttime noise [26]. However, most indications for an exposure-response trend were found for “waking up too early in the morning”, and least indications for “agitated sleep” and “waking phases during the night” in that same study [26].

Whether, or how, the observed pattern with stronger effects for evening and early morning noise translates into chronic health effects is unclear. Separating long-term noise effects regressed on average night noise exposure from time-specific effects in specific time periods is challenging for epidemiological studies, primarily because transportation noise in different time periods within the night is highly correlated, at least if no night curfews are in force. This is especially the case for modeled road traffic noise when traffic input data are based on traffic count samples, which are then extrapolated [2,3]. In reality, as demonstrated in our study, diurnal and day-to-day variation in traffic flows leads to a lower correlation between environmental noise exposures at different time intervals than one would observe between such computed standard metrics [44]. For large-scale



epidemiological studies on long-term risks, individual noise measurements as done here are not feasible. A Swiss cohort study on cardiovascular mortality has evaluated the effects of diurnal noise variation in their analysis by combining road, rail and aircraft noise. Although such a combination introduces additional diurnal variability due to different pattern and night curfews for aircraft noise, the correlation between different exposure time windows remained high ( $\geq 0.94$ ), precluding any firm conclusion. There was a trend that, for all cardiovascular causes combined, exposure during 01:00 to 05:00 was most relevant. For hypertensive related causes of death early morning noise (05:00-06:00) was most relevant; and for ischemic stroke and heart failure early evening and early morning noise were more detrimental than the rest of the night, although strongest association were seen with daytime noise [45].

We hypothesized that we would find stronger associations for estimated indoor noise than measured outdoor noise but could not confirm this with our data. This is in line with results from a survey on self-reported sleep disturbance in the same special issue of this journal [46]. However, our findings do not match the results of an epidemiological study done on indoor noise, which found stronger associations for hypertension with indoor noise compared to modeled noise at the most exposed façade [47].

We did not measure indoor noise, but rather applied an indirect procedure to estimate indoor noise levels from measured outdoor noise because our main interest was outdoor noise penetrating into the building. Direct indoor noise measurements would be heavily affected by the behavior of the participants. For instance, a sleepless person may produce some sound, which would yield a biased correlation between sleeplessness and noise exposure. There is no obvious explanation why indoor noise was less good a predictor than outdoor noise in our study. One may speculate that people who feel disturbed by outdoor noise close their bedroom window and thus have a lower indoor noise estimates, as this is the strongest predictor of indoor noise. This would mean that we deal with reverse causation in the sense that the sleep quality affects the noise exposure and not the other way round. Alternatively, estimated indoor noise levels may be subject to higher exposure misclassification as some levels were low. Estimated indoor noise levels below 20 dB(A) were censored and replaced with 20 dB(A) for the analysis; this was the case for 79 nights in 21 individuals. Finally, we obtained window opening habits for each season from the baseline questionnaire but did not specifically ask about the window position for each night, which may also add to exposure misclassification. Exposure misclassification may have resulted in reduced statistical power, which would explain the observed similar regression coefficients for indoor and outdoor noise but higher p-values for indoor noise.

We also hypothesized that sleep effects depend on the exposure characteristics. In particular, exposure situations with individual noise events clearly standing out from average (background) noise, as quantified with the Intermittency Ratio (IR) [32], were considered to be more detrimental for sleep than exposure to a steady sound level. However, we could not confirm that sleep disturbances are increasing with increasing IR. There was a non-significant non-linear pattern with lowest sleep efficiency for moderate (50-63%) levels of IR, which is in line with findings of the effect of IR in a cohort study on cardiovascular mortality [33]. However, in a cross-sectional survey on arterial stiffness, rather number of events during night was relevant [48]; and in a cohort study on diabetes IR was not associated with the diabetes risk [7]. Thus, it remains open whether IR is contributing to Leq-based metrics for predicting long term health effects of noise.

The strengths of this study include the prospective and detailed data collection with acquisition of objective sleep data (actimetry) and measurement of noise exposure. By measuring instead of calculating outdoor noise exposure, we could adequately record the diurnal variability of noise acting upon study participants. Our measurements allowed us to estimate outdoor noise passing into the sleeping rooms for individual bedroom characteristics. We could also match exposure time windows to the individual bed and rise times. The relatively small sample is a limitation and thus the power of the study rather limited. P-values of the analyses are not adjusted for multiple comparisons because we were interested in the pattern of the effect estimates rather than hypothesis testing. Thus some significant coefficients may in fact be chance findings. Note also

that some of the exposure and outcome measures are correlated and thus analysis should not be considered as mutually independent. This explains, for instance, the paradox result that noise exposure two hours prior to wake up is significant associated with sleep latency.

**5. Conclusions**

Our study provides evidence that matching the nighttime noise exposure time window to the individual’s diurnal sleep-wake pattern results in a better estimate of detrimental nighttime noise effects on sleep. The study suggests that noise exposure in the early morning hours is probably most crucial for a negative impact on objective sleep efficiency and subjective sleep quality. However, evening noise exposure was also associated with longer sleep latency. We could not confirm that noise induced sleep effects are better explained by indoor noise compared to outdoor noise, which might be due to reverse causality. This needs to be confirmed in larger studies.

**Supplementary Materials:** The following are available online at [www.mdpi.com/xxx/s1](http://www.mdpi.com/xxx/s1), Table S1: Pearson correlation matrix of sleep outcomes. Table S2: Summary of IR exposure data, Table S3: Pearson correlation matrix for measured nighttime noise exposure metrics, Table S4: Association between all outcomes and IRnight

**Author Contributions:** conceptualization, MRÖ, CC, JMW, and MB; methodology, MRÖ, MB, CC, FR, MRA, RP, BF, DV, JMW; formal analysis, BF, MRÖ; resources, RP, JMW; data curation, BF, FR, RP; writing—original draft preparation, MRÖ; writing—review and editing, all; visualization, MRÖ; funding acquisition, MRÖ, CC, and JMW.

**Funding:** This research was funded by the Federal Office for the Environment, grant number 00.0066.PZ/O513-1869 and the Swiss National Science Foundation, grant number 324730\_173330.

**Acknowledgments:** We thank Felix Schlatter and Selina Züst for their technical support in deriving the noise metrics and André Piquerez and Jannis Portmann for their engagement in the field work

**Conflicts of Interest:** The authors declare no conflict of interest.

**References**

- Heritier, H.; Vienneau, D.; Foraster, M.; Eze, I.C.; Schaffner, E.; de Hoogh, K.; Thiesse, L.; Rudzik, F.; Habermacher, M.; Kopfli, M., et al. A systematic analysis of mutual effects of transportation noise and air pollution exposure on myocardial infarction mortality: a nationwide cohort study in Switzerland. *European heart journal* **2019**, *40*, 598–603, doi:10.1093/eurheartj/ehy650.
- Sorensen, M.; Andersen, Z.J.; Nordsborg, R.B.; Jensen, S.S.; Lillielund, K.G.; Beelen, R.; Schmidt, E.B.; Tjønneland, A.; Overvad, K.; Raaschou-Nielsen, O. Road traffic noise and incident myocardial infarction: a prospective cohort study. *PLoS One* **2012**, *7*, e39283, doi:10.1371/journal.pone.0039283.
- Cai, Y.; Hodgson, S.; Blangiardo, M.; Gulliver, J.; Morley, D.; Fecht, D.; Vienneau, D.; de Hoogh, K.; Key, T.; Hveem, K., et al. Road traffic noise, air pollution and incident cardiovascular disease: A joint analysis of the HUNT, EPIC-Oxford and UK Biobank cohorts. *Environ Int* **2018**, *114*, 191-201, doi:10.1016/j.envint.2018.02.048.
- Hansell, A.L.; Blangiardo, M.; Fortunato, L.; Floud, S.; de Hoogh, K.; Fecht, D.; Ghosh, R.E.; Laszlo, H.E.; Pearson, C.; Beale, L., et al. Aircraft noise and cardiovascular disease near Heathrow airport in London: small area study. *Bmj* **2013**, *347*, f5432, doi:10.1136/bmj.f5432.
- Foraster, M.; Eze, I.C.; Vienneau, D.; Schaffner, E.; Jeong, A.; Heritier, H.; Rudzik, F.; Thiesse, L.; Pieren, R.; Brink, M., et al. Long-term exposure to transportation noise and its association with adiposity markers and development of obesity. *Environ Int* **2018**, *121*, 879-889, doi:10.1016/j.envint.2018.09.057.

- 387 6. Christensen, J.S.; Hjortebjerg, D.; Raaschou-Nielsen, O.; Ketzel, M.; Sorensen, T.I.; Sorensen, M.  
 388 Pregnancy and childhood exposure to residential traffic noise and overweight at 7 years of age. *Environ*  
 389 *Int* **2016**, *94*, 170–176, doi:10.1016/j.envint.2016.05.016.
- 390 7. Eze, I.C.; Foraster, M.; Schaffner, E.; Vienneau, D.; Heritier, H.; Rudzik, F.; Thiesse, L.; Pieren, R.;  
 391 Imboden, M.; von Eckardstein, A., et al. Long-term exposure to transportation noise and air pollution  
 392 in relation to incident diabetes in the SAPALDIA study. *International journal of epidemiology* **2017**, *46*,  
 393 1115–1125, doi:10.1093/ije/dyx020.
- 394 8. Ohlwein, S.; Hennig, F.; Lucht, S.; Matthiessen, C.; Pundt, N.; Moebus, S.; Jockel, K.H.; Hoffmann, B.  
 395 Indoor and outdoor road traffic noise and incident diabetes mellitus: Results from a longitudinal  
 396 German cohort study. *Environmental Epidemiology* **2019**, *3*, e037.
- 397 9. Sorensen, M.; Andersen, Z.J.; Nordsborg, R.B.; Becker, T.; Tjonneland, A.; Overvad, K.; Raaschou-  
 398 Nielsen, O. Long-Term Exposure to Road Traffic Noise and Incident Diabetes: A Cohort Study.  
 399 *Environmental health perspectives* **2013**, *121*, 217–222, doi:10.1289/ehp.1205503.
- 400 10. Beutel, M.E.; Junger, C.; Klein, E.M.; Wild, P.; Lackner, K.; Blettner, M.; Binder, H.; Michal, M.; Wiltink,  
 401 J.; Braehler, E., et al. Noise Annoyance Is Associated with Depression and Anxiety in the General  
 402 Population- The Contribution of Aircraft Noise. *PLoS One* **2016**, *11*, e0155357,  
 403 doi:10.1371/journal.pone.0155357.
- 404 11. Orban, E.; McDonald, K.; Sutcliffe, R.; Hoffmann, B.; Fuks, K.B.; Dragano, N.; Viehmann, A.; Erbel, R.;  
 405 Jockel, K.H.; Pundt, N., et al. Residential Road Traffic Noise and High Depressive Symptoms after Five  
 406 Years of Follow-up: Results from the Heinz Nixdorf Recall Study. *Environmental health perspectives* **2016**,  
 407 *124*, 578–585, doi:10.1289/ehp.1409400.
- 408 12. Seidler, A.; Hegewald, J.; Seidler, A.L.; Schubert, M.; Wagner, M.; Droge, P.; Haufe, E.; Schmitt, J.; Swart,  
 409 E.; Zeeb, H. Association between aircraft, road and railway traffic noise and depression in a large case-  
 410 control study based on secondary data. *Environmental research* **2017**, *152*, 263–271,  
 411 doi:10.1016/j.envres.2016.10.017.
- 412 13. Clark, C.; Crombie, R.; Head, J.; van Kamp, I.; van Kempen, E.; Stansfeld, S.A. Does traffic-related air  
 413 pollution explain associations of aircraft and road traffic noise exposure on children's health and  
 414 cognition? A secondary analysis of the United Kingdom sample from the RANCH project. *American*  
 415 *journal of epidemiology* **2012**, *176*, 327–337, doi:10.1093/aje/kws012.
- 416 14. Stansfeld, S.A.; Berglund, B.; Clark, C.; Lopez-Barrio, I.; Fischer, P.; Ohrstrom, E.; Haines, M.M.; Head,  
 417 J.; Hygge, S.; van Kamp, I., et al. Aircraft and road traffic noise and children's cognition and health: a  
 418 cross-national study. *Lancet* **2005**, *365*, 1942–1949, doi:10.1016/S0140-6736(05)66660-3.
- 419 15. van Kempen, E.; van Kamp, I.; Lebet, E.; Lammers, J.; Emmen, H.; Stansfeld, S. Neurobehavioral effects  
 420 of transportation noise in primary schoolchildren: a cross-sectional study. *Environ Health* **2010**, *9*, 25,  
 421 doi:10.1186/1476-069X-9-25.
- 422 16. Sygna, K.; Aasvang, G.M.; Aamodt, G.; Oftedal, B.; Krog, N.H. Road traffic noise, sleep and mental  
 423 health. *Environmental research* **2014**, *131*, 17–24, doi:10.1016/j.envres.2014.02.010.
- 424 17. Munzel, T.; Schmidt, F.P.; Steven, S.; Herzog, J.; Daiber, A.; Sorensen, M. Environmental Noise and the  
 425 Cardiovascular System. *J Am Coll Cardiol* **2018**, *71*, 688–697, doi:10.1016/j.jacc.2017.12.015.
- 426 18. McEwen, B.S. Sleep deprivation as a neurobiologic and physiologic stressor: Allostasis and allostatic  
 427 load. *Metabolism* **2006**, *55*, S20–23, doi:10.1016/j.metabol.2006.07.008.

19. Cappuccio, F.P.; Cooper, D.; D'Elia, L.; Strazzullo, P.; Miller, M.A. Sleep duration predicts cardiovascular outcomes: a systematic review and meta-analysis of prospective studies. *European heart journal* **2011**, *32*, 1484–1492, doi:10.1093/eurheartj/ehr007.
20. Schmid, S.M.; Hallschmid, M.; Schultes, B. The metabolic burden of sleep loss. *Lancet Diabetes Endocrinol* **2015**, *3*, 52–62, doi:10.1016/S2213-8587(14)70012-9.
21. DePorter, D.P.; Coborn, J.E.; Teske, J.A. Partial Sleep Deprivation Reduces the Efficacy of Orexin-A to Stimulate Physical Activity and Energy Expenditure. *Obesity (Silver Spring)* **2017**, *25*, 1716–1722, doi:10.1002/oby.21944.
22. Basner, M.; Muller, U.; Elmenhorst, E.M. Single and combined effects of air, road, and rail traffic noise on sleep and recuperation. *Sleep* **2011**, *34*, 11–23.
23. Elmenhorst, E.M.; Griefahn, B.; Rolny, V.; Basner, M. Comparing the Effects of Road, Railway, and Aircraft Noise on Sleep: Exposure(-)Response Relationships from Pooled Data of Three Laboratory Studies. *Int J Environ Res Public Health* **2019**, *16*, doi:10.3390/ijerph16061073.
24. Rudzik, F.; Thiesse, L.; Pieren, R.; Wunderli, J.M.; Brink, M.; Foraster, M.; Heritier, H.; Eze, I.C.; Garbazza, C.; Vienneau, D., et al. Sleep spindle characteristics and arousability from nighttime transportation noise exposure in healthy young and older individuals. *Sleep* **2018**, *41*, doi:10.1093/sleep/zsy077.
25. Elmenhorst, E.M.; Pennig, S.; Rolny, V.; Quehl, J.; Mueller, U.; Maass, H.; Basner, M. Examining nocturnal railway noise and aircraft noise in the field: sleep, psychomotor performance, and annoyance. *The Science of the total environment* **2012**, *424*, 48–56, doi:10.1016/j.scitotenv.2012.02.024.
26. Frei, P.; Mohler, E.; Röösli, M. Effect of nocturnal road traffic noise exposure and annoyance on objective and subjective sleep quality. *International journal of hygiene and environmental health* **2014**, *217*, 188–195, doi:10.1016/j.ijheh.2013.04.003.
27. Evandt, J.; Oftedal, B.; Hjertager Krog, N.; Nafstad, P.; Schwarze, P.E.; Marit Aasvang, G. A Population-Based Study on Nighttime Road Traffic Noise and Insomnia. *Sleep* **2017**, *40*, doi:10.1093/sleep/zsw055.
28. Basner, M.; McGuire, S. WHO Environmental Noise Guidelines for the European Region: A Systematic Review on Environmental Noise and Effects on Sleep. *Int J Environ Res Public Health* **2018**, *15*, doi:10.3390/ijerph15030519.
29. Janssen, S.A.; Centen, M.R.; Vos, H.; van Kamp, I. The effect of the number of aircraft noise events on sleep quality. *Appl Acoust* **2014**, *84*, 9–16.
30. Basner, M.; Samel, A.; Isermann, U. Aircraft noise effects on sleep: Application of the results of a large polysomnographic field study. *J Acoust Soc Am* **2006**, *119*, 2772–2784.
31. Brink, M.; Lercher, P.; Eisenmann, A.; Schierz, C. Influence of slope of rise and event order of aircraft noise events on high resolution actimetry parameters. *Somnologie* **2008**, *12*, 118–128.
32. Wunderli, J.M.; Pieren, R.; Habermacher, M.; Vienneau, D.; Cajochen, C.; Probst-Hensch, N.; Röösli, M.; Brink, M. Intermittency ratio: A metric reflecting short-term temporal variations of transportation noise exposure. *Journal of exposure science & environmental epidemiology* **2016**, *26*, 575–585, doi:10.1038/jes.2015.56.
33. Heritier, H.; Vienneau, D.; Foraster, M.; Eze, I.C.; Schaffner, E.; Thiesse, L.; Rudzik, F.; Habermacher, M.; Kopfli, M.; Pieren, R., et al. Transportation noise exposure and cardiovascular mortality: a nationwide cohort study from Switzerland. *Eur J Epidemiol* **2017**, *32*, 307–315, doi:10.1007/s10654-017-0234-2.

34. Griefahn, B.; Marks, A.; Robens, S. Experiments on the time frame of temporally limited traffic curfews to prevent noise induced sleep disturbances. *Somnologie* **2008**, *12*, 140-148.
35. Brink, M.; Schaffer, B.; Vienneau, D.; Foraster, M.; Pieren, R.; Eze, I.C.; Cajochen, C.; Probst-Hensch, N.; Röösli, M.; Wunderli, J.M. A survey on exposure-response relationships for road, rail, and aircraft noise annoyance: Differences between continuous and intermittent noise. *Environ Int* **2019**, *125*, 277-290, doi:10.1016/j.envint.2019.01.043.
36. **!!! INVALID CITATION !!! [36].**
37. Griefahn, B.; Marks, A.; Gjestland, T.; Preis, A. Annoyance and noise sensitivity in urban areas. In *Proceedings of ICA, Madrid*.
38. Akerstedt, T.; Gillberg, M. Subjective and objective sleepiness in the active individual. *International Journal of Neuroscience* **1990**, *52*.
39. Heutschi, K. SonRoad: new Swiss road traffic noise model. *Acta Acustica United with Acustica*, **2004**, *903*, 548-554.
40. Karipidis, I.; Vienneau, D.; Habermacher, M.; Köpfli, M.; Brink, M.; Probst-Hensch, N.; Röösli, M.; Wunderli, J.-M. Reconstruction of historical noise exposure data for environmental epidemiology in Switzerland within the SiRENE project. *Noise Mapping* **2014**, *1*, 3-14.
41. Schlatter, F.; Piquerez, A.; Habermacher, M.; Ragettli, M.S.; Röösli, M.; Brink, M.; Cajochen, C.; Probst-Hensch, N.; Foraster, M.; Wunderli, J.-M. Validation of large scale noise exposure modeling by long-term measurements. *Noise Mapping* **2017**, *4*, 75-86.
42. Locher, B.; Piquerez, A.; Habermacher, M.; Ragettli, M.; Röösli, M.; Brink, M.; Cajochen, C.; Vienneau, D.; Foraster, M.; Müller, U., et al. Differences between outdoor and indoor sound levels for open, tilted, and closed windows. *International Journal of Environmental Research and Public Health* **2018**, *15*, E149.
43. Halonen, J.I.; Vahtera, J.; Stansfeld, S.; Yli-Tuomi, T.; Salo, P.; Pentti, J.; Kivimäki, M.; Lanki, T. Associations between nighttime traffic noise and sleep: the Finnish public sector study. *Environmental health perspectives* **2012**, *120*, 1391-1396, doi:10.1289/ehp.1205026.
44. Brink, M.; Schaeffer, B.; Pieren, R.; Wunderli, J.M. Conversion between noise exposure indicators Leq(24h), L-Day, L-Evening, L-Night, L-dn and L-den: Principles and practical guidance. *International journal of hygiene and environmental health* **2018**, *221*, 54-63, doi:10.1016/j.ijheh.2017.10.003.
45. Heritier, H.; Vienneau, D.; Foraster, M.; Eze, I.C.; Schaffner, E.; Thiesse, L.; Rudzik, F.; Habermacher, M.; Köpfli, M.; Pieren, R., et al. Diurnal variability of transportation noise exposure and cardiovascular mortality: A nationwide cohort study from Switzerland. *International journal of hygiene and environmental health* **2018**, *221*, 556-563, doi:10.1016/j.ijheh.2018.02.005.
46. Brink, M.; Schäffer, B.; Vienneau, D.; Pieren, R.; Foraster, M.; Eze, I.C.; Rudzik, F.; Thiesse, L.; Cajochen, C.; Probst-Hensch, N., et al. Self-reported sleep disturbance from road, rail and aircraft noise: Exposure-response relationships and effect modifiers in the Swiss SiRENE study. *International Journal of Environmental Research and Public Health*. **submitted**.
47. Foraster, M.; Kunzli, N.; Aguilera, I.; Rivera, M.; Agis, D.; Vila, J.; Bouso, L.; Deltell, A.; Marrugat, J.; Ramos, R., et al. High blood pressure and long-term exposure to indoor noise and air pollution from road traffic. *Environmental health perspectives* **2014**, *122*, 1193-1200, doi:10.1289/ehp.1307156.
48. Foraster, M.; Eze, I.C.; Schaffner, E.; Vienneau, D.; Heritier, H.; Endes, S.; F., R.; L., T.; R., P.; C., S., et al. Exposure to road, railway, and aircraft noise and arterial stiffness in the SAPALDIA study: annual average noise levels and temporal noise characteristics. *Environmental health perspectives* **2017**, *125*, 097004, doi:10.1016/j.envint.2016.03.011.