

Article

Variation in responses of photosynthesis and apparent Rubisco kinetics to temperature in three soybean cultivars

James Bunce ¹

¹ Adaptive Cropping Systems Lab, USDA-ARS, Beltsville, MD, USA, retired; buncejames49@gmail.com

* Correspondence: buncejames49@gmail.com; Tel.: 410-451-7343

Abstract: Recent *in vivo* assays of the responses of Rubisco to temperature in C₃ plants have revealed substantial diversity. Three cultivars of soybean (*Glycine max* L. Merr.), Holt, Fiskeby V, and Spencer, were grown in indoor chambers at 15, 20, and 25 °C. Leaf photosynthesis was measured over the range of 15 to 30 °C, deliberately avoiding higher temperatures which may cause deactivation of Rubisco, in order to test for differences in temperature responses of photosynthesis, and to investigate *in vivo* Rubisco kinetic characteristics responsible for any differences observed. The three cultivars differed in the optimum temperature for photosynthesis (from 15 to 30 °C) at 400 μmol mol⁻¹ external CO₂ concentration when grown at 15 °C, and in the shapes of the response curves when grown at 25 °C. The apparent activation energy of the maximum carboxylation rate of Rubisco differed substantially between cultivars at all growth temperatures, as well as changing with growth temperature in two of the cultivars. The activation energy ranged from 58 to 84 kJ mol⁻¹, compared with the value of 64 kJ mol⁻¹ used in many photosynthesis models. Much less variation in temperature responses occurred in photosynthesis measured at nearly saturating CO₂ levels, suggesting more diversity in Rubisco than in electron transport thermal properties among these soybean cultivars.

Keywords: photosynthesis; Rubisco; temperature; acclimation; soybean

1. Introduction

The temperatures at which crop leaves accomplish photosynthesis vary diurnally, seasonally, and with geographic location. Intraspecific variation in the response and acclimation of photosynthesis to temperature has been studied extensively since the first infra-red CO₂ analyzers came into use in plant physiology [1, 2]. While photosynthetic response and acclimation to temperature has often been studied in crop species e.g. [3], and intraspecific variation in responses to brief, extreme temperature events has been documented [4-7] studies of intraspecific variation in photosynthetic response or acclimation to temperature in crops are rare. Intraspecific variation in response or acclimation of photosynthesis to temperature could prove a useful avenue for crop improvement or for the matching of plant physiological characteristics with climate. That strategy may avoid the inverse relationship between leaf size and photosynthetic rate commonly found in studies of intraspecific variation in photosynthesis in crops, which limits the usefulness of photosynthetic rate itself as a selection criterion [8].

Photosynthesis of C₃ species measured at high light and at the current atmospheric CO₂ concentration has an optimum temperature which may vary with species, and sometimes with growth temperature [9-11]. Growth at different temperatures may also affect the maximum photosynthetic rates without changing the shape of the temperature response curves or the optimal temperature [3, 10, 11]. Several recent studies have found considerable variation in the kinetic properties of Rubisco which affected the response of photosynthesis to temperature [10, 12-16], but none of these studies compared cultivars of soybean.

Soybean (*Glycine max* L. Merr.), while of sub-tropical origin, is grown in North America from southern Canada to the gulf coast states of the southern United States of America. Mean monthly temperatures during soybean growing seasons range from about 15 °C in the north to about 27 °C in the south, while midday mean temperatures during the growing seasons range from a low of about 20 °C in the early season in the north to about 32 °C in mid-season in the southern part of this range. Thus soybeans grown in North America are subject to a wide range of both growth and midday temperatures, even without considering extreme temperature events.

This study examined photosynthetic responses to temperature in three cultivars of soybean when grown at three growth temperatures in order to determine whether significant variation in response exists within soybeans, and to identify which kinetic parameters may be involved in any such variation.

2. Results

The rate of A measured at 400 μmol mol⁻¹ CO₂ was highest at temperatures which ranged from 15 to 30 °C, depending upon the cultivar and the growth temperature (Fig. 1). The 20 °C growth temperature produced leaves with the highest A, for measurement temperatures of 25 and 30 °C, for all three cultivars. The shape of the response of A to measurement temperature was least affected by growth temperature in Fiskeby V, and most affected in Spencer (Fig. 1). Only Spencer had highest A at 15 °C when grown at 15 °C. For the other two cultivars highest A at 15 °C occurred in leaves grown at 20 °C.

Fig. 1




Figure 1. Values of A measured at 400 $\mu\text{mol mol}^{-1}$ external CO_2 over a range of leaf temperatures, for plants of three cultivars, grown at 15, 20, or 25 °C.

When grown at 15 °C, A at a C_i of 200 $\mu\text{mol mol}^{-1}$, increased between 25 and 30 °C in Fiskeby V, and decreased between 25 and 30 °C in the other two cultivars (Fig. 2). Two-way ANCOVA indicated significant effects of cultivar, temperature and their interaction (Table A1). In contrast, values of A obtained at a measurement C_i of 500 $\mu\text{mol mol}^{-1}$, increased from 15 to 30 °C in all three cultivars (Fig. 2). Effects of cultivar and temperature were significant, but the interaction term was not significant (Table A2). For all measurement CO_2 conditions, A was highest in Spencer and lowest in Fiskey V at all temperatures, with Holt intermediate.

Fig. 2




Figure 2. Values of A measured at either C_i = 200 μmol mol⁻¹, or at C_i = 500 μmol mol⁻¹ CO₂ for three soybean cultivars grown at 15 °C.

When grown at 20 °C, A did not differ between cultivars at any temperature or measurement CO₂ condition, and increased between 15 and 30 °C (Fig. 3). For each measurement CO₂ condition,

two-way analysis of variance indicated a significant effect of measurement temperature, but no cultivar effects and no interaction effects (Tables A3 and A4).

Fig. 3




Figure 3. Values of A measured at either C_i = 200 μmol mol⁻¹, or at C_i = 500 μmol mol⁻¹ CO₂ for three soybean cultivars grown at 20 °C.

At the growth temperature of 25 °C, A at the measurement $C_i = 200 \mu\text{mol mol}^{-1}$ changed much less with temperature in Spencer than in Holt or Fiskeby V (Fig. 4). For measurements at $C_i = 200$, effects of cultivar, temperature, and their interaction were all significant (Table A5). At 500 $\mu\text{mol mol}^{-1}$ measurement C_i , all three cultivars had similar increases in A with temperature (Fig. 4), with only the effect of temperature being significant (Table A6).

Fig. 4




Figure 4. Values of A measured at either C_i = 200 μmol mol⁻¹, or at C_i = 500 μmol mol⁻¹ CO₂ for three soybean cultivars grown at 25 °C.

Mesophyll conductance did not vary substantially with either cultivar or growth temperature, but increased strongly with measurement temperature (Table 1).

Table 1. Mesophyll conductance to CO₂ movement from the intercellular airspace to the site of fixation (g_m) in three soybean cultivars grown at three temperatures, and measured over the range of 15 to 30 °C. Values are means for 3 or 4 leaves. Values followed by different letters are different at $P = 0.05$, using analysis of variance on log transformed data.

Cultivar	Growth Temperature (°C)	Mesophyll conductance (mol m ⁻² s ⁻¹)			
		Measurement Temperature (°C)			
		15	20	25	30
Holt	15	0.40d	0.93c	1.6b	2.4a
Fiskeby V	15	0.30d	0.72c	1.3b	2.6a
Spencer	15	0.28d	1.10c	1.5b	2.5a
Holt	20	0.31d	0.93c	1.7b	2.4a
Fiskeby V	20	0.35d	0.77c	1.5b	2.3a
Spencer	20	0.41d	0.95c	1.8b	2.6a
Holt	25	0.38d	0.85c	1.8b	2.3a
Fiskeby V	25	0.45d	0.75c	1.7b	2.3a
Spencer	25	0.33d	0.85c	1.6b	2.5a

The activation energy of $V_{C_{max}}$ was in all cases lower by 2 to 3 kJ mol⁻¹ when based on C_c compared with C_i . The activation energy of $V_{C_{max}}$ based on C_i was consistently lower in Spencer than in the other cultivars (Fig. 5). The activation energy did not change substantially with growth temperature in Spencer, but increased at growth temperatures of 20 and 25 °C in Fiskeby V and Holt (Fig. 5).

Fig. 5




Figure 5. The activation energy (E_a) of $V_{C_{max}}$ of Rubisco based on C_i , for three cultivars of soybean grown at three temperatures (15, 20, 25 °C).

3. Discussion

This work identified a wide range of responses of photosynthesis to growth and measurement temperature within only three commercial cultivars of soybean. Photosynthetic rates at high light and at air levels of CO_2 varied by at least a factor of 1.8 among the three cultivars at all measurement temperatures examined (15 to 30 °C), and the optimum temperatures for photosynthesis at air levels of CO_2 ranged from at least 15 to 30 °C.

The large impact that differences in the activation energy of $V_{C_{max}}$ can have on responses of photosynthesis to temperature is illustrated in Fig. 6, which shows photosynthetic rates at a C_c of 250 $\mu mol\ mol^{-1}$ for a fixed value of $V_{C_{max}}$ at 15 °C combined with different activation energies of $V_{C_{max}}$. At 30 °C, a 1.33x range in activation energy (from 60 to 80 $kJ\ mol^{-1}$) would result in a 1.56x range in A . The range of activation energy values for $V_{C_{max}}$ (58 to 84 $kJ\ mol^{-1}$) observed in this study is comparable to the variation among herbaceous species found by Hikoska et al. [10], and those reviewed by Kattge and Knorr [17], and also to that reported from temperature acclimation experiments with quinoa [18]. Others have also found variation in the activation energy of Rubisco among species from different climates [14], within the Triticeae [15] and within the Paniceae [16].

Fig. 6




Figure 6. Hypothetical rates of A at $C_c = 250 \mu\text{mol mol}^{-1}$, for a single value of $V_{C_{\text{max}}}$ at $15 \text{ }^\circ\text{C}$, combined with different activation energies (E_a , in kJ mol^{-1}) of $V_{C_{\text{max}}}$ of Rubisco.

In addition to differences in the response of $V_{C_{\text{max}}}$ of Rubisco to measurement temperature among the soybean cultivars studied here, approximately two-fold variation in the value of $V_{C_{\text{max}}}$ measured at $15 \text{ }^\circ\text{C}$ also occurred among these three cultivars when grown at 15 and $25 \text{ }^\circ\text{C}$ (Fig. 2 and 4). Variation in the response of $V_{C_{\text{max}}}$ to growth temperature among species and ecotypes is well known [3,19], but the possibility of intraspecific variation in crop species has received little attention to date.

Among these soybean cultivars, intraspecific variation in the temperature dependence of $V_{C_{max}}$ was much larger than differences in the temperature dependence of J_{max} . Consequently, the ratio of $V_{C_{max}}$ to J_{max} varied substantially among the cultivars, and with growth and measurement temperature, rather than being relatively constant, as suggested by some studies [20, 21]. The temperature dependence of photosynthesis at current air levels of CO_2 was much more closely related to $V_{C_{max}}$ than to J_{max} in these soybean cultivars, but that could change with rising atmospheric CO_2 .

Based on photosynthetic rates measured at current air levels of CO_2 , the cultivar Spencer seemed the best adapted to cool growth and measurement temperatures, as it had the highest rates of the three cultivars when grown at 15 °C and measured at 15 and 20 °C. The cultivar Holt seemed the best adapted to warm temperatures, having the highest photosynthetic rate among the three cultivars when measured at 30 °C, for plants grown at 25 °C. It may be useful to consider photosynthetic adaptation to temperature as an additional criterion for developing soybean cultivars for different locations.

4. Materials and Methods

Seeds of three cultivars of soybean (*Glycine max* L. Merr.), Holt, Fiskeby V, and Spencer were obtained from the USDA germplasm collection, and were grown in indoor, controlled environment chambers. Seeds were sown in 15 cm diameter plastic pots filled with a medium grade of vermiculite. Pots were flushed daily with a complete nutrient solution containing 14.5 mM nitrogen. Plants were grown in two M-12 chambers made by Environmental Growth Chambers (Chagrin Falls, Ohio) equipped with metal halide and high pressure sodium lamps. Twelve hours per day had light at 1000 $\mu\text{mol m}^{-2} \text{s}^{-1}$ photosynthetic photon flux density (PPFD). Air temperature was controlled at 15, 20, or 25 °C, with respective dew point temperatures of 8, 13, and 19 °C. All three cultivars were grown together in each chamber, with temperatures randomly assigned to chambers in successive "runs". Each "run" had three pots of each cultivar, with one plant per pot. A total of nine "runs" were grown in order to obtain data on all of the various photosynthetic parameters. Photosynthetic characterization was accomplished using third main stem trifoliolate leaves within a few days after those leaves had reached maximum area, when tests showed that photosynthetic properties were stable over several days.

Basic responses of photosynthesis to temperature were obtained by measuring responses of A to external CO_2 concentrations from 100 to 1200 $\mu\text{mol mol}^{-1}$ at temperatures of 15, 20, 25, and 30 °C. In all cases 1200 $\mu\text{mol mol}^{-1}$ CO_2 was saturating to A . Higher temperatures, which could lead to deactivation of Rubisco [22], were deliberately avoided. These measurements were made on three or four plants of each cultivar for each growth temperature. Gas exchange measurements were made with a CIRAS-3 portable photosynthesis system (PPSystems, Amesbury MA) operated within a controlled environment chamber. During the gas exchange measurements, leaf temperature was controlled to ± 0.3 °C, the PPFD was 1500 $\mu\text{mol m}^{-2} \text{s}^{-1}$, and the leaf to air water vapor pressure deficit ranged from about 0.9 kPa at 15 °C to about 1.4 kPa at 30 °C. The temperature of the controlled environment chamber was set to match the target leaf temperature, and the chamber PPFD was 1000 $\mu\text{mol m}^{-2} \text{s}^{-1}$.

Responses of A to sub-stomatal CO_2 concentration (C_i) were determined using either traditional steady-state measurements at external CO_2 concentrations of 400, 100, 150, 200, 250, 300, 400, 500, 600,

800, 1000, and 1200 $\mu\text{mol mol}^{-1}$ sequentially, or transient measurements during linear ramping of CO_2 concentrations from 100 to 1200 $\mu\text{mol mol}^{-1}$ [23]. The CO_2 ramping technique compares apparent CO_2 fluxes for an empty chamber with those when a leaf is present to obtain values of A at approximately 6 $\mu\text{mol mol}^{-1}$ CO_2 intervals. Because stomatal conductance did not change during the CO_2 ramping, values of C_i could be calculated for each value of A . Details of the CO_2 ramping method using the CIRAS-3 instrument, and examples comparing A vs. C_i curves obtained by ramping and by steady state measurements are given in Bunce (2018) [23]. The advantage of the CO_2 ramping method is that a complete A vs. C_i curve could be obtained in about 5 minutes, compared with about 30 minutes for a steady-state response curve. For each cultivar, growth temperature, and measurement temperature, comparisons were made of A vs. C_i curves obtained on the same leaf by the two methods to verify that photosynthetic parameters obtained by both methods did not differ substantially in this experiment.

Mesophyll conductance (g_m) for CO_2 movement from intercellular airspace to the site of fixation was measured for each growth and measurement temperature in all cultivars. Mesophyll conductance was determined from the oxygen sensitivity of photosynthesis in the Rubisco-limited region [24]. Because that method of measuring g_m depends upon knowing values of respiration in the light (R_l) and the CO_2 concentration at which carboxylation equals photorespiratory CO_2 release (Γ^*), R_l and Γ^* were also measured. Γ^* was measured from the intersection of A vs. C_i curves at high and low PPFD, using the precautions detailed by Walker and Ort (2015) [25]. R_l was determined by extrapolating A vs. C_i curves measured at 2% O_2 to zero C_i . The values of R_l and Γ^* measured for each leaf were then used to calculate g_m by the oxygen sensitivity method [24]. Prior work in soybean indicated that g_m did not vary with C_i [26]. The CO_2 concentration at Rubisco (C_c) was then calculated from A and C_i , using $C_c = C_i - A/g_m$ for each set of values of A and C_i .

The maximum carboxylation capacity of Rubisco ($V_{C_{\max}}$) was then estimated from the initial slopes of A vs. C_i and A vs. C_c curves for each leaf, growth temperature, and measurement temperature. The temperature dependencies of $V_{C_{\max}}$ based on C_i and C_c were summarized by their activation energies over the range of 15 to 30°C. Activation energy was calculated as the slope of $1/V_{C_{\max}}$ vs. $1/T$ (in °K). No deactivation term was used, since high temperatures causing deactivation were not used in this study.

Responses of photosynthesis to CO_2 for each cultivar, growth temperature, and measurement temperature were summarized as A at an external CO_2 (C_a) of 400 $\mu\text{mol mol}^{-1}$, which is approximately the current atmospheric CO_2 , A at $C_i = 200 \mu\text{mol mol}^{-1}$ as an indication of $V_{C_{\max}}$, and A at $C_i = 500 \mu\text{mol mol}^{-1}$, as an indication of the electron transport-limited A (J_{\max}). These parameters were calculated for each leaf from the A vs. C_i response curves, and two-way ANOVA was used to test for effects of cultivar, temperature, and their interaction separately for the three growth temperatures, for each photosynthetic parameter.

Funding: This research received no external funding.

Conflicts of Interest: The author declares no conflict of interest.

Appendix A

Table A1: Analysis of variance for photosynthetic rates measured at 200 $\mu\text{mol mol}^{-1}$ C_i for three soybean cultivars grown at 15 °C.

Source	Degrees of Freedom	Sum of Squares	Mean Square	F-Value	P-Value
Cultivar	2	278	139	196	<0.0001
Temperature	3	8.19	2.73	3.85	0.0221
C x T	6	51.6	8.60	12.2	<0.0001
Residual	24	17.0	0.708		

Table A2: Analysis of variance for photosynthetic rates measured at 500 $\mu\text{mol mol}^{-1}$ C_i for three soybean cultivars grown at 15 °C.

Source	Degrees of Freedom	Sum of Squares	Mean Square	F-Value	P-Value
Cultivar	2	890	445	113	< 0.0001
Temperature	3	1091	364	92.5	<0.0001
C x T	6	66.5	11.1	2.82	0.0320
Residual	24	94.3	3.93		

Table A3: Analysis of variance for photosynthetic rates measured at 200 $\mu\text{mol mol}^{-1}$ C_i for three soybean cultivars grown at 20 °C.

Source	Degrees of Freedom	Sum of Squares	Mean Square	F-Value	P-Value
Cultivar	2	0.945	0.472	0.484	0.626
Temperature	3	418	139	143	<0.0001
C x T	6	7.05	1.18	1.20	0.327
Residual	36	35.2	0.976		

Table A4: Analysis of variance for photosynthetic rates measured at 500 $\mu\text{mol mol}^{-1}$ C_i for three soybean cultivars grown at 20 °C.

Source	Degrees of Freedom	Sum of Squares	Mean Square	F-Value	P-Value
Cultivar	2	41.0	20.5	0.616	0.546
Temperature	3	2143	714.6	21.5	<0.0001
C x T	6	186	30.9	0.929	0.486
Residual	36	1198	33.3		

Table A5: Analysis of variance for photosynthetic rates measured at 200 $\mu\text{mol mol}^{-1} \text{C}_i$ for three soybean cultivars grown at 25 °C.

Source	Degrees of Freedom	Sum of Squares	Mean Square	F-Value	P-Value
Cultivar	2	3.50	1.75	3.00	0.0687
Temperature	3	303	101	173	<0.0001
C x T	6	93.5	15.6	26.7	<0.0001
Residual	36	14.0	0.583		

Table A6: Analysis of variance for photosynthetic rates measured at 500 $\mu\text{mol mol}^{-1} \text{C}_i$ for three soybean cultivars grown at 25 °C.

Source	Degrees of Freedom	Sum of Squares	Mean Square	F-Value	P-Value
Cultivar	2	30.5	15.3	14.1	<0.0001
Temperature	3	3068	1023	944	<0.0001
C x T	6	149.5	24.9	23.0	<0.0001
Residual	24	26.0	1.08		

References

1. Billings, W.D.; Godfrey, P.J.; Chabot, B.F.; Bourque, D.P. Metabolic acclimation to temperature in arctic and alpine ecotypes of *Oxyria digyna*. *Arctic Alpine Res.* **1971**, *3*, 277-299.
2. Mooney, H.A.; Bjorkman O.; Collatz G.J. Photosynthetic acclimation to temperature in the desert shrub *Larrea divaricata*. I. Carbon dioxide exchange characteristic of intact leaves. *Plant Physiol.* **1978**, *61*, 406-410.
3. Bunce, J.A. Acclimation to temperature in eight cool and warm climate herbaceous C_3 species: temperature dependence of parameters of a biochemical photosynthesis model. *Photosynthesis Res.* **2000**, *63*, 59-67.
4. Caulfield, F.; Bunce, J.A. Comparative responses of photosynthesis to growth temperature in soybean (*Glycine max* (L.) Merr.) cultivars. *Can. J. Pl. Sci.* **1988**, *68*, 419-425.
5. Camejo, D., Rodriguez, P., Angeles Morales, M., Dell'Amico, J.M., Torrecillas, A., Alarcon, J.J. High temperature effects on photosynthetic activity of two tomato cultivars with different heat susceptibility. *J. Pl. Physiol.* **2005**, *162*, 281-289.
6. Chovancek, E., Zivcak, M., Botyanszka, L., Hauptvogel, P., Yang, X., Misheva, S., Hussain, S., Brestic, M. Transient heat waves may affect the photosynthetic capacity of susceptible wheat genotypes due to insufficient photosystem I photoprotection. *Plants*, **2019**, *8*, 282.
7. Chen J.; Xu W.; Velten J.; Xin Z.; Stout J. Characterization of maize inbred lines for drought and heat tolerance. *J. Soil Water Conser.* **2012**, *67*, 354-364.
8. Bunce, J. A. Measurements and modeling of photosynthesis in field crops. *Crit. Rev. Plant Sci.* **1986**, *4*, 47-77.
9. Berry, J.A; Bjorkman O. Photosynthetic response and adaptation to temperature in higher plants. *Ann. Rev. Plant Physiol.* **1980**, *31*, 491-543.

10. Hikosaka, H.; Ishikawa, K.; Borjigidal, A.; Muller, O.; Onoda, Y. Temperature acclimation of photosynthesis: mechanisms involved in the changes in temperature dependence of photosynthetic rate. *J. Exp. Bot.* **2006**, *57*, 291-302.
11. Yamori, W.; Hikosaka, H.; Way, D.A. Temperature responses of photosynthesis in C₃, C₄, and CAM plants: temperature acclimation and temperature adaptation. *Photosynthesis Res.* **2014**, *119*, 101-117.
12. Silva-Perez, V.; Furbank, R.T.; Condon A.G.; Evans, J.R. Biochemical model of C₃ photosynthesis applied to wheat at different temperatures. *Plant Cell Environ.* **2017**, *40*, 1552-1564.
13. Galmes, J.; Hermida-Carreia C.; Laanisto, L.; Niinemets, U. A compendium of temperature responses of Rubisco kinetic traits: variability among and within photosynthetic groups and impacts on photosynthesis modelling. *J. Exp. Bot.* **2016**, *67*, 5067-5091.
14. Hermida-Carrera, C.; Kapralov, M.V.; Galmes, J. Rubisco catalytic properties and temperature response in crops. *Plant Physiol.* **2016**, *171*, 2549-2561.
15. Prins, A.; Orr, D.J.; Adralojic, P.J.; Reynolds, M.P.; Carmo-Silva, A.E.; Parry, M.A.J. Rubisco catalytic properties of wild and domesticated relatives provide for improving wheat photosynthesis. *J. Exp. Bot.* **2016**, *67*, 1827-1838.
16. Sharwood, R.E.; Ghannoum, O.; Kapralov, M.V.; Gunn, L.H.; Whitmey, S.M. Temperature responses of Rubisco from Paniceae grasses provide opportunities for improving C₃ photosynthesis. *Nature Plants.* **2016**, *2*, 16186.
17. Kattge J.; Knorr W.: Temperature acclimation in a biochemical model of photosynthesis: a reanalysis of data from 36 species. *Plant Cell Environ.* **2007**, *30*, 1176-1190.
18. Bunce, J.A. Thermal acclimation of the temperature dependence of the V_{Cmax} of Rubisco in quinoa. *Photosynthetica.* **2018**, *56*, 1171-1176.
19. Slatyer, R.O.; Ferrar, P.J. Altitudinal variation in the photosynthetic characteristics of snow gum, *Eucalyptus paucifloa* Sieb. Ex Spring. II. Effect of growth under controlled conditions. *Austral. J. Plant Physiol.* **1977**, *4*, 289-299.
20. Leuning, R. Scaling to a common temperature improves the correlation between the photosynthetic parameters J_{max} and V_{Cmax}. *J. Exp. Bot.* **1997**, *48*, 345-347.
21. Wullschleger, S.D. Biochemical limitations to carbon assimilation in C₃ plants – a retrospective analysis of A/Ci curves from 109 species. *J. Exp. Bot.* **1993**, *44*, 907-920.
22. Crafts-Brandner, S.J.; Salvucci, M.E. Rubisco activase constrains the photosynthetic potential of leaves at high temperature CO₂. *Proc. Nat. Acad. Sci.* **2000**, *97*, 13430-13435.
23. Bunce, J. Three methods of estimating mesophyll conductance agree regarding its CO₂ sensitivity in the Rubisco-limited C_i range. *Plants.* **2018**, *7*, 62.
24. Bunce, J.A. Use of the response of photosynthesis to oxygen to estimate mesophyll conductance to carbon dioxide in water-stressed soybean leaves. *Plant Cell Environ.* **2009**, *32*, 2875-881.
25. Walker, B.I.; Ort, D.R. Improved method for measuring the apparent CO₂ photocompensation point resolves the impact of multiple internal conductances to CO₂ to net gas exchange. *Plant Cell Environ.* **2015**, *38*, 2462-2474.
26. Bunce, J.A. Variable responses of mesophyll conductance to substomatal carbon dioxide concentration in common bean and soybean. *Photosynthetica.* **2010**, *48*, 507-512.