

Article

Functional Mobility and Basic Motor Skills in Patients with Multiple Sclerosis and its Relation to the Anthropometrical Status and Body Composition Parameters

Edyta Matusik^{1,2*}, Aldona Augustak² and Jacek Durmala^{1,2}

¹ School of Health Science in Katowice, Chair and Department of Rehabilitation, Medical University of Silesia, Katowice, Poland; email: ematusik@sum.edu.pl

² Upper-Silesian Medical Centre, Department of the Therapeutic Rehabilitation, Katowice, Poland; email: reh@gcm.pl

* Correspondence: ematusik@wp.pl; Tel.: +48-32-3598241 (E.M.)

Abstract: *Background and objectives:* Patients with multiple sclerosis (MS) have many potential factors (spasticity, immobilization, glucocorticoids use) which can deteriorate the anthropometrical status and body composition and may have the potential impact on the functional mobility and basic motor skills improvement after physiotherapy. The aim of the study was to assess the functional mobility and basic motor skills in patients with MS and to correlate them with disability and anthropometrical status and body composition parameters. *Materials and Methods:* Timed Up-and-Go test (TUG), and six-minute walk test (6MWT) were performed in 36 patients with MS before and after 4 weeks of physiotherapy. Body mass index (BMI), waist to height ratio (W/HtR), and waist-to-hip ratio (WHR) were assessed in this group. Body composition was evaluated by bioelectrical impedance analysis (BIA) and fat mass (FAT), fat free mass (FFM), total body water (TBW) and predicted muscle mass (PMM) were expressed as percentage of body mass. Clinical status was assessed by EDSS and AI scales. *Results:* After physiotherapy, there was a significant improvement in functional mobility and basic motor skills assessed by total distance in 6MWT ($p < 0.001$) and in TUG trials ($p < 0.001$). Positive significant correlations were found between the results obtained in both tests (either before and after physiotherapy) vs. FFM, TBW and PMM, whilst worse results in functional mobility and basic motor skills correlated significantly with higher WHtR, WHR and FAT ($p < 0.05$). Clinical status (EDSS) were significantly related to the WHtR and body composition parameters with the same manner as the results in the either 6MWT and TUG. However, there were no significant relationships between BMI vs. either clinical status (EDSS, AI) and functional mobility tests results in patients with MS. *Conclusions:* Functional mobility and basic motor skills may be significantly improved during the physiotherapy, but they are related to the anthropometrical status and body composition of MS patients. Moreover, disability status is also significantly related to this parameters. Body composition deterioration seems to be the important target for the therapeutic intervention in MS patients. For proper nutritional status assessment in patients with MS, body composition analysis or WHtR instead BMI should to be used.

Keywords: multiple sclerosis; physical fitness; body composition; functional mobility; physiotherapy

1. Introduction

The introduction Multiple sclerosis (MS) is an immune-modulated disease characterized by inflammation and demyelination of the central nervous system [1]. Symptoms of MS include weakness and fatigue, spasticity, abnormal sensation, impaired coordination, mobility and ambulation, and depression. The etiology of MS is not fully understood, but the most popular theory is that MS is provoked by the complex interaction of genetic influences, and environmental or modifiable factors which are causative agents in the disease process. One of the recently underlined modifiable factors, is obesity [2,3]. Large scale population based study conducted in Swedish MS patients showed that subjects whose BMI exceeded 27 kg/m² at age 20 had a two-fold increased risk of developing MS compared with normal weight subjects [4]. To be able to properly assess nutritional status of MS patients either standard anthropometry or body composition analysis (by dual energy X-ray absorptiometry (DXA) or bioelectrical impedance analysis (BIA)) have to be performed. However, body composition assessment in people with MS have not been extensively studied. Limited studies have reported that body mass index (BMI) assessment may underestimate adiposity in patients with multiple sclerosis [5]. On the other hand, in the group of MS patients with BMI > 25 kg/m² there was significantly lower percentage of no evidence of disease activity (NEDA) after beta-interferon therapy than in the normal weight group [6]. Moreover, patients with multiple sclerosis (MS) have many other potential factors (spasticity, immobilization, glucocorticoids use) which can deteriorate the anthropometrical status and body composition and may have the potential impact on the walking ability and other basic motor skills in these persons. Available data showed that only few patients with MS may achieve recommended daily physical activity levels, and their walking ability (measured as the steps count by 30 minutes) was most strongly related to gait and balance measures [7]. The most widely use therapeutic methods for the improvement of functional mobility in MS patients is physiotherapy. Little is known about the potential relationships between the nutritional status of MS patients and the outcomes of physiotherapy. That's why, the aim of this cross-sectional study was to assess the functional mobility and basic motor skills in patients with MS before and after the 4 weeks rehabilitation and physical activity programme, and to correlate them with disability and anthropometrical status and body composition parameters as well.

2. Materials and Methods

2.1. Studied population

Included subjects (n=36, 24 females) had a definite diagnosis of MS according to the McDonald criteria [8] and preservation of at least some ambulatory function (Expanded Disability Status Scale [EDSS] 1.5–6.5, median score 4.5; age 53.83 ± 11.21 years); of these, 9 required a walking aid (i.e. a cane or crutches). Subjects had not experienced an exacerbation in the 30 days prior to rehabilitation and testing, and had no other medical conditions that interfered with walking (i.e. cardiological diseases, endocrine disorders, diseases of the musculoskeletal system, orthopaedic surgery). A routine medical examination was carried out on each patient on the day of admission. The assessment of neurological status on the EDSS and Ambulatory Index (AI) scale was performed by one neurologist (EM).

2.2. Physiotherapy

All subjects included in the study were subjected to rehabilitation lasting 6 days/week for four weeks. Every patients started and finished the exercises at the same time. The rehabilitation program conducted in an individual form was dependent on the clinical condition of the patient, but always included exercises: equivalent, motor coordination, active trunk and limbs. Each patient underwent training on a treadmill. The intensity of the exercises was individually regulated for each patient and carried out under the supervision of a physiotherapist. Duration, speed, exercise parameters varied depending on the individual possibilities covered by the study. The total exercise time was 60 to 90 minutes. In the afternoon, relaxation exercises, stretching, and exercises lasting 30 to 45 minutes were

carried out as part of group classes. The duration of individual exercises, the number of rests varied depending on the patient's exercise tolerance.

2.3. Anthropometric measurements and body composition analysis

A set of anthropometric measurements was recorded at the first day of physiotherapy. Standing height was measured by a wall-mounted Harpenden Stadiometer to the nearest 0.1 cm. Weight (in underwear) was measured with an electronic scale with readings accurate to 0.1 kg. Body mass index (BMI) was then calculated, using the standard formula (kilograms per meter squared). Waist and hip circumferences were also and either, waist to hip (WHR) and waist to height ratio (W/HtR) were calculated. Body composition parameters: fat mass (FAT), fat-free mass (FFM), predicted muscle mass (PMM) and total body water (TBW) were assessed (in kilograms [kg] or as percentage of body weight [%]) based on bioelectrical impedance using a segmental body composition analyzer (BC-420MA Tanita Europe BV, Hoofddorp, The Netherlands).

2.4. Functional mobility and basic motor skills testing

To assess the functional mobility and basic motor skills the six-minute walk test (6MWT) and Timed Up-and-Go test (TUG) were administered by the same investigator (AA). 6MWT was made according the methodology described by Goldman et al. [9]. Subjects walked, at maximal effort, back and forth in a 30-meter hallway turning round cones, and were allowed to use their habitual assistive devices. Total walked distance and heart rate (HR) at baseline and at the end of the attempt were registered. In the TUG the timed performance of getting up from a chair, walking 3 meters, turning around, and walking back to sit down again was assessed [10]. Three trials of the TUG test were performed. The Time was recorded in seconds, with the fastest of the three trials used for analysis. The both tests were carried out twice before and after the rehabilitation program and were performed under the same conditions.

2.5. Ethical considerations

The study was approved by the Ethics Committee of the Medical University of Silesia (KNW/0022/KB/179/17). All participants gave informed consent. Patient rights were also approved according to the Helsinki Declaration.

2.6. Statistical analysis

All data were distributed normally (assessed by Kolmogorov-Smirnov test). Differences in continuous variables between before vs. after physiotherapy were assessed by unpaired Student's t-test for independent variables with non-equal variances. One-way analysis of variance (ANOVA) was used to analyze any significant difference among the four nutritional status subgroups, i.e. underweight, normal weight, overweight, and obesity. Correlations between continues parametrical were based on linear Pearson's correlation coefficient. All statistical analysis was made with the Statistica™ 12 PL software and p value less than 0.05 was considered statistically significant.

3. Results

Baseline characteristics and anthropometric measurements of all studied MS patients are reported in Table 1.

Table 1. Clinical and anthropometrical characteristics of the study population

	<i>Studied population N = 36 (F/M 24/12)</i>			
	Mean	Minimum	Maximum	SD
<i>EDSS</i>	4.4	1.5	6,5	1.5
<i>AI</i>	2.7	0	9	2.5
<i>Height [cm]</i>	167.4	152.1	190.1	9.5
<i>Weight [kg]</i>	72.4	44.6	94.1	12.5
<i>BMI [kg/m²]</i>	25.89	17.2	39.2	4.6
<i>Waist c. [cm]</i>	93.8	70.2	117.3	11.8
<i>Hip c. [cm]</i>	102.4	85.1	127.3	8.3
<i>WHtR</i>	0.56	0.42	0.69	0.08
<i>WHR</i>	0.92	0.73	1.03	0.08
<i>FAT [kg]</i>	22.7	6.3	41.4	8.5
<i>FAT [%]</i>	30.9	12.9	45.2	8.9
<i>FFM [kg]</i>	49.9	36.5	70.5	9.7
<i>FFM [%]</i>	69.2	54.8	87.1	8.9
<i>TBW [kg]</i>	35.7	26	51.5	6.7
<i>TBW [%]</i>	49.4	38.8	60.6	5.9
<i>PMM [kg]</i>	47.5	34.7	67.3	9.3
<i>PMM [%]</i>	65.8	51.9	82.7	8.6

Abbreviations: EDSS – Expanded Disability Status Scale , AI – Ambulatory Index, BMI – body mass index, W/Ht – waist to height ratio, WHR – waist to hip ratio, FAT – fat mass, FFM – fat-free mass, TBW – total body water, PMM – predicted muscle mass

Functional mobility and basic motor skills in patients with MS, assessed by total distance at 6MTW and timed performance in TUG, improved significantly after 4 weeks of physiotherapy ($p < 0.001$), see Figures 1. & 2. The mean difference of total walked distance in 6MTW after physiotherapy was 14.05 ± 22.43 meters, while mean TUG difference was recorded as $-0.58 \pm 1,11$ seconds.




Figure 1. Total walking distance of the 6-minute walk test before (6MWT1) vs. after (6MWT2) physiotherapy




Figure 2. Timed Up-and-go test (TUG) results [s] before (TUG 1) vs. after (TUG 2) physiotherapy

For the next stage of the analysis, the study group was further divided according to the nutritional status i.e. underweight (BMI < 18.5 kg/m², N =2), normal weight (BMI = 18.5 – 24.9 kg/m², N=17), overweight (BMI = 25 – 29.9 kg/m², N=10), obesity (BMI ≥ 30 kg/m², N=7).

One-way analysis of variance (ANOVA) did not reveal significant differences in clinical status of patients with multiple sclerosis between four anthropometrical subgroups, neither for EDSS (Figure 3A) and AI (Figure 3B). However, a trend of gradual deterioration in the clinical status (assessed either as EDSS and AI) of patients with MS was observed along with an increase in BMI value, but these differences did not reach statistical significance.




Figure 3. Clinical status of MS patients expressed as EDSS [A] and AI [B] after the stratification to the different nutritional status subgroups

However, further analysis revealed the significant correlations between EDSS vs. WHtR and body composition parameters as FAT, FFM and PMM, while there were no significant correlations between BMI, waist and hip circumferences, WHR vs. neither EDSS and AI (Table 2.).

Table 2. Correlations of clinical status with individual anthropometric and body composition parameters in studied MS population (*p<0.05)

	<i>BMI</i>	<i>Waist c.</i>	<i>Hip c.</i>	<i>WHtR</i>	<i>WHR</i>	<i>FAT [%]</i>	<i>FFM [%]</i>	<i>PPM [%]</i>
EDSS	0.314	0.245	0.250	0.423*	0.114	0.408*	-0.407*	-0.398*
AI	0.328	0.165	0.296	0.330	-0.040	0.308	-0.307	-0.301

Abbreviations: EDSS – Expanded Disability Status Scale , AI – Ambulatory Index, BMI – body mass index, W/Ht – waist to height ratio, WHR – waist to hip ratio, FAT – fat mass, FFM – fat-free mass, TBW – total body water, PMM – predicted muscle mass

Table 3 summarizes the relationship between the age and clinical status of MS patient with the results obtained in 6MTW and TUG trials before and after physiotherapy. The age of MS patients correlates with the significant expected manner (negatively) with 6MTW total distance either before and after physiotherapy as well as with the TUG trials (positively). The same significant relationships were found in case of EDSS and AI. Moreover, the clinical status expressed as either EDSS and AI was significantly related to the heart rate at baseline and at the end of the every 6MTW trials.

Table 3. Correlations of age and clinical status of MS patients with results obtained in individual 6MTW and TUG trials before (1) and after (2) physiotherapy (*p<0.05;**p<0.001; #p<0.0001; ##p<0.00001)

	<i>6MTW 1</i>	<i>6MTW 1</i>	<i>6MTW 2</i>	<i>6MTW 2</i>	<i>6MTW 1</i>	<i>6MTW 2</i>	<i>TUG 1</i>	<i>TUG 2</i>
	<i>HR</i>	<i>HR</i>	<i>HR</i>	<i>HR</i>	<i>distance</i>	<i>distance</i>		
	<i>at baseline</i>	<i>at the end</i>	<i>at baseline</i>	<i>at the end</i>				
age	0.169	0.139	0.114	0.010	-0.561**	-0.558**	0.380*	0.389*
EDSS	0.357*	0.489*	0.254	0.264	-0.911##	-0.890##	0.645#	0.644#
AI	0.273	0.532**	0.153	0.216	-0.816##	-0.783##	0.555**	0.562**

Abbreviations: EDSS – Expanded Disability Status Scale , AI – Ambulatory Index, 6MTW – 6 minute walk test, HR – heart rate TUG – Timed Up-and-Go test

Table 4 is a summary of the relationship between the anthropometrical standard measurements (BMI, waist and hip circumferences, WHtR and WHR) with the results obtained in 6MTW and TUG trials before and after physiotherapy. BMI and hip c. did not correlate significantly with neither parameters from functional mobility tests performed. However, there were significant positive correlation between waist c. and HR at the end of either before and after physiotherapy performed 6MTW test. There were also significant correlation between abdominal obesity diagnosed as WHtR and 6MTW distance (negative) as well as TUG trials (positive).

Table 4. Correlations of individual anthropometric parameters with the results obtained in individual 6MTW and TUG trials before (1) and after (2) physiotherapy (*p<0.05)

	6MTW 1	6MTW 1	6MTW 2	6MTW 2	6MTW 1	6MTW 2	TUG 1	TUG 2
	HR	HR	HR	HR	distance	distance		
	at baseline	at the end	at baseline	at the end				
BMI	0.164	0.162	0.129	0.139	0.148	0.115	0.162	0.156
Waist c.	0.323	0.358*	0.294	0.375*	0.210	0.182	0.289	0.272
Hip c.	0.118	0.089	0.169	0.198	0.219	0.197	0.251	0.246
WHtR	0.228	0.251	0.199	0.224	-0.348*	-0.322*	0.351*	0.342*
WHR	0.323	0.410*	0.239	0.329*	0.095	0.076	0.182	0.164

Abbreviations: 6MTW – 6 minute walk test, HR – heart rate, TUG – Timed Up-and-Go test, BMI – body mass index, W/Ht – waist to height ratio, WHR – waist to hip ratio

Body composition parameters (FAT, FFM, TBW and PMM) findings were similar to the WHtR findings and reach the same significance. Larger adiposity has significant negative impact on the mobility functioning and basic motor skills in the studied MS population (Table 5.).

Table 5. This is a table. Tables should be placed in the main text near to the first time they are cited.

	6MTW 1	6MTW 1	6MTW 2	6MTW 2	6MTW 1	6MTW 2	TUG 1	TUG 2
	HR	HR	HR	HR	distance	distance		
	at baseline	at the end	at baseline	at the end				
FAT [%]	0.145	0.061	0.186	0.136	0.417*	0.415*	0.383*	0.372*
FFM [%]	-0.145	0.060	-0.186	0.136	0.412*	0.414*	-0.382*	-0.372*
TBW [%]	-0.145	0.009	-0.179	0.102	0.380*	0.377*	-0.352*	-0.341*
PMM [%]	-0.141	0.053	-0.182	0.133	0.414*	0.412*	-0.379*	-0.369*

Abbreviations: 6MTW – 6 minute walk test, HR – heart rate, TUG – Timed Up-and-Go test, FAT – fat mass, FFM – fat-free mass, TBW – total body water, PMM – predicted muscle mass

4. Discussion

The results of this cross-sectional study confirmed the significant correlations between clinical disability (expressed as EDSS) vs. nutritional status (expressed as WHtR and FAT, FFM and PMM), while there were no significant correlations between routinely performed BMI, waist and hip circumferences, and WHR. Our finding confirm date coming from the study by Pilutti et al, which showed that BMI assessment may cause underestimation of adiposity in patients with MS comparing with body composition assessed by DXA [4]. Moreover, in our study one-way analysis of variance (ANOVA) did not reveal significant differences in clinical status of patients with multiple sclerosis

between four anthropometrical subgroups (stratified by BMI). There was only a trend of gradual deterioration in the clinical status (assessed either as EDSS and AI) of patients with MS that was observed along with an increase in BMI value, but these differences did not reach statistical significance.

Areal BMD (aBMD) measured by dual-energy X-ray absorptiometry (DXA) is currently the gold standard not only for the diagnosis of osteoporosis but also for the body composition evaluation. However, a noninvasive body composition assessment technique is currently available, based on bioelectrical impedance analysis (BIA). A good correlation between BIA and DXA has been reported in estimating adiposity in the different groups of patients [11,12]. BIA is relatively simple, quick, non-invasive and readily accessible compared to other more sophisticated methods, such as quantitative computed tomography (qCT) or DXA. In the present study body composition parameters as FAT, FFM, TBW and PPM were significantly related to the total walked distance assessed in the 6MTW and with the level of functional mobility based on the timed performance of the TUG trials. These relationships persisted also after 4 weeks of the rehabilitation programme. Body composition analysis by BIA in MS patients has only been used in study conducted by Bromley et al [13] in only 20 MS patients. Their work focused mainly on the nutritional intake rather than nutritional status, and they found significant correlation between the diet comprising saturated fat (positive) and carbohydrates (negative) with distance covered during the 6MWT. However, body fat percentage assessed by BIA was only showed as a mean result for the studied population, but the authors did not show any detailed results related to BIA. Another aspect of body composition in MS patients is the possible higher risk for sarcopenia related to the level of disability. Our study showed significant positive correlation of all lean mass related parameters (FFM, TBW and PMM) to the longer total walking distance in 6MWT and shorter TUG trials. It is also important to realize that lean mass is strongly related to the bone mineral density (BMD) either in whole body and lumbar spine projection [14]. To summarize, our study strongly suggest that the larger adiposity has significant negative impact on the mobility functioning and basic motor skills in the studied MS population and may influence the physiotherapy outcomes in this patients group. However, in our study MS patients there were no significant relations between the level of functional mobility and basic motor skills improvement (expressed as delta of total distance in 6MWT and delta TUG) and the baseline disability level (EDSS, AI) and the baseline nutritional status. On the other hand, our data proved the efficacy of the 4 weeks rehabilitation programme on the functional capacity in the MS patients. Recent study by Feys et al. showed that exercise therapy in persons with MS not only improve functional mobility and aerobic capacity but also a positive impact on visuospatial memory, fatigue and quality of life [15].

Based on our study only anthropometrical parameter seems to be useful in nutritional status assessment in MS patients is the waist to height ratio (WHtR). In our study, WHtR correlated significantly with either total walked distance in 6MWT and TUG trials with the same manner as percentage of fat mass (FAT) assessed by BIA. WHtR is now widely studied with the aim to find relatively simple parameters of fat tissue distribution in connection with visceral obesity and its comorbidities. A recent analysis showed that WHtR is better than the waist to hip ratio (WHR) for the prognosis of visceral obesity and its comorbidities [16]. In our study WHR was only significantly related to the heart rate (HR) at the end of the 6MWT attempt independently from physiotherapy. The above findings confirm great importance of the body fat distribution, with the special risk for abdominal/visceral obesity, which can't be properly assessed by BMI itself.

Limitations of this study must be acknowledged. First, use of a cross-sectional design with relatively small number of participants (especially with underweight), can provide only correlations and not direct causative findings. Second, participants with better clinical status (lower EDSS scores) may have more social (dietary and behavioral) opportunities to change their body composition, by the interaction with friends and family and at work, which allows for more frequent episodes of dining outside of the house, and have still quite normal daily physical activity. In contrast,

participants with higher EDSS scores may be dependent on a caretaker for meal preparation and have definitely lower physical activity and are probably with the higher risk for the body composition deterioration (adiposity or sarcopenia). That's why, the future studies will have to be focused on the specific MS severity subgroups of the patients taking into account to other potentially confounding factors.

5. Conclusions

Functional mobility and basic motor skills may be significantly improved during the physiotherapy, but they are related to the anthropometrical status and body composition of MS patients. Moreover, disability status is also significantly related to this parameters. Body composition deterioration seems to be the important target for the therapeutic intervention in MS patients. For proper nutritional status assessment in patients with MS, body composition analysis or WHtR instead BMI should to be used.

Author Contributions: E.M. was responsible for conception and design of the study/experiment, choice of the patients, acquisition of the material and data, participation in the diagnostics procedures, creating the database, statistical analysis and interpretation of data, drafting the manuscript, elaborating references, and critical revision of the manuscript. A.A. was responsible for choice of the patients, participation in the diagnostics procedures, acquisition of the material and data, critical revision of the manuscript J.D. was responsible for drafting and critical revision of the manuscript, and supervision of the project on the basis of merit

Funding: This research was supported by the Medical University of Silesia grant (No. KNW-1-122/K/7/Z) founded by the Polish Ministry of Science.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Lassmann, H.; Bruck, W.; Lucchinetti, C.F. The immunopathology of multiple sclerosis: an overview. *Brain Pathology* **2007**, *17*, 210-218.
2. Ascherio, A.; Munger, K.L. Environmental risk factors for multiple sclerosis, part II: noninfectious factors. *Ann Neurol* **2007**, *61*, 504-513.
3. Palavra, F.; Almeida, L.; Ambrosio, A.F.; Reis, F. Obesity and brain inflammation: a focus on multiple sclerosis. *Obes Rev.* **2016**, *17*, 211-224.
4. Hedström, A.K.; Olsson, T.; Alfredsson, L. High body mass index before age 20 is associated with increased risk for multiple sclerosis in both men and women. *Multiple Sclerosis Journal* **2012**, *18*, 1334-1336.
5. Pilutti, L.A.; Motl, R.W. Body Mass Index Underestimates Adiposity in Persons With Multiple Sclerosis. *Archives of Physical Medicine and Rehabilitation* **2016**, *97*, 405-412.
6. Kvistad, S.S.; Myhr, K.M.; Holmøy, T.; Šaltytė Benth, J.; Wergeland, S.; Beiske, A.G.; Bjerve, K.S.; Hovdal, H.; Lilleås, F.; Midgard, R.; Pedersen, T.; Bakke, S.J.; Michelsen, A.E.; Aukrust, P.; Ueland, T.; Sagen, J.V.; Torkildsen, Ø. Body mass index influence interferon-beta treatment response in multiple sclerosis. *Journal of Neuroimmunology* **2015**, *288*, 92-97.
7. Cavanaugh, J.T.; Gappmaier, V.O.; Dibble, L.E.; Gappmaier, E. Ambulatory Activity in Individuals With Multiple Sclerosis. *JNPT* **2011**, *35*, 26-33.
8. Polman, C.H.; Reingold, S.C.; Banwell, B.; Clanet, M.; Cohen, J.A.; Filippi, M.; Fujihara, K.; Havrdova, E.; Hutchinson, M.; Kappos, L.; Lublin, F.D.; Montalban, X.; O'Connor, P.; Sandberg-Wollheim, M.; Thompson, A.J.; Waubant, E.; Weinschenker, B.; Wolinsky, J.S. Diagnostic criteria for multiple sclerosis: 2010 revisions to the McDonald criteria. *Ann Neurol.* **2011**, *69*, 292-302. doi: 10.1002/ana.22366.
9. Goldman, M.D.; Marrie, R.A.; Cohen, J.A. Evaluation of the six-minute walk in multiple sclerosis subjects and healthy controls. *Multiple Sclerosis* **2008**, *14*, 383-390.
10. Podsiadlo, D.; Richardson, S. The timed "Up & Go": a test of basic functional mobility for trail elderly persons. *J Am Geriatr Soc* **1991**, *39*, 142-148.

11. de Lorenzo, A.; Sorge, S.P.; Iacopino, L.; Andreoli, A.; de Luca, P.P.; Stasso, G.F. Fat-free mass by bioelectrical impedance vs. dual-energy X-ray absorptiometry (DXA). *Applied Radiation and Isotopes* **1998**, *49*, 739-741.
12. Thomson, R.; Brinkworth, G.D.; Buckley, J.D.; Noakes, M.; Clifton, P.M. Good agreement between bioelectrical impedance and dual-energy X-ray absorptiometry for estimating changes in body composition during weight loss in overweight young women. *Clinical Nutrition* **2007**, *26*, 771-777.
13. Bromley, L.; Horvath, P.J.; Bennett, S.E.; Weinstock-Guttman, B.; Ray, A.D. Impact of Nutritional Intake on Function in People with Mild-to-Moderate Multiple Sclerosis. *Int J MS Care* **2019**, *21*, 1-9. doi: 10.7224/1537-2073.2017-039.
14. Mojtahedi, M.C.; Snook, E.M.; Motl, R.W.; Evans, E.M. Bone health in ambulatory individuals with multiple sclerosis: Impact of physical activity, glucocorticoid use, and body composition. *Journal of Rehabilitation Research & Development* **2008**, *45*, 851-862.
15. Feys, P.; Moundjian, L.; Van Halewyck, F.; Wens, I.; Eijnde, B.O.; Van Wijmeersch, B.; Popescu, V.; Van Asch, P. Effects of an individual 12-week community located "start-to-run" program on physical capacity, walking, fatigue, cognitive function, brain volumes, and structures in persons with multiple sclerosis. *Multiple Sclerosis Journal* **2019**, *25*, 92-103. DOI: 10.1177/1352458517740211.
16. Browning LM, Hsieh SD, Ashwell M.: A systematic review of waist-to-height ratio as a screening tool for the prediction of cardiovascular disease and diabetes: 0-5 could be a suitable global boundary value. *Nutrition Research Reviews* **2010**, *23*, 247-269.