

## Article

# Salinity Tolerance in *Fraxinus Angustifolia* Vahl. Seed Emergence in the Field and Germination Trials

Sabrina Raddi <sup>1\*</sup>, Barbara Mariotti <sup>1</sup>, Sofia Martini <sup>1</sup> and Alberto Pierguidi <sup>1</sup>

<sup>1</sup> DAGRI, Università degli Studi di Firenze; sabrina.raddi@unifi.it

\* Correspondence: sabrina.raddi@unifi.it

**Abstract:** The effect of salinity on seed germination/emergence in narrow-leaved ash (*Fraxinus angustifolia*) was studied both under field and laboratory conditions, in order to detect critical values to NaCl exposure. *Research Highlights:* Novel statistical methods in germination ecology has been applied (i) to determine the effects of chilling length and salinity (up to 150 mM NaCl) on *Fraxinus angustifolia* subsp. *oxycarpa* seed emergence, and (ii) to estimate threshold limits treating germination response to salinity as a biomarker. *Background and Objectives:* Salinity cut values at germination stage had relevant interest for conservation and restoration aims of Mediterranean floodplain forests in coastal areas subjected to salt spray exposure and/or saline water introgression. *Results:* Salinity linearly decreased germination/emergence both in the field and laboratory tests. Absence of germination was observed at 70 mM NaCl in the field and at 150 mM NaCl for 4-week (but not for 24-week) chilling. At 50 mM NaCl germination percentage was 50% (or 80%) of control for 4-week (or 24-week) chilling. Critical values for salinity were estimated between freshwater and 50 (75) mM NaCl for 4-week (24-week) chilling by Bayesian analysis. After 7-week freshwater recovery, critical cut-off values included all tested salinity levels up to 150 mM NaCl, indicating a marked resumption of seedling emergence. *Conclusions:* *Fraxinus angustifolia* is able to germinate at low salinity and to tolerate temporarily moderate salinity conditions for about two months. Prolonged chilling widened salinity tolerance.

**Keywords:** Mediterranean wetland; NaCl salinity; *Fraxinus angustifolia*; seed germination

---

## 1. Introduction

*Fraxinus angustifolia* Vahl. (narrow-leaved ash) is a typical tree of hygrophilous mixed-hardwood forests of seasonally inundated wetlands along rivers and alluvial plains [1,2]. This tree species has been usefully employed in remediation [3,4] and urban greening programs [5–7] for its high ability to resist waterlogging [8], adverse conditions of soil contamination and for its moderate tolerance to salinity at seedling and tree stage [9–11], but nevertheless information on seed germination behavior under salinity conditions are still lacking. Floodplain ashwoods are included in "interest" (Habitat 91F0 – Riparian mixed forests along the great rivers) or "priority" (Habitat 91E0\* – Alluvial forests) Natura 2000 lists under the EU Habitats Directive (Annex I 92/43/EEC [12]), both relevant for conservation of vulnerable temporary water ecosystems in Mediterranean basin [13,14]. In this region and despite conservation efforts, contraction and degradation risk of wetlands and swamps are still ongoing processes, accounting more than 50% loss since 1900, with 10% decrease in recent times, from 1975 to 2005 [15–18]. The global conservation awareness for these ecosystems – which nowadays resulted in a protected status under the international biodiversity-related Multilateral Environmental Agreements [19] for more than one third of swamps, flooded forests and coastal wetlands – might not be sufficient to prevent degradation risks arisen from interactions with large-scale processes, land use changes and agricultural land use intensification.

Human pressure, altered water flows, contaminants from internal urban and agricultural lands, and increased soil salinity [20–22] were particularly relevant in coastal floodplains. *F. angustifolia* coastal alluvial forests were mainly judged under 'unfavorable' (habitat 91F0) or 'inadequate' (habitat

91E0\*) state, following the 2007-2012 EU Habitat Directive Art. 17 assessment [23]. High fragmentation with connectivity loss among sites [24,25], potential susceptibility to diseases (e.g. ash dieback by *Hymenoscyphus fraxineus* [26]), changes in water regime, availability and quality due to pollution, salinity [27,28] and saltwater intrusion [29] were considered the major treats for this tree species.

International processes, such as the Convention on Biological Diversity (CBD) Aichi Targets and the UN Sustainable Development Goals (SDGs), pointed out the importance of undertaking practice and policy actions to prevent wetland degradation and loss. Consequently, management options that take advantage from science-based decision-making to face adverse climate-related impacts exacerbated by increasing human-induced pressures [30–32] were highly recommended and deemed urgent [33], particularly in coastal areas.

Within this perspective and for ensuring natural regeneration of *F. angustifolia*, our aim was to evaluate seed germination/emergence under increasing salinity conditions (in field and laboratory) to estimate NaCl tolerance threshold values at this life stage. Moreover, novel statistical methods for germination ecology (based on survival analysis and hierarchical Bayesian analysis) have been applied to infer cut off values for salinity threshold.

## 2. Materials and Methods

### 2.1. Germination in natural environment

Within the coastal hygrophilous mixed-hardwood forest (43.69° N, 10.29° E; Natural Reserve “Lame di fuori” – a complex of 655 ha of swamps and marshes [34–37] in San Rossore Park, Tuscany, Italy), three sites were sampled at different distance from the seaside, which differed in soil salinity (Table 1). Sampled sites were dominated by *F. angustifolia* subsp. *oxycarpa* both in number ( $n_{tree} = 54$ ) and basal area (BA = 71%), associated with field elm (*Ulmus minor*,  $n_{tree} = 38\%$ ; BA = 25%), black alder (*Alnus glutinosa*,  $n_{tree} = 1\%$ , BA = 2%), common hawthorn and common pear (together,  $n_{tree} = 7\%$ , BA = 1%). Average ( $\pm$  SD, standard deviation) for forest stand traits were: 18 ( $\pm 1$ ) m dominant height, 18 ( $\pm 1$ ) cm breast tree diameter, 51 ( $\pm 1$ ) years of age, 1104 ( $\pm 267$ ) trees  $ha^{-1}$  and 35 ( $\pm 7$ )  $m^2 ha^{-1}$  basal area. Narrow-leaved ash seedlings at cotyledon stage were counted at end June 2013 within 3 randomly transects (40 m x 1 m) per site. Ten soil cores per transect (0 - 20 cm, litter layer removed, spaced every 4 m) were sampled and soil salinity determined by ECe (electrical conductivity of soil saturation extract at 25°C), then NaCl concentration derived by  $NaCl_{(mM)} = 9.5 \cdot ECe_{(ds/m)} - 1.1327$  calibration equation. Soil samples had a sandy texture and pH (water) of 6.7 (CV=10%).

All statistical analyses were performed in R environment [38]. Normality of data distribution was tested (Shapiro-Wilk W-test [39] by shapiro.test() R function [40]). As most of the variables were not normally distributed, comparisons among groups were performed by nonparametric Wilcoxon rank-sum test by wilcox.test() R function [41]. Mean, standard error (SE) and 95% confidence interval range (95% CI) were calculated by bootstrapping applying boot.ci() function in 'boot' R package [42] along with the adjusted bootstrap percentile (BCa) method [43].

### 2.2. Germination trials

A bulk of narrow-leaved ash samaras was tested for seed vitality by TZ (tetrazolium) test [44,45]. Embryo development was assessed by embryo length to seed length ratio (E:S ratio) on 50 seeds. Samaras (intact pericarp) were manually graded to remove aborted seeds or empty samaras, soaked in water overnight, then moist-stratified at 4°C (chilling effect, C) for four (C<sub>4</sub>) or 24 (C<sub>24</sub>) weeks. Germination took place at 15°C/25°C (16 h dark/ 8 h light) for a period of 8 weeks, testing a range of salinity up to 150 mM NaCl (nacl effect, N<sub>x</sub> where x is NaCl in mM). Freshwater (control, N<sub>1</sub>) had naturally dissolved 1 mM NaCl together with other cations (1.5 mM Ca<sup>2+</sup>, 0.5 mM Mg<sup>2+</sup> and 0.1 mM K<sup>+</sup>). The salinity levels were obtained by adding NaCl to freshwater up to the concentration (in mM) indicated by N<sub>x</sub> subscript, that is N<sub>10</sub>, N<sub>50</sub> and N<sub>150</sub> for 4-week chilling (C<sub>4</sub>) and N<sub>50</sub>, N<sub>75</sub>, N<sub>100</sub> and N<sub>150</sub> for 24-week chilling (C<sub>24</sub>), respectively. About two hundred samaras per treatment (224) were

distributed in 8 germination boxes ( $\varnothing = 15$  cm [46]). Germination was assessed weekly as 'visible germination' (i.e. radicle protrusion  $> 1$  cm [47]). At the end of the germination test, C<sub>4</sub> samaras underwent a recovery germination test; samaras were rinsed three times for 10 minutes in flowing water, placed in N<sub>1</sub>, then followed for further 7 weeks.

**Table 1.** Descriptive statistics for soil salinity (0–20 cm) and number of *Fraxinus angustifolia* seedlings in "Lame di fuori" sampling sites (S1–S3). Distance from seaside (m); N = number of samples (soil) or transects (seedlings). W = *p*-value for Shapiro-Wilk normality test. ECe = soil ECe min–max range in dS/m). % = fraction (in percent) on the total of saline (ECe  $\geq 4$  dS/m) soil samples.

Site (m)	N	min – max	mean (SE)	95% CI		W	ECe	%
soil (mM NaCl)								
S1 890	30	32 – 95	55 (2.6)	a <sup>1</sup>	50 – 61	0.345	3.1 – 10.2	96
S2 1210	30	3 – 102	38 (5.6)	b	28 – 50	0.008	4.1 – 11.2	45
S3 1650	30	18 – 41	27 (0.8)	b	25 – 29	0.009	2.7 – 4.4	7
Total 1250	90	3 – 102	40 (2.4)		35 – 45	0.001	2.7 – 11.2	48
seedlings (ha <sup>-1</sup> )								
S1	3	0 – 500	167 (96)	b <sup>1</sup>	0 – 333	0.001		
S2	3	0 – 1500	667 (226)	ab	167 – 1000	0.415		
S3	3	0 – 2500	1250 (307)	a	583 – 1750	0.751		
Total	9	0 – 2500	694 (168)		389 – 1028	0.010		

<sup>1</sup> Wilcoxon test (different letters for *p* < 0.05).

### 2.3. Statistical analysis

Seed germination was assessed by survival analysis [48], where the response in terms of germination was defined as time-to-event [49,50]. Respect to the classic approach of calculating germination capacity and other germination indexes, this method offered the advantage of a proper statistical handling for time-repeated measures (namely the weekly counts of germinated seeds) together with an evaluation of treatment effects over the entire trial period [51]. An event was defined as the change from ungerminated (value = 0) to germinated status (value = 1); only viable seeds (assessed by cut test) were included in the counts. For each samara, the response variable *T* 'time until an event occurred' was counted in weeks since the beginning of the trial (germination- or recovery-test, respectively). The probability of germination per unit of time, *f*(*t*), was calculated by *T* probability density function (PDF); while the probability of changing status (germinating) within time *t* (*T*  $\leq$  *t*), *F*(*t*), was given by the cumulative distribution function (CDF). Here, the survival function, *S*(*t*) = 1 – *F*(*t*), returned the probability of not germinating within time *t*, that is the probability that a seed had to remain in its original (ungerminated) state for a time longer than *t* (*T* > *t*).

The hazard rate of germinating, *h*, was defined as the probability that an ungerminated seed (until the beginning of a time interval) had to germinate in the successive time interval (i.e. the conditional rate of germinating). It followed that the hazard function, *h*(*t*) = *f*(*t*)/*S*(*t*), returned the rate at which events occur for those seeds, that have not yet germinated at time *t*. The integral of *h*(*t*) over time gave the cumulative hazard rate function, *H*(*t*); the accumulated risk that a seed would face up to time *t*. Non parametric methods were used for estimating both survival function  $\hat{S}(t)$  (Kaplan and Meier method, KM [52]) and cumulative hazard rate function  $\hat{H}(t)$  (Nelson-Aalen method, NA [53–55]). These non-parametric estimators did not assume any underlying distribution and they returned non-parametric maximum likelihood estimators (NPMLE) for survival and hazard functions, resp., in right-censored data [56]. A 95% confidence interval (CI) for  $\hat{S}(t)$  was obtained by minimizing the likelihood function [57] within 'km.ci' R package [58], then it was derived for  $\hat{H}(t)$  by

$-\log \hat{S}(t)$  of 95% CI values. Hazard ratio, HR, could be interpreted as a relative measure of effects, as it compared survival between two treatments by  $S_0(t) = S_1(t)^{HR}$ , where  $S_0$  and  $S_1$  are the survival probabilities in control and in an alternative treatment, respectively. Therefore,  $HR = 1$  indicated no difference between treatments; HR values greater (or lower) than 1 indicated a higher (or lower) hazard of germinating in the control respect to the alternative treatment.

Chilling, salinity and their interaction effects on germination were tested by semi-parametric Cox Proportional Hazards regression model, Cox PH [59], applying `coxph()` function in 'survival' R package [60]. Cox PH model had no assumption about the shape of the hazard function, but it assumed proportional hazards rates among treatments, no time change in tested variables, lack of influential observations and linearity. Assumptions were verified by plotting scaled Schoenfeld residuals [61] against time [62]. The significance of model terms (*nacl*, *chilling*, *nacl*  $\times$  *chilling*) was assessed by Wald test; differences among treatment levels were compared by Peto-Peto-Prentice generalized Wilcoxon test – this latter did not require a consistent hazard ratio over time, but only that  $H(t)$  curves did not cross [50].

Backward stepwise regression using `drop1()` R function and likelihood ratio (LRT) were used for model reduction. Differences in goodness of fit for each model was then performed by log-likelihood, Akaike information criterion (AIC) and Bayesian information criterion (BIC), using `logLik()` and `stepAIC()` functions in 'MASS' R package, respectively [63]. In order to select the best model (most parsimonious; the simpler model with a high explanatory predictive power), the lowest (with difference exceeding 2) AIC (or BIC) and the highest likelihood values (LLV) were used. Cut off values for salinity in germination trials were determined fitting biomarker threshold models by 'bhm' R package version 1.15 [64,65], treating germination/emergence response to salinity (*nacl* levels) as biomarker. The aim of this hierarchical Bayesian method was to identify within the dataset a range of salinity (and its credible intervals), which were most likely to give a similar response to the best treatment in terms of germination probability.


### 3. Results

#### 3.1. Seed germination in natural environment

Soil salinity had an average ECe of 4.1 ( $\pm 0.3$  SE) dS m<sup>-1</sup> and 40 ( $\pm 2.4$  SE) mM NaCl with a wide variability in S1 and S2 sites, both including values higher than 10 dS m<sup>-1</sup> (Table 1). In transects, newly germinated seedling were found up to 53 mM NaCl soil salinity, and no seedling was found above 60 mM NaCl. Seedling density decreased linearly with soil salinity ( $r^2_{adj} = 0.466$ ,  $p < 0.026$ ,  $n$  transects = 9). Following this relationship, the ratio between intercept (1532.7 ha<sup>-1</sup>) and slope (-22.0 ha<sup>-1</sup> / mM NaCl) estimated the absence of seedlings at 70 mM NaCl. By *leave-one-out* validation procedure (i.e. removing one transect per time from the analysis), this estimate ranged between 65 and 73 mM NaCl and it had a bootstrapped 95% confidence interval range of 67-71 mM NaCl (Figure 1).

#### 3.2. Seed viability and embryo development

Seed lot had 70% viable, 22% non-viable, and 8% unclassified seeds (TZ test). Normality was observed only for embryo length in non-viable seeds and E:S ratio in viable seeds (W test,  $p > 0.14$ ;  $p > 0.32$ , resp.). The two viability classes did not differ for seed and embryo development (Wilcoxon test: seed length,  $p > 0.14$ ; embryo length,  $p > 0.63$ ; E:S ratio,  $p > 0.77$ ). Overall, seed and embryo length had an average of 17 mm ( $\pm 0.27$  SE; min-max: 10-22) and 12 mm ( $\pm 0.23$  SE; min-max: 6-15), resp., with E:S ratio of 0.7 ( $\pm 0.01$  SE; min-max 0.4-0.9). Prediction of embryo length by seed length was poor, even if significant ( $r^2_{adj} < 0.14$ ,  $p < 0.004$ ). E:S ratio was more influenced by embryo length increase (slope = 0.038,  $r^2_{adj} = 0.39$ ,  $r = +0.63$ ,  $p < 0.000$ ), than by seed length decrease (slope = -0.023,  $r^2_{adj} = 0.19$ ;  $r = -0.45$ ,  $p < 0.000$ ).




**Figure 1.** Salinity and seedling density at the three sites (S1–S3, see Table 1) in “Lame di fuori”.


### 3.3. Chilling and salinity effect in germination

After 8 weeks, germination was higher in C<sub>24</sub> than in C<sub>4</sub> ( $\hat{F}$ , 15% vs 9%,  $p < 0.01$ ; Figure 2a). Longer chilling treatment showed a marked higher germination hazard rate in the first week, while, thereafter, similar hazard rates indicated comparable germination between chilling treatments (Figure 2b). Final germination percentage ( $\hat{F}$ ) decreased with salinity, ranging from 0% to 12% in C<sub>4</sub>, and from 6% to 23% in C<sub>24</sub> (Figures 3a and 3b). Highest and promptest germination was observed in C<sub>24</sub> at low salinity ( $\leq 50$  mM NaCl). Complete absence of germination occurred only for C<sub>4</sub> at 150 mM NaCl. C<sub>24</sub> strongly germinated during the first-week, when hazard rate was more than 10 times higher respect to what observed in C<sub>4</sub> or at any other time in the trial (Figures 4a, 4b and 2). C<sub>24</sub> first week germination was linearly and positively related to final  $\hat{F}$  values (Figure 5a), and it had a marked effect on final  $\hat{F}$  value, particularly at high salinity (up to 89% of  $\hat{F}$  at 150 mM NaCl), representing almost all observed events, while at low salinity ( $\leq 50$  mM NaCl) it contributed only 53% of  $\hat{F}$  (Figure 5a inset).


Differently, the quicker chilling length, C<sub>4</sub>, showed germination during the first week only in control (hazard rate,  $h = 1.5\%$ ), and about 3 to 4 weeks were needed to reach at least 50% of  $\hat{F}$  for low salinity levels up to 50 mM NaCl. Apart from this initial difference, since second week forward hazard rates did not significantly differ between chilling treatments at any salinity level (paired Wilcoxon test,  $p > 0.43$ ), showing a ratio not different from unit both for control (N<sub>1</sub>,  $h_{C_4}/h_{C_{24}} \pm \text{SE}$ :  $1.04 \pm 0.38$ ,  $p > 0.43$ ) and common salinity levels, obtained by pooling N<sub>1</sub>, N<sub>50</sub> and N<sub>150</sub> together ( $h_{C_4}/h_{C_{24}} \pm \text{SE}$ :  $1.27 \pm 0.33$ ,  $p > 0.59$ ).



**Figure 2.** Germination test. Chilling treatments (C<sub>24</sub> and C<sub>4</sub>) averaged over common salinity levels (N<sub>1</sub>, N<sub>50</sub> and N<sub>150</sub>), mean  $\pm$  95% CI bands or bars for (a) germination probability,  $F(t)$  and (b) hazard rate,  $h(t)$ .



**Figure 3.** Cumulative germination probability,  $F(t)$ , with salinity ( $N_x$ ;  $x = \text{NaCl}$  in mM) during germination (a, b) and recovery (c). Mean  $\pm$  95% CI band for  $N_1$  (light gray),  $N_{50}$  (diagonal stripes) and  $N_{150}$  (dark gray). Same symbols in all plots.



**Figure 4.** Hazard function,  $h(t)$ , with salinity during germination (a, b) and recovery (c). Mean  $\pm$  95% CI bars. Same symbols in all graphs.

Salinity (*nacl*) and chilling length (*chilling*) tested by Cox PH model were significant, without a significant interaction effect (Table 2). Hierarchical Bayesian model '*bhm*' and LRT analysis confirmed these results (Tables 2 and 3). BIC (but not AIC) indicated a better goodness of fit if interaction was omitted from the statistical model. When individually considered, *nacl* contributed more than *chilling* to model goodness of fit (Table 3).

In Cox PH model the assumption of proportional hazard held for *nacl* ( $p > 0.93$ ) and interaction effect ( $p > 0.53$ ), but not for *chilling* ( $p < 0.004$ ), mainly because of the high hazard rate during the first week in C<sub>24</sub> for any of the salinity level (Figures 4a and 4b; Table 4). PH assumption rejection for *chilling* suggested to test salinity effect under conditions that met such assumptions; either by excluding first week germination from data analysis, or by separately evaluating *nacl* effect within C<sub>4</sub> and C<sub>24</sub>. The former analysis showed that, after the first week, hazard became proportional for all effects. In doing so, only *nacl* (but not *chilling* nor interaction) remained significant under Cox PH model ( $p < 0.030$ , Table 2). 'bhm', LRT, AIC and BIC analyses confirmed this result, indicating that most of the differences between chilling treatments were in the first week (Tables 2 and 3), as also shown by time trends for survival and hazard rate (Figure 2).

$C_{24}$  showed germination also at 150 mM NaCl (6.2%), with cumulative germination curve,  $F(t)$ , statistically not significantly different from moderate-high salinity ( $N_{75}$ ,  $p > 0.44$ ;  $N_{100}$ ,  $p > 0.08$ ), but highly different from low salinity levels ( $N_{50}$ ,  $p < 0.003$ ; control,  $N_1$ ,  $p < 0.000$ ). At low salinity ( $N_1$  and

$N_{50}$ ) no difference in germination curves was observed between these two levels ( $p > 0.29$ ), both showing higher germination respect to moderate and high salinities ( $p < 0.023$ , Table 4, Figure 3).

**Table 2.** Summary table for Cox PH regression model and cut off values for *nacl* biomarker threshold ('*bhm*') in germination (G) and recovery (R) trials. Cox proportional hazard assumption (PH  $p$ ) rejected if  $p < 0.05$  (in **bold**). Effects ( $Z_i$ ) tested by Wald and '*bhm*' tests ( $p < 0.05$ , in **bold**). Cut off values for *nacl* biomarker threshold (in mM NaCl) by '*bhm*' analysis and their 95% CI (in brackets).

Trial	$Z_i$	PH	' <i>bhm</i> '		cut off values (95% CI)
			$p$	Wald <sup>1</sup>	
<i>Complete dataset</i>					
G	<i>nacl</i>	0.930	<b>0.016</b>	<b>0.011</b>	1 – 50 (1-1) – (50-50)
	<i>chilling</i>	<b>0.004</b>	<b>0.011</b>	<b>0.008</b>	
	<i>interaction</i>	0.526	0.155	0.218	
<i>1<sup>st</sup> week excluded</i>					
G	<i>nacl</i>	0.559	<b>0.030</b>	<b>0.022</b>	1 – 50 (1-1) – (50-50)
	<i>chilling</i>	0.383	0.830	0.585	
	<i>interaction</i>	0.904	0.639	0.644	
<i>C<sub>24</sub></i>					
G	<i>nacl</i>	<b>0.014</b>	<b>0.000</b>	<b>0.000</b>	1 – 75 (1-1) – (75-75)
<i>C<sub>4</sub></i>					
G	<i>nacl</i>	0.683	<b>0.009</b>	<b>0.005</b>	1 – 50 (1-1) – (50-50)
R	<i>nacl</i>	0.104	0.057	<b>0.021</b>	10 – 50 (1-10) – (50-150)
G + R	<i>nacl</i>	<b>0.010</b>	<b>0.001</b>	<b>0.000</b>	1 – 50 (1-1) – (10-150)

<sup>1</sup>  $p < 0.01$  for Cox model (by LRT, Wald and logrank tests) in all trials except recovery (R: LRT,  $p < 0.050^*$ ; Wald,  $p < 0.057^{ns}$ ; logrank,  $p < 0.052^{ns}$ )

Salinity negatively influenced germination independently from chilling treatment, being *nacl* effect highly significant ( $p < 0.010$ ) over and within each chilling level (Table 2 and 3). Comparing the two chilling lengths ( $C_{24}$  vs.  $C_4$ ), longer chilling resulted in an overall higher germination (generalized Wilcoxon test,  $p < 0.004$ ), also confirmed within each salinity level ( $N_1$ ,  $p < 0.006$ ;  $N_{50}$ ,  $p < 0.020$ ;  $N_{150}$ ,  $p < 0.040$ ). The salinity levels (in common over chilling treatments) showed similar germination curves for  $N_1$  and  $N_{50}$  ( $p > 0.58$ ), both higher than  $N_{150}$  ( $p < 0.000$ ). Within  $C_4$ , 150 mM NaCl failed to germinate and significantly ( $p < 0.05$ ) differed from all the other salinity levels; while at low salinity ( $\leq 50$  mM NaCl) no statistically significant difference was observed between *nacl* levels ( $N_1$  vs.  $N_{10}$ ,  $p > 0.52$ ;  $N_1$  vs.  $N_{50}$ ,  $p > 0.17$ ;  $N_{10}$  vs.  $N_{50}$ ,  $p > 0.09$ ; Table 4, Figure 3). Aggregating  $C_{24}$  data in three salinity ranges of 50 mM NaCl – *low*, *moderate*, and *high*; (0, 50], (50, 100] and (100, 150] – germination in *low* salinity was higher ( $p < 0.001$ ) respect each of the other two classes, being the last two ones not statistically different ( $p > 0.15$ ). These results suggested (similarly to what observed in  $C_4$  and in overall analysis) that germination probability decreased above 50 mM NaCl. Applying '*bhm*' model in order to determine cut off values for conditions that could be interpreted as homogenous, the range between


1 mM to 50 mM NaCl was inferred for (i) all data pooled together and (ii) C<sub>4</sub>, while C<sub>24</sub> had a wider salinity range from 1 to 75 mM NaCl (Table 2).

### 3.4. Recovery from salt stress

In recovery, C<sub>4</sub> final germination was highest under 10 mM NaCl ( $\hat{F} = 29\%$ ) followed by control (16%), 50 mM NaCl (10%) and 150 mM NaCl (Table 4, Figure 3). Samaras in high salinity (N<sub>150</sub>) did not germinate until the third week in freshwater, with a peak of hazard rate in the fourth week ( $h = 4.6\%$ ). Salinity was not significant (but close to critical value) in survival analysis, differently from 'bhm' ( $p < 0.021$ , Table 2), AIC and BIC (chi square  $p < 0.037$ , Table 3) analysis, even if AIC and BIC values were only slightly higher than the threshold value of 2 (AIC, 2.3; BIC, 2.7).

Salinity at 10 mM NaCl had a higher germination probability curve respect to any other *nacl* treatment (Wilcoxon test, Figure 3c) without statistically significant difference among these latter ones (i.e. 1, 50 and 150 mM NaCl, Table 4). Cut off values for salinity during recovery (determined by 'bhm' and including confidence intervals) extended over the full salinity range tested – from control up to 150 mM NaCl.

An overall evaluation, which included the 15 weeks of germination and recovery tests (G+R), again showed the highly significant effect of salinity (by Cox PH model, 'bhm', AIC, BIC and LRT; Tables 2 and 3). Even if Cox PH results should be interpreted with prudence (as salinity changed between germination and recovery trial for most treatments, infringing assumptions), highest germination ( $\hat{F}$ ) was achieved at 10 mM NaCl (N<sub>10</sub>, 39%), followed by control (N<sub>1</sub>, 26%), low (N<sub>50</sub>, 15%) and high (N<sub>150</sub>, 9%) salinity (Table 4). Overall, salinity at 10 mM NaCl did not differ ( $p < 0.05$ ) from control, while it was statistically higher from N<sub>50</sub> and N<sub>150</sub> salinity levels. Control  $\hat{F}$  was higher only respect to high salinity (N<sub>150</sub>), while N<sub>50</sub> and N<sub>150</sub> did not differ (Table 4). The cut off values inferred by 'bhm' did not substantially deviate from the results obtained by the recovery test, indicating a marked and positive influence of the period in freshwater, which reduced, at least in part, the negative effects of salinity. Since germination-test beginning, two peaks of germination had been observed in control and 10 mM NaCl treatment, namely at the 4<sup>th</sup> and 11<sup>th</sup> week (i.e. the 3<sup>rd</sup> week of recovery, Figures 4b and 4c).



**Figure 5.** (a) Salinity and C<sub>24</sub> first-week germination probability,  $F_{(week=1)}$ ;  $y = -0.04^{**}x + 11.5^{**}$ ,  $r^2_{adj} = 0.633$ ,  $p > 0.067$ ,  $n = 5$ . Inset: Salinity and relative germination risk for 1<sup>st</sup> week relative to final germination (in percent), % final =  $100 \cdot F_{(week=1)} / \hat{F}$ ; for 50-150 mM NaCl salinity range,  $y = 0.32^*x + 40.3^{**}$ ,  $r^2_{adj} = 0.915$ ,  $p < 0.029$ ,  $n_{obs} = 4$ . (b) Linear regression between salinity and  $\hat{F}$  in the trials (G = germination, R = recovery): solid bold, C<sub>24</sub>(G),  $r^2_{adj} = 0.77$ ,  $p < 0.031^*$ ; dashed, C<sub>4</sub>,  $r^2_{adj} = 0.76$ ,  $p < 0.034^*$ ; solid, C<sub>4</sub>(R),  $r^2_{adj} = 0.37$ ,  $p < 0.105^{ns}$ ; densely dashed, C<sub>4</sub>(G+R),  $r^2_{adj} = 0.47$ ,  $p < 0.197^{ns}$ .

The negative trend between salinity and  $\hat{F}$  had comparable slopes among trials. It should be noted that relationship between salinity and  $\hat{F}$  was significant only for C<sub>24</sub> and C<sub>4</sub> germination ( $r^2_{adj} > 0.77$ ; 0.76, resp., both  $p < 0.05$ ), but not for recovery ( $r^2_{adj} = 0.37$ ,  $p > 0.10$ ), nor germination + recovery ( $r^2_{adj} = 0.47$ ,  $p > 0.20$ ) trials. In all salinity levels, final  $\hat{F}$  for C<sub>4(G+R)</sub> was greater than C<sub>4</sub> and similar to C<sub>24</sub> (as inferred by 95% CI range, Table 4). In control (intercept in Figure 5b) C<sub>24</sub> > C<sub>4</sub> and C<sub>24</sub> ≈ C<sub>4(R)</sub> < C<sub>4(G+R)</sub>, highlighting the effects of chilling and recovery on germination.

**Table 3.** Model selection (best models in **bold**) by log-likelihood value (LLV), Akaike information criterion (AIC), Bayesian information criterion (BIC); df, degree of freedom. Likelihood ratio (LRT)  $p$ -value listed following the order of terms in model formula (significant terms in **bold**,  $p < 0.05$ ).

Model	df	LLV <sup>1</sup>	AIC	BIC	LRT $p$ -value
<i>Complete dataset (Germination)</i>					
<i>nacl + chilling + nacl × chilling</i>	3	<b>-906 a</b>	<b>1818</b> <sup>2</sup>	1833 <sup>2</sup>	<b>0.001, 0.011, 0.092</b>
<i>nacl + chilling</i>	2	<b>-907 a</b>	1819 <sup>3</sup>	<b>1829</b> <sup>3</sup>	<b>0.000, 0.000</b>
<i>nacl</i>	1	-916 b	1833	1838	<b>0.000</b>
<i>chilling</i>	1	-923 d	1848	1853	<b>0.047</b>
<i>nacl × chilling</i>	1	-922 c	1846	1850	<b>0.015</b>
<i>none</i>	0	-925 e	1850	1850	
<i>1<sup>st</sup> week excluded (Germination)</i>					
<i>nacl + chilling + nacl × chilling</i>	3	<b>-473 a</b>	952 <sup>2</sup>	967 <sup>2</sup>	<b>0.003, 0.830, 0.626</b>
<i>nacl + chilling</i>	2	<b>-473 a</b>	951 <sup>3</sup>	960 <sup>3</sup>	<b>0.000, 0.605</b>
<i>nacl</i>	1	<b>-473 a</b>	<b>949</b>	<b>954</b>	<b>0.000</b>
<i>chilling</i>	1	-489 c	981	986	<b>0.034</b>
<i>nacl × chilling</i>	1	-479 b	959	964	<b>0.000</b>
<i>none</i>	0	-492 d	983	983	
<i>C<sub>24</sub> (Germination)</i>					
<i>nacl</i>	1	<b>-609 a</b>	<b>1220</b>	<b>1225</b>	<b>0.000</b>
<i>none</i>	0	-618 b	1236	1236	
<i>C<sub>4</sub> (Germination)</i>					
<i>nacl</i>	1	<b>-218 a</b>	<b>437</b>	<b>442</b>	<b>0.000</b>
<i>none</i>	0	-215 b	451	451	
<i>C<sub>4</sub> (Recovery)</i>					
<i>nacl</i>	1	<b>-317 a</b>	<b>635.1</b>	<b>637.4</b>	<b>0.037</b>
<i>none</i>	0	-319 b	637.4	640.1	
<i>C<sub>4</sub> (Germination + Recovery)</i>					
<i>Nacl</i>	1	<b>-559 a</b>	<b>1120</b>	<b>1125</b>	<b>0.000</b>
<i>none</i>	0	-567 b	1134	1134	

<sup>1</sup> Best model selection by analysis of deviance; different letters for  $p < 0.05$ .

<sup>2</sup> AIC (BIC) values after single term drop from model. *Complete dataset (Germination)*: *nacl × chilling*, 1819 (1829); *chilling*, 1823 (1833); *nacl*, 1823 (1838). *1<sup>st</sup> week excluded (Germination)* dataset: *nacl × chilling*, 951 (961); *chilling*, 950 (960); *nacl*, 959 (960).

<sup>3</sup> AIC (BIC) values after single term drop from model. *Complete dataset (Germination)*: *chilling*, 1833 (1838); *nacl*, 1848 (1852). *1<sup>st</sup> week excluded (Germination)* dataset: *chilling*, 949 (954); *nacl*, 981 (986).

**Table 4.** Cumulative germination probability ( $\hat{F}$ ) and cumulative hazard rate ( $\hat{H}$ ) in percent along with their 95% CI range (in brackets). Differences between C<sub>24</sub> and C<sub>4</sub> for the same salinity levels (*nacl*, in mM) in **bold** (Wilcoxon  $p < 0.05$ ); within each trial different letters for Wilcoxon  $p < 0.05$ . RR (=  $\hat{F}_{Nx}/\hat{F}_{N1}$ ) = relative risk respect to control.

<i>chilling</i>	<i>nacl</i>	$\hat{F}$ (CI range)	$\hat{H}$ (CI range)	RR	
<i>Germination</i>					
C <sub>24</sub>	1	<b>22.9</b> (16.5 - 30.8)	<b>25.0</b> (17.1 - 35.9)	<i>a</i>	1
	50	<b>17.6</b> (12.3 - 24.5)	<b>18.7</b> (12.5 - 27.5)	<i>ab</i>	0.8
	75	8.5 (5.0 - 14.0)	8.7 (5.0 - 14.9)	<i>c</i>	0.4
	100	12.2 (7.8 - 18.7)	12.6 (7.7 - 20.4)	<i>bc</i>	0.5
	150	<b>6.2</b> (3.3 - 11.3)	<b>6.2</b> (3.2 - 19.1)	<i>c</i>	0.3
	1+50+150	<b>15.3</b> (12.2 - 19.0)	<b>16.1</b> (12.5 - 20.6)		
C <sub>4</sub>	1	12.1 (8.3 - 17.4)	12.8 (8.5 - 19.0)	<i>a</i>	1
	10	<b>15.2</b> (8.4 - 25.7)	<b>16.0</b> (8.5 - 29.4)	<i>a</i>	1.3
	50	6.1 (2.4 - 14.6)	6.2 (2.4 - 15.7)	<i>a</i>	0.5
	150	0.0	0.0.	<i>b</i>	0
	1+50+150	8.5 (5.9 - 12.0)	8.8 (6.1 - 12.7)		
	<i>Recovery</i>				
C <sub>4</sub>	1	15.7 (11.0 - 21.9)	16.7 (11.3 - 24.3)	<i>b</i>	1
	10	<b>28.6</b> (18.4 - 41.5)	<b>32.2</b> (19.1 - 52.3)	<i>a</i>	1.8
	50	9.7 (4.5 - 19.6)	10.0 (4.5 - 21.6)	<i>b</i>	0.6
	150	9.1 (4.2 - 18.5)	9.4 (4.2 - 20.3)	<i>b</i>	0.6
	1+50+150	<b>13.0</b> (9.7 - 17.3)	<b>13.4</b> (9.9 - 18.7)		
	<i>Germination + Recovery</i>				
C <sub>4</sub>	1	26.0 (20.4 - 32.6)	29.6 (22.2 - 38.9)	<i>ab</i>	1
	10	<b>39.4</b> (28.5 - 51.5)	<b>48.0</b> (31.7 - 70.4)	<i>a</i>	1.5
	50	15.2 (8.4 - 25.7)	16.2 (8.6 - 29.5)	<i>bc</i>	0.6
	150	9.1 (4.2 - 18.4)	9.4 (4.2 - 20.3)	<i>c</i>	0.3
	1+50+150	<b>20.4</b> (16.4 - 25.1)	<b>22.5</b> (17.6 - 28.6)		

#### 4. Discussion

The comparison of seed germination response to salinity in natural environment and germination tests resulted useful to infer tolerance levels over a wide range of environmental conditions. It was often pointed out that salt tolerance of plants is difficult to quantify, because it may vary with environmental conditions, plant life stage, and genotype [66,67]. Therefore estimates obtained by standard laboratory germination tests may not reflect seedling emergence in the field [68], therefore extrapolation of monosaline response (such as NaCl) to field salinity conditions might be only speculative [69]. On the other hand, although sampling newly germinated seedling in the forest had the advantage to return results based on effective environmental conditions, such trials had several *cons* in identifying cause-effect relationships (e.g. [70]). Time and space variability in abiotic or biotic factors and conditions (apart from salinity) might strongly influence results; here, site vicinity (below 1 km), similar soil and stand conditions could in part minimize this problem, leaving salinity as the main source of variability.

Salinity reduced seed germination in plants [71–74], even in halophytes [75–77], delaying or preventing germination beyond tolerance limits [78]. Although the issue of critical values determination for salinity is complex and controversial – due to the diversity of saline environments and responses induced by salts, such as: osmotic and oxidative stresses, ion-toxicity and/or nutritional disorder [79] – generalized relationships between germination curves and salinity were suggested. Following Läuchli and Grattan classification [80], a delay in germination onset (without differences in rate and thus in final germination percentage) sorted 'moderate' from 'low' salinity levels, while a further delay along with significant final germination percent reduction characterized 'high' salinity levels. This scheme might not be confirmed by C<sub>24</sub> results (which did not show any delay in germination onset; or at least it was limited to 1 week), where only a reduction of final germination with 75 mM NaCl and beyond was observed. On the other hand, a 'moderate' salinity effect up to 50 mM NaCl resulted in C<sub>4</sub> treatment, suggesting the important role of chilling to overcome dormancy and saline stress.

The comparison of the two chilling lengths confirmed that northern Tuscany (Italy) *F. angustifolia* population had an intermediate physiological seed dormancy (PD, following Baskin and Baskin definition [81,82]), and took advantage from a prolonged chilling period. The time required to break germination has a central role not only for physiological dormancy classification, but also to prevent freshly matured seed from germinating under otherwise favourable conditions, such as under warm autumn days in Mediterranean climate [83,84]. The two chilling lengths tested in this study were placed to the either sides of recommended values (4–26 weeks) for *F. angustifolia* [82] and intermediate PD species (1–6 months [81]). In control, 24-week long chilling resulted in nearly double germination probability (23%) respect to 4-week chilling (12%), and this latter treatment needed almost a double period of time (15 instead to 8 weeks) to reach similar values to the former one. Seed dormancy was mainly attributed to seed coat, as *F. angustifolia* isolated embryos could grow and develop into seedling [85]. The observed low germination, even in freshwater conditions, could partly be explained by the choice of not having applied two dormancy-breaking procedures, that is (i) after-ripening freshly harvested mature seeds in a relatively dry state (effective also for this species [86]) and (ii) to remove pericarp before dormancy-release treatments [82,87]. Both choices were justified by the purpose of investigating germination response to chilling and salinity in conditions similar to those experienced by seed after dispersal in natural environment.

*F. angustifolia* populations showed a decrease of dormancy from northern to southern Italian provenances [88], without chilling requirement in Sicily [86]. At the northern geographical range (41–43° Lat. N), lack of cold stratification resulted in low germination percentage (below 25% after 7–8 weeks) for Spanish [89], French [90], and Turkish [91] provenances, suggesting that origin site temperature or other environmental cues might play a role in dormancy levels [92,93]. Assuming a linear increase germination with chilling length for our seed lot in freshwater conditions, germination probability would be reduced to  $\hat{F} = 10\%$  without chilling (intercept) with a germination increase of 0.54% per chilling week (slope), not so much lower than what inferred from Turkish provenances dataset (0.8–1.4% per week) [91].

Chilling promotes germination, as soon as temperature (and light) conditions returned favourable (as evidenced by C<sub>24</sub> high germination rate during the first-week), allowing emergence also in the sub-optimal conditions given by salinity. Most of the chilling effect was detected in the first week. Excluding first-week emergence from data analysis both Bayesian and survival analysis showed comparable germination between the two chilling lengths. A week of high germination rate could result in an effective mechanism to escape the negative effect of salinity for a higher fraction of seeds. In several plant species (salt-intolerant glycophytes included), cold stratification under freshwater conditions induced emergence under saline conditions [94]. Cold stratification might induce several physiological changes involved in germination under salty conditions, such as higher seed imbibition [95,96], changes in enzyme kinetics or activities [97,98], and/or storage protein degradation [99,100].

In our dataset, seed germination was progressively reduced under salinity conditions and marked reduction of seed germination over 50 (75) mM NaCl was highlighted both under laboratory

(Cox PH and hierarchical Bayesian analysis) and natural environment conditions, indicating that this species might complete germination under 'low' to 'moderate' salt stress [101]. In *F. angustifolia*, osmotic and ion-toxicity of NaCl effects were evidenced on embryogenic callus culture growth at 100 mM NaCl [102]. The critical values found in our study were similar to those detected in *F. rhynchophylla* [103] and slightly lower than what reported for *F. chinensis*, which is able to retain 60% of control germination at 100 mM NaCl [104].

Germination resumption under freshwater conditions might have adaptive consequences, as it prevents emergence under harsh environmental conditions, ensuring a higher chance to proper seedlings establishment [105,106]. This species can tolerate up to 150 mM NaCl salt stress (at least for two months), taking advantage of a prolonged chilling or recovery period, even if in both cases hazard rate for seed germination in high salinity remains low and about 25% and 50% of control values. Low salt-stress exposure (10 mM NaCl) – a process known as salt acclimation [107] – improved germination particularly during recovery. In *Arabidopsis thaliana*, exposure to these same NaCl concentration was found to trigger several physiological responses such as reactive oxygen species (ROS) increase, ROS-scavenging-enzyme activation, compatible solutes accumulation, and Na<sup>+</sup>, Cl<sup>-</sup>, and K<sup>+</sup> ion-content changes [108,109]. Germination kinetics during recovery from saline stress was similar among NaCl concentrations, and germination probability decrease with salinity was not significant for none of the applied statistical approaches, i.e. survival analysis, critical value determination by hierarchical Bayesian, and regression between final  $\hat{F}$  values vs. salinity. Moreover, the similar behaviour among salt concentrations during recovery suggested that NaCl had a mainly osmotic effect (at least up to 150 mM and 2-month exposure). This was also confirmed by the observation that 4-week chilled seeds reached (at the end of recovery) similar final values to C<sub>24</sub> for germination probability ( $\hat{F}$ ) and hazard rate ( $\hat{H}$ ) under all tested NaCl conditions. Thus, moderate-high salinity limited emergence, retaining seed viability. Seed germination resumed under favorable freshwater conditions, highlighting an escape mechanism [110], that might be positive to ensure seedling survival soon after emergence [68], and thus widening regeneration opportunities.

## 5. Conclusions

*F. angustifolia* moderate tolerance to salt spray and soil salinity at seedling and plant stage [9–11] could be also be confirmed in germination/emergence stage. Even if a progressively reduction was observed in seed germination with salinity for *F. angustifolia*, emergence took place up to 50 (70) mM NaCl without a significant differences respect to control. The potential toxicity of high salt levels, hyperosmotic stress, and related secondary stresses, such as oxidative damage [111,112] could be recovered up to the highest NaCl tested levels (150 mM NaCl) after about two months in freshwater conditions. Hence, from an ecological point of view, this study confirmed the importance for overcoming dormancy and complete germination in sub-optimal saline conditions by a prompt germination after (i) a prolonged cold and humid season or (i) rainy periods such as to reduce soil salinity below threshold critical values.

**Author Contributions:** S.R. coordinated the overall project. A.P. followed germination tests. S.R., B.M., and S.M. analyzed data and wrote the manuscript.

**Funding:** This work was funded by Italian PRIN-MIUR [grant number 2012E3F3LK].

**Acknowledgments:** The authors are indebted to Paolo Cherubini (WSL, Swiss) and Andrea Bertacchi (Università di Pisa, Italy) for field trips in study areas and to "Ente Parco Regionale Migliarino, San Rossore, Massaciuccoli" for providing access. We also thank Leonora Di Gesualdo for field help.

**Conflicts of Interest:** The authors declare no conflict of interest.

## References

1. Caudullo, G.; Houston Durrant, T. *Fraxinus angustifolia* in Europe: distribution, habitat,

usage and threats. In *European Atlas of Forest Tree Species*; San-Miguel Ayanz, J., de Rigo, D., Caudullo, G., Houston Durrant, T., Mauri, A., Eds.; Publ. Off. EU: Luxembourg, 2016; p. 97 ISBN 978-92-79-36740-3.

- 2. Raddi, S. *Fraxinus angustifolia*. *Enzyklopädie der Holzgewächse Handb. und Atlas der Dendrol.* 2014, 1–18 doi: 10.1002/9783527678518.ehg2010001.
- 3. Pinheiro, J.C.; Marques, C.R.; Pinto, G.; Bouguerra, S.; Mendo, S.; Gomes, N.C.; Gonçalves, F.; Rocha-Santos, T.; Duarte, A.C.; Roembke, J.; et al. The performance of *Fraxinus angustifolia* as a helper for metal phytoremediation programs and its relation to the endophytic bacterial communities. *Geoderma* **2013**, 202–203, 171–182.
- 4. Rosselli, W.; Keller, C.; Boschi, K. Phytoextraction capacity of trees growing on a metal contaminated soil. *Plant Soil* **2003**, 256, 265–272.
- 5. Baycu, G.B.; Tolunay, D.; Özden, H.; Günebakan, S. Ecophysiological and seasonal variations in Cd, Pb, Zn, and Ni concentrations in the leaves of urban deciduous trees in Istanbul. *Environ. Pollut.* **143** *2006*, 143, 545–554.
- 6. Semenzato, P.; Cattaneo, D.; Dainese, M. Growth prediction for five tree species in an Italian urban forest. *Urban For. Urban Green.* **2011**, 10, 169–176.
- 7. Roloff, A.; Korn, S.; Gillner, S. The Climate-Species-Matrix to select tree species for urban habitats considering climate change. *Urban For. Urban Green.* **2009**, 8, 295–308.
- 8. Jaeger, C.; Gessler, A.; Biller, S.; Rennenberg, H. Differences in C metabolism of ash species and provenances as a consequence of root oxygen deprivation by waterlogging. *J. Exp. Bot.* **2009**, 60, 4335–4345.
- 9. Miyamoto, S. *Site Suitability Assessment for Irrigating Urban Landscapes with Water of Elevated Salinity in the Southwest Consolidated Final Report Part I. Water Quality and Plant Salt Tolerance*; TWRI TR-416. Texas A&M University, AgriLife Research Center at El Paso Texas Water Resources Institute, 2011; pp. 1-91.
- 10. Wu, L.; Guo, X.; Hunter, K.; Zagory, E.; Waters, R.; Brown, J. Studies of salt tolerance of landscape plant species and California native grasses for recycled water irrigation. *Slosson Rep. 2000-2001* **2001**, 1–14.
- 11. Xie, Y.J.; Xu, S.; Han, B.; Wu, M.Z.; Yuan, X.X.; Han, Y.; Gu, Q.; Xu, D.K.; Yang, Q.; Shen, W.B. Evidence of *Arabidopsis* salt acclimation induced by up-regulation of HY1 and the regulatory role of RbohD-derived reactive oxygen species synthesis. *Plant J.* **2011**, 66, 280–292.
- 12. European Commission Interpretation Manual of European Union Habitats - EUR28. DG Environment B.3, Nature ENV. 2013, pp. 1-144.
- 13. Schwartz, S.S.; Jenkins, D.G. Temporary aquatic habitats: Constraints and opportunities. *Acquat. Ecol.* **2000**, 34, 3–8.
- 14. Viciani, D.; Lastrucci, L.; Dell'Olmo, L.; Ferretti, G.; Foggi, B. Natura 2000 habitats in Tuscany (central Italy): Synthesis of main conservation features based on a comprehensive database. *Biodivers. Conserv.* **2014**, 23, 1551–1576.
- 15. Gardner, R.C.; Barchiesi, S.; Beltrame, C.; Finlayson, C.; Galewski, T.; Harrison, I.; Paganini, M.; Perennou, C.; Pritchard, D.; Rosenqvist, A.; et al. *State of the World's Wetlands and Their Services to People: A Compilation of Recent Analyses*; Gland, Switzerland Ramsar Convention Secretariat, 2015; pp. 1-19.
- 16. MedWet Secretariat *Wetlands for Sustainable Development in the Mediterranean Region. A*

*Framework for Action 2016-2030*; MedWet Secretariat: Palais de la Porte Dorée, Paris, France, 7-11 February 2016, 2016; ISBN 978-2-9554-6210-2.

- 17. Ramsar Convention on Wetlands *Global Wetland Outlook: State of the World's Wetlands and their Services to People*; Dudley, N., Ed.; Ramsar Convention Secretariat: Gland, Switzerland, 2018; pp. 1-86.
- 18. Davidson, N.C. How much wetland has the world lost? Long-term and recent trends in global wetland area. *Mar. Freshw. Res.* **2014**, *65*, 934–941.
- 19. Juffe-Bignoli, D.; Burgess, N.D.; Bingham, H.; Belle, E.M.; De Lima, M.G.; Deguignet, M.; Bertzky, B.; Milam, A.N.; Martinez-Lopez, J.; Lewis, E.; et al. *Protected Planet Report 2014*; UNEP-WCMC: Cambridge, UK, 2014; ISBN 978-92-807-3416-4.
- 20. Rundel, P.W. Landscape disturbance in Mediterranean-type ecosystems: An overview. In *Landscape Disturbance and Biodiversity in Mediterranean-Type Ecosystems*; Rundel, P.W., Montenegro, G., Jaksic, F.M., Eds.; Springer Berlin Heidelberg: Berlin, Heidelberg, 1998; pp. 3–22 ISBN 978-3-662-03543-6.
- 21. Herbert, E.R.; Boon, P.; Burgin, A.J.; Neubauer, S.C.; Franklin, R.B.; Ardón, M.; Hopfensperger, K.N.; Lamers, L.P.M.; Gell, P. A global perspective on wetland salinization: Ecological consequences of a growing threat to freshwater wetlands. *Ecosphere* **2015**, *6*, art206.
- 22. Geijzendorffer, I.R.; Beltrame, C.; Chazee, L.; Gaget, E.; Galewski, T.; Guelmami, A.; Perennou, C.; Popoff, N.; Guerra, C.A.; Leberger, R.; et al. A more effective Ramsar Convention for the conservation of Mediterranean wetlands. *Front. Ecol. Evol.* **2019**, *7*, 1–6.
- 23. EEA Natura 2000 Network Viewer Available online: <http://natura2000.eea.europa.eu/#> (accessed on Aug 10, 2019).
- 24. Levin, N.; Elron, E.; Gasith, A. Decline of wetland ecosystems in the coastal plain of Israel during the 20th century: Implications for wetland conservation and management. *Landsc. Urban Plan.* **2009**, *92*, 220–232.
- 25. Cosentino, B.; Colleges, W.S. Dispersal and wetland fragmentation. In *The Wetland Book: I: Structure and Function, Management, and Methods*; Finlayson, C.M., Everard, M., Irvine, K., McInnes, R.J., Middleton, B.A., van Dam, A.A., Davidson, N.C., Eds.; Springer Netherlands: Dordrecht, 2018; pp. 105–111 ISBN 9789048196593.
- 26. Ghelardini, L.; Migliorini, D.; Santini, A.; Pepori, A.L.; Maresi, G.; Vai, N.; Montuschi, C.; Carrari, E.; Feducci, M.; Capretti, P.; et al. From the Alps to the Apennines: possible spread of ash dieback in Mediterranean areas. In *Dieback of European Ash (*Fraxinus spp.*): Consequences and Guidelines for Sustainable Management. The Report on European Cooperation in Science & Technology (COST) Action FP1103 FRAXBACK*; Vasaitis, R., Enderle, R., Eds.; SLU Service/Repro: Uppsala, Sweden, 2017; pp. 140–149 ISBN 978-91-576-8696-1.
- 27. Ortega, M.; Velasco, J.; Millán, A.; Guerrero, C. An ecological integrity index for littoral wetlands in agricultural catchments of semiarid mediterranean regions. *Environ. Manage.* **2004**, *33*, 412–430.
- 28. Schuerch, M.; Spencer, T.; Temmerman, S.; Kirwan, M.L.; Wolff, C.; Lincke, D.; McOwen, C.J.; Pickering, M.D.; Reef, R.; Vafeidis, A.T.; et al. Future response of global coastal wetlands to sea-level rise. *Nature* **2018**, *561*, 231–234.
- 29. Tsanis, I.K.; Daliakopoulos, I.N.; Koutoulis, A.G.; Karatzas, G.P.; Varouchakis, E.; Kourgialas, N. Soil salinization. In *Soil threats in Europe. JRC Technical Report*. EUR 27607; Stolte, J., Tesfai,

M., Øygarden, L., Kværnø, S., Keizer, J., Verheijen, F., Panagos, P., Ballabio, C., Hessel, R., Eds.; 2015; pp. 104–117.

30. Hoepffner, N.; Dowell, M.D.; Edwards, M.; Fonda-Umano, S.; Green, D.R.; Greenaway, B.; Hansen, B.; Heinze, C.; Leppänen, J.-M.; Lipiatou, E.; et al. *Marine and Coastal Dimension of Climate Change in Europe Marine and Coastal Dimension of Climate Change in Europe*; Hoepffner, N., Ed.; Institute for Environment and Sustainability, EUR 22554 EN, 2006; ISBN 9279037471.

31. IPCC *Summary for policymakers*. In: *Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part A: Global and Sectoral Aspects. Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*; Field, C.B., Barros, V.R., Dokken, D.J., Mach, K.J., Mastrandrea, M.D., Bilir, T.E., Chatterjee, M., Ebi, K.L., Estrada, Y.O., Genova, R.C., Girma, B., Kissel, E.S., Levy, A.N., MacCracken, S., Mastrandrea, P.R., White, L.L., Eds.; Cambridge University Press: Cambridge, United Kingdom and New York, NY, USA, 2014; pp. 1–32.

32. Amores, M.J.; Verones, F.; Raptis, C.; Jurasko, R.; Pfister, S.; Stoessel, F.; Antón, A.; Castells, F.; Hellweg, S. Biodiversity impacts from salinity increase in a coastal wetland. *Environ. Sci. Technol.* **2013**, *47*, 6384–6392.

33. Zedler, J.B. What's new in adaptive management and restoration of coasts and estuaries? *Estuaries and Coasts* **2017**, *40*, 1–21.

34. Barducci, A.; Guzzi, D.; Marcoionni, P.; Pippi, I.; Raddi, S. PROBA contribution to wetland monitoring in the coastal zone of San Rossore Natural Park. In Proceedings of the European Space Agency, (Special Publication) ESA SP; 2007.

35. Bertacchi, A.; Lombardi, T.; Tomei, P.E. Le aree umide salmastre della Tenuta di San Rossore (PI): zonazione e successione delle specie vegetali in relazione alla salinità del suolo. *INTER NOS* **2007**, *1*, 1–14.

36. Sani, A.; Tomei, P.E. La vegetazione psammofila del litorale di San Rossore (Toscana settentrionale) e la sua importanza conservazionistica. *Parlatore* **2006**, *VIII*, 99–119.

37. Bertacchi, A. Dune habitats of the Migliarino - San Rossore - Massaciuccoli Regional Park (Tuscany-Italy). *J. Maps* **2017**, *13*, 322–331.

38. RStudio Team. RStudio: Integrated Development for R. RStudio, Inc., Boston, MA URL <http://www.rstudio.com/>. 2015.

39. Shapiro, S.S.; Wilk, M.B. An analysis of variance test for normality (complete sample). *Biometrika* **1965**, *52*, 591–611.

40. Royston, P. The W-test for Normality. *J. R. Stat. Soc. Ser. C (Applied Stat.)* **1995**, *44*, 547–551.

41. Wilcoxon, F. Individual comparisons by ranking methods. *Biom. Bull.* **1945**, *1*, 80–83.

42. Canty, A.; Ripley, B. Bootstrap Functions. Package 'boot' R - Project. 2019, 1–117.

43. Canty, A.J. Resampling Methods in R: The boot Package. *Newsl. R Proj.* **2002**, *2/3*, 2–7.

44. ISTA. *ISTA working sheets on tetrazolium testing Volume II. 1 st edition 2003 including supplements* 2011; International Seed Testing Association, Eds.; Bassersdorf, Switzerland, 2011; ISBN 3906549402 9783906549408.

45. ISTA. *International Rules for Seed Testing*; International Seed Testing Association.: Bassersdorf, Switzerland, 2017; ISBN 2310-3655.

46. Magini, E. Forest seed handling, equipment and procedures. *Unasylva* **1962**, *64*, 1–22.

47. Bewley, J.D.; Bradford, K.J.; Hilhorst, H.W.M.; Nonogaki, H. *Seeds: Physiology of development*,

germination and dormancy; 3rd ed.; Springer: New York, Heidelberg, Dordrecht, London, 2013; ISBN 9781461446934.

- 48. Klein, J.P.; van Houwelingen, H.C.; Ibrahim, J.G.; Scheike, T.H. *Handbook of Survival Analysis*; CRC Press, Taylor and Francis Group, 2016; ISBN 9781466555679.
- 49. Muenchow, G. Ecological use of Failure Time Analysis. *Ecology* **1986**, *67*, 246–250.
- 50. Kalbfleisch, J.D.; Prentice, R.L. *The Statistical Analysis Failure Time Data*; 2nd ed.; John Wiley & Sons, 2011; ISBN 9781118031230.
- 51. McNair, J.N.; Sunkara, A.; Frobish, D. How to analyse seed germination data using statistical time-to-event analysis: non-parametric and semi-parametric methods. *Seed Sci. Res.* **2012**, *22*, 77–95.
- 52. Kaplan, E.L.; Meier, P. Nonparametric estimation from incomplete samples. *J. ASA* **1958**, *53*, 457–481.
- 53. Nelson, W. Hazard plotting for incomplete failure data. *J. Qual. Technol.* **1969**, *1*, 27–52.
- 54. Nelson, W. Theory and applications of hazard plotting for censored failure data. *Technometrics* **1972**, *14*, 945–966.
- 55. Aalen, O. Nonparametric inference for a family of counting processes. *Ann. Stat.* **1978**, *6*, 701–726.
- 56. Zhou, M. *Empirical Likelihood Method in Survival Analysis*; Zhou, M., Ed.; Chapman and Hall/CRC: New York, 2015; ISBN 9781466554931.
- 57. Thomas, D.R.; Grunkemeier, G.L. Confidence interval estimation of survival probabilities for censored data. *J. Am. Stat. Assoc.* **1975**, *70*, 865–871.
- 58. Strobl, R.; Verbeke, T. Confidence intervals for the Kaplan-Meier estimator. Package 'km.ci' for RStudio 2015, 1–8.
- 59. Cox, D.R. Regression models and life-tables. *J. R. Stat. Soc. Ser. B* **1972**, *34*, 187–220.
- 60. Therneau, T.M. Survival Analysis. RStudio Package 'survival' 2019, 1–161.
- 61. Grambsch, P.M.; Therneau, T.M. Proportional hazards tests and diagnostics based on weighted residuals. *Biometrika* **1994**, *81*, 515–526.
- 62. Hess, K.R. Graphical methods for assessing violations of the proportional hazards assumption in Cox regression. *Stat. Med.* **1995**, *14*, 1707–1723.
- 63. Ripley, B.; Venables, B.; Bates, D.M.; Hornik, K.; Gebhardt, A.; Firth, D. Package 'MASS.' **2019**, 1–169.
- 64. Chen, B.E.; Jiang, W.; Tu, D. A hierarchical Bayes model for biomarker subset effects in clinical trials. *Comput. Stat. Data Anal.* **2014**, *71*, 324–334.
- 65. Fang, T.; Mackillop, W.; Jiang, W.; Hildesheim, A.; Wacholder, S.; Chen, B.E. A Bayesian method for risk window estimation with application to HPV vaccine trial. *Comput. Stat. Data Anal.* **2017**, *112*, 53–62.
- 66. Kozlowski, T.T. Responses of woody plants to flooding and salinity. *Tree Physiol. Monogr.* **1997**, 1–29.
- 67. Kozlowski, T.T. Responses of woody plants to human-induced environmental stresses: Issues, problems, and strategies for alleviating stress. *CRC. Crit. Rev. Plant Sci.* **2000**, *19*, 91–170.
- 68. Esechie, H.A.; Al-Saidi, A.; Al-Khanjari, S. Effect of sodium chloride salinity on seedling emergence in chickpea. *J. Agron. Crop Sci.* **2002**, *188*, 155–160.
- 69. Sosa, L.; Llanes, A.; Reinoso, H.; Reginato, M.; Luna, V. Osmotic and specific ion effects on the

germination of *Prosopis strobulifera*. *Ann. Bot.* **2005**, *96*, 261–267.

70. Gotelli, N.J.; Ellison, A.M. *A Primer of Ecological Statistics*; Sinauer, 2013; ISBN 1605350648, 9781605350646.

71. Peel, M.D.; Waldron, B.L.; Jensen, K.B.; Chatterton, N.J.; Horton, H.; Dudley, L.M. Screening for salinity tolerance in alfalfa: A repeatable method. *Crop Sci.* **2004**, *44*, 2049–2053.

72. Rajjou, L.; Duval, M.; Gallardo, K.; Catusse, J.; Bally, J.; Job, C.; Job, D. Seed germination and vigor. *Annu. Rev. Plant Biol.* **2012**, *63*, 507–33.

73. Zedler, J.B. Wetlands at your service: reducing impacts of agriculture at the watershed scale. *Front. Ecol. Environ.* **2003**, *1*, 65–72.

74. Shumway, S.W.; Bertness, M. Salt stress limitation of seedling recruitment in a salt marsh plant community. *Oecologia* **1992**, *92*, 490–497.

75. Ungar, I.A. Halophyte seed germination. *Bot. Rev.* **1978**, *44*, 233–264.

76. Grigore, M.-N. *Romanian Salt Tolerant Plants: Taxonomy and Ecology*; TEHNOPRESS: Iasi, 2012; ISBN 978-973-702-923-2.

77. González, M.B. Adaptation of halophytes to different habitats. In *Seed Dormancy and Germination*; IntechOpen, 2019; pp. 1–23.

78. Grigore, M.-N.; Toma, C. *Anatomical Adaptations of Halophytes*; Springer International Publishing, 2017; ISBN 978-3-319-66480-4.

79. Liang, W.; Ma, X.; Wan, P.; Liu, L. Plant salt-tolerance mechanism : A review. *Biochem. Biophys. Res. Commun.* **2018**, *495*, 286–291.

80. Läuchli, A.; Grattan, S.R. Plant growth and development under salinity stress. In *Advances in Molecular Breeding Toward Drought and Salt Tolerant Crops*; Jenks, M.A., Hasegawa, P.M., Jain, S.M., Eds.; Springer, 2007; pp. 1–32.

81. Baskin, C.C.; Baskin, J.M. *Seeds - Ecology, Biogeography, and, Evolution of Dormancy and Germination*; 2nd ed.; Academic Press: San Diego, CA, 2014; ISBN 978-0-12-416677-6.

82. Piotto, B.; Di Noi, A. *Seed propagation of mediterranean trees and shrubs*; APAT, Ed.; IGER srl: Roma, Italy, 2003; ISBN 88-448-0081-0.

83. Carta, A.; Bedini, G.; Mu, J.V. Comparative seed dormancy and germination of eight annual species of ephemeral wetland vegetation in a Mediterranean climate. *Plant Biol.* **2013**, *214*, 339–349.

84. Carta, A.; Hanson, S.; Müller, J. V. Plant regeneration from seeds responds to phylogenetic relatedness and local adaptation in Mediterranean Romulea (Iridaceae) species. *Ecol. Evol.* **2016**, *6*, 4166–4178.

85. Raquin, C.; Jung-Muller, B.; Dufour, J.; Frascaria-Lacoste, N. Rapid seedling obtaining from European ash species *Fraxinus excelsior* (L.) and *Fraxinus angustifolia* (Vahl.). *Ann. For. Sci.* **2002**, *59*, 219–224.

86. Lombardo, G.; Scialabba, A.; Schicchi, R. Seed cryopreservation of *Fraxinus angustifolia* Vahl. *African J. Biotechnol.* **2013**, *12*, 1930–1936.

87. French, D.; Meilan, R. Germination trials for Asian and North American ash species. *Tree Plant. Notes* **2013**, *56*, 27–34.

88. Piotto, B.; Piccini, C. Influence of pretreatment and temperature on the germination of *Fraxinus angustifolia* seeds. *Seed Sci. Technol.* **1998**, *26*, 799–812.

89. Cabra-Rivas, I.; Castro-Díez, P. Potential germination success of exotic and native trees

coexisting in central Spain riparian forests. *Int. J. Ecol.* **2016**, *ID* 7614683, 1–10.

90. Gérard, P.R.; Fernández-Manjarrés, J.F.; Bertolino, P.; Dufour, J.; Raquin, C.; Frascaria-Lacoste, N. New insights in the recognition of the European ash species *Fraxinus excelsior* L. and *Fraxinus angustifolia* Vahl as useful tools for forest management. *Ann. For. Sci.* **2006**, *63*, 733–738.

91. Tilki, F.; Çiçek, E. Effects of stratification, temperature and storage on germination in three provenances of *Fraxinus angustifolia* subsp. *oxycarpa* seeds. *Turkish J. Agric. For.* **2005**, *29*, 323–330.

92. He, H.; De Souza Vidigal, D.; Basten Snoek, L.; Schnabel, S.; Nijveen, H.; Hilhorst, H.; Bentsink, L. Interaction between parental environment and genotype affects plant and seed performance in *Arabidopsis*. *J. Exp. Bot.* **2014**, *65*, 6603–6615.

93. Née, G.; Xiang, Y.; Soppe, W.J.J. The release of dormancy, a wake-up call for seeds to germinate. *Curr. Opin. Plant Biol.* **2017**, *8*–14.

94. Zhang, L.; Tian, C.; Wang, L. Cold stratification pretreatment improves salinity tolerance in two wheat varieties during germination. *Seed Sci. Technol.* **2018**.

95. Meng, H.; Wang, X.; Tong, S.; Lu, X.; Hao, M. Seed germination environments of *Typha latifolia* and *Phragmites australis* in wetland restoration. **2016**, *96*, 194–199.

96. Bungard, R.A.; McNeil, D.; Morton, J.D. Effects of chilling, light and nitrogen-containing compounds on germination, rate of germination and seed imbibition of *Clematis vitalba* L. *Ann. Bot.* **1997**, *79*, 643–650.

97. Chen, S.; Chou, S.; Tsai, C.; Hsu, W.; Baskin, C.C.; Baskin, J.M.; Chien, C.; Kuo-Huang, L. Plant physiology and biochemistry effects of moist cold stratification on germination, plant growth regulators, metabolites and embryo ultrastructure in seeds of *Acer morrisonense* (Sapindaceae). *Plant Physiol. Biochem.* **2015**, *94*, 165–173.

98. Miransari, M.; Smith, D.L. Plant hormones and seed germination. *Environ. Exp. Bot.* **2014**, *99*, 110–121.

99. Gai, Y.P.; Li, X.Z.; Ji, X.L.; Wu, C.A.; Yang, G.D.; Zheng, C. Chilling stress accelerates degradation of seed storage protein and photosynthetic protein during cotton seed germination. *J. Agron. Crop Sci.* **2008**, *194*, 278–288.

100. Einali, A.L.I.R.; Sadeghipour, H.R. Alleviation of dormancy in walnut kernels by moist chilling is independent from storage protein mobilization. *Tree Physiol.* **2007**, *27*, 519–525.

101. Polle, A.; Rennenberg, H. Physiological responses to abiotic and biotic stress in forest trees. *Forests* **2019**, *10*, 711, 1–5.

102. Tonon, G.; Kevers, C.; Faivre-Rampant, O.; Graziani, M.; Gaspar, T. Effect of NaCl and mannitol iso-osmotic stresses on proline and free polyamine levels in embryogenic *Fraxinus angustifolia* callus. *J. Plant Physiol.* **2004**, *161*, 701–708.

103. Hong, L.; Ping, C.; ZhaoHui, Z.; ShouWen, S.; PiJun, L.; BaoQing, W. Seeds germination of three common afforestating trees under salt and drought stress in Xinjiang. *Acta Bot. Boreali-Occidentalia Sin.* **2011**, *31*, 1466–1473.

104. Zhang, Z.; Yu, F. Effects of salt stress on seed germination of four ornamental non- halophyte species. *Intern. J. Agr. & Biol.* **2019**, *21*, 47–53.

105. Llanes, A.; Andrade, A.; Masciarelli, O.; Alemano, S.; Luna, V. Drought and salinity alter endogenous hormonal profiles at the seed germination phase. *Seed Sci. Res.* **2016**, *26*, 1–13.

106. Macovei, A.; Pagano, A.; Leonetti, P.; Carbonera, D.; Balestrazzi, A.; Araújo, S.S. Systems biology and genome-wide approaches to unveil the molecular players involved in the pre-germinative metabolism: implications on seed technology traits. *Plant Cell Rep.* **2017**, *36*, 669–688.
107. Shen, X.; Wang, Z.; Song, X.; Xu, J.; Jiang, C.; Zhao, Y.; Ma, C.; Zhang, H. Transcriptomic profiling revealed an important role of cell wall remodeling and ethylene signaling pathway during salt acclimation in *Arabidopsis*. *Plant Mol. Biol.* **2014**, *86*, 303–317.
108. Sivritepe, N.; Sivritepe, H.O.; Eris, A. The effects of NaCl priming on salt tolerance in melon seedlings grown under saline conditions. *Sci. Hortic. (Amsterdam)*. **2003**, *97*, 229–237.
109. Bajji, M.; Kinet, J.; Lutts, S. Osmotic and ionic effects of NaCl on germination, early seedling growth, and ion content of *Atriplex halimus* (Chenopodiaceae). *Can. J. Bot.* **2002**, *80*, 297–304.
110. Ievinsh, G. Biological basis of biological diversity: Physiological adaptations of plants to heterogeneous habitats along a sea coast. *Acta Univ. Latv.* **2006**, *710*.
111. Zhu, J.K. Salt and drought stress signal transduction in plants. *Annu. Rev. Plant Biol.* **2002**, *53*, 247–273.
112. Zhu, J.K. Abiotic stress signaling and responses in plants. *Cell* **2016**, *167*, 313–324.