

Article

Cognitive Fusion Mediates the Relationship between Dispositional Mindfulness and Anxiety and Other Negative Emotions: A Study in a Sample of Spanish Primary-School Students

María García-Gómez¹, Joan Guerra², Víctor M. López-Ramos² and José M. Mestre^{3*}

¹ Departamento de Psicobiología, Universidad de Murcia, 30001 Murcia, Spain; mgg99158@um.es

² Departamento de Psicología, Universidad de Extremadura, Cáceres, 10071, Spain; joangb@unex.es; vmlopez@unex.es

³ Instituto de Investigación y Desarrollo Social Sostenible (INDESS), Universidad de Cádiz, 11405 Jerez de la Frontera, Cádiz, Spain; josemi.mestre@uca.es; <https://orcid.org/0000-0002-6822-4970>

* Correspondence author: josemi.mestre@uca.es

Abstract: Nowadays, mindfulness-based interventions (MBI) have experimented a remarkable development of studies among childhood and adolescent interventions. For this reason, dispositional mindfulness (DM) measures for children and adolescents have been developed to determine the effectiveness of MBI at this age stage. However, little is known about how key elements of DM (f. e., cognitive defusion or experiential avoidance that both conform psychological inflexibility) are involved in the mechanisms of the children and adolescents' mental health outcomes. This research examined the mediating effect of cognitive fusion between DM and anxiety and other negative emotional states in a sample of 318 Spanish primary-school students (aged between 8 and 16 years, $M=11.24$, $SD=2.19$, 50.8% males). Participants completed the AFQ-Y, which is a measure of psychological inflexibility that encompasses cognitive defusion and experiential avoidance; CAMM (DM for children and adolescents), PANAS-N (positive and negative affect measure for children, Spanish version of PANASC), and STAIC (an anxiety measure for children). The study accomplished ethical standards. As MBI relevant literature has suggested, cognitive defusion was a significant mediator between DM and symptoms of both negative emotions and anxiety in children and adolescents. However, experiential avoidance did not show any significant mediating relationship. Probably, it is needed an improvement of the assessment of experiential avoidance. MBI programs for children and adolescents may include more activities for reducing effects of the cognitive defusion on their emotional distress.

Keywords: dispositional mindfulness; cognitive defusion; anxiety; mindfulness based on interventions; mental health; experiential avoidance; children and adolescents

1. Introduction

Activities for children and adolescents based on mindfulness interventions are being used more and more by researchers, instructors, and therapists. Recent reviews have reported that mindfulness-based interventions (MBI) are feasible for young participants [1]. MBI for children and adolescents are promising initiatives as there is evidence of their usefulness in treating childhood anxiety [2], reducing symptoms of obsessive-compulsive disorder [3] and their pathological concerns [4].

On the one hand, the most sought-after goals of MBIs and Acceptance and Commitment Therapy (ACT) interventions for children and adolescents is the promotion of psychological flexibility [5]. Psychological flexibility (PF) "is about being aware of thoughts and feelings that unfold in the present moment without needless defence, and depending on what the situation affords, persisting or changing behaviour to pursue central interests and goals" (p.868, [6]). For this reason,

and from ACT assumptions, one way to increase psychological flexibility of individuals could be to promote producing better mental health outcomes [7, 8]. Therefore, psychological inflexibility (PI) would be the opposite pole of the same dimension and, consequently, would have the opposite outcomes to those reported above [9, 10].

On the other hand, dispositional mindfulness (DM) has been defined as a trait, in which individuals differ in how they accept their current contingencies and live with a commitment [11]. Therefore, DM is a predisposition or trait for living in a mindfulness way [12]. Considering any psychological intervention as a mechanism of change (f. i., MBIs), DM as a trait might be included as a mechanism as well [12].

However, little is known about how the combined nature of the relationship between PI and DM regarding their potential predictions of mental-wellbeing outcomes in children and adolescents. For instance, some studies stated that the mechanisms through which being mindful can influence positive outcomes have only recently been explored, hence, it is necessary to explore these mechanisms [13]. Our review of the literature found studies on how DM and/or PI are related to mental health outcomes. However, we have not found any important studies on whether PI can play a mediating role between DM and emotional-wellbeing criteria. We found no studies on what aspect of PI (cognitive fusion and experiential avoidance) may have a mediating effect on some indicators of mental health criteria. Therefore, there is a shortage of studies on PI and DM in children and adolescents [14, 15]. Given the relevance of PI, the Avoidance and Fusion Questionnaire for Youth (AFQ-Y) was developed to assess this construct [16]. The instrument measures the two main interrelated processes which produce psychological inflexibility: cognitive fusion (CF) and experiential avoidance (EA) [16]. Cognitive fusion means responding as if private events were true, and so increase the tendency to avoid the internal experience, the opposite of a mindful attitude, such as present moment by moment awareness [17, 18].

There are some promising antecedents with samples of children and adolescents using this PI measure. A longitudinal study with adolescents, experiential acceptance (the opposite of avoidance) predicted increasing positive affect and decreasing fear and sadness [14]. In a Dutch sample, PI (using a shorter version of the AFQ-Y and only one factor) and anxiety symptoms were positively correlated [9]. Using a sample of Spanish, the total score of AFQ-Y was related negatively to depression symptoms and positively to satisfaction with life [16]. In this last study, however, CF predicted depression and satisfaction with life, while experiential avoidance did not.

Although the AFQ-Y has been validated in various languages -Spanish [16], Italian [19], Swedish [20], or Dutch [9], we found some controversies concerning the factor structure of the measure. Some authors described a single loaded factor (just psychological flexibility), while others found two factors (cognitive fusion and experiential avoidance). It is important to try to clarify this debate because it has consequences for the interpretation of the AFQ-Y. We believe that both cognitive fusion and experiential avoidance might have different natures for the same construct: psychological inflexibility (PI). Therefore, potentially, there could be different mediating effects of CF and EA between DM and negative/positive emotional outcomes.

A brief review of both AFQ-Y factors (cognitive fusion and experiential avoidance) helps to understand our point of view described above. Psychological flexibility (using an adaptive-positive term) has been defined as the ability to contact with present-moment awareness, and adjust behaviour to valued aims and interests [21, 22], hence, PI is a lack of this ability. Psychological flexibility is a dynamic process which could be observed when a person: (1) adapts to a changing situational request, (2) redefines mental resources, (3) transforms perspective, and (4) balances needs, competing desires, and life domains [22, 23]. For this reason, a lack of this ability, called psychological inflexibility, has been associated with different diagnostic categories [23-25] and is a predictor of emotional distress, anxiety, attenuation of positive life appraisals and emotions [26].

Besides, two processes have been included as factors of this psychological inflexibility topic: cognitive fusion and experiential avoidance. On the one hand, cognitive fusion (CF) is a process by which the individual becomes entangled with memories, thoughts, judgments, and evaluations and adjust behaviour to the internal experiences [27]. Therefore, CF should have a mediating role between

DM and some psychological outcomes [24]. On the other hand, experiential avoidance (EA) has been defined as a set of behaviours in which a person refuses to remain in contact with the internal experience such as thoughts, feelings, images, and sensations, labelling as negative events [25]. Experiential avoidance can be considered adaptive if it is used as self-control strategies in short term [27]. Hence, EA becomes problematic if used continuously and rigidly. According to this last point of view, EA has been related to psychological symptoms and problem behaviours [23] as well as anxiety, depression, emotional distress [28], and obsessive-compulsive symptoms [29].

At this point, it is necessary to understand how each of the two components of psychological flexibility contributes to the effects of DM on key criteria in the emotional wellbeing of children and adolescents. According to the antecedents previously reported, DM and psychological flexibility have significant relationships with positive and negative emotions, especially, with both state and trait anxiety [24].

Regarding potential mediating effects of all this PI and its elements, it has been verified that psychological inflexibility fully mediated the effect of early maladaptive schemes on psychopathology in a non-clinical undergraduate sample [30]. Cognitive fusion had a mediating association with catastrophizing coping strategy in a chronic-pain community sample of young people [31]. Regarding experiential avoidance, some authors have suggested that it is important to understand models of psychopathology by assessing the role of experiential avoidance as a mediator and/or moderator [32, 33]. Beyond these studies, little has been explored about the mediating effect of psychological flexibility, cognitive fusion, or experiential avoidance between DM and some positive or negative emotional states, especially in children and adolescents.

Our cross-sectional study was designed to understand better how DM and PI are present in the mechanisms of emotional wellbeing among children and adolescents. We hypothesised that cognitive fusion and experiential avoidance (as components of the psychological inflexibility) have a mediating role between DM and emotional health criteria (positive/negative affectivity and both state and trait anxiety).

2. Materials and Methods

2.1. Participants

We randomly selected 327 children and adolescents from nine primary schools in southern Spain. The age of participants ranged from eight to 16 years old ($M=11.24$, $SD=2.19$, 50.8% males). Study information and consent forms were distributed to all parents (according to the Spanish Organic Law of Data Protection). Students were asked to fill out the questionnaires in one session during the class by supervising psychologists (M. G-G, C.S., and J. T., some authors of this article). Nine children were excluded due to errors or omissions in their responses or because their parents did not provide informed consent.

In conformance with the Research and Ethical Commission of INDESS (University Research Institute on Social and Sustainable Development, University of Cadiz, Spain), we adhered to the following ethical recommendations: (a) all participants had to bring an informed consent from their parents, particularly those under 14 years old; (b) we had to inform and receive permission from every single Parents' Association, and (c) the study had to be approved by an external ethical board (Ethical board of University of Murcia, Spain). Consequently, this research is subject to compliance with ethical standards: the 1964-Helsinki declaration and its later amendments or comparable ethical standards and. According to article 13.1 of the Spanish Organic Law of Data Protection, the "data of persons over fourteen years of age may be processed with their consent, except in those cases in which the Law requires the assistance of the holders of parental authority or guardianship. In the case of minors under 14 years of age, the consent of the parents or guardians will be required".

2.2. Measures

2.2.1. Psychological inflexibility: cognitive fusion and experiential avoidance

Psychological inflexibility (PI) was measured using the Spanish adaptation [16] of Avoidance and Fusion Questionnaire for Youth (AFQ-Y, [15]). This tool was developed to measure psychological inflexibility, and its components - cognitive fusion and experiential avoidance - in children and adolescents. It was adapted from the Acceptance and Action Questionnaire [34], which is a self-report measure for psychological inflexibility in adults. The AFQ-Y consists of 17 items that can be rated on a Likert scale ranging from 0 (not at all true) to 4 (very true). Example items include "I am afraid of my feelings" and "the bad things I think about myself must be true". Total scores ranged from 0 to 68. High scores of AFQ-Y indicate higher psychological inflexibility, experiential avoidance and cognitive fusion. AFQ-Y total, cognitive fusion and experiential avoidance were promediated (ranged from 0 to 4) due to every single subscale have a different number of items.

2.2.2. Dispositional Mindfulness

To assess DM of participants, we used the validated Spanish version [35] of the Child and Adolescent Mindfulness Measure (CAMM; [36]). The CAMM consists of 10 items, answered on a 5-point Likert scale, ranging from 0 (never true) to 4 (always true). An example of a CAMM item was 'I keep myself busy so I don't notice my thoughts or feelings'. However, the Ten-Spanish CAMM version had to be erased items 5 and 10 due to cultural biases among Spanish-children samples. Spanish CAMM-8 had better internal consistency and reliabilities than CAMM-10. Previous research using CAMM-8 has been widely reported in a pilot study with gifted children [37]. Recently, another Spanish validation of the CAMM in a larger sample of Chilean and Spanish children adolescents with similar outcomes regarding items 5 and 10, however, authors recommended a shorter version of a 7-item CAMM (removing # 2, 5, and 10 from Greco's original CAMM), [38]. Nonetheless, we used the total score of Spanish CAMM-8 [35] and it was promediated (ranging from 0 to 4) for this study. The DM measure was also recoded into positive total scores, so higher scores reflected better dispositional mindfulness rather than the opposite.

2.2.3. Positive and negative affect

The Positive and Negative Affect Schedule for Children (Spanish validation 'PANASN', [39]). This measure was based on the original version of the instrument called PANAS-C [40]. This tool consists of a 30-item measure for children and young adolescents, which assesses several positive affects (PA; e.g., joyful) and negative affects (NA; e.g., lonely) with 15 items in each subscale. Besides, it returned a score on Balance (PA - NA). Through a 3-point Likert scale, ranging from 1 (slightly or never) to 3 (much), participants valued how they felt during the last few weeks. The minimum score for each scale is 15 and the maximum score is 45. Internal consistency, convergent and discriminant validity for the Spanish PANAS-C (named as PANASN) had been previously obtained [39]. Our research used an average score of PANASN (from 1 to 3). We used the acronym PANASN for this study. Positive affect (PA), negative affect (NA), and balance were used as criteria.

2.2.4. Anxiety

The State-Trait Anxiety Inventory for Children (STAIC; [41]). This self-report scale has been frequently used to measure state and trait anxiety in children and adolescents. The instrument was created to measure the propensity to anxiety both as a transient state and as a relatively stable individual difference. To measure trait and state anxiety, this questionnaire incorporates two separate 20-item self-report rating scales. The participant has to evaluate the degree to which he or she experiences a particular symptom (e.g., I feel 1-not scared, 2-scared, and 3-very scared). By adding up scores for each scale, the state-anxiety and trait-anxiety are obtained. Total scores for situational and baseline questions range, separately, from 20 to 60, with higher scores denoting higher levels of anxiety.

3. Results

Before studying the mediation effects of psychological flexibility and its components, we show the results of descriptive statistics, reliabilities and correlations of all variables in Table 1.

Table 1. Descriptive and intercorrelations values for the empirical study (N=318, 49.1% females). Cronbach reliabilities (α) for the study is also reported.

	Min-Max	M (SD)	α	1	2	3	4	5	6	7	8	9	10
1 Age	8 - 16	11.25 (2.2)	-	*									
2 Gender	(0: male, 1: female)		-	-.02	*								
3 CF_AFQ_Y	0 - 3.38	1.03 (.70)	.65	-.12*	.02	*							
4 EA_AFQ_Y	0 - 4	1.87 (.89)	.76	-.36**	.00	.62**	*						
5 AFQ_Y_Total	0 - 3.53	1.47 (.72)	.82	-.29**	.01	.87**	.93**	*					
6 DM_CAMM	0.4 - 4	2.66 (.27)	.73	.23**	.00	-.55**	-.60**	-.64**	*				
7 PA_PANASN	1 - 3	2.35 (.37)	.72	-.05	.15*	-.07	-.02	-.04	-.04	*			
8 NA_PANASN	1 - 2.6	1.61 (.39)	.78	-.11*	.13*	.40**	.33**	.40**	-.47**	-.15**	*		
9 BALANCE	-0.8 - 2	.74 (.58)	.65	.04	.01	-.31**	-.24**	-.30**	.29**	.75**	-.77**	*	
10 ANX_S	20 - 55	29.97 (6.41)	.82	-.09	-.15**	.34**	.24**	.31**	-.31**	-.29**	.37**	-.44**	*
11 ANX_T	20 - 54	33.37 (7.19)	.85	-.18*	.07	.50**	.49**	.55**	-.67**	-.00	.62**	-.42**	.34**

* $p < .05$; ** $p < .01$.

3 FC: Fusion cognitive; 4 Experiential avoidance; 5 8 AFQ: Cognitive Inflexibility Total Average; 6 Dispositional Mindfulness (DM) of CAMM; 7 Positive Affect PANASN; 8

Negative Affect PANASN; 9 Balance (POS - NEG) PANASN; 10 ANX_S: Anxiety State STAI-C; and 11 ANX_T: Anxiety Trait STAI-C

Our primary goal was to determine the mediating roles of the two PI factors (experiential avoidance and cognitive fusion). Regarding *experiential avoidance*, despite negative and moderate significance relationships with negative affect, trait, and state anxiety, EA did not show significant influences on the different mediation analysis (negative affect, anxiety trait, and anxiety state). Neither did *age* significantly covariate in such mediation analyses. Likewise, *gender* had no moderating role in further analyses. However, we did find a mediating role of *cognitive fusion* between DM and negative affect and both state and trait anxiety. We decided not report mediation analysis with balance criterion due to its being the result of the subtracted difference between the positive affection and the negative affection. This subtraction does not reflect a balance but the difference between the totals of the positive and negative items of the PANASN.

Regarding DM, CF, and negative affect of PANASN mediation analysis, there was a significant indirect effect of DM on Negative Affect through Cognitive Fusion, $b = .051$, 95% BC CI (Confidence Interval) [.019, .086], as shown in Figure 1.




Figure 1. Model of DM as predictor of Negative Affect, mediated by Cognitive Fusion.

The confidence interval for the indirect effect is a bootstrapped CI based on 5000 samples. Similar findings appeared regarding anxiety state (STAIC). There was a significant indirect effect of DM on anxiety state through Cognitive Fusion, $b = .83$, 95% CI [.24, 1.44], as shown in Figure 2.




Figure 2. Model of Mindfulness as predictor of Anxiety-estate, mediated by Cognitive Fusion.

The confidence interval for the indirect effect is a bootstrapped CI based on 5000 samples. Finally, there was also a significant indirect effect of DM on anxiety trait through Cognitive Fusion, $b = .84$, 95% CI [.29, 1.43], as shown on Figure 3.




Figure 3. Model of DM as predictor of anxiety trait, mediated by Cognitive Fusion.

The confidence interval for the indirect effect is a bootstrapped CI based on 5000 samples.

4. Discussion

The purpose of the present study was to assess the potential mediating role of cognitive fusion and experiential avoidance between DM and emotional wellbeing criteria such as negative affect and anxiety. However, our arguments should be considered tentatively for two main reasons.

On one hand, although the benefits of MBIs for school-aged children have been reported (see three systematic reviews: [1, 42, 43]), most children-and-adolescents MBI studies use DM measures rather than a systematic observation of how mindful mechanisms work for better mental and emotional health outcomes after MBIs. Hence, little is known about mindful mechanisms, so any mediation analysis could provide tentative but not, as yet, confirmatory ideas. In our study, we did not conduct an MBI, just a correlational study. However, at earlier stages of mediation analysis, these types of correlational studies serve useful purposes [44-47]. Furthermore, we used a DM measure, the Spanish eight-item CAMM [47]. We recoded the total score for a better understanding since CAMM assesses indispositional mindfulness rather than DM. The original CAMM [36] seems to be cultural biased when is translated and validated in Spanish samples [48, 49] despite good reliabilities with the 10-item CAMM, it fits better when some items were removed (#5 and 10 for Turanzas's version [35]; #2, 5, and 10 for the García-Rubio et al. version [48]; and #2, 3, 5, 6, and 10 for the Guerra et al. version [49]). Yet, our arguments about the predictor variable are tentative because DM shows a trait [12] rather than a well-trained set of thoughts and behaviours based on the principles of mindfulness training.

We used Spanish AFQ-Y [20]. This measure provided our two potential mediators (CF and EA) between DM and emotional criteria. Our findings regarding cognitive fusion mediating role were consistent. However, our concerns were addressed to experiential avoidance assessment issues. We have found some articles that report the role of the construct of experiential avoidance used by clinical researchers [40, 50]. A systematic review of the EA has revealed its relationship with various psychopathological symptoms [32], this review sustains that EA is a factor in the aetiology of maladaptive behaviours related to some mental disorders (e.g., generalized anxiety disorder). Although the current literature suggests that EA may be involved in various forms of psychopathology, some authors also alert to a "lack of theoretical integration and refinement

concerning operationalizing and assessing experiential avoidance" [40]. Therefore, in the assessment of the construct for adults, adolescents, and children doubts remain as to whether AFQ-Y correctly measures the experiential avoidance construct. We believe that EA is a negative reinforcement process [51]. Avoid and escape are two key verbs related to understanding this process [52] and its relationship with psychopathologic symptoms [23]. In targeting experiential avoidance behaviour, the central premise of behavioural theory and activation treatments is that increasing avoidance and escape behaviour will decrease exposure to positive reinforcement for healthy outcomes (e.g., increasing depression symptoms, see [52]). We agree with the arguments that the role of negative reinforcement processes (like EA) in mental health (for instance, depression) depends more on temporality terms than on the certainty or not of avoidance-or-escape behaviours [53]. Therefore, we cannot confirm whether EA has a mediating role or not due to assessment issues with AFQ-Y regarding this construct (involved Spanish AFQ items are 6 to 9, 11, 12, 14, 15, and 17). Instead of responding about the degree of certainty in an EA item ('I push away thoughts and feelings that I don't like'), we suggest including items some key information about the temporality of such information in these EA items ('I usually push away thoughts and feelings that I don't like').

With these caveats in mind, Table 1 reports high moderate and significant relationships between DM and AFQ-Y factors –CF and EA- (from $r=.57$ to $r=.93$, all $p<.001$), and moderate relationships between DM, EA, and CF and negative affect and anxiety trait, and a low moderate relationship with anxiety state. There are potential casualties between a positive DM, low CF and EA regarding negative affectivity and anxiety. Reliabilities showed a good index for all measures (predictor, mediators, and criteria). Age and sex variables had no relevant effects on the measures implemented in this study. However, only cognitive fusion seems to be a mediating role when we performed mediation and moderation analyses.

According to Kazdin's earlier mediation analysis [54], we fulfilled the four relations conditions for a mediation analysis (basically, all involved variables are related to each other, and by controlling the mediator predictor-criterion, the relationship has to change).

Cognitive fusion was positively correlated with anxiety and depression. Other studies have found a moderate association between CF and anxiety that was not due to negative affect and, besides, this association was independent of EA [55, 56]. Furthermore, a study [57] examined the mediating effect of CF in a cross-sectional design. Authors measured negative affect, depressive symptoms, rumination, cognitive fusion and mindfulness in depressed outpatients and normative individuals. Their model showed that CF was the only significant mediator of the relationship between negative affect and depressive symptoms (accounting for 61% of the total variance) [57]. Besides, a longitudinal study found that CF predicted and mediated the effect of mindfulness on negative affect, depression and posttraumatic stress symptoms [58]. Our results are in line with these studies, which highlighted the potential mediator role of CF in mental health.

Some authors hold that mindfulness and cognitive defusion could be overlap constructs. Thinking about this relationship, it could be that when an individual pays attention to his thoughts as mental events, this increases the separation or defusion of his thoughts, seeing them with a perspective [59]. In fact, in some exercise of the ACT model, mindfulness is used to increased cognitive defusion [5]. Although both terms are related, mindfulness is a concept that encompasses aspects beyond cognitive fusion. However, studies seeking to address the mechanism of mindfulness and mental health find inconsistencies [60]. Previous literature suggested that the relation between mindfulness and mental health outcomes could be a function of the effect of mindfulness on cognitive fusion [17, 61]. In our study, we found a strong relationship between both concepts and the mediator effect of CF on negative affect and anxiety. One possible explanation of our model is that CF could be proximal processes underlying the effect of mindfulness on mental health [62]. The non-judge perspective towards thought, being aware of them, without reaction, allows individuals to achieve a psychological distance from their thoughts [63]. These individuals change their old pattern of interaction with their thoughts. Consequently, the soaring effects from the sad mood to severe affective disorders could be broken [64]. Thus, the defuse perspective, which is given by mindfulness practice, may contribute to less dysregulated emotional outcomes. In conclusion, the cognitive aspect

of DM could be the key factor in the contribution of mindfulness to mental health. This finding has an important impact on treatments to improve mental health. Interventions should take into account the main role of cognitive fusion in the promotion of mental health through mindfulness approaches. As such, this study underlines the role of mindfulness and helps to understand how mindfulness is beneficial to mental health and the mediator role of CF.

As for the relationship between mindfulness and EA, although research has documented the role of EA in anxiety [65], our study does not find support for the important role of EA in psychopathology, probably because AFQ-Y does not encompass the Experiential Avoidance construct adequately. However, another study found that EA was not a predictor of depressive symptoms and negative-emotions states, being the DM a better protector factor in depression than EA [66]. These results are consistent with our findings and suggest that avoidance behaviour (towards internal or external events) it is not the main factor contributing to negative outcomes, rather it is cognitive aspects that seem to play a mediating role between DM and negative affect.

Author Contributions: MGG, JG, VML, and JMM designed the study and they had full access to all the data in the study. MGG and JG conducted all the field research. J. M. and VML performed all statistical analyses and the interpretation of the data. J.M, J.C. and MGG wrote the paper. VML supervised the whole process. All authors have read and approved the final manuscript.

Funding: Proyecto Regional de Investigación "EMOROBOTIC: Emotional Management through Robot Programming in Primary Education" (Junta de Extremadura IB16090/18HJ01) y 2. Consolidation of Research Groups (Junta de Extremadura GR18091/18.HJ.11)

Conflicts of Interest: The authors declare no conflict of interest.

References

- Tomlinson E, Yousaf O, Vittersø A, Jones L. Dispositional Mindfulness and Psychological Health: a Systematic Review. *Mindfulness* (N Y). 2017; doi:10.1007/s12671-017-0762-6
- Semple R, Lee J. Treating anxiety with mindfulness: Mindfulness-based cognitive therapy for children. In: Greco L, Hayes S, editors. *Acceptance and mindfulness treatments for children and adolescents*. New Harbinger Publications; 2008. pp. 63–68.
- Armstrong A, Morrison K, Twohig M. A Preliminary Investigation of Acceptance and Commitment Therapy for Adolescent Obsessive-Compulsive Disorder. *J Cogn Psychother*. 2013;27: 175–190. doi:10.1891/0889-8391.27.2.175
- Meagher R, Chessor D, Fogliati V. Treatment of Pathological Worry in Children With Acceptance-Based Behavioural Therapy and a Multisensory Learning Aide: A Pilot Study. 2017; doi:10.1111/ap.12288
- Hayes SC, Pistorello J, Levin ME. Acceptance and Commitment Therapy as a Unified Model of Behavior Change. *Couns Psychol*. SAGE PublicationsSage CA: Los Angeles, CA; 2012;40: 976–1002. doi:10.1177/0011000012460836
- Kashdan T, Rottenberg J. Psychological flexibility as a fundamental aspect of health. *Clin Psychol Rev*. NIH Public Access; 2010;30: 865–78. doi:10.1016/j.cpr.2010.03.001
- Keng S, Smoski M, Robins C. Effects of mindfulness on psychological health: a review of empirical studies. *Clin Psychol Rev*. NIH Public Access; 2011;31: 1041–56. doi:10.1016/j.cpr.2011.04.006
- A-Tjak J, Davis M, Morina N, Powers M, Smits J, Emmelkamp P. A Meta-Analysis of the Efficacy of Acceptance and Commitment Therapy for Clinically Relevant Mental and Physical Health Problems. *Psychother Psychosom*. 2015;84: 30–36. doi:10.1159/000365764
- Simon E, Verboon P. Psychological Inflexibility and Child Anxiety. *J Child Fam Stud*. 2016;25. doi:10.1007/s10826-016-0522-6
- Van der Gucht K, Griffith JW, Hellemans R, Bockstaele M, Pascal-Claes F, Raes F. Acceptance and Commitment Therapy (ACT) for Adolescents: Outcomes of a Large-Sample, School-Based, Cluster-Randomized Controlled Trial. *Mindfulness* (N Y). 2017;8: 408–416. doi:10.1007/s12671-016-0612-y
- Goodall K, Trejnowska A, Darling S. The relationship between dispositional mindfulness, attachment security and emotion regulation. *Pers Individ Dif*. Elsevier Ltd; 2012;52: 622–626. doi:10.1016/j.paid.2011.12.008
- Petrides K, García-Gómez M, Pérez-González JC. Pathways into psychopathology: Modeling the effects of trait emotional intelligence, mindfulness, and irrational beliefs in a clinical sample. *Clin Psychol Psychother*. 2017;24: 1130–1141. doi:10.1002/cpp.2079

13. Bluth K, Blanton P. Mindfulness and Self-Compassion: Exploring Pathways to Adolescent Emotional Well-Being. *J Child Fam Stud*. NIH Public Access; 2014;23: 1298–1309. doi:10.1007/s10826-013-9830-2
14. Ciarrochi J, Kashdan T, Leeson P, Heaven P, Jordan C. On being aware and accepting: A one-year longitudinal study into adolescent well-being. *J Adolesc*. Elsevier Ltd; 2011;34: 695–703. doi:10.1016/j.adolescence.2010.09.003
15. Greco LA, Lambert W, Baer RA. Psychological inflexibility in childhood and adolescence: Development and evaluation of the Avoidance and Fusion Questionnaire for Youth. *Psychol Assess*. 2008;20: 93–102. doi:10.1037/1040-3590.20.2.93
16. Valdivia-Salas S, Martin-Albo J, Zaldivar P, Lombas AS, Jimenez TI. Spanish Validation of the Avoidance and Fusion Questionnaire for Youth (AFQ-Y). *Assessment*. 2016; doi:10.1177/1073191116632338
17. Hayes S, Wilson K. Mindfulness: Method and Process. *Clin Psychol Sci Pract*. 2006;10: 161–165. doi:10.1093/clipsy.bpg018
18. Shonin E, Van Gordon W, Griffiths MD. Mindfulness-based interventions: Towards mindful clinical integration. *Front Psychol*. 2013;4: 1–4. doi:10.3389/fpsyg.2013.00194
19. Schweiger M, Ristallo, A., Oppo, A., Pergolizzi, T., Presti, G., and Moderato, P. Ragazzi in lotta con emozioni e pensieri: la validazione della versione italiana dell'Avoidance and Fusion Questionnaire for Youth (I-AFQ.Y). *Psicoterapia Cognitiva e Comportamentale* 2017, 23 (2), 141-162. Available at <http://web.b.ebscohost.com/ehost/pdfviewer/pdfviewer?vid=0&sid=5e9eac3-0515-4319-9665-b97af5fdb883%40pdc-v-sessmgr03>
20. Livheim F, Tengström A, Bond F, Andersson G, Dahl J, Rosendahl I. Psychometric properties of the Avoidance and Fusion Questionnaire for Youth: A psychological measure of psychological inflexibility in youth. *J Context Behav Sci*. 2016;5: 103–110. doi:10.1016/j.jcbs.2016.04.001
21. Davis D, Hayes J. What are the benefits of mindfulness? A practice review of psychotherapy-related research. *Psychotherapy*. 2011;48: 198–208. doi:10.1037/a0022062
22. Kashdan TB, Rottenberg J. Psychological flexibility as a fundamental aspect of health. *Clin Psychol Rev*. 2010;30: 865–878. doi:10.1016/j.cpr.2010.03.001
23. Hayes S, Wilson K, Gifford E, Follette V, Strosahl K. Experiential avoidance and behavioral disorders: A functional dimensional approach to diagnosis and treatment. *J Consult Clin Psychol*. 1996;64: 1152–1168. doi:10.1037/0022-006X.64.6.1152
24. Hayes S, Luoma J, Bond F, Masuda A, Lillis J. Acceptance and Commitment Therapy: Model, processes and outcomes. *Behav Res Ther*. 2006;44: 1–25. doi:10.1016/j.brat.2005.06.006
25. Kashdan T, Barrios V, Forsyth J, Steger M. Experiential avoidance as a generalized psychological vulnerability: Comparisons with coping and emotion regulation strategies. *Behav Res Ther*. Pergamon; 2006;44: 1301–1320. doi:10.1016/J.BRAT.2005.10.003
26. Lee JK, Orsillo SM. Investigating cognitive flexibility as a potential mechanism of mindfulness in Generalized Anxiety Disorder. *J Behav Ther Exp Psychiatry*. 2014;45: 208–16. doi:10.1016/j.jbtep.2013.10.008
27. Ruiz F, Langer Herrera Á, Luciano C, Cangas A, Beltrán I. Measuring experiential avoidance and psychological inflexibility: The Spanish version of the Acceptance and Action Questionnaire - II. *Psicothema*. 2013;25: 123–129.
28. Feldner M, Zvolensky M, Eifert G, Spira A. Emotional avoidance: an experimental test of individual differences and response suppression using biological challenge. *Behav Res Ther*. Pergamon; 2003;41: 403–411. doi:10.1016/S0005-7967(02)00020-7
29. Abramowitz J, Lackey G, Wheaton M. Obsessive-compulsive symptoms: The contribution of obsessional beliefs and experiential avoidance. *J Anxiety Disord*. Pergamon; 2009;23: 160–166. doi:10.1016/J.JANXDIS.2008.06.003
30. Fischer TD, Smout MF, Delfabbro PH. The relationship between psychological flexibility, early maladaptive schemas, perceived parenting and psychopathology. *J Context Behav Sci*. 2016;5: 169–177. doi:10.1016/j.jcbs.2016.06.002
31. Solé E, Tomé-Pires C, de la Vega R, Racine M, Castarlenas E, Jensen MP, et al. Cognitive Fusion and Pain Experience in Young People. *Clin J Pain*. 2016;32: 602–608. doi:10.1097/AJP.0000000000000227
32. Chawla N, Ostafin B. Experiential avoidance as a functional dimensional approach to psychopathology: An empirical review. *J Clin Psychol*. 2007;63: 871–890. doi:10.1002/jclp.20400
33. Bowen S, Chawla N, Collins S, Witkiewitz K, Hsu S, Grow J, et al. Mindfulness-based relapse prevention for substance use disorders: a pilot efficacy trial. *Subst Abuse*. NIH Public Access; 2009;30: 295–305.

- doi:10.1080/08897070903250084
34. Hayes S, Strosahl K, Wilson K, Bissett R, Pistorello J, Toarmino D, et al. Measuring experiential avoidance: A preliminary test of a working model. *Psychol Rec.* 2004;54: 553–578. doi:10.1007/BF03395492
 35. Turanzas J. Adaptación Transcultural de la Escala CAMM (Child and Adolescent Mindfulness Measure) y Estudio preliminar de sus características psicométricas. VIU Valencian International University. 2013. doi:10.1002/jclp
 36. Greco L, Baer R, Smith G. Assessing mindfulness in children and adolescents: development and validation of the Child and Adolescent Mindfulness Measure (CAMM). *Psychol Assess.* 2011;23: 606–14. doi:10.1037/a0022819
 37. Turanzas J, Cordón JR, Choca JP, Mestre JM. Evaluating the APAC (Mindfulness for Giftedness) Program in a Spanish Sample of Gifted Children: a Pilot Study. *Mindfulness*; 2018; doi:10.1007/s12671-018-0985-1
 38. García-Rubio C, Rodríguez-Carvajal R, Langer A, Paniagua D, Steinebach P, Andreu C, et al. Validation of the Spanish Version of the Child and Adolescent Mindfulness Measure (CAMM) with Samples of Spanish and Chilean Children and Adolescents. *Mindfulness (N Y)*. 2019; doi:10.1007/s12671-019-01108-8
 39. Sandín B. Escalas Panas de afecto positivo y negativo para niños y adolescentes (PANASN). *Rev Psicopatología y Psicol Clínica.* 2003;8: 173–182. doi:10.5944/rppc.vol.8.num.2.2003.3953
 40. Laurent J, Catanzaro SJ, Joiner TE, Rudolph KD, Potter KI, Lambert S, et al. A measure of positive and negative affect for children: Scale development and preliminary validation. *Psychol Assess.* 1999;11: 326–338. doi:10.1037/1040-3590.11.3.326
 41. Spielberger C, Gorsuch R, Lushene R. Cuestionario de Ansiedad Estado - Rasgo, STAI. [Questionnaire State - Trait Anxiety]. Madrid: TEA Ediciones; 1982.
 42. Burke C. Mindfulness-Based Approaches with Children and Adolescents: A Preliminary Review of Current Research in an Emergent Field. *J Child Fam Stud.* Springer US; 2010;19: 133–144. doi:10.1007/s10826-009-9282-x
 43. Harnett PH, Dawe S. The contribution of mindfulness-based therapies for children and families and proposed conceptual integration. *Child Adolesc Ment Health.* Blackwell Publishing Ltd; 2012;17: 195–208. doi:10.1111/j.1475-3588.2011.00643.x
 44. Goodall K, Trejnowska A, Darling S. The relationship between dispositional mindfulness, attachment security and emotion regulation. *Pers Individ Dif.* Elsevier Ltd; 2012;52: 622–626. doi:10.1016/j.paid.2011.12.008
 45. Kazdin AE, Nock MK. Delineating mechanisms of change in child and adolescent therapy: methodological issues and research recommendations.
 46. Kraemer HC, Stice E, Kazdin A, Offord D, Kupfer D. How Do Risk Factors Work Together? Mediators, Moderators, and Independent, Overlapping, and Proxy Risk Factors. *Am J Psychiatry.* 2001;158: 848–856. doi:10.1176/appi.ajp.158.6.848
 47. Breslin F, Zack M, McMain S. An Information-Processing Analysis of Mindfulness: Implications for Relapse Prevention in the Treatment of Substance Abuse. *Clin Psychol Sci Pract.* Blackwell Publishing Ltd; 2002;9: 275–299. doi:10.1093/clipsy.9.3.275
 48. García-Rubio C, Luna T, Castillo R, Rodríguez-Carvajal R. Impacto de una intervención breve basada en mindfulness en niños: un estudio piloto. *Rev Interuniv Form del Profr.* 2016;87: 3.
 49. Guerra, García-Gómez, Turanzas, Cordón, Suárez-Jurado, Mestre. A Brief Spanish Version of the Child and Adolescent Mindfulness Measure (CAMM). A Dispositional Mindfulness Measure. *Int J Environ Res Public Health.* Multidisciplinary Digital Publishing Institute; 2019;16: 1355. doi:10.3390/ijerph16081355
 50. Muris P, Meesters C, Herings A, Jansen M, Vossen C. Inflexible Youngsters: Psychological and Psychopathological Correlates of the Avoidance and Fusion Questionnaire for Youths in Nonclinical Dutch Adolescents. *Mindfulness*; 2017; doi:10.1007/s12671-017-0714-1
 51. Biglan A, Hayes S, Pistorello J. Acceptance and Commitment: Implications for Prevention Science. *Prev Sci.* 2008;9: 139–152. doi:10.1007/s11121-008-0099-4
 52. Carvalho JP, Hopko DR. Journal of Behavior Therapy and Behavioral theory of depression : Reinforcement as a mediating variable between avoidance and depression. *J Behav Ther Exp Psychiatry.* Elsevier Ltd; 2011;42: 154–162. doi:10.1016/j.jbtep.2010.10.001
 53. Manos RC, Kanter JW, Busch AM. Author ' s personal copy Clinical Psychology Review A critical review of assessment strategies to measure the behavioral activation model of depression. *Clin Psychol Rev.* 2010;30: 547–561. doi:10.1016/j.cpr.2010.03.008

54. Kazdin A. Mediators and Mechanisms of Change in Psychotherapy Research. *Annu Rev Clin Psychol.* 2007;3: 1–27. doi:10.1146/annurev.clinpsy.3.022806.091432
55. Fergus TA. I Really Believe I Suffer From a Health Problem: Examining an Association Between Cognitive Fusion and Healthy Anxiety. *J Clin Psychol.* 2015;71: 920–934. doi:10.1002/jclp.22194
56. Dyck MJ, Denver E. Can the Emotion Recognition Ability of Deaf Children be Enhanced? A Pilot Study. *J Deaf Stud Deaf Educ.* 2003;8: 348–56. doi:10.1093/deafed/eng019
57. Costa J, Pinto-Gouveia J, Marôco J. The Role of Negative Affect, Rumination, Cognitive Fusion and Mindfulness on Depressive Symptoms in Depressed Outpatients and Normative Individuals. *Int J Psychol Psychol Ther.* 2018;18: 207–220.
58. Nitzan-Assayag Y, Yuval K, Tanay G, Aderka IM. Reduced Reactivity to and Suppression of Thoughts Mediate the Effects of Mindfulness Training on Recovery Outcomes Following Exposure to Potentially Traumatic Stress. *Mindfulness.* 2017; 920–932. doi:10.1007/s12671-016-0666-x
59. Gillanders D, Bolderston H, Bond F, Dempster M, Flaxman P, Campbell L, et al. The Development and Initial Validation of the Cognitive Fusion Questionnaire. *Behav Ther. Elsevier.* 2014;45: 83–101. doi:10.1016/J.BETH.2013.09.001
60. Rau HK, Williams PG. Dispositional mindfulness: A critical review of construct validation research. *Pers Individ Dif.* 2016;93: 32–43. doi:10.1016/j.paid.2015.09.035
61. Hayes S, Wilson K. Acceptance and commitment therapy: Altering the verbal support for experiential avoidance. *Behav Anal. Association for Behavior Analysis International.* 1994;17: 289–303.
62. Nitzan-Assayag Y, Aderka I, Bernstein A. Dispositional mindfulness in trauma recovery: Prospective relations and mediating mechanisms. *J Anxiety Disord.* 2015;36: 25–32. doi:10.1016/j.janxdis.2015.07.008
63. Luoma J, Hayes S, Walser R. *Learning ACT: An Acceptance & Commitment Therapy Skills-Training Manual for Therapists.* Oakland: New Harbinger Publications; 2007.
64. Coffman S, Dimidjian S, Baer R. Mindfulness-Based Cognitive Therapy for prevention of depressive relapse. In: Baer R, editor. *Mindfulness-Based Treatment Approaches Clinician's Guide to Evidence Base and Applications.* Burlington: Elsevier Academic press; 2006. pp. 31–50.
65. Kämpfe C, Gloster A, Wittchen H, Helbig-Lang S, Lang T, Gerlach A, et al. Experiential avoidance and anxiety sensitivity in patients with panic disorder and agoraphobia: Do both constructs measure the same?. *Int J Clin Heal Psychol.* 2012;12: 5–22.