

Review

Chemical Constituents, Antimicrobial Activity and Food Preservative Characteristics of *Aloe vera* Gel

İbrahim Kahramanoğlu ¹, Chuying Chen ², Jinyin Chen ^{2,3} and Chunpeng Wan ^{2,*}

¹ European University of Lefke, Gemikonagi, Northern Cyprus, via Mersin 10 Turkey; ikahramanoglu@eul.edu.tr (İ.K.)

² Jiangxi Key Laboratory for Postharvest Technology and Nondestructive Testing of Fruits & Vegetables, College of Agronomy, Jiangxi Agricultural University, Nanchang 330045, China; cy.chen@jxau.edu.cn (C.C.); jinyinchen@126.com (J.C.)

³ Pingxiang University, Pingxiang 337055, China

* Correspondence: chunpengwan@jxau.edu.cn (C.W.); Tel.: +86-791-83813158

Abstract: Edible coating gels developed from the *Aloe vera* plant have been used as a traditional medicine for about 3000 years. *Aloe vera* contains 110 potentially active constituents from six different classes: chromone and its glycoside derivatives; anthraquinone and its glycoside derivatives; flavonoids; phenylpropanoids and coumarins; phenylpyrone and phenol derivatives; and phytosterols and others. Apart from medicinal uses, *Aloe* gels have an important role in food preservation as edible coatings. They provide an edible barrier for atmospheric gases and moisture, and help to reduce the respiration and transpiration of fresh produce, which helps to preserve its postharvest quality. To date, numerous studies have been conducted on the postharvest use of *Aloe vera* gel. The present review article summarizes and discusses existing available information about the chemical constituents, antimicrobial activity, and food preservative characteristics of *Aloe vera*.

Keywords: *Aloe vera*; chemical constituents; antimicrobial activity; postharvest storage; biodegradable; edible coating

1. Introduction

Food quality mainly refers to three attributes: external (size, colour, appearance, etc.), internal (taste, colour, juicy, texture, seedless, etc.) and hidden (food safety and nutritional contents). External and internal quality attributes were of greatest importance to consumers for many years. However, since the occurrence of food-derived health problems has begun to increase, consumers have started to pay more attention to the hidden quality attributes of fresh produce, and are asking for food to be free of chemical residues [1]. Fungicides and other agrochemicals are of great importance in controlling postharvest diseases and have crucial role for the preservation of the postharvest quality, but misuse and/or excessive use of them might cause negative impacts on human health [2,3]. There is an increasing effort in postharvest studies to develop natural preservatives and antimicrobials to extend the storage duration of foods without chemical preservatives [4,5]. So far, many storage techniques and natural preservatives have been developed to extend the postharvest life of foods. The currently utilized natural preservatives are chitosan [6,7], essential oils [8,9], propolis extract [10], plant extracts [11,12], edible coatings [13,14], and organic salts [15]. Among these, edible coatings have been receiving more attention in recent years due to their potential for developing edible packaging materials [16]. Weight loss, changes in textural quality, changes in chemical structure, and microbial pathogens (mostly fungus) are the most important postharvest problems for foods [17,18].

Aloe belongs to the family of Xanthorrhoeaceae, which consists of about 420 species, and has been used as a traditional medicine for about 3000 years [19]. The perennial plant known as *Aloe vera* is *Aloe barbadensis* Miller, which is a well-known pharmaceutical herb that has long been used

in traditional Chinese medicine for the treatment of various diseases. It is widely distributed in the semitropical regions, and cultivated in many provinces of China.

Aloe gels have an important role in food preservation as edible coatings. Edible coatings generally provide a thin layer on the fruit surface, which acts as a barrier to atmospheric gases and moisture [20,21]. *Aloe* gels help to reduce the respiration and transpiration of fresh produce and delay postharvest deterioration of foods, promoting food preservation [17]. Edible coatings are generally applied by dipping the foods, spraying, or brushing. To date, numerous studies have been conducted into the postharvest use of *Aloe vera* gel as an edible coating. The following sections of the present review article aim to summarize and discuss the existing available information regarding the use of *Aloe vera* gel as a food preservative. First, however, it is important to mention the chemical constituents and antimicrobial activity of *Aloe vera*, which are summarized herein.

2. Chemical Constituents of *Aloe vera*

The two main class active constituent of the *Aloe vera* plant extract are chromone and anthraquinone and its glycoside derivatives, alongside others such as phenylpyrone derivatives, flavonoids, phenylpropanoids, coumarins, phytosterols, naphthalene analogs, lipids and vitamins.

2.1. Chromone and its Glycoside Derivatives

Approximately 29 chromone derivatives were isolated and identified from *Aloe vera* (Table 1, Figure 1). Aloesin (**1**, formerly called aloeresin B), aloeresin A (**23**), isoaloeresin D (**13**) and aloeresin E (**9**) are the most significant active constituents of *Aloe vera*. Three aloediols (**7**, **8** and **9**) were isolated and identified from *Aloe vera*, but the absolute configuration has not yet been determined.




Figure 1. Chemical structure of chromone and its glycoside derivatives from *Aloe vera*.

Table 1. Chromone and its glycoside derivatives isolated and identified from *Aloe vera*.

No	Constituents	Molecular formula	Exact Mass	References
1	aloesin	C ₁₉ H ₂₂ O ₉	394.1264	[22,23]
2	neoaloesin A	C ₁₉ H ₂₂ O ₉	394.1264	[24]
3	8-C-glucosyl-(R)-aloesol	C ₁₉ H ₂₄ O ₉	396.142	[22]
4	8-C-glucosyl-7-methoxy-(R)-aloesol	C ₂₀ H ₂₆ O ₉	410.1577	[22]
5	8-C-glucosyl-(S)-aloesol	C ₁₉ H ₂₄ O ₉	396.142	[25]
6	8-C-glucosyl-7-methoxy-(S)-aloesol	C ₂₀ H ₂₆ O ₉	410.1577	[25,26]
7	8-C-glucosyl-7-O-methylaloediol	C ₂₀ H ₂₆ O ₁₀	426.1526	[22,25]
8	8-glucosyl-(2'-O-cinnamoyl)-7-O-methylaloediol A	C ₂₉ H ₃₂ O ₁₂	572.1894	[27]
9	8-glucosyl-(2'-O-cinnamoyl)-7-O-methylaloediol B	C ₂₉ H ₃₂ O ₁₂	572.1894	[27]
10	C-2'-decoumaroyl-aloesin G	C ₂₀ H ₂₄ O ₈	392.1471	[22]
11	aloesin E	C ₂₉ H ₃₂ O ₁₀	540.1995	[26]
12	isoaloesin D	C ₂₉ H ₃₂ O ₁₁	556.1945	[26,28,29]
13	iso-rabaichromone	C ₂₉ H ₃₂ O ₁₂	572.1894	[28]
14	8-[C-β-D-[2-O-(E)-cinnamoyl]glucopyranosyl]- 2-[(R)-2-hydroxypropyl]-7-methoxy-5-methylchromone	C ₂₉ H ₃₂ O ₁₀	540.1995	[30]
15	aloesin D	C ₂₉ H ₃₂ O ₁₁	556.1945	[22,30]
16	rabaichromone	C ₂₉ H ₃₂ O ₁₂	572.1894	[22]
17	allo-aloesin D	C ₂₉ H ₃₂ O ₁₁	556.1945	[22]
18	aloesin K	C ₃₁ H ₃₄ O ₁₂	598.205	[29]
19	aloesin J	C ₃₀ H ₃₄ O ₁₁	570.2101	[29]
20	8-C-glucosyl-noreugenin	C ₁₆ H ₁₈ O ₉	354.0951	[27]
21	4'-O-glucosyl-isoaloesin DI	C ₃₅ H ₄₂ O ₁₆	718.2473	[27]
22	4'-O-glucosyl-isoaloesin DII	C ₃₅ H ₄₂ O ₁₆	718.2473	[27]
23	aloesin A	C ₂₈ H ₂₈ O ₁₁	540.1632	[23]
24	7-O-methyl-aloesin A	C ₂₉ H ₃₀ O ₁₁	554.1788	[23,31]
25	9-dihydroxyl-2'-O-(Z)-cinnamoyl-7-methoxy-aloesin	C ₂₉ H ₃₀ O ₁₂	570.1737	[31]
26	6'-O-coumaroyl-aloesin	C ₂₈ H ₂₈ O ₁₂	556.1581	[32]
27	7-methoxy-6'-O-coumaroyl-aloesin	C ₂₉ H ₃₀ O ₁₂	570.1737	[33]
28	aloeveraside B	C ₂₈ H ₂₈ O ₁₂	556.1581	[32,34]
29	aloeveraside A	C ₂₉ H ₃₀ O ₁₂	570.1737	[32,34]

2.2. Anthraquinone and its Glycoside Derivatives

Approximately 32 anthraquinones and their glycoside derivatives were isolated and identified from *Aloe vera* (Table 2, Figure 2). The isomers of aloin A (**30**) and aloin B (**31**), two anthraquinone glucosides, are the most abundant active constituents of *Aloe vera*. However, chrysophanol (**52**), emodin (**53**), physcione (**54**), alo-emodin (**55**) are four major anthraquinone aglycones. Six anthraquinone dimmers (**45–50**) were also identified from *Aloe vera*.

Table 2. Anthraquinone and its glycoside derivatives isolated and identified from *Aloe vera*.

No	Constituents	Molecular formula	Exact Mass	References
30	aloin A	C ₂₁ H ₂₂ O ₉	418.1264	[29]
31	aloin B	C ₂₁ H ₂₂ O ₉	418.1264	[29]
32	6'-O-acetyl-aloin A	C ₂₃ H ₂₄ O ₁₀	460.1369	[29]
33	6'-O-acetyl-aloin B	C ₂₃ H ₂₄ O ₁₀	460.1369	[29]
34	10-hydroxyaloin A	C ₂₁ H ₂₂ O ₁₀	434.1213	[28,32]
35	10-hydroxyaloin B	C ₂₁ H ₂₂ O ₁₀	434.1213	[28,32]
36	aloinoside A	C ₂₇ H ₃₂ O ₁₃	564.1843	[29]
37	aloinoside B	C ₂₇ H ₃₂ O ₁₃	564.1843	[29]
38	7-hydroxyaloin A	C ₂₁ H ₂₂ O ₁₀	434.1213	[23]
39	7-hydroxyaloin B	C ₂₁ H ₂₂ O ₁₀	434.1213	[23]
40	7-hydroxy-8-O-methylaloin A	C ₂₂ H ₂₄ O ₁₀	448.1369	[23,28]
41	7-hydroxy-8-O-methylaloin B	C ₂₂ H ₂₄ O ₁₀	448.1369	[23,28]
42	6'-malonylnataloin A	C ₂₄ H ₂₄ O ₁₂	504.1268	[23]
43	6'-malonylnataloin B	C ₂₄ H ₂₄ O ₁₂	504.1268	[23]
44	homonataloside B	C ₂₈ H ₃₄ O ₁₄	594.1949	[23]
45	elgonica dimer A	C ₃₆ H ₃₀ O ₁₄	686.1636	[29,35,36]
46	elgonica dimer B	C ₃₆ H ₃₀ O ₁₄	686.1636	[29,35,36]
47	aloindimer A	C ₄₂ H ₄₂ O ₁₈	834.2371	[29]
48	aloindimer B	C ₄₂ H ₄₂ O ₁₈	834.2371	[29]
49	aloindimer C	C ₄₂ H ₄₂ O ₁₈	834.2371	[29]
50	aloindimer D	C ₄₂ H ₄₂ O ₁₈	834.2371	[29]
51	aloe-emodin-11-O-rhamnoside	C ₂₁ H ₂₀ O ₉	416.1107	[33]
52	chrysophanol	C ₁₅ H ₁₀ O ₄	254.0579	[17]
53	emodin	C ₁₅ H ₁₀ O ₅	270.0528	[33]
54	physcione	C ₁₆ H ₁₂ O ₅	284.0685	[37]
55	aloe-emodin	C ₁₅ H ₁₀ O ₅	270.0528	[33]
56	nataloeemodin	C ₁₅ H ₁₀ O ₅	270.0528	[23]
57	aloesaponarin I	C ₁₇ H ₁₂ O ₆	312.0634	[38]
58	aloesaponarin II	C ₁₅ H ₁₀ O ₄	254.0579	[38]
59	madagascine	C ₂₀ H ₁₈ O ₅	338.1154	[39]
60	3-Geranyloxyemodin	C ₂₄ H ₂₄ O ₅	392.1624	[39]
61	rhein	C ₁₅ H ₈ O ₆	284.0321	[37]





Figure 2. Chemical structure of anthraquinone and its glycoside derivatives from *Aloe vera*.

2.3. Flavonoids

Approximately 13 flavonoids and their glycoside derivatives were isolated and identified from *Aloe vera* (Table 3, Figure 3), including three types; namely flavone (62–67), flavonol (68–72) and flavan-3-ol (73,74).

**Figure 3.** Chemical structure of flavonoids from *Aloe vera*.**Table 3.** Flavonoids isolated and identified from *Aloe vera*.

No	Constituents	Molecular formula	Exact Mass	References
62	apigenin	C ₁₅ H ₁₀ O ₅	270.0528	[40]
63	luteolin	C ₁₅ H ₁₀ O ₆	286.0477	[41]
64	isovitexin	C ₂₁ H ₂₀ O ₁₀	432.1056	[41]
65	isorientin	C ₂₁ H ₂₀ O ₁₁	448.1006	[41]
66	saponarin	C ₂₇ H ₃₀ O ₁₅	594.1585	[41]
67	lutonarin	C ₂₇ H ₃₀ O ₁₆	610.1534	[41]
68	kaempferol	C ₁₅ H ₁₀ O ₆	286.0477	[40]
69	quercetin	C ₁₅ H ₁₀ O ₇	302.0427	[40]
70	myricetin	C ₁₅ H ₁₀ O ₈	318.0376	[40]
71	quercitrin	C ₂₁ H ₂₀ O ₁₁	448.1006	[40]
72	rutin	C ₂₇ H ₃₀ O ₁₆	610.1534	[40]
73	catechin	C ₁₅ H ₁₄ O ₆	290.0790	[40]
74	epicatechin	C ₁₅ H ₁₄ O ₆	290.0790	[40]

2.4. Phenylpropanoids and Coumarins

Approximately 12 phenylpropanoid acids and their ester derivatives (75–86), and four coumarins (87–90), were isolated and identified from *Aloe vera* (Table 4, Figure 4).

Table 4. Phenylpropanoids and coumarins isolated and identified from *Aloe vera*.

No	Constituents	Molecular formula	Exact Mass	References
75	cinnamic acid	C ₉ H ₈ O ₂	148.0524	[42]
76	p-coumaric	C ₉ H ₈ O ₃	164.0473	[40]
77	caffeic acid	C ₉ H ₈ O ₄	180.0423	[40]
78	ferulic acid	C ₁₀ H ₁₀ O ₄	194.0579	[40]
79	sinapic acid	C ₁₁ H ₁₂ O ₅	224.0685	[40]

80	5-p-coumaroylquinic	C ₁₆ H ₁₈ O ₈	338.1002	[41]
81	chlorogenic	C ₁₆ H ₁₈ O ₉	354.0951	[40]
82	5-feruloylquinic	C ₁₇ H ₂₀ O ₉	368.1107	[41]
83	caffeoylshikimic	C ₁₆ H ₁₆ O ₈	336.0845	[41]
84	5-p-cis-coumaroylquinic	C ₁₆ H ₁₈ O ₈	338.1002	[41]
85	3-(4-hydroxyphenyl)propanoic acid	C ₉ H ₁₀ O ₃	166.063	[32]
86	methyl 3-(4-hydroxyphenyl)propionate	C ₁₀ H ₁₂ O ₃	180.0786	[32]
87	7-demethylsiderin	C ₁₁ H ₁₀ O ₄	206.0579	[32]
88	feralolide	C ₁₈ H ₁₆ O ₇	344.0896	[36]
89	dihydrocoumarin	C ₂₂ H ₁₈ O ₇	394.1053	[43]
90	dihydrocoumarin ethyl ester	C ₂₄ H ₂₄ O ₈	440.1471	[43]




Figure 4. Chemical structure of phenylpropanoids and coumarins from *Aloe vera*.

2.5 Phenylpyrone and Phenol Derivatives

Approximately three phenylpyrone derivatives (**91–93**), one triglucosylated naphthalene derivative named aloverside A (**94**), and one 1-methyltetralin derivative feroxidin (**95**) were isolated and identified from *Aloe vera* (Table 5, Figure 5). Nine phenol derivatives (**96–104**) and vitamin C (**105**) were also isolated from *Aloe vera*.

Table 5. Phenylpyrone and phenol derivatives isolated and identified from *Aloe vera*.

No	Constituents	Molecular formula	Exact Mass	References
91	aloenin A	C ₁₉ H ₂₂ O ₁₀	410.1213	[44]
92	aloenin B	C ₃₄ H ₃₈ O ₁₇	718.2109	[35,44]
93	p-coumaroyl aloenin	C ₂₈ H ₂₈ O ₁₂	556.1581	[35]
94	aloverside A	C ₃₀ H ₄₀ O ₁₇	672.2265	[35]
95	feroxidin	C ₁₁ H ₁₄ O ₃	194.0943	[32]
96	1-(2,4-dihydroxy-6-methylphenyl)ethanone	C ₉ H ₁₀ O ₃	166.0630	[32]
97	p-anisaldehyde	C ₈ H ₈ O ₂	136.0524	[32]
98	salicylaldehyde	C ₇ H ₆ O ₂	122.0368	[32]
99	p-cresol	C ₇ H ₈ O	108.0575	[32]
100	pyrocatechol	C ₆ H ₆ O ₂	110.0368	[42]

101	gentisic acid	$C_7H_6O_4$	154.0266	[40]
102	gallic acid	$C_7H_6O_5$	170.0215	[40]
103	vanillic acid	$C_8H_8O_4$	168.0423	[40]
104	syringic acid	$C_9H_{10}O_5$	198.0528	[40]
105	ascorbic acid	$C_6H_8O_6$	176.0321	[40]




Figure 5. Chemical structure of phenylpyrone and phenol derivatives from *Aloe vera*.

2.6 Phytosterols and Others

Five phytosterols (Table 6, Figure 6) were isolated from *Aloe vera* gel, including cycloartanol (**106**), 24-methylene-cycloartanol (**107**), lophenol (**108**), 24-methyl-lophenol (**109**), and 24-ethyl-lophenol (**110**). Some polar and nonpolar lipids, as well as prostanoids, were also isolated from *Aloe vera* leaves [45]. Chemical investigation of the major constituents in *Aloe vera* leaves revealed moisture, ash, fiber, protein, lipids, minerals, organic acids, free sugars, and polysaccharides. Glucose, fructose, and sucrose were the main free sugars. Oxalic, L-Malic, isocitric, lactic, acetic, isocitric, lactone, citric, and fumaric acid were the main organic acids [43].

Table 6. Phytosterols isolated and identified from *Aloe vera*.

No	Constituents	Molecular formula	Exact Mass	References
106	cycloartanol	$C_{30}H_{52}O$	428.4018	[46]
107	24-methylene-cycloartanol	$C_{31}H_{52}O$	440.4018	[46]
108	lophenol	$C_{28}H_{48}O$	400.3705	[46]
109	24-methyl-lophenol	$C_{29}H_{50}O$	414.3862	[46]
110	24-ethyl-lophenol	$C_{30}H_{52}O$	428.4018	[46]




Figure 6. Chemical structure of phytosterols from *Aloe vera*.

3. Antimicrobial Activity of *Aloe vera*

Aloe vera plant extracts have antimicrobial characteristics that kill microorganisms (including bacteria [anti-bacterial activity], fungi [anti-fungal activity] and viruses [anti-viral activity]) or stop their growth. Fruit decay is an important parameter influencing the postharvest quality of fresh produce. Previous studies have shown that the use of *Aloe vera* gel as an edible coating has positive effects on the prevention of fruit decay and microbial spoilage. The inhibitory effects of *Aloe vera* gel on the growth of mycelium (*Penicillium digitatum* and *Aspergillus niger*) was reported by Nabigol and Asghari [47], who performed a range of laboratory tests. They suggested that the inhibition of the mycelium growth rate increased with gel concentration. The 500 ml/L dose of *Aloe vera* gel was found to cause 100% inhibition of *P. digitatum* and 64% of *A. niger*. According to the findings of Kator et al. [48], 20%, 60% and 100% concentrations of *Aloe vera* gel are effective in preventing the occurrence of decay in tomato fruits for seven days of storage. However, these authors suggested that a 100% concentration has significantly higher effects, and the positive impact may continue for 16 days of storage. Benitez et al. [49] reported that *Aloe vera* gel provides higher efficacy for the prevention of mesophilic bacteria and yeasts and moulds than alginate and chitosan for kiwifruit slices. In another study, the shelf life of guava was reported to be increased by about one more week with the application of an *Aloe vera* gel coating, due to the fact that the edible coating prevents microbial growth [50]. Sitara et al. [51] conducted a comprehensive study regarding the antifungal activity of *Aloe vera* gel at three different doses against five plant pathogenic fungi: *A. niger*, *Aspergillus flavus*, *Alternaria alternata*, *Drechslera hawaiiensis* and *P. digitatum*. The highest test dose (0.35%) of *Aloe vera* gel was reported to completely inhibit the growth of *Drechslera hawaiiensis* and *Alternaria alternata*. In another study, the minimum fungicidal concentrations of *Aloe vera* against *Botrytis gladiolorum*, *Fusarium oxysporum* f.sp. *gladioli*, *Heterosporium pruneti* and *Penicillium gladioli* were reported to vary between 80 and 100 µl/ml, depending on the fungal species [52].

Previous studies have also shown that the combination of *Aloe vera* gel with some homogenizers, such as glycerol starch (0.15 g), improves the efficacy in controlling fungal decay and weight loss in cherry tomatoes [53]. The specific mechanism of action is still unknown but it is known that saponins, acemannan and anthraquinone derivatives, which are found in *Aloe vera*, have antibacterial activity [54]. Navarro et al. [55] performed a study with *Aloe vera* gel alone or in combination with thymol on nectarines and reported that the *Aloe vera* gel alone is more efficient in prevention of the decay caused by *Rhizopus stolonifer*, *B. cinerea* and *P. digitatum*. *Aloe vera* gel coatings were previously tested against decay and found to significantly lower counts for moulds, yeast, and mesophilic aerobics in different fruits and vegetables, including tomatoes [56,57], citrus fruits [58,59], raspberry fruits [60], blueberries [61], strawberries [62], and ready-to-eat pomegranate arils [63]. The preharvest application of *Aloe vera* gel treatment was also previously tested and found to be effective in postharvest storage; specifically, it was found to reduce the decay incidence of table grapes [64,65].

In a different study [66], *Aloe vera* leaf gel was found to inhibit the growth of two bacteria: *Shigella flexneri* and *Streptococcus progenes*. The anti-bacterial activities of *A. vera* gel was also reported by Wang et al. [67] and Cellini et al. [68] against *Helicobacter pylori*. Moreover, anti-viral activities of *A. vera* have also been of interest to many researchers, wherein its positive influence has been reported against herpes simplex virus (HSV) type 2 strains by Zandi and Rastian [69] and against influenza A virus replication by Li et al. [70].

4. Food Preservative Characteristics of *Aloe vera*

Numerous previous studies and the current extensive review have concluded that *Aloe vera* gel coatings have been used to preserve the postharvest quality of numerous fruits (Table 7). Herein, we summarized the effect of *Aloe vera* gel coatings on the respiration rate, weight loss, soluble solids concentration (SSC), titratable acidity (TA), firmness, ethylene production, overall appearance and colour, aroma volatile biosynthesis, and ascorbic acid content of preserved postharvest fruits.

1

Table 7. Use of *Aloe vera* gel as an edible coating on different fruits.

Fruits	Dose of <i>A. vera</i>	Incorporate with	Fruit storage conditions	Acceptable storage duration for		Ref
				untreated fruits	treated fruits	
Pineapples	100%	Ascorbic acid (1.9 - 2.0g L ⁻¹) and citric acid (4.5 - 4.6g L ⁻¹)	Ambient temperature (27 + 2 °C) and 50–60% RH	21 days	49 days	[73]
Pistachio	50% and 100%	Chitosan (0.5% and 1.0%)	At 4 °C	-	30 days	[82]
Nectarine cv ‘Arctic Snow’	2.5 g/L	0.05% Tween-20	At 0 ± 0.5 °C and 90 ± 5% RH	21 days	42 days	[71]
Tomato cv. ‘Ruchi 618’	2%	0.3% antioxidant rich herb, Glycerol (2 %) and oleic acid (0.6%)	N/A	20 days	39 days	[84]
Tomato cv. ‘Dafni’	10% and 10%	-	At 11 °C and 90% RH in darkness	7 days	14 days	[57]
Tomato var. ‘Roma’ and ‘UTC’	20%, 60%, 100%	-	N/A	7 days	13–16 days	[48]
Table grape ‘Crimson Seedless’	33.3%	-	At 2 °C in controlled chambers with 85-90% RH	5–10 days	15–20 days	[64]
Grape ‘Thompson’	5% and 10%	-	In air tight plastic container and at 15 °C, 96-98% RH	15 days	40 days	[65]
Raspberry (grown naturally in Iran)	25%, 50%, 75%	-	At 4 °C	4 days	8 days	[80]
Peach (from Iran)	25%	-	Air dried and stored at 1 °C and 95% RH	10 days	20-30 days	[86]
Nectarine cv. ‘Flavela’ and ‘Flanoba’		1 mL/L. Thymol (99.5% purity)	At 25 °C and 85% RH	-	6 days	[55]
Sweet cherry cv. ‘Star King’	25%	-	Air-dried and stored at 1 °C and 95% RH	2–6 days	9–16 days	[88]
Plum cv. ‘President’	100%	Rosehip oil (2%)	In a controlled chamber at 20 °C and 85% RH or at 2 °C and 90% RH	14 days	28 days	[80]
Bell pepper cv. ‘Cardio’	30%	Gum tragacanth (20% w/w)	At 4, 10, 15 and 23 °C	6 days	18–22 days	[79]
Strawberry cv. ‘Bari’	100%	1% (w/v) CMC	At 6 ± 1 °C and 50 ± 5% RH	3–6 days	12–15 days	[62]
Mango var. ‘Ngowe’	25%, 50%, 75%	-	At 13 and 15–22 °C			[85]
Fresh-cut kiwifruit cv. ‘Hayward’	5%	-	At 4 ± 1 °C and 75% RH	6 days	11 days	[49]
Fresh-cut papaya cv. Pusa delicious	100%	1.5% glycerol	At 4 ± 1 °C and 95% RH	6 days	12 days	[75]
Ready-to-eat pomegranate arils cv. Mollar de Elche	50% and 100%	Ascorbic acid and citric acid (0.5% and 1.0%)	At 3 °C and 90% RH	4 days	8–12 days	[63]
Minimally processed pomegranate arils cv. ‘Malas Saveh’	60, 125, 250, 500 ml/L	-	At 5 °C and 95% RH	-	-	[47]
Fresh-cut oranges	50% and 100%	Gelatin	At 4 °C	9 days	17 days	[81]
Fresh-cut apples cv. ‘Hongro’	50%	0.5% Cysteine	At 4 °C	8 days	16 days	[89]

4.1. Respiration Rate and Weight Loss

Respiration is an important characteristic for the postharvest quality of fresh produce. The higher the respiration rate during storage, the lower the storage duration of fruits, and vice versa. Weight loss is an important parameter for determination of the postharvest quality of fresh produce. It is mainly a result of respiration (loss of carbon reserves) and transpiration (loss of water). The gel of *Aloe vera* acts as a barrier, thereby restricting water transfer and preventing weight loss. Ahmed et al. [71] reported that *Aloe vera* gel reduces the respiration rate, retarding fruit softening and reduce weight loss in 'Arctic Snow' nectarine fruits kept in ambient and cold storage conditions. Researchers used *Aloe vera* gel dried powder (200:1) in a ratio of 2.5/1 (g/L, w/v) to prepare the coating material, and Tween-20 surfactant was also used at 0.05% concentration. Another study by Achipiz et al. [72] showed that the *Aloe vera*-based edible coating is successful for extending the shelf life of guava (*Psidium guajava*). The researchers suggested that the positive effect of *Aloe vera* in preventing weight loss is the result of the reducing effect of the gel on the respiratory rate of the fruit. The positive effects of *Aloe vera* gel as an edible coating on the prevention of weight loss in pineapple fruits stored at ambient temperatures (27±2 °C) and 55–60% relative humidity was also discovered by Adetunji et al. [73]. The researchers reported that coating pineapple fruits with *Aloe vera* gel extends the storability of the fruits by seven weeks. Positive effects of *Aloe vera* gel on the prevention of weight loss were also noted for tomatoes. Kator et al. [48] noted that the use of 20%, 60% and 100% concentrations of *Aloe vera* gel resulted in 58.20, 59.20 and 65.80 g tomatoes after 16 days of storage, while the untreated tomato fruits were found to have only 25.90 g of fruit. The initial weights of the tested tomato fruits were between 82.40–83.80 g for all treatments. In another study [57] it was reported that the different concentrations of *Aloe vera* show different efficacies in the postharvest quality of fresh tomato fruits, and a 15% concentration provides higher efficacy than 5% and 10% concentrations. The positive influence of *Aloe vera* gel on the prevention of weight loss for pomegranate arils was reported by Nabigol and Asghari [47]. It was found that the combination of *Aloe vera* gel with calcium or citric acid improves the efficacy in the prevention of weight loss of grape fruits [65,74]. The positive influence of *Aloe vera* gel on the reduction of the respiration rate was previously reported for kiwifruit slices. Benitez et al. [49] noted that the O₂ concentration in the headspace of packages decreases in control fruits, but stays higher in fruits treated with *Aloe vera* gel. Contrary to O₂, the CO₂ level is reported to be lower in the treated fruits. The reason for this influence was attributed to the permeability properties of *Aloe vera* gel by the researchers. Similar findings were reported by Martínez-Romero et al. [63] for ready-to-eat pomegranate arils. The researchers noted that the application of *Aloe vera* alone or in combination with citric acid led to lower CO₂ and higher O₂ concentrations inside the packages. The potential postharvest effects of *Aloe vera* gel were also tested under ambient conditions. In one such study, it was found that an *Aloe vera* gel coating prevents weight loss in papaya fruits under ambient conditions [75]. It was also reported that the *Aloe vera* gel coating retards the ethylene production rate, and delays ripening. The impact of *Aloe vera* gel on the postharvest quality attributes is also reported to improve with the combination of some other materials, including chitosan [76], plantain flour [77,78], glycerol starch [53], gum tragacanth [79], rosehip oil [80], and gelatin [81]. Previous studies have also shown that *Aloe vera* gel prevents weight loss in pistachios [82], strawberries [62,83], tomatoes [56,57,84], bell peppers [79], raspberry fruits [60], mangoes [85], blueberries [61], peaches [86], nectarines [71], sour cherries [87], sweet cherries [88], plums [80], fresh-cut papaya fruits [75], fresh-cut oranges [81], and fresh-cut apples [89].

4.2. Soluble Solids Concentration (SSC) and Titratable Acidity (TA)

The soluble solids concentration is a measure of the total soluble solids in fruit juices. It plays a crucial role in the taste of fresh produce, and alterations in the SSC are of utmost important for postharvest storage. The SSC content is highly related to the total amount of soluble solids and the water content. Thus, respiration and transpiration have a large influence on the SSC content of fruits. Therefore, *Aloe vera* gel has the potential to prevent the loss of SSC through reducing

respiration and transpiration. Nabigol and Asghari [47] reported that different doses of *Aloe vera* gel (60, 125 and 250 ml/L) provide favourable conditions for the storage of pomegranate arils, and help to maintain the SSC for 21 days of storage when compared with untreated pomegranate arils. Similarly, *Aloe vera* gel has been reported to reduce the speed of changes in the soluble solids concentration of strawberries [83], tomatoes [57,84], raspberry fruits [60], blueberries [61], peaches [86], fresh-cut papaya fruits [75], sour cherries [87], sweet cherries [88], and fresh-cut oranges [81].

Researchers reported that the titratable acidity (TA) of fruits generally decreases during storage, but edible coating materials would provide favourable conditions for reducing the speed of the decrease in TA. It was found that the application of *Aloe vera* gel prevents the loss of acidity in grapefruits, and this effect increases when the *Aloe vera* gel is incorporated with calcium or citric acid [74]. Some other previous studies have also noted that the use of an *Aloe vera* gel coating helps to maintain TA of fruits during storage, including tomatoes [57], raspberry fruits [60], mangoes [85], peaches [86], fresh-cut papaya [75], and fresh-cut oranges [81].

4.3. Firmness

Fruit texture and firmness are important characteristics which affect consumer preferences for fresh produce. Previous studies have shown that the application of *Aloe vera* gel as an edible coating during postharvest storage helps to maintain fruit firmness, or reduce the speed of firmness loss. For example, previous studies have shown that the fruit firmness of pineapples stored at ambient temperatures (27±2 °C) and 55–60% relative humidity for seven weeks with an *Aloe vera* gel coating was about 600 N, while the firmness of uncoated fruits was found to be less than 100 N at the same time [73]. *Aloe vera* gel at a concentration of 20%, 60% and 100% was reported to maintain the fruit firmness of tomatoes for about 16 days of storage [48]. Castillo et al. [64] noted similar results for table grapes, while *Aloe vera* gel was found to keep berry firmness higher than the control. Hazrati et al. [86] conducted a study with *Aloe vera* gel coatings on peach fruits and noted that they protect against enzymes which have cell wall degradation activity. The *Aloe vera* gel coated fruits were reported to maintain the turgor pressure of the cell walls and incorporate texture enhancers to reduce firmness loss. An *Aloe vera* coating has previously been reported to promote firmness retention in different kinds of fruits and/or ready-to-eat fruits, including pomegranate arils [63], tomatoes [84], peaches [86], strawberries [62], sour cherries [87], sweet cherries [88], plums [80] and fresh-cut apples [89].

4.4. Ethylene Production

Ethylene (C₂H₄) is a natural phytohormone responsible for regulation of the growth and senescence of plants [90–92]. It governs the development of leaves, flowers, and especially fruits. Due to its special characteristics, it is used to accelerate the ripening process of climacteric fruits, such as bananas, mangoes, apples and similar, just before they are transferred to markets. This also allows the harvest of climacteric fruits before ripening, and increases the storage duration of the fruits. It was found that the *Aloe vera* gel coating prevents ethylene production by peach and plum fruits, thereby protecting the postharvest quality of the fruits. Guillén et al. [93] reported 70% and 50% inhibition of ethylene production, respectively.

4.5. Overall Appearance and Colour

Although food safety and nutrition are of great importance for consumers, visual appearance is still the first impression and a key characteristic in the choice of fruits. Moreover, external colour is one of the most important visual characteristic for fruits. Vanaei et al. [82] noted that the application of *Aloe vera* gel protects the brightness of pistachio fruits, and this positive influence increases when the gel is incorporated with chitosan. The researchers noted that the bright colour of pistachio bark darkened during storage with all treatments, but the degree of darkening was lowest in the fruits treated with the combination of *Aloe vera* gel (50%) and chitosan (1%). Similarly, a delay in changes in the external colour of peach and plum fruits was reported when treatments with *Aloe*

vera was undertaken, whereby the chroma index significantly decreased in untreated fruits but the decrease was lower in treated fruits [93]. Similar results were also noted by Benitez et al. [49], who reported that *Aloe vera* gel treated kiwifruit slices have a higher chroma index as compared to untreated fruits. Sharmin et al. [75] conducted a study of papaya fruits and reported that the fruit peel color changed from green to greenish yellow in untreated control fruits; *Aloe vera* treated papayas showed retarded colour and physiological changes for up to 12 days during storage. Studies with fresh-cut apples also showed that an *Aloe vera* gel coating 75% (v/v) alleviates browning and helps to maintain surface colour [94]. *Aloe vera* gel alone or in combination with gum tragacanth has been reported to delay colour changes in bell peppers during long-term storage [79]. Prevention of colour changes by the application of an *Aloe vera* gel coating was also reported for mangoes [85], fresh-cut oranges [81], strawberries [62], plums [80] and fresh-cut apples [89].

4.6. Aroma Volatile Biosynthesis

Flavour composition is a complex attribute for fruits, which includes sugars, acids, and volatiles. Aroma has an important influence on consumers' preferences by significantly affecting the four basic flavours (sweetness, sourness, saltiness, and bitterness). Fruit aroma develops from a complex mixture of many volatile compounds, including alcohols, aldehydes, and esters. Storage conditions are known to significantly affect the synthesis, transport, and/or degradation of volatile compounds. Although knowledge of the biosynthesis mechanisms of regulation or modulation is still limited, it is known that postharvest practices significantly affect aroma [95]. Previous studies have suggested that *Aloe vera*-based coatings reduce aroma volatile biosynthesis in the fruit pulp of mangoes [13]. The researchers noted that this influence was characterized by the suppression of respiration.

4.7. Ascorbic Acid (AsA)

Ascorbic acid (Vitamin C) is a natural water-soluble vitamin and antioxidant which has the potential to fight bacterial infections, maintain skin, bone and teeth health, and protect body cells from damage. It is known as an antioxidant and a toxicide. Previous studies have reported that during postharvest storage, ascorbic acid is lost due to the activities of phenoloxidase [96]. Studies have also suggested that lowering the oxygen content around fresh produce during storage reduces the activities of phenoloxidase, and thus the loss of ascorbic acid [97]. *Aloe vera* gel coatings were found to be effective in reducing the loss of ascorbic acid content in pineapple fruits during storage [73]. Wounding of fruits is known to speed up the loss of nutritional properties (i.e. ascorbic acid) of minimally processed fruits by initiating enzymatic browning, which involves the mixing of PPO with phenolic compounds. Ascorbic acid as an antioxidant is mainly involved in oxidative reduction reactions in fresh-cut fruits, and is converted to dehydroascorbic acid [98]. An increase in storage duration, high temperatures, low relative humidity and physical damage have been reported to enhance loss of AsA [99]. Edible coatings of *Aloe vera* gel are reported to prevent the loss of AsA, and retention of AsA in the fruit is important in judging the efficacy of coating materials [75]. The addition of calcium chloride (2%) and citric acid (1%) in *Aloe vera* gel has been reported to retain AsA more effectively in table grapes [74]. Similar positive effects for *Aloe vera* gel were reported for grapefruits [74], strawberries [83]) and fresh-cut oranges [81].

5. Conclusions: Current Use and Future Trends

Bio-materials derived from plants have a long history in the use of human health and as therapeutic agents, and also for the postharvest storage of fresh and processed fruits and vegetables. *Aloe vera* gel coatings are effective and safe alternative to postharvest chemical treatments. The biological effects and the mechanisms of action responsible for their effects have been extensively studied and reviewed in the last decades; however, the mechanisms remain unclear. Further research into the compounds responsible for antimicrobial activity is of the utmost important, and

will broaden the research field of *Aloe vera*. On the other hand, the effects of *Aloe vera* gel coatings on the enzymatic actions, such as CAT, SOD, POD and PPO, are highly important for the understanding of the preservative mechanisms [60].

Critical discussion of the existing literature has made it possible to find huge differences among the tested doses and preparation methods of *Aloe vera* gel coatings. These two factors are among the most important factors which determine the success of the preservative materials. The next factor is the product itself, and the selection of the correct dose and suitable preparation method are highly important for each different product. According to the existing literature, lower doses of *Aloe vera* ($\leq 25\%$) are effective preservatives for thin-shelled products such as grapes, nectarines, raspberries, tomatoes and sweet cherries [65,71,80,84,88]; moderate doses (25–50%) are effective for medium-shelled products such as peppers and mangoes [79,85]; and high doses ($\geq 50\%$) are required for thick-shelled fresh products such as pineapples, plums and pistachios [73,80,82]. The current literature also shows that the incorporation of *Aloe vera* with ascorbic acid, glycerol, gum tragacanth, rosehip oil, chitosan, oleic acid and cysteine [63,75,79,80,82,84,89] improves the preservative characteristics of coatings. Incorporation of *Aloe vera* with other edible materials has mainly been tested for minimally processed ready-to-eat fruits and vegetables. Some of these materials were reported to be applied for pH adjustment, and the recommended pH is between 4.0 and 5.6, depending on the characteristics of the fresh produce.

Apart from the dose, additives and pH, pasteurisation is very important for stabilisation of the gel. The pasteurisation temperature is highly important for optimum stabilisation and for obtaining good film thickness, water solubility, and swelling behaviour. The most used temperature and duration combinations for pasteurisation are 65 °C for 30 min [61]; 70 °C for 45 min [62,65,75,85]; 75 °C for 45 min [50]; 80 °C for 10 s [55]; 80 °C for 10 min [81]; and 90 °C for 30 min [77]. However, some of the previously published articles did not mention pasteurisation [83,89]. Among the tested temperature and duration combinations for pasteurisation, the most preferred and successful method was found to be 70 °C for 45 min. However, the effectiveness of the temperature and duration combination for pasteurisation has not been adequately explored, and further studies should be carried out on this topic. The dipping duration of the fresh produce into the *Aloe vera* gel is another critical point for improving the efficiency. The literature review showed that there is not a uniform duration for fresh produce, even for the same products in different studies. The most tested dipping durations are 1 min [62,81]; 2 min [87,89]; 5 min [74,75,82,83,86]; 10 min [49,55,88,93]; and 25 min [85]. The “lack of uniformity” in the preparation and application methods would not be a problem if they had been tested to compare the effectiveness of the different methodologies. However, the literature search showed no studies for such a comparison. Thus, further studies are needed to test the effects of dipping durations, for determination of the optimum duration.

To conclude the criticism, there is not a clear methodology (or uniformity) for the preparation and application methods of *Aloe vera* gel, and further studies are required to compare the effectiveness of different methods on the biochemical and physical effectiveness of gel coatings. Discussion of the existing literatures also made it possible to conclude that the incorporation of *Aloe vera* gel with some other effective bio-degradable materials (chitosan, essential oils, propolis, plant extracts, etc.) is important, in order to increase the effectiveness and reduce the application doses of *Aloe vera*.

Author Contributions: conceptualization, İ.K. and C.W.; investigation, İ.K.; C.C. and C.W.; resources, İ.K. and J.C.; reviewing, İ.K. and C.W.; discussion, İ.K. and C.W.; writing—original draft preparation, İ.K. and C.W.; writing—review and editing, İ.K. and C.W.; project administration and funding acquisition, J.C.

Funding: This research was funded by Natural Science Foundation in Jiangxi Province (20181BCB24005).

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Cordenunsi, B.R.; Nascimento, J.R.O.; Lajolo, F.M. Physicochemical changes related to quality of five strawberry fruit cultivars during cool-storage. *Food Chem.* **2003**, *83*, 167-173. [https://doi.org/10.1016/S0308-8146\(03\)00059-1](https://doi.org/10.1016/S0308-8146(03)00059-1)
2. Sharma, R.R.; Singh, D.; Singh, R. Biological control of postharvest diseases of fruits and vegetables by microbial antagonists: a review. *Biol. Cont.* **2009**, *50*(3), 205-221. <https://doi.org/10.1016/j.biocontrol.2009.05.001>
3. Coulibaly, O.; Nouhoheiflin, T.; Aitchedji, C.C.; Cherry, A.J.; Adegbola, P. Consumers' Perceptions and Willingness to Pay for Organically Grown Vegetables. *Int. J. Veg. Sci.* **2011**, *17*(4), 349-362. <https://doi.org/10.1080/19315260.2011.563276>
4. Lin, D.; Zhao, Y. Innovations in the development and application of edible coatings for fresh and minimally processed fruits and vegetables. *Comprehensive Reviews in Food Science and Food Safety-CRFSFS* **2007**, *6* (3), 60-75. 2007. <https://doi.org/10.1111/j.1541-4337.2007.00018.x>
5. Silvestre, C.; Duraccio, D.; Cimmino, S. Food packaging based on polymer nanomaterials. *Progress in Polymer Science* **2011**, *36* (12), 1766-1782. <https://doi.org/10.1016/j.progpolymsci.2011.02.003>
6. Sharif, R.; Mujtaba, M.; Ur Rahman, M.; Shalmani, A.; Ahmad, H.; Anwar, T.; Tianchan, D.; Wang, X. The Multifunctional Role of Chitosan in Horticultural Crops; A Review. *Molecules* **2015**, *23*(4), 872. <https://doi.org/10.3390/molecules23040872>
7. Adiletta, G.; Pasquariello, M.S.; Zampella, L.; Mastrobuoni, F.; Scortichini, M.; Petriccione, M. Chitosan coating: A postharvest treatment to delay oxidative stress in loquat fruits during cold storage. *Agronomy* **2018**, *8*(4), 54. <https://doi.org/10.3390/agronomy8040054>
8. Prakash, B.; Kedia, A.; Mishra, P.K.; Dubey, N.K. Plant essential oils as food preservatives to control moulds, mycotoxin contamination and oxidative deterioration of agri-food commodities—Potentials and challenges. *Food Cont* **2015**, *47*, 381-391. <https://doi.org/10.1016/j.foodcont.2014.07.023>
9. Kahramanoğlu, İ. Effects of lemongrass oil application and modified atmosphere packaging on the postharvest life and quality of strawberry fruits. *Sci Hort* **2019**, *256*, <https://doi.org/10.1016/j.scienta.2019.05.054>
10. Kahramanoğlu, İ.; Aktaş, M.; Gündüz, Ş. Effects of fludioxonil, propolis and black seed oil application on the postharvest quality of “Wonderful” pomegranate. *PLoS ONE* **2018**, *13*(5): e0198411. <https://doi.org/10.1371/journal.pone.0198411>
11. Obagwu, J.; Korsten, L. Control of citrus green and blue molds with garlic extracts. *Eur J Plant Pathol* **2003**, *109*(3), 221-225. <https://doi.org/10.1023/A:1022839921289>
12. Gatto, M.A.; Sergio, L.; Ippolito, A.; Di Venere, D. Phenolic extracts from wild edible plants to control postharvest diseases of sweet cherry fruit. *Postharvest Biol Technol* **2016**, *120*, 80-187. <https://doi.org/10.1016/j.postharvbio.2016.06.010>
13. Dang, K.T.H.; Singh, Z.; Swinny, E.E. Edible coatings influence fruit ripening, quality, and aroma biosynthesis in mango fruit. *J Agric Food Chem* **2008**, *56*(4):1361-1370. <https://doi.org/10.1021/jf072208a>
14. Saucedo-Pompa, S.; Rojas-Molina, R.; Aguilera-Carbó, A.F.; Saenz-Galindo, A.; de La, G.H.; Jasso-Cantú, D.; Aguilar, C.N. Edible film based on candelilla wax to improve the shelf life and quality of avocado. *Food Res Int* **2009**, *42*(4), 511-515. <https://doi.org/10.1016/j.foodres.2009.02.017>
15. Troyo, R.D.; Acedo, A.L. Effects of calcium ascorbate and calcium lactate on quality of fresh-cut pineapple (*Ananas comosus*). *Int J Agric For Life Sci* **2019**, *3*(1), 143-150.
16. Misir, J.; Brishti, F.H.; Hoque, M.M. Aloe vera gel as a novel edible coating for fresh fruits: A Review. *Am J Food Sci Technol* **2014**, *2*(3), 93-97. <https://doi.org/10.12691/ajfst-2-3-3>
17. Kahramanoğlu, İ. Introductory chapter: postharvest physiology and technology of horticultural crops. In: *Postharvest Handling*, Ed.: İ. Kahramanoğlu, InTech Open, London, **2017**, pp. 1-5, <https://doi.org/10.5772/intechopen.69466>
18. Singh, D.; Sharma, R.R. Postharvest diseases of fruits and vegetables and their management. In: *Postharvest Disinfection of Fruits and Vegetables*. Elsevier, **2018**. p. 1-52.
19. Dagne, E.; Bisrat, D.; Viljoen, A., & Van Wyk, B. E. Chemistry of Aloe Species. *Current Organic Chemistry* **2000**, *4*, 1055-1078.
20. Mchugh, T.H.; Senesi, E. Apple wraps: A novel method to improve the quality and extend the shelf life of fresh-cut apples. *Journal of Food Science* **2000**, *65*(3), 480-485. <https://doi.org/10.1111/j.1365-2621.2000.tb16032.x>

21. Michailides, T.J.; Manganaris, G.A. Harvesting and handling effects on postharvest decay. *Stewart Postharvest Rev* **2009**, *5*(2), 1-7. <https://doi.org/10.2212/spr.2009.2.3>
22. Lv, L., Yang, Q. Y., Zhao, Y., Yao, C. S., Sun, Y., Yang, E. J. Fang, W. S. BACE1 (β -secretase) inhibitory chromone glycosides from Aloe vera and Aloe nobilis. *Planta medica* **2008**, *74*, 540-545.
23. Lee, S.; Do, S.-G.; Kim, S.Y.; Kim, J.; Jin, Y.; Lee, C.H. Mass Spectrometry-Based Metabolite Profiling and Antioxidant Activity of Aloe vera (Aloe barbadensis Miller) in Different Growth Stages. *Journal of Agricultural and Food Chemistry* **2012**, *60*, 11222-11228, <http://dx.doi.org/10.1021/jf3026309>
24. Park, M.K., Park, J. H., Shin, Y. G., Kim, W. Y., Lee, J. H., & Kim, K. H. Neoaloesin A_ A New C-Glucosyl Chromone from Aloe barbadensis. *Planta medica* **1996**, *64*, 363-365.
25. Okamura, N.; Hine, N.; Tateyama, Y.; Nakazawa, M.; Fujioka, T.; Mihashi, K.; Yagi, A. Three chromones of Aloe vera leaves. *Phytochemistry* **1997**, *45*, 1511-1513, [http://dx.doi.org/10.1016/s0031-9422\(97\)00148-9](http://dx.doi.org/10.1016/s0031-9422(97)00148-9)
26. Okamura, N.; Hine, N.; Harada, S.; Fujioka, T.; Mihashi, K.; Yagi, A. Three chromone components from Aloe vera leaves. *Phytochemistry* **1996**, *43*, 495-498, [http://dx.doi.org/10.1016/0031-9422\(96\)00306-8](http://dx.doi.org/10.1016/0031-9422(96)00306-8)
27. Okamura, N.; Hine, N.; Tateyama, Y.; Nakazawa, M.; Fujioka, T.; Mihashi, K.; Yagi, A. Five chromones from Aloe vera leaves. *Phytochemistry* **1998**, *49*, 219-223, [http://dx.doi.org/10.1016/s0031-9422\(97\)01071-6](http://dx.doi.org/10.1016/s0031-9422(97)01071-6)
28. Okamura, N.; Hine, N.; Harada, S.; Fujioka, T.; Mihashi, K.; Nishi, M.; Miyahara, K.; Yagi, A. Diastereomeric C-glucosylanthrones of Aloe vera leaves. *Phytochemistry* **1997**, *45*, 1519-1522, [http://dx.doi.org/10.1016/s0031-9422\(97\)00149-0](http://dx.doi.org/10.1016/s0031-9422(97)00149-0)
29. Zhong, J.S.; Huang, Y.Y.; Zhang, T.H.; Liu, Y.P.; Ding, W.J.; Wu, X.F.; Xie, Z.Y.; Luo, H.B.; Wan, J.Z. Natural phosphodiesterase-4 inhibitors from the leaf skin of Aloe barbadensis Miller. *Fitoterapia* **2015**, *100*, 68-74, <http://dx.doi.org/10.1016/j.fitote.2014.11.018>
30. Hutter, J.A., Salman, M., Stavinoha, W. B., Satsangi, N., Williams, R. F., Streeper, R. T., & Weintraub, S. T. Antiinflammatory C-Glucosyl Chromone from Aloe barbadensis. *Journal of natural products* **1996**, *59*, 541-543.
31. Kim, J.H.; Yoon, J.-Y.; Yang, S.Y.; Choi, S.-K.; Kwon, S.J.; Cho, I.S.; Jeong, M.H.; Kim, Y.H.; Choi, G.S. Tyrosinase inhibitory components from Aloe vera and their antiviral activity. *Journal of Enzyme Inhibition and Medicinal Chemistry* **2017**, *32*, 78-83, <http://dx.doi.org/10.1080/14756366.2016.1235568>
32. Rehman, N.U.; Al-Riyami, S.A.; Hussain, H.; Ali, A.; Khan, A.L.; Al-Harrasi, A. Secondary metabolites from the resins of Aloe vera and Commiphora mukul mitigate lipid peroxidation. *Acta Pharmaceutica* **2019**, *69*, 433-441, <http://dx.doi.org/10.2478/acph-2019-0027>
33. Rehman, N.U.; Hussain, H.; Khiat, M.; Khan, H.Y.; Abbas, G.; Green, I.R.; Al-Harrasi, A. Bioactive chemical constituents from the resin of Aloe vera. *Zeitschrift Fur Naturforschung Section B-a Journal of Chemical Sciences* **2017**, *72*, 955-958, <http://dx.doi.org/10.1515/znbs-2017-0117>
34. Rehman, N.U.; Hussain, H.; Khiat, M.; Al-Riyami, S.A.; Csuk, R.; Khan, H.Y.; Abbas, G.; Al-Thani, G.S.; Green, I.R.; Al-Harrasi, A. Aloeverasides A and B: Two Bioactive C-Glucosyl Chromones from Aloe vera Resin. *Helvetica Chimica Acta* **2016**, *99*, 687-690, <http://dx.doi.org/10.1002/hlca.201600126>
35. Yang, Q.-Y.; Yao, C.-S.; Fang, W.-S. A new triglucosylated naphthalene glycoside from Aloe vera L. *Fitoterapia* **2010**, *81*, 59-62, <http://dx.doi.org/10.1016/j.fitote.2009.07.006>
36. Choi, J.S.; Lee, S.K.; Sung, C.K.; Jung, J.H. Phytochemical study on Aloe vera. *Archives of Pharmacal Research* **1996**, *19*, 163-167, <http://dx.doi.org/10.1007/bf02976854>
37. Tan, Z.-j.; Li, F.-f.; Xu, X.-l. Extraction and purification of anthraquinones derivatives from Aloe vera L. using alcohol/salt aqueous two-phase system. *Bioprocess and Biosystems Engineering* **2013**, *36*, 1105-1113, <http://dx.doi.org/10.1007/s00449-012-0864-4>
38. Borges-Argaez, R.; Chan-Balan, R.; Cetina-Montejo, L.; Ayora-Talavera, G.; Sansores-Peraza, P.; Gomez-Carballo, J.; Caceres-Farfan, M. In vitro evaluation of anthraquinones from Aloe vera (Aloe barbadensis Miller) roots and several derivatives against strains of influenza virus. *Industrial Crops and Products* **2019**, *132*, 468-475, <http://dx.doi.org/10.1016/j.indcrop.2019.02.056>
39. Epifano, F.; Fiorito, S.; Locatelli, M.; Taddeo, V.A.; Genovese, S. Screening for novel plant sources of prenyloxanthraquinones: Senna alexandrina Mill. and Aloe vera (L.) Burm. F. *Natural Product Research* **2015**, *29*, 180-184, <http://dx.doi.org/10.1080/14786419.2014.971792>
40. Lopez, A.; Suarez de Tangil, M.; Vega-Orellana, O.; Ramirez, A.S.; Rico, M. Phenolic Constituents, Antioxidant and Preliminary Antimycoplasmic Activities of Leaf Skin and Flowers of Aloe vera (L.) Burm. f. (syn. A. barbadensis Mill.) from the Canary Islands (Spain). *Molecules* **2013**, *18*, 4942-4954, <http://dx.doi.org/10.3390/molecules18054942>

41. Keyhanian, S.; Stahl-Biskup, E. Phenolic constituents in dried flowers of Aloe vera (Aloe barbadensis) and their in vitro antioxidative capacity. *Planta Medica* **2007**, *73*, 599-602, <http://dx.doi.org/10.1055/s-2007-967202>
42. Lawrence, R.; Tripathi, P.; Jeyakumar, E. Isolation, purification and evaluation of antibacterial agents from Aloe vera. *Brazilian Journal of Microbiology* **2009**, *40*, 906-915, <http://dx.doi.org/10.1590/s1517-83822009000400023>
43. Zhang, Y.; Bao, Z.; Ye, X.; Xie, Z.; He, K.; Mergens, B.; Li, W.; Yacilla, M.; Zheng, Q. Chemical Investigation of Major Constituents in Aloe vera Leaves and Several Commercial Aloe Juice Powders. *Journal of Aoac International* **2018**, *101*, 1741-1751, <http://dx.doi.org/10.5740/jaoacint.18-0122>
44. Speranza, G.; Dadá, G.; Lunazzi, L.; Gramatica, P., & Manitto, P. Aloenin B, a New Diglucosylated 6-Phenyl-2-pyrone from Kenya Aloe. *Journal of Natural Products* **1986**, *49*, 800-805.
45. Afzal, M.; Ali, M.; Hassan, R.A.H.; Sweedan, N.; Dhami, M.S.I. Identification of Some Prostanoids in Aloe vera Extracts. *Planta Medica* **1991**, *57*, 38-40, <http://dx.doi.org/10.1055/s-2006-960012>
46. Tanaka, M.; Misawa, E.; Ito, Y.; Habara, N.; Nomaguchi, K.; Yamada, M.; Toida, T.; Hayasawa, H.; Takase, M.; Inagaki, M., et al. Identification of five phytosterols from aloe vera gel as anti-diabetic compounds. *Biological & Pharmaceutical Bulletin* **2006**, *29*, 1418-1422, <http://dx.doi.org/10.1248/bpb.29.1418>
47. Nabigol, A.; Asghari, A. Antifungal activity of Aloe vera gel on quality of minimally processed pomegranate arils. *International Journal of Agronomy and Plant Production* **2013**, *4*(4), 833-838.
48. Kator, L.; Hosea, Z.Y.; Ene, O.P. The Efficacy of Aloe-vera coating on postharvest shelf life and quality tomato fruits during storage. *Asian Research Journal of Agriculture* **2018**, *8*(4), 1-9. <https://doi.org/10.9734/ARJA/2018/41540>
49. Benitez, S.; Achaerandio, I.; Pujol, M.; Sepulcre, F. Aloe vera as an alternative to traditional edible coatings used in freshcut fruits: A case of study with kiwifruit slices. *LWT - Food Science and Technology* **2015**, *61*, 184-193. <http://dx.doi.org/10.1016/j.lwt.2014.11.036>
50. Krishnan, S.A.; Ullas, A.; Sagarika, N.; Oommen, T.E.; Sunaila, K. Development of Aloevera Based Edible Coating. *Int J Pure App Biosci* **2017**, *5*(5), 796-801. <http://dx.doi.org/10.18782/2320-7051.3026>
51. Sitara, U.; Hassan, N.; Naseem, J. Antifungal activity of Aloe vera gel against plant pathogenic fungi. *Pak J Bot* **2011**, *43*(4), 2231-2233.
52. Rosca-Casian, O.; Parvu, M.; Vlase, L.; Tamas, M. Antifungal activity of Aloe vera leaves. *Fitoterapia* **2007**, *78*, 219-222. <https://doi.org/10.1016/j.fitote.2006.11.008>
53. Ortega-Toro, R.; Collazo-Bigliardi, S.; Roselló, J.; Santamarina, P.; Chiralt, A. Antifungal starch-based edible films containing Aloe vera. *Food Hydrocolloids* **2017**, *72*, 1-10. <https://doi.org/10.1016/j.foodhyd.2017.05.023>
54. Serrano, M.; Miguel, J.; Guillen, F.; Castillo, S.; Martinez-Romero, D.; Valero, D. Use of Aloe vera gel coating preserves the functional properties of table grapes. *J Agri Food Chem* **2006**, *54*, 3882-3886. <https://doi.org/10.1021/jf060168p>
55. Navarro, D.; Díaz-Mula, H.M.; Guillén, F.; Zapata, P.J.; Castillo, S.; Serrano, M.; Valero, D.; Martínez-Romero, D. Reduction of nectarine decay caused by *Rhizopus stolonifer*, *Botrytis cinerea* and *Penicillium digitatum* with Aloe vera gel alone or with the addition of thymol. *Int J Food Microbiol* **2011**, *151*, 241-246. <https://doi.org/10.1016/j.ijfoodmicro.2011.09.009>
56. Garcia, M.A.; Ventosa, M.; Diaz, R.; Falco, S.; Casariego, A. Effects of Aloe vera coating on postharvest quality of tomato. *Fruits* **2014**, *69*, 117-126. <https://doi.org/10.1051/fruits/2014001>
57. Chrysargyris, A.; Nikou, A.; Tzortzakis, N. Effectiveness of Aloe vera gel coating for maintaining tomato fruit quality. *New Zealand J Crop Hort Sci* **2016**, <http://dx.doi.org/10.1080/01140671.2016.1181661>
58. Saks, Y.; Barkai-Golan, R. Aloe vera gel activity against plant pathogenic fungi. *Postharvest Biol Technol.* **1995**, *6*, 159-165. [https://doi.org/10.1016/0925-5214\(94\)00051-S](https://doi.org/10.1016/0925-5214(94)00051-S)
59. Jhalegar, J.; Sharma, R.R.; Singh, D. Antifungal efficacy of botanicals against major postharvest pathogens of Kinnow mandarin and their use to maintain postharvest quality. *Fruits* **2014**, *69*, 223-237. <https://doi.org/10.1051/fruits/2014012>
60. Hassanpour, H. Effect of Aloe vera gel coating on antioxidant capacity, antioxidant enzyme activities and decay in raspberry fruit. *LWT - Food Science and Technology* **2015**, *60*(1), 495-501. <https://doi.org/10.1016/j.lwt.2014.07.049>
61. Vieira, J.M.; Flores-López, M.L.; de Rodríguez, D.J.; Sousa, M.C.; Vicente, A.A.; Martins, J.T. Effect of chitosan-Aloe vera coating on postharvest quality of blueberry (*Vaccinium corymbosum*) fruit. *Postharvest Biol Technol* **2016**, *116*, 88-97. <http://dx.doi.org/10.1016/j.postharvbio.2016.01.011>

62. Nasrin, T.A.A.; Rahman, M.A.; Hossain, M.A.; Islam, M.N.; Arfin, M.S. Postharvest quality response of strawberries with aloe vera coating during refrigerated storage. *The J Hort Sci Biotechnol* **2017**, *https://doi.org/10.1080/14620316.2017.1324326*
63. Martínez-Romero, D.; Castillo, S.; Guillén, F.; Díaz-Mula, H.M.; Zapata, P.J.; Valero, D.; Serranoba, M. *Aloe vera* gel coating maintains quality and safety of ready-to-eat pomegranate arils. *Postharvest Biol Technol* **2013**, *86*, 107-112. <http://dx.doi.org/10.1016/j.postharvbio.2013.06.022>
64. Castillo, S.; Navarro, D.; Zapata, D.J.; Guillén, F.; Valero, D.; Martínez-Romero, D.; Serrano, M. Using *Aloe vera* as a preharvest treatment to maintain postharvest organic table grape quality. In: Proc. XXVIIIth IHC – IS on Organic Horticulture: Productivity and Sustainability, Eds.: I. Mourão and U. Aksoy, Acta Hort. 933, ISHS **2012**; pp. 621-626. <https://doi.org/10.17660/ActaHortic.2012.933.81>
65. Chauhan, S.; Gupta, K.C.; Agrawal, M. Application of Biodegradable *Aloe vera* gel to control post harvest decay and longer the shelf life of Grapes. *Int J Curr Microbiol App Sci*, **2014**, *3*(3), 632-642.
66. Ferro, V.A.; Bradbury, F.; Cameron, P.; Shakir, E.; Rahman, S.R.; Stimson, W.H. In vitro susceptibilities of *Shigella flexneri* and *Streptococcus pyogenes* to inner gel of *Aloe barbadensis* Miller. *Antimicrobial agent and Chemotherapy* **2003**, 1137-1139.
67. Wang, H.H.; Chung, J.G.; Ho, C.C.; Wu, L.T.; Chang, S.H. Aloe-emodin effects on arylamine N-Acetyltransferase activity in the bacterium *heliobacter pylori*. *Planta Medica* **1998**, *64*, 176-178
68. Cellini, L.; Di Bartolomeo, S.; Di Campli, E.; Genovese, S.; Locatelli, M.; Di Giulio, M. In vitro activity of *Aloe vera* inner gel against *Helicobacter pylori* strains. *Letters in Applied Microbiology* **2014**, *59*, 43-48
69. Zandi, K.; Rastian, Z. Antiviral activity of *Aloe vera* against herpes simplex virus type 2: An in vitro study. *African Journal of Biotechnology* **2007**, *6*, 1770-1773.
70. Li, S. W.; Yang, T.C.; Lai, C.C.; Huang, S.H.; Liao, J.M.; Wan, L. et al. Antiviral activity of aloe-emodin against influenza A virus via galectin-3 up-regulation. *European Journal of Pharmacology* **2014**, *738*, 125-132
71. Ahmed, M.J.; Singh, Z.; Khan, A.S. Postharvest *Aloe vera* gel-coating modulates fruit ripening and quality of 'Arctic Snow' nectarine kept in ambient and cold storage. *Int J Food Sci Technol* **2009**, *44*, 1024-1033. <https://doi.org/10.1111/j.1365-2621.2008.01873.x>
72. Achipiz, S.M.; Castillo, A.E.; Mosquera, S.A.; Hoyos, J.L.; Navia, D.P. Efecto de Recubrimiento a Base de Almidón Sobre la maduración de la Guayaba (*Psidium guajava*). *Biotechnol en el Sect Agropecu y Agroind* **2013**, *2*(2), 92-101.
73. Adetunji, C.O.; Fawole, O.B.; Arowora, K.A.; Nwaubani, S.I.; Ajayi, E.S.; Oloke, J.K.; Majolagbe, O.M.; Ogundele, B.A.; Aina, J.A.; Adetunji, J.B. Effects of edible coatings from *Aloe vera* gel on quality and postharvest physiology of *Ananas comosus* (L.) fruit during ambient storage. *Global Journal of Science Frontier Research Bio-Tech & Genetics* **2012**, *12* (5), 39-43
74. Shahkoomahally, S.; Ramezani, A. Effect of Natural *Aloe Vera* Gel Coating Combined with Calcium Chloride and Citric Acid Treatments on Grape (*Vitis vinifera* L. Cv. Askari) Quality during Storage. *American Journal of Food Science and Technology* **2014**, *2* (1), 1-5. <https://doi.org/10.12691/ajfst-2-1-1>
75. Sharmin, M.R.; Islam, M.N. and Alim, M.A. Shelf-life enhancement of papaya with aloe vera gel coating at ambient temperature. *J Bangladesh Agril Univ* **2015**, *13*(1), 131-136.
76. Khoshgozaran-Abrasa, S.; Azizia, M.H.; Hamidya, Z.; Bagheripoor-Fallah, N. Mechanical, physicochemical and color properties of chitosan based-films as a function of *Aloe vera* gel incorporation. *Carbohydrate Polymers* **2012**, *87*, 2058-2062. <https://doi.org/10.1016/j.carbpol.2011.10.020>
77. Gutierrez, T.J.; Alvarez, K. Physico-chemical properties and *in vitro* digestibility of edible films made from plantain flour with added *Aloe vera* gel. *Journal of Functional Foods* **2016**, *26*, 750-762. <http://dx.doi.org/10.1016/j.jff.2016.08.054>
78. Gutiérrez, T.J.; González, G. Effect of cross-linking with *Aloe vera* gel on surface and physicochemical properties of edible films made from plantain flour. *Food Biophysics* **2016**, <https://doi.org/10.1007/s11483-016-9458-z>
79. Mohebbi, M.; Hasanpour, N.; Ansarifard, E.; Amiryousefi, M.R. Physicochemical properties of bell pepper and kinetics of its color change influenced by *Aloe vera* and gum tragacanth coatings during storage at different temperatures. *J Food Process Preserv* **2014**, *38*(2), 684-693. <https://doi.org/10.1111/jfpp.12018>
80. Martínez-Romero, D.; Zapata, P.J.; Guillén, F.; Paladines, D.; Castillo, S.; Valero, D.; Serrano, M. The addition of rosehip oil to *Aloe* gels improves their properties as postharvest coatings for maintaining quality in plum. *Food Chem* **2017**, *217*, 585-592. <http://dx.doi.org/10.1016/j.foodchem.2016.09.035>

81. Radi, M.; Firouzi, E.; Akhavan, H.; Amiri, S. Effect of gelatin-based edible coatings incorporated with *Aloe vera* and black and green tea extracts on the shelf life of fresh-cut oranges. *Hindawi Journal of Food Quality* **2017**, ID 9764650, 10 p. <https://doi.org/10.1155/2017/9764650>
82. Vanaei, M.; Sedaghat, N.; Abbaspour, H.; Kaviani, M.; Azarbad, H.R. Novel edible coating based on *Aloe Vera* gel to maintain pistachio quality. *International Journal of Scientific Engineering and Technology* **2018**, 3 (8), 1016-1019.
83. Sogvar, O.B.; Saba, M.K.; Emamifar, A. Aloe vera and ascorbic acid coatings maintain postharvest quality and reduce microbial load of strawberry fruit. *Postharvest Biol Technol* **2016**, 114, 29-35. <http://dx.doi.org/10.1016/j.postharvbio.2015.11.019>
84. Athmaselvi, K.A.; Sumitha, P.; Revathy, B. Development of *Aloe vera* based edible coating for tomato. *Int Agrophys* **2013**, 27, 369-375. <https://doi.org/10.2478/intag-2013-0006>
85. Sophia, O.; Robert, G.M.; Ngwela, W.J. Effect of Aloe vera gel coating on postharvest quality and shelf life of mango (*Mangifera indica* L.) fruits var. 'Ngowe'. *J Hort For* **2015**, 7(1), 1-7. <https://doi.org/10.5897/JHF2014.0370>
86. Hazrati, S.; Kashkooli, A.B.; Habibzadeh, F.; Tahmasebi-Sarvestani, Z.; Sadeghi, A.R. Evaluation of *Aloe vera* gel as an alternative edible coating for peach fruits during cold storage period. *Gesunde Pflanzen* **2017**, 69, 131-137. <https://doi.org/10.1007/s10343-017-0397-5>
87. Ravanfar, R.; Niakousari, M.; Maftoonazad, N. Postharvest sour cherry quality and safety maintenance by exposure to Hot- water or treatment with fresh *Aloe vera* gel. *J Food Sci Technol* **2012**, <https://doi.org/10.1007/s13197-012-0767-z>
88. Martinez-Romero, D.; Alburquerque, N.; Valverde, J.M.; Guillen, F.; Castillo, S.; Valero, D.; Serrano, M. Postharvest sweet cherry quality and safety maintenance by *Aloe vera* treatment: A new edible coating. *Postharvest Biol Technol* **2006**, 39, 93-100. <https://doi.org/10.1016/j.postharvbio.2005.09.006>
89. Song, H.Y.; Jo, W.S.; Song, N.B.; Min, S.C.; Song, K.B. Quality change of apple slices coated with *Aloe vera* gel during storage. *J Food Sci* **2013**, 78(6), 817-822. <https://doi.org/10.1111/1750-3841.12141>
90. Reid, M.S. Ethylene in plant growth, development, and senescence. In: *Plant Hormones*, Ed.: P.J. Davis, Dordrecht: Springer, **1995**. https://doi.org/10.1007/978-94-011-0473-9_23
91. Pierik, R.; Tholen, D.; Poorter, H.; Visser, E.J.; Voesenek, L.A. The Janus face of ethylene: growth inhibition and stimulation. *Trends Plant Sci* **2006**, 11(4), 176-83. <https://doi.org/10.1016/j.tplants.2006.02.006>
92. Nazar, R.; Khan, M.I.; Iqbal, N.; Masood, A.; Khan, N.A. Involvement of ethylene in reversal of salt-inhibited photosynthesis by sulfur in mustard. *Physiol Plant* **2014**, 152(2), 331-344. <https://doi.org/10.1111/pp1.12173>
93. Guillén, F.; Díaz-Mula, H.M.; Zapata, P.J.; Valero, D.; Serrano, M.; Castillo, S.; Martínez-Romero, D. Aloe arborescens and Aloe vera gels as coatings in delaying postharvest ripening in peach and plum fruit. *Postharvest Biology and Technology* **2013**, 83, 54-57. <http://dx.doi.org/10.1016/j.postharvbio.2013.03.011>
94. Supapvanich, S.; Mitsrang, P.; Srinorkham, P.; Boonyariththongchai, P.; Wongs-Aree, C. Effects of fresh Aloe vera gel coating on browning alleviation of fresh cut wax apple (*Syzygium samarangense*) fruit cv. Taaptimjaan. *J Food Sci Technol* **2016**, 53(6), 2844-2850. <https://doi.org/10.1007/s13197-016-2262-4>
95. Defilippi, B.G.; Manríquez, D.; Luengwilai, K.; González-Agüero, M. Chapter 1 Aroma Volatiles: Biosynthesis and Mechanisms of Modulation During Fruit Ripening. *Advances in Botanical Research* **2009**, 50, 1-37. [https://doi.org/10.1016/S0065-2296\(08\)00801-X](https://doi.org/10.1016/S0065-2296(08)00801-X)
96. Salunkhe, D.K.; Boun, H.R.; Rddy, N.R. Storage Processing and Nutritional Quality of Fruits and Vegetables, vol. 1. Fresh Fruits and Vegetables. Boston: CRC Press Inc, **1991**.
97. Weichmann, J. Postharvest Physiology of Vegetables. New York: Marcel Dekker, **1985**.
98. Deutsch, J.C. Dehydroascorbic acid: Review. *J. Chromatogr A* **2000**, 881, 299-307. [https://doi.org/10.1016/S0021-9673\(00\)00166-7](https://doi.org/10.1016/S0021-9673(00)00166-7)
99. Lee, S.K.; Kader, A.A. Preharvest and postharvest factors influencing vitamin C content of horticultural crops. *Postharvest Biol Technol* **2000**, 20, 207-220. [https://doi.org/10.1016/S0925-5214\(00\)00133-2](https://doi.org/10.1016/S0925-5214(00)00133-2)