

Establishment and Characterization of Canine Mammary Gland Carcinoma Cell Lines with Vasculogenic Mimicry Ability *in vitro* and *in vivo*

Patrícia de Faria Lainetti^{1,‡}, Andressa Brandi^{1,‡}, Antonio Fernando Leis Filho¹, Maria Carolina Mangini Prado¹, Priscila Emiko Kobayashi¹, Renée Laufer-Amorim¹, Carlos Eduardo Fonseca-Alves^{1,2,*}

¹ School of Veterinary Medicine and Animal Science, Sao Paulo State University – UNESP, Botucatu, Brazil.

² Institute of Animal Health, Universidade Paulista-UNIP, Bauru, Brazil.

[‡] Both authors have equal contribution.

* **Correspondence:**

Carlos Eduardo Fonseca-Alves

carlos.e.alves@unesp.br

ABSTRACT

Mammary neoplasms affect a population of intact and elderly female dogs and 50% of these neoplasms are malignant. In order to study this disease, cell culture is as a promising preclinical model, creating the opportunity to deposit cell lines at a cell bank, allowing a great reproducibility of the assays and making the validation of the results more reliable. Another important aspect is the possibility to establish models for the better understanding of tumour characteristics, such as vasculogenic mimicry. VM is related to cancer cells able to generate vascular-like structures from epithelial cells, mimicking the vasculogenic process and it is closely related to patient's prognosis. Due the importance of cancer cell lines in preclinical models, this study aimed to establish and characterize ten cell lines from canine mammary gland tumours according to immunophenotype and tumorigenicity, and with its ability to form vasculogenic mimicry-like structures *in vitro* and *in vivo*. Cell cultures were evaluated for morphology, phenotype, vasculogenic mimicry and tumorigenicity abilities. All cell lines presented spindle shape morphology and expressed concomitant pan-cytokeratin and cytokeratin 8/18. Four cell lines had vasculogenic mimicry ability *in vitro* and two of them showed *in vivo* tumorigenic potential and forming VM in the xenotransplant tumour. Cell characterization of those lines will help to create a database for more knowledge of

mammary carcinomas in dogs, including studies of tumor behavior and new therapeutic targets.

Keywords: female dog; cell culture; mammary cancer; veterinary; oncology

BACKGROUND

Intact and old female dogs are frequently affected by mammary gland tumours and more than 50% of the cases are malignant [1, 2]. In women, breast cancer is the main cause of mortality and the most common cancer type diagnosed [3]. The occurrence of neoplasms in dogs are spontaneous and similar to breast cancer in women, as histological appearance, molecular targets and biological behaviour. For these reasons, canine mammary gland tumour is a natural model for human breast cancer [4].

Female dogs with mammary carcinoma do not have the same therapeutic outcome as women, since chemotherapy treatment is not as effective as in humans and doesn't increase the patient's survival time or influence local disease control [5, 6]. In dogs, the golden standard treatment is radical mastectomy and chemotherapy is based on the human breast cancer literature [7, 8]. Thus, even performing chemotherapy in female dogs affected by mammary gland tumours, there is no standardized protocol [6]. Besides that, we lack information regarding markers that can predict antitumor response, as occurs in human breast cancer treatment [9, 10]. In this scenario, cancer cell lines represent a great opportunity for the evaluation of antitumor response.

Cell lines are an alternative and an experimental model *in vitro* to human breast cancer and canine mammary gland tumours, allowing the investigation of carcinogenesis process as proliferation, apoptosis and migration [11]. Cell culture is considered an excellent preclinical model, essential for the identification and evaluation of drugs action mechanism, identification of genes involved in carcinogenesis, such as oncogenes and tumour suppressors, defining cell signalling pathways and how they contribute to tumour pathogenesis, discovering of new drugs and for the development process of antitumor drugs [12].

In canine and human patients affected by highly aggressive mammary neoplasms, neoplastic cells can be able to form vascular-like structures or channels, used to conduct

plasma, red cells and neoplastic cells during epithelial mesenchymal transition [13]. Thus, vasculogenic mimicry (VM) has been widely studied as a mechanism of tumour nutrition and metastasis [13, 14, 15, 16]. In canine mammary gland tumours, the phenomenon has been studied in inflammatory mammary carcinomas and other tumour subtypes. Thus, this study aimed to establish and characterize nine cell lines from canine mammary gland tumour according to immunophenotype, tumorigenicity and ability to form vascular-like structures *in vitro* and *in vivo*.

MATERIALS AND METHODS

Reagents

All reagents used had high purity and were purchased from companies GE Healthcare (Uppsala, Sweden), Sigma-Aldrich (São Paulo, Brazil), Merck SA (São Paulo, Brazil), otherwise, they were cited. In addition, cell culture medium used was *Mammary Epithelial Cell Growth Medium* (MEGM™, Lonza Inc., Allendale, NJ, USA), medium *Dulbecco's Modified Eagle Medium:F12* (DMEM/F12, Lonza Inc., Allendale, NJ, USA), fetal bovine serum (FBS, LGC Biotecnologia, Cotia, SP, Brazil), *Dulbecco's Phosphate Buffered Saline* (DPBS) (Sigma Aldrich, St. Louis, MO, USA), antibiotic/antimycotic solution (TermoFischer Scientific, Waltham, MA, USA) and trypsin (0.25%, GBICO TermoFischer Scientific, Waltham, MA, USA).

Tumour samples and cell isolation

Fifteen samples of mammary gland tumours were collected at the veterinary teaching hospital from Veterinary Medicine and Animal Science School, UNESP (between December 2016 and March 2017) for histopathological examination and the samples were expanded in cell culture. Sample of the tumour was formalin fixed and paraffin-embedded and part placed in media and brought immediately to cell culture. From the fifteen samples, twelve were primary tumours and three were respective metastasis from canine mammary carcinoma. Cell lines were obtained using enzymatic dissociation as previously described in the literature [17]. Briefly, tumour fragments with approximately 1 cm² were collected and dissociated with type IV collagenase (Sigma-Aldrich, St. Louis, MO, United States) for four hours at 37 °C, in a humidified

atmosphere containing 5% CO₂. After this process the material was separated using a 75 µm mesh filter, centrifuged and washed with Dulbecco's Phosphate-Buffered Saline DPBS (Sigma-Aldrich, St. Louis, MO, United States) to remove excess collagenase. After cell isolation process, they were counted in a Neubauer chamber and the cell viability was evaluated by the Trypan blue technique. Plating was performed at a concentration of 10⁴ cells / mL in 25 mL culture bottles with filter.

Selected samples, histological subtypes, and cell obtention technique are described in Supplementary Table 1. Identification of histological subtypes and tumour grade were based on the international classification of mammary gland tumours [18].

Molecular phenotype of primary tumours and respective metastasis

Immunohistochemistry (IHC) was performed with the tumour's samples from paraffin block, wherein 4 µm-thick sections were placed onto positively charged slides (StarFrost, Braunschweig, Germany) and then, deparaffinized. To antigen retrieval, citrate buffer (pH 6.0) was used in a pressure cooker (Pascal, Dako, Agilent Technologies, Santa Clara, CA, USA) and endogenous peroxidase was blocked with 8% hydrogen peroxide (Dinâmica Química Contemporânea, Indaiatuba, SP, Brazil) in methanol (Dinâmica Química Contemporânea, Indaiatuba, SP, Brazil) for 20 min. Nonspecific binding was blocked with 8% skim milk for 60 min, both at room temperature. The antibodies for HER2, ER α , PR, Ki-67, CK5/6 and EGFR and respective clones were diluted and samples remained incubated according to Nguyen et al. [19] (Supplementary Table 1). Antibody detection was achieved using a polymer system (EnVision, Agilent Technologies, Santa Clara, CA, USA). 3,3'-Diaminobenzidine (DAB) (EnVision, FLEX, High pH, Dako, Agilent Technologies, Santa Clara, CA, USA) was used as a chromogen and tissue counterstaining was performed with Harris Haematoxylin. Positive [19] and negative [20] controls were performed according to the previous literature.

ER α and PR were considered positive when $\geq 10\%$ of the nuclei were stained, CK8/18 and EGFR receptors when cytoplasmic staining was $\geq 10\%$ and Ki-67 when more than 33.3% of the cells had stained nuclei [19]. For the evaluation, a group of more than 500 cells were counted HER-2 staining was scored for labelling distribution (0: unlabelled

or <10% labelled tumour cells; 1+:> 10% of labelled tumour cells with incomplete membrane labelling; 2+: moderate to > 10% strong tumour cells; 3+: complete and strong membrane labelling of > 10% tumour cells). In this classification, 0 and 1+ were considered negative and, 2+ and 3+ positive.

The classification of different molecular phenotypes of canine mammary carcinoma is described as Luminal A, Luminal B, Triple negative or HER2 overexpressing. The Luminal A contemplates tumours HER2-, ER or PR + and Ki67 less than 33%. The Luminal B type is HER2-, ER or PR + and Ki67 more than 33%. The triple negative is a tumour negative for HER 2, ER and PR, and the tumour HER2 overexpressing is when it's positive for HER2 [20].

Cell Expansion

Cell culture was established from tumour fragments in MEGM™ medium containing 1% antibiotic/antimycotic solution and 10% FBS kept in a humid atmosphere containing 5% CO₂ at 37° C. When the cells reached 80% confluence, the medium was discarded and the bottles were washed with sterile DPBS pH 7.2 to eliminate residual FBS. For detachment of the cells from the bottle, trypsin 0.25% was added at 37 °C, followed by a 5 min incubation period in a 5% CO₂ humid atmosphere and 37 °C. Cells were cultured until 10th passage and used for cell phenotyping, karyotype, morphology and western blot.

Elimination of contaminating fibroblasts

To eliminate fibroblasts from primary cultures, selective cell trypsinization with cold trypsin (4°C) was performed according to the previous literature [21] at passage 5 (P5). Briefly, cells were washed with DPBS at 4 °C to avoid direct thermal shock by the cold trypsin. Subsequently, 2 mL of cold trypsin (4 °C) were used at room temperature (27 °C) for 2 min. Then, supernatant was inactivated with complete medium containing 10% FBS in a 1: 1 ratio and trypsin was collected and discharged. Then, the flasks were washed with DPBS buffer to remove residual trypsin. Subsequently, cells were washed with DPBS solution at room temperature twice, and then washed with DPBS buffer at 37 °C once. After this procedure, 800µl of trypsin at 37 °C were

added to the bottle and the cells were kept in a humid atmosphere at 37 °C in a humidified atmosphere containing 5% CO₂ for 5 min. After this period, the remaining cells were detached with manual mechanical impact and the trypsin was inactivated with complete medium (containing 10% FBS) in a 1:1 ratio. Then, cells were centrifuged for 5 min at 1200 rpm and resuspended in 5 ml of MEGM™ medium containing 10% FBS and 1% antibiotic and antifungal.

Cell Karyotype

Karyotype analysis was performed according to Moorhead et al. [22]. Initially the cells at P10 were cultured in DMEN/F12 medium, supplemented with FBS (10%) and phytohemagglutinin for 72 hours. After this period, the cells were evaluated under the inverted microscope and the mitotic spindle was interrupted by the addition of colchicine (16 µg / ml). Subsequent washes and centrifugations were performed at 4 °C to then fix the slide material and stain by the Wright-Giemsa staining method for karyotype assembly. After karyotype preparation and staining, 70 different images were captured for each cell culture (n=10) and at least 20 metaphases of each culture were analysed, according to the previously described by Gouveia et al. [25].

Cell morphology and doubling time

The morphology of each culture was evaluated at P10 to characterize the cell clone that grew in each culture. For this, 12 sterile well plates with sterile circular coverslips were used and placed in the centre of each well. After plate preparation, 500 µl of complete culture medium were added to each well and subsequently 10³ cells were pipetted into the middle of the well for 72 hours. After this period cell density in each coverslip was verified. When the coverslips presented confluence above 50% they were removed for cell morphology.

For the morphology technique the medium was removed, the coverslips washed with DPBS three times and then fixed with cold methanol absolute (4 °C) for 30 min in a refrigerator (8 °C). After methanol was removed, the cells were washed three times with PBS and immersed in 0.1% Triton-X solution for 10 min at room temperature for cell permeabilization. Subsequently, the permeabilizing solution was removed and the cells

washed three times with DPBS. After that, cells were stained with hematoxylin and eosin (HE).

Cells were also evaluated for cell doubling time. The cell medium was discarded and the bottles were washed with sterile DPBS pH 7.2 to eliminate residual FBS. For detachment of the cells from the bottle, trypsin was added at 37 °C, followed by a 5 min incubation period in a 5% CO₂ humid atmosphere and after this period, trypsin was inactivated with cell culture medium supplemented with 10% FBS and 1% antibiotic/antimycotic solution. Cells were centrifuged (450 g, 5 min) and after discarding the supernatant, resuspended in 1 mL DMEM/F12 medium with 10% FBS and 1% antibiotic/antimycotic solution. Cells were diluted in Trypan blue (Trypan blue solution, cod. T8154, Sigma-Aldrich, St. Louis, MO, USA) in a 1: 1 ratio and counted in a Neubauer chamber. For doubling time, the protocol described by Caceres et al. [23] was followed. Briefly, 1x10⁵ were plated in 25cm² flasks and maintained with DMEN/F12 medium supplemented with 10% FBS and 1% antibiotic/antimycotic solution in triplicate. Every 24 hours cells were trypsinized and counted. This procedure was performed for 5 consecutive days to evaluate the exponential growth curve. The final amount of cells for each culture was obtained by averaging three counts.

Cell Phenotype

The phenotype of each culture was evaluated at P10 to characterize the cell clone that grew in each culture. For this, 12 sterile well plates with sterile circular coverslips were used and placed in the centre of each well. After plate preparation, 500 µl of complete culture medium were added to each well and subsequently 10³ cells were pipetted into the middle of the well for 72 hours. After this period cell density in each coverslip was verified. When the coverslips presented confluence above 50% they were removed for immunofluorescence (IF).

For the IF technique the medium was removed, the coverslips washed with DPBS three times and then fixed with cold methanol absolute (4 °C) for 30 min in a refrigerator (8 °C). After methanol was removed, the cells were washed three times with PBS and

immersed in 0.1% Triton-X solution for 10 min at room temperature for cell permeabilization. Then, cells were blocked with commercial solution (Protein block, Dako, CA, USA) for 30 min at room temperature and primary antibodies were added to each well. We investigated pan-cytokeratin, cytokeratin 8/18 and vimentin expression. The information regarding the antibodies can be found in Supplementary Table 3. Cells were incubated with goat anti-mouse IgG secondary antibody (Alexa Fluor 647, Life Technologies, Corporation, Carlsbad, CA, EUA) and counterstained with DAPI (Sigma Aldrich, St. Louis, MO, EUA) at 1:10.000 dilution. As a negative reaction control, the primary antibodies tested during the procedure were omitted and replaced with TRIS buffer solution.

Tubular formation *in vitro*

Tubular formation *in vitro* it is an assay to identify vasculogenic mimicry ability in culture conditions. For tubular formation assay, the cells were cultured in three-dimensional (3D) conditions. After each cell culture achieved more than 80% of confluency, they were trypsinized and 50×10^3 viable cell were cultured with DMEM in a 24-well plate with 250 μ L of Matrigel (Matrigel® Growth Factor Reduced (GFR) Basement Membrane Matrix, *LDEV-Free, Corning, New York, NY, USA). The Matrigel was added in each well and air-dried for 30 min, at room temperature. The cells were incubated in a humidified atmosphere with 5% of CO₂ at 37°C. Then, the cells were evaluated in an inverted microscopy every hour to determine tubular formation. The experiment was performed in triplicate for each cell culture.

Tumour growth in immunodeficient mice

To evaluate cell culture tumorigenicity, a total of 12 nude mice (BALB/c nude, C.Cg-Foxn1nu line) were acquired from the Institute of Biomedical Sciences, University of São Paulo – USP and housed in individually ventilated caging. All procedures regarding feeding, humidity, temperature and light control were performed according to the literature [24]. For *in vivo* tumorigenicity evaluation, 1×10^6 cell were inoculated on

subcutaneous mammary gland region and the subjects were accompanied once a week, for at least 60 days. After tumour growth, the visits occurred daily to measure tumour volume. After achieving 3 cm², the subjects were submitted to humanely euthanasia and the material from the tumour was collected to confirm malignancy with histological evaluation. Then, pan-cytokeratin and vimentin immunohistochemistry was performed to confirm tumour phenotype. Immunohistochemistry analysis was performed as described above, using the antibodies mouse monoclonal anti-vimentin (Clone V3, Santa Cruz Biotechnology, Dallas, TX, USA) and mouse monoclonal anti-cytokeratin (Clone AE1/AE3, Santa Cruz Biotechnology, Dallas, TX, USA) at 1:300 dilution, overnight. The secondary antibodies, chromogen, counterstaining and negative controls were performed as described above. The epithelial component of a normal skin was used as positive control for pan-cytokeratin and the endodermis was used as positive control for vimentin.

RESULTS

Molecular phenotype of primary tumours and cell isolation

Of all the fifteen tumour samples, only ten were able to grow on cell culture were evaluated by IHC. Six were classified as HER-2 overexpressing (CM1, CM9, CM11, CM60, MM3 and MM4) and 3 as triple negative (CM5, CM61 and MM1).

Ten out 15 samples grew *in vitro* and the different tumour classification, graduation, acquisition technique, time exposed to collagenase and passage are described in table 1. The samples CM2, CM3, CM6, CM7, CM8, CM10 and CM12-15 did not show *in vitro* expansion. The cell cultures CM2 and CM3 were cultured using enzymatic dissociation overnight and had no cellular growth after 72 hours in culture condition. CM6 and CM7 culture cells were made using explants and for both cell lines, fungal contamination discontinued the cellular growth. Thus, cells submitted enzymatic dissociation with type IV collagenase showed the best *in vitro* expansion.

Contaminating Fibroblast Elimination

Cell cultures prior P5 showed mixed morphology (spindle cells, polygonal cells, cells growing in groups, and rounded cells) (Figure 1A). After selective trypsinization at 4 °C, cells became more homogeneous and showed slow growth with spindle morphology (Figure 1B) and cells did not show confluence above 50% after 30 days of cultivation and were discharged. Cells from trypsinization at 37 °C showed homogeneous morphology with some cultures presenting uniform spindle morphology (CM1, CM4 and MM1) or polygonal morphology (CM5, CM9 and CM11) (Figure 1C). Cells were grown to passage 10 (P10), showing approximately 90% confluence 48 hours after passage.

Cell Karyotype

Of the 70 images captured, the best ones for chromosome counting were selected (Supplementary Figure 2). It was possible to identify aneuploidies in different metaphases of different cell cultures. In cell cultures CM4 and CM11, hypoploidy of metaphase was identified, as well as hyperploid of cultures CM1, CM60 and MM3. The representation of chromosomal alterations in each culture can be seen in Supplementary Figure 2.

Cell morphology, phenotype and doubling time

In the morphological evaluation (Figure 3I and 3J), the primary and metastatic adenosquamous carcinoma tumour cells showed growth in monolayers, in fusiform pattern, basophilic nucleus and eosinophilic cytoplasm, with presence of mitosis. Grade III solid carcinoma cells showed multinucleated cells and colony formation (Figure 3K and 3L). All samples evaluated showed strong pan-cytokeratin (Figure 4A) and CK8/18 (Figure 4B) staining and were negative for p63 (Figure 4C). All cell cultures also showed vimentin positive cells (Figure 4D). The concomitant pan-cytokeratin and CK8/18 expression for all cell lines confirmed their epithelial phenotype. The primary culture cells, CM1, CM4, CM5, CM9, CM60 and CM61 had doubling time at 6.31, 5.1, 7.5, 8.95, 11.79 and 13.06, respectively. The MM1, MM3 and MM4 metastasis cultures presented their doubling time at 25.41, 34.17 and 10.28, respectively.

Vasculogenic mimicry *in vitro* and *in vivo*

The *in vitro* tubular formation was identified in 4 out of 10 cell cultures. The cell lines CM1, CM9, CM60 and MM4 presented tubular formation at four hours (Figure 5) until six hours and tubules were disrupted from 6 hours.

One primary cell culture (CM60) and its respective metastasis (MM4) showed *in vivo* tumorigenicity (2 out of 10 cell cultures). The macroscopic growth was evident after 50 days after cell application and histology revealed a tumour with highly vasculogenic mimicry formation. Both cell lines presented neoplastic cells with evident nucleoli forming tubular structures like capillaries (6A and 6D). These capillaries-like structures were positive for both pan-cytokeratin and vimentin, indicating (6B, 6C, 6D, 6E and 6F). Interestingly, the metastatic cell line (MM4), also showed an intravascular growth (6A). It was possible to observe several blood vessels in tumour periphery with intravascular growth of cancer cells positive for pan-cytokeratin (6B).

DISCUSSION

Canine mammary gland tumours represent one of the most important tumours in intact female dogs and can be considered a model for studying human disease. In human medicine, several cell lines have been used as a preclinical model for understanding breast cancer (BC) development and progression or studying the antitumor effect of new drugs [26, 27, 28, 29]. Comparing with human BC, there is less canine mammary gland tumour cell line developed and studied [24, 30, 31, 32]. To establish the primary cell cultures, we tried different methodologies aiming to identify the most effective in isolating the neoplastic epithelial mammary neoplastic cells. The cultures obtained through explant samples presented several problems in their establishment, mainly related to contamination during cultivation. Thus, we tried protocols based on the use of enzymatic dissociation. In this step we tried collagenase type I and II in different incubation time (3, 4, 12, 24 and 48 hours) with no success (data not shown). On the other hand, using 0.05% type IV collagenase we had better results. For type IV collagenase 0.05% with 24- and 48-hours incubation, we identified cell damage and death, with no cell growth (data not shown). For samples incubated using 0.05% type IV collagenase

within 4 hours there was no cell damage and satisfactory cultures expansion. Therefore, we standardized the time to up to 4 hours for enzymatic dissociation.

After the different cell cultures were established, they presented a heterogeneous morphology, so selective trypsinization of cells was performed to eliminate fibroblasts and stimulate cell clone formation [33]. After initial expansion, cells were used for characterization around passage 10 (P10), since from lower passages they tend to have a heterogeneous morphology and in higher passages can present a higher number of chromosomal alterations related to cell culture conditions [34, 35]. In addition, at passage 10 or above, we did not find contaminating fibroblasts in the cultures. To eliminate fibroblasts, we used two different strategies, using a culture medium that show specific epithelial cell growth factors and also did a selective trypsinization. Together, both strategies were effective in eliminating fibroblasts and selecting epithelial cells during cell expansion, that can be confirmed based on the expression of epithelial markers by neoplastic cells.

Cell lines presented a monolayer growth, similar size and morphology. They showed spindle-shape, high nucleus/cytoplasm ratio and a tight cell-cell adhesion. The high ratio between nucleus and cytoplasm size can usually be associated with malignant tumour behaviour [36]. In haematoxylin and eosin staining, it was possible to observe basophilic nucleus and eosinophilic cytoplasm, with presence of mitosis and some cells appeared multinucleated and with colony formation. Besides that, our cell lines also expressed vimentin, same results found in MCF7 and HeLa cells cultured *in vitro* [37, 38]. Since in culture conditions cells need to change their cytoskeleton for the flask attachment, it is common to observe vimentin expression either in epithelial cells [39]. This expression was previous explained by post translational modification during cell culture conditions [23]. Doubling time of each culture was analysed and the shortest was from CM4 (5.1) and the longest from cell MM3 (34.17) at passage 10. This is important, considering that the cell doubling time should vary with passages. The difference in cell doubling time is expected, as these are different tumours. These results are similar to those found by Cordeiro et al. [24] in which the doubling times of two different cells lines cultured *in vitro* were 26 and 42 hours and, in this case, one cell line was more malignant than other.

An interesting finding in this study was the *in vitro* tubular formation in four different cell lines that can be related to vasculogenic mimicry (VM). Tumours with VM capacity can promote metastasis of tumour cells through these new channels. Also,

tumour cell lines are highly malignant and are capable of penetrate endothelium, developing tumour invasion and metastasis [15]. Among these cells, two also showed vasculogenic mimicry ability *in vivo* (tumour from xenotransplant). The ability of VM formation is associated with aggressive melanoma cells and not with non-aggressive cells [16], and high tumour grade, invasion, metastasis and poor clinical prognosis in hepatocellular carcinomas [14]. In inflammatory mammary tumours, both human and canine, VM is commonly described, possibly related to tumour aggressiveness and metastasis capacity [40, 41]. The VM results corroborate with the results from tumorigenicity assay where two cellular types, that were capable of VM formation, produced tumours *in vivo*. When analysed in HE, the tumours *in vivo* had highly vasculogenic mimicry formation, showing the aggressivity of this neoplasia, *in vivo* and *in vitro*. The ability to form vasculogenic mimicry *in vitro* and *in vivo*, reinforce the use of these cells as a preclinical model for canine mammary gland tumours.

Among our tumour cell lines, four out of 10 showed tumorigenicity *in vivo*. This result was similar with previously described by Cordeiro et al. [24], wherein only one in two different cell lines have grew *in vivo*. Although the cells had resembling epithelial characteristics, as morphology, growth pattern, phenotype, protein expression and tumoursphere formation, the tumour did not grow *in vivo*. The tumour growth *in vivo* depends on many factors, as the local of the application, cell concentration, tumour heterogeneity and the immune system of the animal [42]. In our case, this result may be explained by the animal model used in the study. The BALB-c nude mice lack thymus and is unable to produce T-cells and is considered immunodeficient mice. However, this nude model shows other immune cells, such as B-cells and natural killers that can affect the tumour growth *in vivo*. In our xenograft study, the histological evaluation revealed a high inflammatory infiltrate in all tumours (N=4), indicating that other immune cells were infiltrating the tumours. Thus, a nude mice with no B-cell and natural killer cells can be accurate to evaluate *in vivo* tumorigenicity [43]. However, for using a more immunocompromised mice, it is necessary an accurate pathogen-free laboratory system. Unfortunately, our institution doesn't have the necessary structure for accommodating these immunocompromised mice. Thus, the major limitation of our study is the lack of structure to evaluate the cell tumorigenicity in a less immunogenic mice.

CONCLUSION

In summary, this study had established and characterized 10 cell lines and xenografts from canine mammary gland carcinomas. The cells cultured *in vitro* demonstrated morphological and phenotype similarities, but had tumorigenicity differences. Furthermore, four cell lines were capable of vasculogenic mimicry ability and two cell lines showed *in vivo* tumorigenicity, both related to malignancy and aggressiveness. Thus, described cell lines will be able to be used in the future for clinical investigations, therapeutic targets, and also for studying genes targets and pathways. The results showed similarities with BC in human demonstrating that the results obtained with these cell lines could be used for comparative oncology studies.

Acknowledgement

We would like to thank Professor Heidge Fukumasu and Dr. Yonara de Gouveia Cordeiro from University of São Paulo (USP) for its help during karyotyping experiments.

Disclosure statement

The authors report no conflict of interest

Funding

This work was supported by São Paulo Research Foundation (FAPESP) for providing the scholarship for one co-author (Grant No. 2016/21474-9). This study was financed in part by the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior – Brasil (CAPES) – Finance Code 001.

REFERENCES

1. Sorenmo K. Canine mammary gland tumors. *Vet Clin North Am - Small Anim Pract.* 2003;33(3):573–96. DOI: 10.1016/S0195-5616(03)00020-2
2. Sleenckx N, de Rooster H, Veldhuis Kroeze EJB, et al. Canine mammary tumours, an Overview. *Reprod Domest Anim.* 2011;46(6):1112–31. DOI: 10.1111/j.1439-0531.2011.01816.x

3. Salehiniya H, Ghoncheh M, Pournamdar Z. Incidence and Mortality and Epidemiology of Breast Cancer in the World. *Asian Pacific J Cancer Prev.* 2016;17:43–6. DOI: 10.7314/APJCP.2016.17.S3.43
4. Abdelmegeed SM, Mohammed S. Canine mammary tumors as a model for human disease (Review). *Oncol Lett.* 2018;15(6):8195–205. DOI: 10.3892/ol.2018.8411
5. Tran CM, Moore AS, Frimberger AE. Surgical treatment of mammary carcinomas in dogs with or without postoperative chemotherapy. *Vet Comp Oncol.* 2016;14(3):252–62. DOI: 10.1111/vco.12092
6. Levi M, Brunetti B, Sarli G, et al. Immunohistochemical Expression of P-glycoprotein and Breast Cancer Resistance Protein in Canine Mammary Hyperplasia, Neoplasia and Supporting Stroma. *J Comp Pathol.* 2016;155(4):277–85. DOI: 10.1016/j.jcpa.2016.07.008
7. Hernandez-Aya LF, Gonzalez-Angulo AM. Adjuvant Systemic Therapies in Breast Cancer. *Surg Clin North Am.* 2013;93(2):473–91. DOI: 10.1016/j.suc.2012.12.002
8. Cassali GD, de Campos CB, Bertagnolli AC, et al. Consensus for the diagnosis, prognosis and treatment of feline mammary tumors. *Brazilian J Vet Res Anim Sci.* 2019;55(2):153–80.
9. Sawyers, C. Targeted cancer therapy. *Nature.* 2004;432(7015):294-297. DOI: 10.1038/nature03095
10. Blows FM, Driver KE, Schmidt MK, et al. Subtyping of breast cancer by immunohistochemistry to investigate a relationship between subtype and short and long term survival: a collaborative analysis of data for 10,159 cases from 12 studies. *PLoS Med.* 2010;7(5):e1000279. DOI: 10.1371/journal.pmed.1000279
11. Visan S, Balacescu O, Berindan-Neagoe I, et al. In vitro comparative models for canine and human breast cancers. *Clujul Med.* 2016;89(1):38–49. DOI: 10.15386/cjmed-519
12. Ruggeri BA, Camp F, Miknyoczki SA. Animal models of disease: pre-clinical animal models of cancer and their applications and utility in drug discovery. *Biochem Pharmacol.* 2014; 87(1):150-161. DOI: 10.1016/j.bcp.2013.06.020
13. Ge H, Luo H. Overview of advances in vasculogenic mimicry – a potential target for tumor therapy. *Cancer Manag Res.* 2018;2(10):2429-2437. DOI: 10.2147/cmar.s164675
14. Sun B, Zhang S, Zhang D, et al. Vasculogenic mimicry is associated with high tumor grade, invasion and metastasis, and short survival in patients with hepatocellular carcinoma. *Oncol Rep.* 2006;16(4):693-8
15. Zhang S, Zhang D, Sun B. Vasculogenic mimicry: current status and future prospects. *Cancer Lett.* 2007;254(2):157-164. DOI: 10.1016/j.canlet.2006.12.036
16. Delgado-Bellido D, Serrano-Saenz S, Fernández-Cortés M, et al. Vasculogenic mimicry signaling revisited: focus on non-vascular VE-cadherin. *Mol Cancer.* 2017;16(1):65. DOI: 10.1186/s12943-017-0631-x
17. Costa CD, Justo AA, Kobayashi PE, et al. Characterization of OCT3/4, Nestin, NANOG, CD44 and CD24 as stem cell markers in canine prostate cancer. *Int J Biochem Cell Biol.* 2019;108(1):21–8. DOI: 10.1016/j.biocel.2019.01.002
18. Goldschmidt MH, Peña L, Rasotto R, et al. Classification and grading of canine mammary tumors. *Vet Pathol.* 2011;48:117–31. DOI: 10.1177/0300985810393258
19. Nguyen F, Peña L, Ibsch C, et al. Canine invasive mammary carcinomas as models of human breast cancer. Part 1: Natural history and prognostic factors. *Breast Cancer Res Treat.* 2018;167(3):635–48. DOI: 10.1007/s10549-017-4548-2
20. Dos Anjos DS, Vital AF, Lainetti PF, et al. Deregulation of VEGFR-2 and PDGFR Expression and Microvascular Density in a Triple-Negative Model of Canine Malignant Mammary Tumors with Lymph Node or Lung Metastasis. *Vet Sci.* 2019;6(1):E3. DOI: 10.3390/vetsci6010003

21. Kisselbach L, Merges M, Bossie A, et al. CD90 expression on human primary cells and elimination of contaminating fibroblasts from cell cultures. *Cytotechnology*. 2009;59(1):31–44. DOI: 10.1007/s10616-009-9190-3
22. Moorhead PS, Nowell PC, Mellman WJ, et al. Chromosome preparations of leukocytes cultured from human peripheral blood. *Exp Cell Res*. 1960;20(3):613–6. DOI: 10.1016/0014-4827(60)90138-5
23. Caceres S, Peña L, DeAndres PJ, et al. Establishment and characterization of a new cell line of canine inflammatory mammary cancer: IPC-366. *PLoS One*. 2015;10(3). DOI: 10.1371/journal.pone.0122277
24. Cordeiro YG, Xavier PLP, Rochetti AL, et al. Transcriptomic profile reveals molecular events associated to focal adhesion and invasion in canine mammary gland tumour cell lines. *Vet Comp Oncol*. 2018;16(1):E89–98. DOI: 10.1111/vco.12339
25. Gouveia JG, Wolf IR, Vilas-Boas LA, et al. Repetitive DNA in the Catfish Genome: rDNA, Microsatellites, and Tc1-Mariner Transposon Sequences in *Imparfinis* Species (Siluriformes, Heptapteridae). *J Hered*. 2017;108(6):650–7. DOI: 10.1093/jhered/esx065
26. De AK, Muthiyar R, Mondal S, et al. A Natural Quinazoline Derivative from Marine Sponge *Hyrtios erectus* Induces Apoptosis of Breast Cancer Cells via ROS Production and Intrinsic or Extrinsic Apoptosis Pathways. *Mar Drugs*. 2019;17(12):E658. DOI: 10.3390/md17120658
27. Lefort S, Tan S, Balani S, et al. Initiation of human mammary cell tumorigenesis by mutant KRAS requires YAP inactivation. *Oncogene*. 2019;1-12. DOI: 10.1038/s41388-019-1111-0
28. Muhammed E, Chen L, Gao Y, et al. Chemo-treated 4T1 breast cancer cells radiation response measured by single and multiple cell ionization using infrared laser trap. *Sci Rep*. 2019;9(1):17547. DOI: 10.1038/s41598-019-53821-y
29. Comşa Ş, Cîmpean AM, Raica M. The Story of MCF-7 Breast Cancer Cell Line: 40 years of Experience in Research. *Anticancer Res*. 2015;35(6):3147-54
30. Gentile LB, Nagamine MK, Biondi LR, et al. Establishment of primary mixed cell cultures from spontaneous canine mammary tumors: Characterization of classic and new cancer-associated molecules. *PLoS One*. 2017;12(9):e0184228. DOI: 10.1371/journal.pone.0184228
31. Osaki T, Sunden Y, Sugiyama A, et al. Establishment of a canine mammary gland tumor cell line and characterization of its miRNA expression. *J Vet Sci*. 2016;17(3):385-90. DOI: 10.4142/jvs.2016.17.3.385
32. Hellmén E, Moller M, Blankenstein MA, et al. Expression of different phenotypes in cell lines from canine mammary spindle-cell tumours and osteosarcomas indicating a pluripotent mammary stem cell origin. *Breast Cancer Res Treat*. 2000;61(3):197-210. DOI: 10.1023/a:1006410020384
33. Geng XF, Fang M, Liu SP, et al. Quantum dot-based molecular imaging of cancer cell growth using a clone formation assay. *Mol Med Rep*. 2016;14(4):3007-3012. DOI: 10.3892/mmr.2016.5632
34. Turin I, Schiavo R, Maestri M, et al. In vitro efficient expansion of tumor cells deriving from different types of human tumor samples. *Med Sci*. 2014;2(2):70-81. DOI: 10.3390/medsci2020070
35. Muñoz JJ, Drigo SA, Kuasne H, et al. A comprehensive characterization of cell cultures and xenografts derived from a human verrucous penile carcinoma. *Tumor Biol*. 2016;37(8):11375-11384. DOI: 10.1007/s13277-016-4951-z
36. Chang CY, Pinwen PC, Wei-Jung C, et al. Assessment of the tumorigenesis and drug susceptibility of three new canine mammary tumor cell lines. *Res Vet Sci*. 2010;88(2):285-293. DOI: 10.1016/j.rvsc.2009.08.006

37. Messica Y, Laser-Azogui A, Volberg T, et al. The role of vimentin in regulating cell invasive migration in dense cultures of breast carcinoma cells. *Nano Lett.* 2017;17(11):6941-6948. DOI: 10.1021/acs.nanolett.7b03358
38. Saunus JM, Smart CE, Kutasovic JR, et al. Multidimensional phenotyping of breast cancer cell lines to guide preclinical research. *Breast Cancer Res Tr.* 2018;167(1):289-301. DOI: 10.1007/s10549-017-4496-x
39. Liu CY, Lin HH, Tang MJ, et al. Vimentin contributes to epithelial-mesenchymal transition cancer cell mechanics by mediating cytoskeletal organization and focal adhesion maturation. *Oncotarget.* 2015;6(18), 15966. DOI: 10.18632/oncotarget.3862
40. Clemente M, Perez-Alenza MD, Illera JC, et al. Histological, immunohistological, and ultrastructural description of vasculogenic mimicry in canine mammary cancer. *Vet Pathol.* 2010;47(2):265-274. DOI: 10.1177/0300985809353167
41. Barreno L, Cáceres S, Alonso-Diez Á, et al. Vasculogenic mimicry-associated ultrastructural findings in human and canine inflammatory breast cancer cell lines. *BMC Cancer.* 2019;19(1), 750. DOI: 10.1186/s12885-019-5955-z
42. Mollard S, Mousseau Y, Baaj Y, et al. How can grafted breast cancer models be optimized?. *Cancer Biol Ther.* 2011;12(10):855-864. DOI: 10.4161/cbt.12.10.18139
43. Kariya R, Matsuda K, Gotoh K, Vaeteewoottacharn K, Hattori S, Okada S. Establishment of nude mice with complete loss of lymphocytes and NK cells and application for in vivo bio-imaging. *In Vivo.* 2014;28(5):779-84