

*Review*

# HDAC6 – An Emerging Target against Chronic Myeloid Leukemia?

**Hélène Losson<sup>1</sup>, Michael Schnekenburger<sup>1</sup>, Mario Dicato<sup>1</sup> and Marc Diederich<sup>2,\*</sup>**

<sup>1</sup> Laboratoire de Biologie Moléculaire et Cellulaire du Cancer, Hôpital Kirchberg 9, rue Edward Steichen, L-2540 Luxembourg, Luxembourg; helene.losson@lbmcc.lu (H.L.) ; michael.schnekenburger@lbmcc.lu (M.S.) ; dicato.mario@chl.lu (M. Dic.)

<sup>2</sup> College of Pharmacy, Seoul National University, 1 Gwanak-ro, Gwanak-gu, Seoul 08826, Korea; marcdiederich@snu.ac.kr (M. Die.)

\* Correspondence: marcdiederich@snu.ac.kr; Tel.: +82 2 880 8919 (M.Die.)

**Abstract:** Imatinib became the standard treatment for chronic myeloid leukemia (CML) about 20 years ago, which was a major breakthrough in stabilizing the pathology and improving the quality of life of patients. However, the emergence of resistance to imatinib and other tyrosine kinase inhibitors leads researchers to characterize new therapeutic targets. Several studies have highlighted the role of histone deacetylase 6 (HDAC6) in various pathologies, including cancer. This protein effectively intervenes in cellular activities by its primarily cytoplasmic localization. In this review, we will discuss the molecular characteristics of the HDAC6 protein, as well as its overexpression in CML leukemic stem cells, which make it a promising therapeutic target for the treatment of CML.


**Keywords:** histone deacetylase 6 inhibitor; personalized treatment; heat shock protein 90 $\alpha$ ; leukemia stem cells; imatinib resistance; targeted therapy

## 1. Introduction

Chronic myeloid leukemia (CML) is a well-known hematological malignancy that is characterized in most patients by the translocation t(9; 22)(q34; q11), leading to the formation of the Philadelphia chromosome carrying the chimeric oncogene *breakpoint cluster region-Abelson (BCR-ABL)*. The slow progression of CML can be stopped if the disease is diagnosed and managed early enough with the reference treatment, imatinib, a first-generation tyrosine kinase inhibitor (TKI). Imatinib acts to block the BCR-ABL protein, which has constitutive tyrosine kinase activity, in its inactive form. However, resistances to imatinib and the second and third generation TKIs frequently develop in patients, leading researchers to synthesize molecules targeting other proteins involved in this pathology, such as histone deacetylase 6 (HDAC6). In this review, we will discuss the characteristics and implication of the HDAC6 protein in CML.

## 2. Chronic myeloid leukemia

CML is a myeloproliferative disorder affecting myeloid progenitors, and is considered as a rare cancer with an incidence of approximately 1 in 100000 cases each year in Europe and the United States [1,2]. There are an estimated 6370 new cases each year in Europe [3] and 8430 new cases have been reported in 2018 in the United States [2]. CML is characterized by an excessive proliferation of leukocytes in the bone marrow, most of which are still immature, and called blasts when they pass into the blood [4]. This pathology comprises three distinct phases: the chronic, accelerated, and blast phases [5] (Figure 1).




**Figure 1.** Progression of chronic myeloid leukemia. Chronic myeloid leukemia (CML) is characterized by an excessive proliferation of non-functional leukocytes (dotted lines) or blasts in the bone marrow and then in the blood. The chronic phase remains generally asymptomatic, during which 5% blasts are found in the bloodstream. After 5 to 6 years without treatment, the pathology progresses towards the accelerated phase (5 to 20% blasts) and symptoms including fatigue and loss of appetite and weight begin to appear. Without treatment, the accelerated phase progresses in 6 to 9 months to the blast phase (more than 20% blasts), causing rapid deterioration of the general condition of the patient and death after only a few months without management. Symptoms and resistance to therapy increase during the course of chronic myeloid leukemia, while the probability of survival decreases.

2.1. Chromosomal rearrangement during chronic myeloid leukemia

A chromosomal anomaly, the t(9;22)(q34;q11) translocation between chromosome 9, which carries the *ABL* gene, and chromosome 22, which carries the *BCR* gene, is found in 95% of patients with CML. This translocation leads to the formation of the Philadelphia chromosome carrying the *BCR-ABL* chimeric gene (Figure 2A), and thus to the expression of the corresponding fusion oncoprotein, which has a constitutive tyrosine kinase activity. The Philadelphia or Ph<sup>+</sup> chromosome was identified for the first time in 1960 by the researchers Hungerford and Nowell in the city of Philadelphia, for which it was named [6,7].

2.2. *BCR-ABL* mRNA and protein

There are multiple breakpoints on the *BCR* and *ABL* genes that lead to the formation of different transcripts (Figure 2B). These transcripts encode *BCR-ABL* proteins of different sizes that have been found in patients (Table 1) [4].



**Figure 2.** Breakpoints in the *ABL1* and *BCR* genes result in the formation of different transcripts encoding the BCR-ABL chimeric protein. (A) Structure of the *ABL* (*Abelson*) 1 and *BCR* (*breakpoint cluster region*) genes composed of 11 and 23 exons, respectively. (B) Diagram of ABL transcripts (1130 amino acids), BCR (1271 amino acids) and BCR-ABL hybrids. The hybrid mRNA eXaY is produced after cleavage occurs after exon X of BCR and before exon Y of ABL. (C) ABL protein (145 kDa) has tyrosine kinase activity and predominantly nuclear localization. It comprises a long lipid chain, myristate (Myr), conferring a capacity for self-inhibition, SH2 and SH3 domains of Src homology, allowing interactions with proline-rich regions and with phosphorylated tyrosines, respectively, a tyrosine kinase (TK) domain, and finally an ABD binding domain to DNA and  $\beta$ -actin. ABL is involved in cytoskeletal dynamics (adhesion, polarity, cell migration, and cellular protrusions), proliferation and cell survival, membrane transport (endocytosis, macropinocytosis, phagocytosis, and caveolae formation) by phosphorylating receptors such as epidermal growth factor receptor, platelet-derived growth factor beta receptor and muscle specific tyrosine kinase receptor [8], and finally in autophagy [9,10]. ABL protein is involved in neurodegenerative diseases and neuroinflammation [9,10]. The BCR protein (160 kDa) has serine / threonine kinase activity and is localized in the cytoplasm [11]. BCR comprises an oligomerization domain (OLI). Mutations in this domain cause a decrease in the tyrosine kinase activity of the BCR-ABL protein and prevent the interaction of BCR and BCR-ABL proteins. It also has a domain of homology to the protein DBL (guanosine diphosphate exchange factor (GDP) in guanosine triphosphate (GTP)), a domain of pleckstrin homology (PH), a C2 domain for binding of calcium-dependent phospholipids, and finally a GTPase (GTPase activating protein) domain (GTPase hydrolysis in GDP and phosphate ion) for binding to Rac proteins.

**Table 1.** Human BCR-ABL transcripts and proteins. The name and composition of the various human BCR-ABL hybrid transcripts identified in patients are described. The size of the corresponding proteins, their frequency of detection, and the cell lines expressing them are also indicated.

Hybrid mRNA name	BCR-ABL mRNA Composition		Size of the corresponding protein (kDa)	Examples of CML cell lines	References
	BCR exons <sup>§</sup>	ABL exons <sup>£</sup>			
e13a2 or b2a2	1 - 13	2 - 11	210	MEG-01, KBM-7, KYO-1, CML-T1, KCL-22	[12-15]
e14a2 or b3a2	1 - 14	2 - 11	210	K-562, KBM-5, LAMA-84, EM-3, TK-6, EM-2	[12,13,15,16]
e13a3 or b2a3	1 - 13	3 - 11	210	NA	[17]
e14a3 or b3a3	1 - 14	3 - 11	210	NA	[17]
e18a2	1 - 18	2 - 11	225	AR-230	[13,18]
e19a2	1 - 19	2 - 11	230	AR-230	[13,17]
e1a2	1	2 - 11	190	SUP-B15*, Z-33*, SD-1*, TOM-1*, Z-119*	[13,14]
e6a2	1 - 6	2 - 11	185	NA	[19]

<sup>§</sup> Of the 23 exons that compose the *BCR* gene. <sup>£</sup> Of the 11 exons that compose the *ABL* gene. \* Acute lymphocytic leukemia cell lines. ABL: Abelson; BCR: breakpoint cluster region; NA: not applicable.

The BCR-ABL protein activates many substrates (Table 2) and signaling pathways, including some involved in cell proliferation and survival, through increased activity or expression of a series of anti-apoptotic proteins including the signal transducer and transcriptional activator 5 (STAT5), Akt, phosphoinositide 3-kinase, or B-cell lymphoma-extra-large [20].

**Table 2.** BCR-ABL substrates.

Substrates	Phosphorylation site	Function	References
Abi 1 and 2	ND	Proliferation	[21]
BAP-1	Serine and tyrosine residues	Proliferation	[22,23]
Cbl	Tyr-674	Unknown	[22,24,25]
CK2	ND	Cell cycle, apoptosis, transcription, viral infection	[26-28]
Crk	Tyr-221	Migration and cellular adhesion	[22,29]
CrkL	Tyr-207	Migration and cellular adhesion	[22,29]

Dok1	Tyr-361	Negative regulation of signaling pathways mediated by tyrosine kinase proteins	[8,22,30]
Fes	ND	Myeloid differentiation	[22,31]
GAP-associated proteins	ND	Ras activation	[22]
GCKR	ND	SAPK activation	[24,32]
Grb2	Tyr-209	Ras activation	[24,33,34]
LASP1	Tyr-171	Interaction with the cytoskeleton, migration and cell survival	[35,36]
Lyn	ND	Cell survival	[37]
Paxillin	ND	Focal adhesion, signaling and cell migration	[22,38]
PLC $\gamma$	Tyr-69/Tyr-74	Actin rearrangement and cell migration	[22,39]
PI3-K p85	ND	Proliferation, survival and cellular motility	[22,40]
PKD	Tyr-463	Proliferation, migration and cell survival, angiogenesis, regulation of gene expression	[22,41]
P27Kip1	Tyr-88	Cell proliferation	[42]
p73	ND	Activation of transcription	[22]
Rad9	Tyr-28	DNA damage repair	[22]
Rad51	Tyr-54	DNA damage repair	[22]
Ras-GAP	ND	Apoptosis, proliferation and cell migration	[22,43]
RNA-polymerase II	C-terminal	Transcription	[22]
RAFT1	ND	Cell proliferation, autophagy, cytoskeletal reorganization	[22,44]
Shc	Tyr-427	Migration, angiogenesis	[22,25,45]
SHIP1, SHIP2	Tyr-986 et Tyr-1135 (SHIP2)	Signal transduction, macrophage programming, phagocytosis, migration	[24,25,46]
STAT5	Tyr-694	Signal transduction, transcription activation	[47]
Syp	ND	Unknown	[22]
Talin	ND	Signal transmission between the extracellular matrix and the cytoskeleton	[22,48]
TERT	ND	Genomic integrity	[22,49]
VAV p95	ND	Hematopoietic differentiation	[22]

Abi: Abelson interactor; BAP-1: breakpoint cluster region-associated protein 1; Cbl: Casitas-B-lineage protein; CK: casein kinase; CrkL: Crk like protein; Dok: docking protein; GAP: guanosine triphosphate (GTP)ase-activating proteins; GCKR: germinal-center kinase related protein; Grb2: growth factor receptor-bound protein 2; LASP1: LIM and Src homology (SH)3 protein 1; ND: not determined; PLC $\gamma$ : phospholipase C- $\gamma$ ; PI3-K: phosphatidylinositol 3-kinase; PKD: protein kinase D; RAFT1: rapamycin and FK506-binding protein (FKBP)-target 1; Ras-GAP: RAS GTPase activating proteins; Shc: Src homology 2 domain containing; SHIP: SH2-containing inositol polyphosphate 5-phosphatase; Syp: SH2-containing phosphotyrosine phosphatase; TERT: telomerase reverse transcriptase; Tyr: tyrosine.

### 3. CML treatments and associated resistance mechanisms

#### 3.1. Targeted therapy with a tyrosine kinase inhibitor

In the 1960s, CML was treated with chemotherapy using busulfan and hydroxyurea. Two different therapeutic strategies emerged in the 1970s: allogeneic stem cell transplantation and the use of interferon  $\alpha$  [5]. A new therapy was approved in 2001 by the Food and Drug Administration (FDA) using the TKI imatinib (Gleevec®) [50]. TKIs represent a major advance in the management of CML patients by enabling targeted therapy for the first time.

There are two types of TKIs: type I TKIs target the active and inactive conformation of the BCR-ABL protein and bind to the ATP binding site on the tyrosine kinase domain of the protein, whereas type II inhibitors target the inactive conformation of the protein and bind to a site adjacent to the available ATP binding site when the activation loop is not phosphorylated [51,52].

Imatinib is type II TKI. Its action therefore prevents the binding of ATP, and thus blocks the protein in its inactive form, preventing its autophosphorylation, and therefore preventing the phosphorylation of its substrates [53]. Imatinib is the first-line treatment for CML with daily oral intake. According to the IRIS study, including 1106 new patients with CML, 76% of patients treated with imatinib also lost the Philadelphia chromosome, also called a complete cytogenetic response. This treatment with a TKI thus allowed stabilization of the pathology in the chronic phase, allowing patients to live normally. However, this treatment has numerous side effects including fatigue, headaches, anemia and myalgia, which are more common, and sometimes even leading to organ toxicity in the liver or heart, which can cause QT prolongation associated with a high risk of rhythm disturbances and serious ventricular disease, which are uncommon and occur in less than 1% of patients treated with imatinib [5].

Patients treated with imatinib should be closely monitored to assess their response to treatment. There are three types of responses: the hematological, cytogenetic, and molecular responses. A complete hematologic response corresponds to normalization of blood levels and spleen size. An examination is performed every two weeks until a complete hematologic response is obtained, followed by a control every three months. Then, an examination is performed every 6 months until reaching a complete cytogenetic response, then every 12 to 18 months. Finally, a complete molecular response corresponds to the total disappearance of the BCR-ABL transcript, which is measured by reverse transcription followed by real-time PCR amplification. This test is performed every 3 months according to the recommendations of the European leukemia net [54].

#### 3.2. Imatinib resistances

Although imatinib is the gold standard for CML treatment, 35 to 40% of patients develop resistances to this drug [55]. It is possible to distinguish between primary resistances, defined as a complete lack of hematological response after 3 months, cytogenetic response after 6 months, or a major cytogenetic response after one year of treatment; and secondary resistances, also called acquired resistances, when the previous hematologic or cytologic response is lost. Resistances can also be classified according to whether or not they depend on BCR-ABL [56,57]. The emergence of resistances has led to the development of new TKIs (Table 3).

**Table 3.** Tyrosine kinase inhibitors approved by the US Food and Drug Administration for the treatment of chronic myeloid leukemia.

Generation	Drug name (and others)	Pharmaceutical company and marketing authorization year by FDA	Targets	Daily dosage in adults	References
First	Imatinib <sup>°</sup> (Gleevec, STI571, CGP57148B)	Novartis, 2001	BCR-ABL, c-KIT, PDGFR	400 mg single dose	[58,59]
Second	Dasatinib* (Sprycel, BMS-354825)	Bristol-Myers Squibb, 2006	BCR-ABL, Src family, c-KIT, PDGFR	100 mg single dose	[58-60]
	Nilotinib <sup>°</sup> (Tasigna, AMN107)	Novartis, 2007	BCR-ABL, c-KIT, PDGFR	300 mg in two doses	[58,59,61]
Third	Ponatinib <sup>°</sup> (Iclusig, AP24534)	ARIAD Pharmaceuticals, 2012	BCR-ABL, FTL3, Src family, RET	45 mg single dose	[58,59]
	Bosutinib* (Bosulif, SKI-606)	Pfizer, 2012	BCR-ABL, Src family	500 mg single dose	[58,59,62]


<sup>°</sup> Type II inhibitor binding to the inactive conformation of BCR-ABL. \* Type I inhibitor binding to the active and inactive conformation of BCR-ABL. BCR-ABL: breakpoint cluster region-Abelson; FTL3: Fms-like tyrosine kinase 3; PDGFR: platelet-derived growth factor receptor; RET: rearranged during transfection.

### 3.2.1. BCR-ABL-dependent resistance mechanisms

BCR-ABL-dependent resistance can be caused by duplication and mutation of the *BCR-ABL* gene (Figure 3). Duplication of the gene has been identified in the cells of imatinib-resistant patients and could be a possible source of drug resistance [63]. Although overexpression of BCR-ABL has been reported in patients with accelerated and blastic phase CML who became resistant to imatinib, several studies have shown that only 3% of imatinib-resistant patients have amplification of *BCR-ABL* gene [64].

Mutations in *BCR-ABL* are more common than duplications and occur in 40 to 90% of imatinib-resistant patients, depending on the sensitivity of the detection method used and the stage of CML [65]. To date, more than a hundred have been discovered [66], which can explain the recently observed decrease in the effectiveness of imatinib treatment [63]. The first mutation described, which is also the most common, represents 14% of all mutations detected [64], and corresponds to the nucleotide substitution of a cytosine by a thymine at position 944 of the *ABL* gene. This mutation results in the substitution of the amino acid 315, initially threonine, with an isoleucine (T315I). This results in the loss of an oxygen molecule that is necessary for the hydrogen bond between imatinib and the tyrosine kinase domain, and also creates steric hindrance, preventing binding and drastically reducing treatment efficacy [67-69].





**Figure 3.** BCR-ABL-dependent and -independent imatinib resistances. BCR-ABL-dependent (purple) and -independent (blue) resistances can be explained by *BCR-ABL* duplication and mutation mechanisms, co-medication, interindividual variety, decreased import proteins, increased export proteins, binding of imatinib to plasma proteins, and the presence of imatinib-insensitive leukemic stem cells (LSCs). CYP3A4: cytochrome 3A4.

The seven most common mutations are: G250A/E, Y253F/H, and E255D/K/R/V located in the ATP binding P loop, T315I located at the imatinib binding site, M351T and F359C/L/V/R located in the catalytic loop, and H396P located at the activation loop A [64]. Mutations at the P-loop represent 38 to 46% of all mutations and result in a conformational change that prevents imatinib from binding to BCR-ABL [54]. Mutations occurring at loop A prevent BCR-ABL from attaining its active conformation, thus also preventing binding to imatinib [64].



It is interesting to note that the frequency of mutations is higher in patients who have developed secondary resistance, and that the site of mutation varies according to the progression of the pathology. Mutations of amino acids at position 244, 250, and 351 are more frequent in patients in the chronic phase, whereas mutations of amino acids at position 253, 255, and 315 are more frequently encountered in patients in the accelerated or blast phases [64].

### 3.2.2. BCR-ABL-independent resistance mechanisms

BCR-ABL-independent resistances can be explained by interindividual variability, increased export protein, decreased import protein, and also by binding of imatinib to plasma proteins [67].

Interindividual variability may underlie differences in drug metabolism, and thus a different drug response in patients. The metabolism of imatinib to its main circulating metabolite, the N-desmethyl piperazine derivative [55], progresses *via* cytochrome (CYP) P450, and in particular to the CYP3A4 isoform, which has variable activity. This may explain the observed lower levels of imatinib in some patients despite a similar dose. Cytochrome CYP3A4 can be activated or inhibited by many drugs. The metabolism of imatinib by CYP3A4 may therefore be important during co-medication [65].

A reduced cellular concentration of imatinib can be explained by decreased levels of import proteins or by increased levels of export proteins (Figure 3B). The organic cation transporter (OCT)1 is responsible for the influx of imatinib into leukemic cells, and a polymorphism of this transporter is associated with imatinib resistance in K-562 CML cells [64]. Glycoprotein G, also known as multi-drug resistance protein 1, is an export protein that has been associated with failed leukemia treatment by chemotherapy. Recent studies have supported the notion that imatinib is also a substrate for glycoprotein G [70].

Although this hypothesis remains controversial in the literature, the binding of imatinib to plasma proteins such as alpha-1-acid glycoprotein may explain the decrease in free imatinib concentration described in some patients with little or no response to treatment [54,64].

Finally, the existence of quiescent leukemic stem cells (LSC), which are unaffected by current drug therapies, would also explain the emergence of resistance to imatinib. LSCs are derived from a group of myeloid progenitors that gain the ability to self-renew, become quiescent, and finally survive in specific microenvironments called hematopoietic niches [4]. These allow interactions between the LSCs and the cells of the microenvironment, which can promote the development of resistance mechanisms. The persistence of LSCs in the presence of TKIs could be the cause of the molecular minimal residual disease that can promote the long-term development of resistance, making them potentially responsible for the clonal evolution and progression of CML [71].

### 3.3. Development of novel tyrosine kinase inhibitors

The emergence of resistances to imatinib has led to the development of new TKIs with an activity against specific mutations in BCR-ABL protein. Dasatinib and nilotinib are second-generation TKIs (Table 3) used in cases of imatinib resistance, allowing a complete, lasting cytogenetic response in approximately 40% of these cases. Despite promising results, these new molecules are nevertheless not effective in patients with the T315I mutation, and lead to the appearance of new mutations generating resistance and subsequently to a decrease in their effectiveness. In addition, the benefit-risk ratio is important to consider as the side effects caused by second-generation TKIs are numerous, and for the most part similar to those generated by imatinib [5].

Ponatinib and bosutinib are third generation TKIs (Table 3) that are effective in second generation TKI refractory patients [72]. Only ponatinib is effective in patients with imatinib resistance associated with the T315I mutation. A complete cytogenetic response was observed in 46% of chronic phase patients receiving ponatinib who had already been treated with other TKIs. Data from a phase II study suggest an increased incidence of arterial thrombotic events in patients receiving ponatinib, thus ending a phase III study. In addition, third generation TKIs do not escape the appearance of resistance phenomena [5,62]. Third generation TKIs also include radotinib, a TKI with structural

similarities to imatinib and nilotinib, which was approved in Korea in 2012 for the second-line treatment of CML [73]. This TKI is one of the most effective treatments for chronic phase CML [74], but has side effects including pigment changes such as eruptive melanocytic nevi [75].


4. Histone deacetylase 6

The HDAC6 protein is part of the HDAC family, which are enzymes catalyzing the deacetylation of proteins, which corresponds to the removal of an acetyl group from lysine residues [76]. The 18 HDACs found in mammals are divided into four classes according to their sequence homology. For classes I (HDAC1, 2, 3 and 8), IIa (HDAC4, 5, 7 and 9), IIb (HDAC6 and 10), and IV (HDAC11), the deacetylation of lysine occurs through a transfer of charge, and their essential component is a zinc ion (Zn<sup>2+</sup>) present at the bottom of the catalytic pocket of HDAC enzymes [77]. For class III [sirtuins (SIRT) 1-7] HDACs, the presence of a cofactor, nicotinamide adenine dinucleotide (NAD<sup>+</sup>), is essential for the reaction [78,79].

Class I HDACs are ubiquitously present in many human cell lines and tissues, while class II HDACs exhibit a specific expression profile in certain human tissues such as the heart (HDAC5), the breast (HDAC6), the ovary (HDAC7 and 9), and the kidney (HDAC10) [80,81].

4.1. Structure

Here, we will focus more specifically on HDAC6 belonging to class IIb. This enzyme is the only HDAC to possess two functional active catalytic sites, and has a nuclear localization sequence, a nuclear export sequence, and a repetitive region of eight consecutive serine-glutamic acid tetradecapeptides, a cytoplasmic retention signal, and is mainly present in the cytoplasm [82]. HDAC6 also has a C-terminal ubiquitin binding domain required when binding to poly-ubiquitinated proteins (Figure 4).



**Figure 4.** Protein structure of HDAC6. The HDAC6 protein consists of 1215 amino acids. It has a nuclear localization sequence (NLS), a nuclear export sequence (NES), two functional active catalytic (SC) sites, a cytoplasmic retention signal of eight consecutive serine-glutamic acid tetradecapeptides (SE14), and a ubiquitin binding domain (BUZ) at the C-terminal.

4.2. Function

The HDAC6 protein deacetylates many substrates [83] (Table 4) including  $\alpha$ -tubulin, cortactin, and heat shock protein (HSP)90 $\alpha$ , and is thus involved in many cell processes, some of which are described below [84].

**Table 4.** List of substrates specifically deacetylated by HDAC6.

Substrates	Localization of the substrate	Deacetylated lysine(s)	Function of the deacetylated substrate	Interaction domains of HDAC6	Reference
14-3-3 $\zeta$	Cytoplasm and nucleus	49, 120	Regulation of protein binding Bad and AS160	ND	[85]
$\beta$ -catenin	Cytoplasm and nucleus	49	Epidermal growth factor-induced nuclear localization and	ND	[83]

			decreased expression of c-Myc		
Cortactin*	Cytoplasm	87, 124, 161, 189, 198, 235, 272, 309, 319	Regulation of cell migration and actin filament binding	DD1 and DD2	[83]
DNAJA1	Cytoplasm	ND	Protein folding	ND	[86]
ERK1	Cytoplasm and nucleus	72	Proliferation, mobility and cell survival		[87]
Foxp3*	Nucleus	ND	ND	ND	[88]
HDAC9	Cytoplasm and nucleus	ND	Modulation of cell survival and arrest of cellular movement	DD2	[89]
HDAC11	Nucleus	ND	Transcriptional activation of interleukin 10	ND	[90]
HMGN2	Nucleus	2	Increased transcription of STAT5	ND	[91]
HSC70	Cytoplasm	ND	Protein folding	ND	[86]
HSPA5	Cytoplasm	353	Ubiquitination of HSPA5 mediated by GP78	ND	[92]
HSP90α	Cytoplasm	294	Degradation and elimination of misfolded proteins and regulation of glucocorticoid receptors	DD1, DD2 et BUZ	[83]
K-RAS*	Cytoplasm	104	Cell proliferation	ND	[93]
Ku70	Cytoplasm	539, 542	Suppression of apoptosis	ND	[83]
LC3B-II*	Cytoplasm	ND	Regulation of autophagy	ND	[94]
MSH2	Cytoplasm and nucleus	845, 847, 871, 892	Reduced cellular sensitivity to DNA damaging agents and reduced DNA mismatch repair activities by downregulation of MSH2	DD1	[95]
MYH9	Cytoplasm	ND	Regulation of binding to actin filaments	ND	[86]
PrxI	Cytoplasm and nucleus	197	Antioxidant activity	ND	[96 ,97]


PrxII	Cytoplasm and nucleus	196	Antioxidant activity	ND	[96 ,97]
RIG-I	Cytoplasm	858, 909	Recognition of viral RNA	ND	[98]
Sam68	Nucleus	ND	Alternative splicing	ND	[99]
Survivin	Nucleus	129	Anti-apoptotic function	DD2	[83]
Tat	Cytoplasm	28	Suppression of HIV transactivation	DD2 and BUZ	[100]
$\alpha$ -tubulin*	Cytoplasm	40	Formation of immune synapses, viral infection, cell migration and chemotaxis	DD1 or DD2	[83,101 ]

\* Cortactin and LC3B-II are also deacetylated by SIRT1, K-RAS and  $\alpha$ -tubulin are also deacetylated by SIRT2, and Foxp3 is also deacetylated by HDAC9 and SIRT1. AS160: Akt substrate of 160 kDa; Bad: Bcl-2 associated agonist of cell death; BUZ: binding-of-ubiquitin zinc; DD: deacetylase domain; DNAJA1: dnaJ homolog subfamily A member 1; ERK1: extracellular signal-regulated kinase 1; Foxp3: forkhead box P3; GP: glycoprotein; HDAC: histone deacetylase; HIV: human immunodeficiency virus; HMGN2: high mobility group nucleosomal binding domain 2; HSC: heat-shock cognate; HSP (A): heat shock protein [family A (HSP70) member 5]; LC3B-II: microtubule-associated protein 1 light chain 3; MSH2: MutS protein homolog 2; MYH9: myosin heavy chain 9; ND: non determined; Prx: peroxiredoxin; RIG-I: retinoic acid-inducible gene I protein; Sam: Src-associated substrate in mitosis; STAT: signal transducer and transcriptional activator; Tat: twin-arginine translocation protein.

The HDAC6 protein plays an important role in the dynamism of two components of the cytoskeleton, actin filaments (or F-actin) and microtubules,  $\alpha$ - and  $\beta$ -tubulin polymers, which are involved in particular in cell mobility and division. Cortactin, which improves the polymerization of actin filaments, and  $\alpha$ -tubulin, a constituent of microtubules, are substrates of HDAC6. The deacetylation of cortactin by HDAC6 and SIRT1 leads to its binding to F-actin, improving its polymerization, and thus contributing to cytoskeletal dynamics (Figure 5A). The deacetylation of  $\alpha$ -tubulin by HDAC6 and SIRT2 is associated with microtubule depolymerization, thus contributing to the dynamism of microtubules (Figure 5B), and to proteasome-independent protein degradation. When the proteasome is degraded, the polyubiquitinated misfolded proteins are transported to the microtubule organizing center and are supported by HDAC6 *via* its ubiquitin binding domain, leading to the formation of aggresomes through deacetylation of cortactin. The aggresomes thus formed are subsequently removed after autophagosome fusion with lysosomes *via* autophagy (Figure 5C). A decrease in the acetylation of microtubule-associated protein 1 light chain 3 by HDAC6 was observed during autophagic degradation [68,102]. The HDAC6 protein is also involved in proteasome-dependent protein degradation *via* its interaction with HSP90 $\alpha$ , a chaperone that stabilizes other proteins when deacetylated by HDAC6. In its acetylated form, HSP90 $\alpha$  loses its chaperone function, which leads to the degradation of its client proteins by the proteasome (Figure 5D). An accumulation of misfolded proteins causes dissociation of the complex containing HSP90 $\alpha$ , heat shock factor 1 (HSF) 1, chaperone valosin-containing protein/ATPase, and HDAC6. In complex in inactive form, during dissociation, the release of HSF1 induces the transcription of many HSPs (Figure 5E), and HDAC6 will allow its binding to misfolded proteins [68].

HDAC6 is also involved in apoptosis by deacetylating the Ku70 protein, which then forms a complex with BAX, a proapoptotic protein, allowing the inhibition of apoptosis (Figure 5F). Similarly, inhibition of the catalytic activity of HDAC6 promotes the dephosphorylation of AKT and ERK, associated with decreased cell proliferation and death of cancer cells [68].

Furthermore, HDAC6 regulates endocytosis and exocytosis vesicles. When the EGFR receptor is bound to its ligand, it interacts with HDAC6 and inactivates it by phosphorylation, which then leads to the hyperacetylation of microtubules and finally the internalization of the receptor. The inhibition of HDAC6 induces the increase of the acetylation of peroxiredoxins 1 and 2, which are antioxidant enzymes, increasing their activity and causing a reduction in cell resistance to chemotherapy [68]. HDAC6 is involved in the process of autophosphorylation of tau protein, giving it the ability to form aggregates called neurofibrillary tangles that can cause neurotoxicity [103].



**Figure 5.** The HDAC6 protein is involved in many cellular processes. HDAC6 is involved in F-actin polymerization, microtubule dynamics, anti-apoptotic activity, proteasome-dependent and -independent degradation, transcriptional activation of chaperone proteins, and autophagy. Ac: acetylated; HAT: histone acetyltransferase; HDAC: histone deacetylase; HSF: heat shock factor; HSP: heat shock protein; MTOC: microtubule organizing center; SIRT: sirtuin; VCP: valosin-containing protein / ATPase.

#### 4.3. Post-transcriptional regulation

There is a lack of current data explaining the post-transcriptional regulation of HDAC6 protein. Nevertheless, some microRNAs stimulating cancer cell proliferation and metastasis formation (miR-22, miR-221, miR-433 and miR-548) [84], and stem cell differentiation (miR-26a) [104], are predicted to interact with HDAC6 protein, thus inducing a destabilization or repression of the translation of its mRNA.

#### 4.4. Post-translational regulation

Post-translational modifications such as phosphorylation and acetylation have a significant impact on HDAC6 functions. Indeed, although EGFR induces an inhibitory phosphorylation of

HDAC6, in the majority of cases it is established that the phosphorylation of HDAC6 improves its deacetylase activity, whereas acetylation decreases its enzymatic activity, preventing the deacetylation of  $\alpha$ -tubulin. Examples of post-translational modifications of the HDAC6 protein influencing its activity are shown in Table 5.

**Table 5.** Post-translational modifications regulating the activity of HDAC6.

Post-translational modification	Enzyme	Target site	Consequences	Reference
Phosphorylation	GSK3 $\beta$	Ser-22	Increased deacetylation activity of $\alpha$ -tubulin	[84]
	ERK1	Ser-1035	Regulation of cellular motility	[84]
	GRK2	ND	Increased deacetylation activity of $\alpha$ -tubulin	[105]
	Aurora	ND	Increased deacetylation activity of $\alpha$ -tubulin	[84]
	PKC $\zeta$	ND	Increased deacetylation activity of $\alpha$ -tubulin	[84]
	CK2	Ser-458	Improved formation and elimination of aggresomes	[84]
	EGFR	Tyr-570	Inhibition of deacetylation activity	[106]
Acetylation	p300	Lys-16	Inhibition of deacetylation activity	[84]

CK2: casein kinase 2; EGFR: epidermal growth factor receptor; ERK1: extracellular signal-regulated kinase; GRK2: G protein-coupled receptor kinase 2; GSK3: glycogen synthase kinase 3; Lys: lysine; ND: non determined; PKC $\zeta$ : protein kinase C isoform  $\zeta$ ; Ser: serine; Thr: threonine.

In addition to these known post-translational modifications, there are proteins interacting directly with the HDAC6 protein and inducing its inhibition by direct interaction (Table 6).

**Table 6.** Proteins that interact directly with the HDAC6 protein.

Protein inhibiting HDAC6 by direct interaction	Protein function	Protein region required for interaction with HDAC6	HDAC6 domain interacting with the protein	Cellular impact	References
--	------------------	--	---	-----------------	------------



CYLD	Deubiquitinase	ND	DD1/DD2	Cell proliferation, ciliogenesis	[84]
Dysferlin	Skeletal muscle membrane repair, myogenesis, cell adhesion, intercellular calcium signaling	Domain C2	ND	Myogenesis	[107]
Mdp3	Stabilization factor of microtubules	Amino-terminal region	ND	Cell motility	[108]
Paxillin	Focal adhesion	Region rich in proline	ND	Polarization and cell migration	[84]
p62	Transport of misfolded proteins	Between the ZZ domain and the TRAF6 link area	DD2	Aggresome formation	[109]
RanBPM	Apoptosis, proliferation and cell migration		ND	Aggresome formation	[110]
Tau	Stabilization factor of microtubules	Tubulin binding region	SE14 domain	Aggresome formation	[109,111]
TPPP1	Polymerization and acetylation of microtubules		ND	Regulation of microtubule acetylation and $\beta$ -catenin expression	[112]

DD: deacetylase domain; Mdp3: microtubule associated protein (MAP) 7 domain-containing protein 3; ND: non determined; RanBPM: Ran-binding protein microtubule-organizing center; tau: tubulin-associated unit; TPPP1: tubulin polymerization-promoting protein-1.

4.5. HDAC6 inhibitors

HDAC6 is overexpressed in many types of cancer (Table 7) and may be implicated in disease progression.

**Table 7.** Deregulation of HDAC6 expression in different types of cancers.

Cancer type	Cancers	Expression of HDAC6 - Comments	References
Solid tumors	Bladder	Overexpressed	[113]
	Melanoma	Overexpressed	[113]
	Lung	Overexpressed	[113]
	Oral squamous cell carcinoma	Overexpressed - Enhanced expression in advanced stages	[68 ,114]



Hematological	Ovarian carcinoma	Overexpressed - Enhanced expression in advanced stages	[68,114 ]
	Breast	Overexpressed - Prediction of a good or bad prognosis	[68,115 ]
	Hepatocytic carcinoma	Overexpressed - Enhanced expression in advanced stages	[68]
		Under-expressed - HDAC6 suggested as a tumor suppressor	[68 ,116]
	Chronic lymphocytic leukemia	Overexpressed - Observation on patient samples, cell lines and a transgenic mouse model	[114]
	Acute myeloid leukemia	Overexpressed	[68 ,114]
	Acute lymphoblastic leukemia	Overexpressed - Enhanced expression in advanced stages	[68]
	Chronic lymphocytic leukemia	Overexpressed - Correlated with longer survival	[68]
	T-cell cutaneous lymphoma	Overexpressed - Correlated with longer survival	[68]
	Chronic myeloid leukemia	Overexpressed - Increased expression in CD34 <sup>+</sup> cells	[117]
	Multiple myeloma	Overexpressed	[118]
	Mantle cell lymphoma	Overexpressed	[118]
	Diffuse large B cell lymphoma	Overexpressed	[118]
	Peripheral T-cell lymphoma	Overexpressed	[118]

CD: cluster of differentiation; HDAC6: histone deacetylase 6.

The ability to specifically target HDAC6 would have valuable clinical utility in the treatment of these cancers. However, despite a large number of pan-HDAC inhibitors, very few compounds are capable of selectively inhibiting HDAC6 (Table 8). This type of inhibitor can be divided into 2 groups according to their chemical structure: benzamides and hydroxamates [68].

Table 8. List of HDAC6 inhibitors.

Class	HDAC6 inhibitor	Binding domain	CI <sub>50</sub> (nM) of the HDAC6 activity <i>in vitro</i>	Inhibition of HDAC6 <i>in cellulo</i> (μM) <sup>s</sup>	Effect on cancer cell lines or cancer type	References
Benzamides	Trithiocarbonate derivative (12ac)	ND	65	10 (lung cancer)	CI <sub>50</sub> = 8.2 μM (cervical cancer)	[119]
	NQN-1 (2-benzyl-amino-naphthoquinone)	ND	5540	4 (chronic myeloid leukemia)	CI <sub>50</sub> = 0.8 μM (leukemia)	[120]
Hydroxamates	Hydroxamic acid containing a phenylalanine (4n)	His215, His216, Tyr386, Phe283, and Tyr255 of DD1 and His610, His611, Tyr782, Phe620, and Phe680 of an HDAC6 homology model	1690	1 (colorectal carcinoma)	IC <sub>50</sub> : 3 to > 50 μM (various cancer cell lines)	[121]
	Hydroxamic acid containing a pyridylalanine (5a)	Phe566 of DD2 of an HDAC6 homology model	3970	ND	IC <sub>50</sub> : 104 μM (breast cancer)	[122]
	ACY-738	ND	1.7	2.5 (neural cells)	ND	[123]
	ACY-775	ND	7.5	2.5 (neural cells)	ND	[123]
	ACY-1083	His573 and His574 of DD2	3	0.03 (neuroblastoma)	ND	[124 ,125]
	Bavarostat	Ser568 of DD2	60	10 (neural progenitor cells derived from induced pluripotent stem cells)	ND	[126]

BRD9757	ND	30	10 (cervical cancer)	ND	[127]
C1A	Phe150 and Phe205 of DD1 and Phe620 and Phe680 of DD2	479	1 (colon cancer)	IC <sub>50</sub> : 0.2 to 7.7 µM (various cancer cell lines)	[128]
Cay10603	His499 of DD2 of an HDAC6 homology model	0.002	<1 to 1 µM (several pancreatic cancer cell lines)	ND	[129 ,130]
Citarinostat (ACY-241)	ND	2.6	0.3 (ovarian cancer)	CI <sub>50</sub> : 4.6 to 6.1 µM (ovarian and breast cancer)	[131]
α3β-cyclic tetrapeptide (23)	ND	39	2 (acute lymphoblastic leukemia)	IC <sub>50</sub> : 9 to > 20 µM (various cancer cell lines)	[132]
Compound containing a phenylisoxazole group as a surface recognition group (7)	His499 of HDAC7	0.002	ND	IC <sub>50</sub> : 0.1 to 1 µM (various prostate cancer cell lines)	[129]
Compound containing a triazolylphenyl group (6b)	ND	1.9	ND	IC <sub>50</sub> : <0.5 to 22 µM (several prostate cancer lines)	[133]
Compound containing a peptoid (2i)	Tyr301 of DD2 of an HDAC6 homology model	1.59	N	IC <sub>50</sub> : 0.34 to 2.7 µM (various cancer cell lines)	[134]
3-aminopyrrolidinone derivative (33)	ND	17	0.3 (multiple myeloma)	Good oral bioavailability	[135]
4-aminomethylaryl acid derivative (1a)	ND	19	0.46 (cervical cancer)	ND	[136]
4-hydroxybenzoic acid derivative (7b)	ND	200	50 (prostate cancer)	IC <sub>50</sub> : 41 to 130 (several prostate and breast cancer cell lines)	[137]

4-hydroxybenzoic acid derivative (13a)	ND	20000	50 (prostate cancer)	IC <sub>50</sub> : 19 to 127 (several prostate and breast cancer cell lines)	[137]
Aminoteraline derivative (32)	Phe620 and Phe680 of an HDAC6 homology model	50	2 (neuroblastoma)	IC <sub>50</sub> = 5.4 µM (neuroblastoma)	[138]
Benzothiophene derivative (39)	ND	14	Same effect as tubastatin A	Does not target NF-κB and AP-1 at the transcriptional level	[139]
2,4-imidazolinedione derivative (10c)	ND	4.4	1.6 (acute myeloid leukemia)	IC <sub>50</sub> : 0.2 to 0.8 µM (various cancer cell lines)	[140]
Mercaptoacetamide derivative (2)	ND	95.3	ND	At 10 µM protects cortical neurons from oxidative stress inducing death	[141]
N-Hydroxycarbonylbenzylamino quinoline derivative (13)	ND	0.29	0.1 (multiple myeloma)	IC <sub>50</sub> : 9.1 to 40.6 µM (multiple myeloma)	[142]
Isoxazole-3-hydroxamate derivative (SS-208)	His463, Pro464, Phe583, and Leu712 of DD2	12	5 (melanoma)	ND	[143]
Phenothiazine derivative (7i)	Phe620 and Phe680 of DD2	5	0.1 (acute myeloid leukemia)	ND	[144]
Phenylhydroxamate derivative (2)	Phe464 and His614 of DD2	3	ND	CI <sub>50</sub> : 0.65 to 2.77 (ovarian cancer and squamous cell carcinoma of the tongue)	[134 ,145]

Phenylsulfonylfuroxan derivative (5c)	ND	7.4	0.013 (acute myeloid leukemia)	IC <sub>50</sub> : 0.4 to 5.8 µM (various cancer cell lines)	[146]
Pyridone derivative (11e)	Phe155 and Phe210 of HDAC2	2.46	ND	IC <sub>50</sub> : 0.14 to 0.38 µM (various cancer cell lines)	[147]
Pyrimidinedione derivative (6)	ND	12.4	ND	Induces arrest of the cell cycle in subG1 phase and death by apoptosis (colon cancer)	[139,148]
Quinazolin-4-one derivative (3f)	ND	29	Increases acetylation levels of α-tubulin and histone H3 at 10 µM	ND	[149]
Sulfone derivative (36)	ND	8	0,01 (unspecified)	ND	[139]
Trichostatine A derivatives (M344, 16b)	ND	88	ND	ND	[150]
Tubacin derivative (WT-161)	Phe200, Phe201, Leu270, Arg194 of HDAC7	0.4	0.3 (multiple myeloma)	IC <sub>50</sub> = 3,6 µM (multiple myeloma)	[151]
Tubastatin A derivative (Marbostat-100)	Asp649, His651 et Asp742 of DD2	0.7	0.05 (acute monocytic leukemia)	CI <sub>50</sub> : 1.5 to 4.7 µM (multiple myeloma cell lines) Non-cytotoxic	[152]
Indolylsulfonylcinnamic hydroxamate (12)	ND	5.2	0.1 (colon cancer)	IC <sub>50</sub> : 0.4 to 2.5 µM (multiple cancer cell lines)	[153]

Hydroxamate phenylsulfonylfuroxane (5c)	ND	7.4	0.0125 (acute myeloid leukemia)	IC <sub>50</sub> : 0.38 to 5.84 µM (multiple cancer cell lines)	[146]
MAIP-032	DD2	58	ND	CI <sub>50</sub> : 3.87 µM (squamous cell carcinoma line of the tongue)	[154]
MPT0G211	ND	0.291	0.1 (neuroblastoma)	ND	[103]
N-hydroxy-4-[(N(2- hydroxyethyl)-2- phenylacetamido)methyl]- benzamide)] (HPB)	His573 and His574 of DD2	31	8 (prostate cancer)	ND	[124 ,155]
N-hydroxy-4-(2-[(2- hydroxyethyl)(phenyl)ami no]- 2-oxoethyl)benzamide (HPOB)	Binding to zinc ion only via its OH group but does not displace the zinc-bound water molecule	56	16 (prostate cancer, adenocarcinoma, glioblastoma)	Increases the effect on cell viability in combination with etoposide, dexamethasone or SAHA	[156 ,157]
N-hydroxy-4-(2-methoxy- 5-(methyl(2- methylquinazolin-4-yl)- amino)phenoxy)butanamid e (23bb)	Tyr298 and Glu255 of an HDAC6 homology model	17	0.051 (cervical cancer)	IC <sub>50</sub> : 14 to 104 nM (various cancer cell lines)	[158]
Nexturastat A	DD2 of an HDAC6 homology model	5	0.01 (murine melanoma)	IC <sub>50</sub> = 14.3 µM (melanoma)	[130,159 ]
Oxazole hydroxamate (4g)	Phe620, Phe680, Leu749, and Tyr782 of DD2 of an HDAC6 homology model	59	10 (cervical cancer)	IC <sub>50</sub> = 10.2 µM (acute myeloid leukemia)	[160]

	Rocilinostat (ACY-1215)	DD2 of an HDAC6 homology model	4.7	0.62 (multiple myeloma)	CI <sub>50</sub> : 2 to 8 µM (multiple myeloma cell lines)	[130,161,162]
	Sahaquine	ND	ND	0.1 (glioblastoma)	CI <sub>50</sub> : 10 µM (glioblastoma)	[163]
	TC24	Ser568, His610, Phe679 and Tyr782 of HDAC6	ND	1 et 10 (gastric cancer)	CI <sub>50</sub> : 10.2 to 17.2 µM (several gastric cancer cell lines)	[164]
	Tetrahydroisoquinoline (5a)	ND	36	0.21 (cervical cancer)	ND	[136]
	Thiophene	ND	19	ND	ND	[136]
	Thiazole	ND	52	ND	ND	[136]
	Tubacin	DD2 of an HDAC6 homology model	4	5 (prostate cancer) 2.5 (acute lymphoblastic leukemia)	IC <sub>50</sub> : 1.2 to 2 µM (acute lymphoblastic leukemia)	[130,165 ,166 ]
	Tubastatine A	His610, His611, Phe679, Phe680 and Tyr782 of HDAC6	15	2.5 (unspecified)	ND	[164,165 ]
	Tubathian A	ND	1.9	0.1 (ovarian cancer)	ND	[167]
Other	3-hydroxypyridine-2-thione (3-HPT)	Tyr306 of HDAC8	681	ND	Inactive against two prostate cancer cell lines and one acute T cell leukemia cell line	[168]
	1-hydroxypyridine-2-thione (1HPT)-6-carboxylic acid	DD	150	ND	CI <sub>50</sub> : 18 to 75 µM (leukemia)	[169]
	Adamantylamino derivative (20a)	ND	82	ND	ND	[170]



Undefined structure	Mercaptoacetamide derivative (2b)	ND	1.3	10 (primary rat cortical culture)	ND	[171]
	Sulfamide derivative (13e)	ND	440	1 (bladder cancer)	ND	[172]
	CKD-506	ND	5	0.03 (Human PBMCs)	ND	[173]

Arg: arginine; Asp: aspartic acid; CI<sub>50</sub>: concentration inhibiting 50% of cell viability; DD: deacetylase domain; Glu: glutamic acid; HDAC: histone deacetylase; His: histidine; IC<sub>50</sub>: concentration inhibiting 50% of cell growth; Leu: leucine; ND: non determined; PBMC: peripheral blood mononuclear cell; Phe: phenylalanine; Pro: proline; Ser: serine; Tyr: tyrosine.

The compounds ACY-241 (Citarinostat) and ACY-1215 (Rocilinostat) are derivatives of hydroxamic acid, which shows a specific inhibitory activity of HDAC6 with IC<sub>50</sub>'s of 2.6 and 5 nM, respectively. They are the only HDAC6 inhibitors in currently clinical trials (Table 9) [131,161]. To date, no HDAC6 inhibitor has yet been approved by the FDA, unlike pan HDAC inhibitors such as romidepsin, SAHA, PXD101, and LBH589 [174].

**Table 9.** HDAC6 Inhibitors in Clinical Trials in Cancer. Clinical studies include four phases. Phase I is performed on healthy volunteers to determine the maximum tolerated dose in humans. Phase II is performed on a limited patient population to determine the optimal dosage. Phase III is performed on several thousand patients and will demonstrate the therapeutic value of the drug and assess its benefit/risk. Phase IV is performed once the drug is marketed and allows to better characterize its adverse effects.

HDAC6 inhibitor	Clinical trial identification	Phase of the clinical trial	Pathology
ACY-241	NCT02400242	Ia/Ib	Multiple myeloma
	NCT02935790	Ib	Stage III and IV unresectable melanoma
	NCT02551185	Ib	Advanced solid tumors
	NCT02635061	Ib	Non-resectable non-small cell lung cancer
ACY-1215	NCT02632071	Ib	Unresectable or metastatic breast cancer
	NCT02787369	Ib	Relapsed chronic lymphocytic leukemia
	NCT02091063	Ib/II	Relapsed or refractory lymphoid malignancies
	NCT01997840	Ib/II	Recurrent and refractory multiple myeloma
	NCT01583283	I/II	Multiple myeloma recurrent or recurrent and refractory
	NCT02189343	Ib	Recurrent and refractory multiple myeloma
	NCT01323751	I/II	Multiple myeloma recurrent or recurrent and refractory
	NCT02856568	Ib	Unresectable or metastatic cholangiocarcinoma
	NCT02661815	Ib	Ovarian cancer, primary peritoneal cancer or platinum-resistant fallopian tubes


It is important to note that inactivation of HDAC6 protein in mice does not result in abnormal development or major organ problems [167], suggesting that HDAC6 inhibition would have few side effects, unlike pan-HDAC inhibitors.

## 5. HDAC6 in CML

Several studies have demonstrated the influence of HDAC6 in neurodegenerative, cardiovascular and renal diseases, as well as in inflammation [175] and viral response [84]. The role of the HDAC6 protein in cancer is also now well better understood. Although its oncogenic or tumor suppressor potential is dependent on the type of cancer [68], its involvement in oncogenic cell transformation, tumor development, and cancer immunity regulation makes a strong therapeutic candidate [113]. The overexpression of HDAC6 in many cancers, and in particular in several types of leukemia, led researchers to test the effects of HDAC6 inhibitors on these cancers. Despite the observation of a moderate overexpression of the HDAC6 protein in urothelial cancerous tissues, the inhibition of the protein had limited efficacy compared to the use of inhibitors targeting several HDACs [176]. On the other hand, the inhibition of HDAC6 has notable anti-cancer properties in the context of multiple myeloma [177], chronic lymphocytic leukemia [114], and acute myeloid leukemia [178], as well as solid cancers including prostate cancer [137], breast cancer [179], melanoma [139], and ovarian cancer [102]. Here, we will focus on the involvement of HDAC6 in CML.

### 5.1. Degradation of BCR-ABL via deacetylation of HSP90 $\alpha$ by HDAC6

Although little research exists on HDAC6 in the context of CML, this protein has a function that makes it particularly interesting in the context of such pathology. HDAC6 deacetylates HSP90 $\alpha$ , which is involved in the stabilization BCR-ABL [180]. In the acetylated form, HSP90 $\alpha$  loses its chaperone function, which leads to the degradation of its client proteins by the proteasome (Figure 6A). The importance of the acetylation status of HSP90 $\alpha$  in the protein degradation of BCR-ABL makes HDAC6 inhibitors potentially promising molecules for the treatment of CML. Pan-HDAC inhibitors are capable of inducing the inhibition of HDAC6, as well as the downregulation of HDAC6 using si-RNA, which increases the acetylation of HSP90 $\alpha$ , and in turn increases the ubiquitination of the BCR-ABL protein, decreasing its expression in K562 cells [181,182].



**Figure 6.** Role of HDAC6 in chronic myeloid leukemia. (A) HDAC6 is implicated in proteasome-dependent protein degradation by its interaction with HSP90 $\alpha$ , a chaperone protein which, when deacetylated by HDAC6, is involved in the stabilization of BCR-ABL. In its acetylated (Ac) form, HSP90 $\alpha$  loses its chaperone function, which leads to ubiquitination (Ub) and subsequent degradation of BCR-ABL by the proteasome. (B) Imatinib-insensitive chronic myeloid leukemia (CML) leukemic stem cells (LSCs) overexpress HDAC6 compared to CML cells.

### 5.2. Overexpression of HDAC6 in CML stem cells

LSCs that are not targeted by TKI and are characterized by a capacity for self-renewal play a crucial role in CML relapse. Although HDAC6 is necessary for the repression of genes involved in the differentiation targeted by the Tip60-p400 complex in embryonic stem cells (ESCs) [183], no study has provided evidence for this in LSCs, more differentiated. In contrast, studies have shown that several proteins in the HDAC family are overexpressed in LSCs of CML. Indeed, SIRT1 is activated by BCR-ABL *via* STAT5 and its expression is increased in LSCs compared to in CML cells [184]. Finally, overexpression of isoforms of HDAC (HDAC1, HDAC2, HDAC3, HDAC4, and HDAC5) and in particular HDAC6 was more frequently observed in LSCs (CD34<sup>+</sup> CD38<sup>-</sup>) isolated from patients with CML than in K562 cells [117] (Figure 6B), making it a protein of interest in the search for treatments to prevent relapse in patients with CML.

## 6. Conclusion

Despite the success of imatinib and TKIs, the emergence of resistance and the presence of LSCs that are unaffected by these treatments and are responsible for relapse mean that the development of new treatments for CML remains urgent. HDACs represent potential therapeutic targets for the treatment of CML [181,182,185]. Inhibitors of the HDAC family have promising results in combination with TKIs [186], some of which also inhibit HDAC6 [187-189]. The combination of HDAC class I inhibitors [190] or HDAC inhibitors including HDAC6 [191] and TKIs induces elimination of the quiescent LSCs that are not eradicated during treatment with imatinib alone. Although a decrease in HDAC6 expression, an increase in HSP90 $\alpha$  acetylation, and a decrease in BCR-ABL expression were observed in imatinib-resistant K562 cells, these cells still showed sensitivity to SAHA, an inhibitor of HDACs including HDAC6 [180]. However, the high toxicity of HDAC inhibitors and their many side effects [177] has necessitated targeting of particular HDAC isoforms.

The targeting of HDAC6 appears promising because it is overexpressed in many types of cancers, including certain leukemia subtypes. A recent study showed that dasatinib inhibits the phosphorylation of BCR-ABL without inducing the death of LSCs, suggesting that LSCs use signaling pathways that are activated by the BCR-ABL protein and are independent of its tyrosine kinase activity. The simple inhibition of the tyrosine kinase activity of the BCR-ABL protein is therefore not sufficient [192]. Since HDAC6 is overexpressed in CML stem cells and that its inhibition can potentially cause BCR-ABL degradation [68], this deacetylase appears as a strong candidate for CML treatment. Accordingly, HDAC6 inhibitors could be further tested in combination with TKIs or other molecules capable of targeting BCR-ABL such as bafetinib, rebastinib, tozasertib, danusertib, HG-7-85-01, GNF-2 and -5 molecules, and 1, 3, 4 thiadiazole derivatives, to potentially reduce resistance to treatment [193]. In the future, further development of selective PROTAC E3 ubiquitin ligase degraders triggering HDAC6 degradation [194] will provide further therapeutic options against various types of cancer, including CML.

**Supplementary Materials:** No supplementary materials.

**Author Contributions:** All authors contributed to writing and edition of the manuscript.

**Funding:** HL was supported by the “Recherche Cancer et Sang” foundation. LBMCC was also supported the “Recherches Scientifiques Luxembourg” association, by the “Een Häerz fir kriibskrank Kanner” association, by the Action LIONS “Vaincre le Cancer” association, and by Télémie Luxembourg. MD was also supported by National Research Foundation (NRF) [Grant Number 019R1A2C1009231] and by a grant from the MEST of Korea for Tumor Microenvironment Global Core Research Center (GCRC) [Grant Number 2011-0030001]. Support from Brain Korea (BK21) PLUS program and Creative-Pioneering Researchers Program at Seoul National University [Funding number: 370C-20160062] are acknowledged.

**Conflicts of Interest:** The authors declare no conflict of interest.

## References

- Hoffmann, V.S.; Baccarani, M.; Hasford, J.; Castagnetti, F.; Di Raimondo, F.; Casado, L.F.; Turkina, A.; Zackova, D.; Ossenkoppele, G.; Zaritskey, A., et al. Treatment and outcome of 2904 CML patients from the EUTOS population-based registry. *Leukemia* **2017**, *31*, 593-601, doi:10.1038/leu.2016.246.
- Siegel, R.L.; Miller, K.D.; Jemal, A. Cancer statistics, 2018. *CA: a cancer journal for clinicians* **2018**, *68*, 7-30, doi:10.3322/caac.21442.
- Hoffmann, V.S.; Baccarani, M.; Hasford, J.; Lindoerfer, D.; Burgstaller, S.; Sertic, D.; Costeas, P.; Mayer, J.; Indrak, K.; Everaus, H., et al. The EUTOS population-based registry: incidence and clinical characteristics of 2904 CML patients in 20 European Countries. *Leukemia* **2015**, *29*, 1336-1343, doi:10.1038/leu.2015.73.
- Apperley, J.F. Part I: mechanisms of resistance to imatinib in chronic myeloid leukaemia. *The Lancet. Oncology* **2007**, *8*, 1018-1029, doi:10.1016/S1470-2045(07)70342-X.
- Apperley, J.F. Chronic myeloid leukaemia. *Lancet* **2015**, *385*, 1447-1459, doi:10.1016/S0140-6736(13)62120-0.
- Nowell, P.C. Discovery of the Philadelphia chromosome: a personal perspective. *The Journal of clinical investigation* **2007**, *117*, 2033-2035, doi:10.1172/JCI31771.
- Kang, Z.J.; Liu, Y.F.; Xu, L.Z.; Long, Z.J.; Huang, D.; Yang, Y.; Liu, B.; Feng, J.X.; Pan, Y.J.; Yan, J.S., et al. The Philadelphia chromosome in leukemogenesis. *Chinese journal of cancer* **2016**, *35*, 48, doi:10.1186/s40880-016-0108-0.
- Colicelli, J. ABL tyrosine kinases: evolution of function, regulation, and specificity. *Science signaling* **2010**, *3*, re6, doi:10.1126/scisignal.3139re6.
- Khatri, A.; Wang, J.; Pendergast, A.M. Multifunctional Abl kinases in health and disease. *Journal of cell science* **2016**, *129*, 9-16, doi:10.1242/jcs.175521.
- Wang, J.Y. The capable ABL: what is its biological function? *Molecular and cellular biology* **2014**, *34*, 1188-1197, doi:10.1128/MCB.01454-13.
- Laurent, E.; Talpaz, M.; Kantarjian, H.; Kurzrock, R. The BCR gene and philadelphia chromosome-positive leukemogenesis. *Cancer research* **2001**, *61*, 2343-2355.
- Drummond, M.W.; Lush, C.J.; Vickers, M.A.; Reid, F.M.; Kaeda, J.; Holyoake, T.L. Imatinib mesylate-induced molecular remission of Philadelphia chromosome-positive myelodysplastic syndrome. *Leukemia* **2003**, *17*, 463-465, doi:10.1038/sj.leu.2402814.
- Schultheis, B.; Wang, L.; Clark, R.E.; Melo, J.V. BCR-ABL with an e6a2 fusion in a CML patient diagnosed in blast crisis. *Leukemia* **2003**, *17*, 2054-2055, doi:10.1038/sj.leu.2403079.
- Vergilio, J.; Bagg, A. Myeloproliferative disorders and myelodysplastic syndromes. In *Molecular Pathology in Clinical Practice*, Leonard, D.G.B., Ed. Springer New York: 2007; pp. 383-396.
- Withey, J.M.; Marley, S.B.; Kaeda, J.; Harvey, A.J.; Crompton, M.R.; Gordon, M.Y. Targeting primary human leukaemia cells with RNA interference: Bcr-Abl targeting inhibits myeloid progenitor self-renewal in chronic myeloid leukaemia cells. *British journal of haematology* **2005**, *129*, 377-380, doi:10.1111/j.1365-2141.2005.05468.x.
- Beran, M.; Pisa, P.; O'Brien, S.; Kurzrock, R.; Siciliano, M.; Cork, A.; Andersson, B.S.; Kohli, V.; Kantarjian, H. Biological properties and growth in SCID mice of a new myelogenous leukemia cell line (KBM-5) derived from chronic myelogenous leukemia cells in the blastic phase. *Cancer research* **1993**, *53*, 3603-3610.

17. Yaghmaie, M.; Ghaffari, S.H.; Ghavamzadeh, A.; Alimoghaddam, K.; Jahani, M.; Mousavi, S.A.; Irvani, M.; Bahar, B.; Bibordi, I. Frequency of BCR-ABL fusion transcripts in Iranian patients with chronic myeloid leukemia. *Archives of Iranian medicine* **2008**, *11*, 247-251, doi:08113/AIM.003.
18. van der Velden, V.H.; Beverloo, H.B.; Hoogeveen, P.G.; Zwaan Ch, M. A novel BCR-ABL fusion transcript (e18a2) in a child with chronic myeloid leukemia. *Leukemia* **2007**, *21*, 833-835, doi:10.1038/sj.leu.2404580.
19. Hochhaus, A.; Reiter, A.; Skladny, H.; Melo, J.V.; Sick, C.; Berger, U.; Guo, J.Q.; Arlinghaus, R.B.; Hehlmann, R.; Goldman, J.M., et al. A novel BCR-ABL fusion gene (e6a2) in a patient with Philadelphia chromosome-negative chronic myelogenous leukemia. *Blood* **1996**, *88*, 2236-2240.
20. O'Hare, T.; Deininger, M.W.; Eide, C.A.; Clackson, T.; Druker, B.J. Targeting the BCR-ABL signaling pathway in therapy-resistant Philadelphia chromosome-positive leukemia. *Clinical cancer research : an official journal of the American Association for Cancer Research* **2011**, *17*, 212-221, doi:10.1158/1078-0432.CCR-09-3314.
21. Dai, Z.; Quackenbush, R.C.; Courtney, K.D.; Grove, M.; Cortez, D.; Reuther, G.W.; Pendergast, A.M. Oncogenic Abl and Src tyrosine kinases elicit the ubiquitin-dependent degradation of target proteins through a Ras-independent pathway. *Genes & development* **1998**, *12*, 1415-1424, doi:10.1101/gad.12.10.1415.
22. Los, M. Tumor Growth and Cell Proliferation. In *The Impact of Tumor Biology on Cancer Treatment and Multidisciplinary Strategies*, Molls, M., Vaupel, P., Nieder, C., Anscher, M.S., Eds. Springer-Verlag Berlin Heidelberg: 2009; 10.1007/978-3-540-74386-6.
23. Reuther, G.W.; Fu, H.; Cripe, L.D.; Collier, R.J.; Pendergast, A.M. Association of the protein kinases c-Bcr and Bcr-Abl with proteins of the 14-3-3 family. *Science (New York, N.Y.)* **1994**, *266*, 129-133, doi:10.1126/science.7939633.
24. Goldman, J.M.; Melo, J.V. Chronic myeloid leukemia--advances in biology and new approaches to treatment. *The New England journal of medicine* **2003**, *349*, 1451-1464, doi:10.1056/NEJMra020777.
25. Goss, V.L.; Lee, K.A.; Moritz, A.; Nardone, J.; Spek, E.J.; MacNeill, J.; Rush, J.; Comb, M.J.; Polakiewicz, R.D. A common phosphotyrosine signature for the Bcr-Abl kinase. *Blood* **2006**, *107*, 4888-4897, doi:10.1182/blood-2005-08-3399.
26. Morotti, A.; Carra, G.; Panuzzo, C.; Crivellaro, S.; Taulli, R.; Guerrasio, A.; Saglio, G. Protein Kinase CK2: A Targetable BCR-ABL Partner in Philadelphia Positive Leukemias. *Advances in hematology* **2015**, *2015*, 612567, doi:10.1155/2015/612567.
27. Shin, S.; Lee, Y.; Kim, W.; Ko, H.; Choi, H.; Kim, K. Caspase-2 primes cancer cells for TRAIL-mediated apoptosis by processing procaspase-8. *The EMBO journal* **2005**, *24*, 3532-3542, doi:10.1038/sj.emboj.7600827.
28. St-Denis, N.A.; Derksen, D.R.; Litchfield, D.W. Evidence for regulation of mitotic progression through temporal phosphorylation and dephosphorylation of CK2alpha. *Molecular and cellular biology* **2009**, *29*, 2068-2081, doi:10.1128/MCB.01563-08.
29. Feller, S.M. Crk family adaptors-signalling complex formation and biological roles. *Oncogene* **2001**, *20*, 6348-6371, doi:10.1038/sj.onc.1204779.
30. Mashima, R.; Hishida, Y.; Tezuka, T.; Yamanashi, Y. The roles of Dok family adapters in immunoreceptor signaling. *Immunological reviews* **2009**, *232*, 273-285, doi:10.1111/j.1600-065X.2009.00844.x.



31. Lionberger, J.M.; Smithgall, T.E. The c-Fes protein-tyrosine kinase suppresses cytokine-independent outgrowth of myeloid leukemia cells induced by Bcr-Abl. *Cancer research* **2000**, *60*, 1097-1103.
32. Shi, C.S.; Tuscano, J.; Kehrl, J.H. Adaptor proteins CRK and CRKL associate with the serine/threonine protein kinase GCKR promoting GCKR and SAPK activation. *Blood* **2000**, *95*, 776-782.
33. Puil, L.; Liu, J.; Gish, G.; Mbamalu, G.; Bowtell, D.; Pelicci, P.G.; Arlinghaus, R.; Pawson, T. Bcr-Abl oncoproteins bind directly to activators of the Ras signalling pathway. *The EMBO journal* **1994**, *13*, 764-773.
34. Li, S.; Couvillon, A.D.; Brasher, B.B.; Van Etten, R.A. Tyrosine phosphorylation of Grb2 by Bcr/Abl and epidermal growth factor receptor: a novel regulatory mechanism for tyrosine kinase signaling. *The EMBO journal* **2001**, *20*, 6793-6804, doi:10.1093/emboj/20.23.6793.
35. Frietsch, J.J.; Kastner, C.; Grunewald, T.G.; Schweigel, H.; Nollau, P.; Ziermann, J.; Clement, J.H.; La Rosee, P.; Hochhaus, A.; Butt, E. LASP1 is a novel BCR-ABL substrate and a phosphorylation-dependent binding partner of CRKL in chronic myeloid leukemia. *Oncotarget* **2014**, *5*, 5257-5271, doi:10.18632/oncotarget.2072.
36. Orth, M.F.; Cazes, A.; Butt, E.; Grunewald, T.G. An update on the LIM and SH3 domain protein 1 (LASP1): a versatile structural, signaling, and biomarker protein. *Oncotarget* **2015**, *6*, 26-42, doi:10.18632/oncotarget.3083.
37. Liang, X.; Hajivandi, M.; Veach, D.; Wisniewski, D.; Clarkson, B.; Resh, M.D.; Pope, R.M. Quantification of change in phosphorylation of BCR-ABL kinase and its substrates in response to Imatinib treatment in human chronic myelogenous leukemia cells. *Proteomics* **2006**, *6*, 4554-4564, doi:10.1002/pmic.200600109.
38. Lopez-Colome, A.M.; Lee-Rivera, I.; Benavides-Hidalgo, R.; Lopez, E. Paxillin: a crossroad in pathological cell migration. *Journal of hematology & oncology* **2017**, *10*, 50, doi:10.1186/s13045-017-0418-y.
39. Wang, Y.; Tomar, A.; George, S.P.; Khurana, S. Obligatory role for phospholipase C-gamma(1) in villin-induced epithelial cell migration. *American journal of physiology. Cell physiology* **2007**, *292*, C1775-1786, doi:10.1152/ajpcell.00420.2006.
40. Kuchay, S.; Duan, S.; Schenkein, E.; Peschiaroli, A.; Saraf, A.; Florens, L.; Washburn, M.P.; Pagano, M. FBXL2- and PTPL1-mediated degradation of p110-free p85beta regulatory subunit controls the PI(3)K signalling cascade. *Nature cell biology* **2013**, *15*, 472-480, doi:10.1038/ncb2731.
41. Roy, A.; Ye, J.; Deng, F.; Wang, Q.J. Protein kinase D signaling in cancer: A friend or foe? *Biochim Biophys Acta Rev Cancer* **2017**, *1868*, 283-294, doi:10.1016/j.bbcan.2017.05.008.
42. Grimmmler, M.; Wang, Y.; Mund, T.; Cilensek, Z.; Keidel, E.M.; Waddell, M.B.; Jakel, H.; Kullmann, M.; Kriwacki, R.W.; Hengst, L. Cdk-inhibitory activity and stability of p27Kip1 are directly regulated by oncogenic tyrosine kinases. *Cell* **2007**, *128*, 269-280, doi:10.1016/j.cell.2006.11.047.
43. Pamonsinlapatham, P.; Hadj-Slimane, R.; Lepelletier, Y.; Allain, B.; Toccafondi, M.; Garbay, C.; Raynaud, F. p120-Ras GTPase activating protein (RasGAP): a multi-interacting protein in downstream signaling. *Biochimie* **2009**, *91*, 320-328, doi:10.1016/j.biochi.2008.10.010.
44. Kim, D.H.; Sarbassov, D.D.; Ali, S.M.; King, J.E.; Latek, R.R.; Erdjument-Bromage, H.; Tempst, P.; Sabatini, D.M. mTOR interacts with raptor to form a nutrient-sensitive complex that signals to the cell growth machinery. *Cell* **2002**, *110*, 163-175, doi:10.1016/s0092-8674(02)00808-5.
45. Audero, E.; Cascone, I.; Maniero, F.; Napione, L.; Arese, M.; Lanfranccone, L.; Bussolino, F. Adaptor ShcA protein binds tyrosine kinase Tie2 receptor and regulates migration and sprouting but not



- survival of endothelial cells. *The Journal of biological chemistry* **2004**, *279*, 13224-13233, doi:10.1074/jbc.M307456200.
46. Freeburn, R.W.; Wright, K.L.; Burgess, S.J.; Astoul, E.; Cantrell, D.A.; Ward, S.G. Evidence that SHIP-1 contributes to phosphatidylinositol 3,4,5-trisphosphate metabolism in T lymphocytes and can regulate novel phosphoinositide 3-kinase effectors. *Journal of immunology (Baltimore, Md. : 1950)* **2002**, *169*, 5441-5450, doi:10.4049/jimmunol.169.10.5441.
  47. Berger, A.; Hoelbl-Kovacic, A.; Bourgeais, J.; Hoefling, L.; Warsch, W.; Grundschober, E.; Uras, I.Z.; Menzl, I.; Putz, E.M.; Hoermann, G., et al. PAK-dependent STAT5 serine phosphorylation is required for BCR-ABL-induced leukemogenesis. *Leukemia* **2014**, *28*, 629-641, doi:10.1038/leu.2013.351.
  48. Rahikainen, R.; von Essen, M.; Schaefer, M.; Qi, L.; Azizi, L.; Kelly, C.; Ihalainen, T.O.; Wehrle-Haller, B.; Bastmeyer, M.; Huang, C., et al. Mechanical stability of talin rod controls cell migration and substrate sensing. *Scientific reports* **2017**, *7*, 3571, doi:10.1038/s41598-017-03335-2.
  49. La, S.H.; Kim, S.J.; Kang, H.G.; Lee, H.W.; Chun, K.H. Ablation of human telomerase reverse transcriptase (hTERT) induces cellular senescence in gastric cancer through a galectin-3 dependent mechanism. *Oncotarget* **2016**, *7*, 57117-57130, doi:10.18632/oncotarget.10986.
  50. Zitvogel, L.; Rusakiewicz, S.; Routy, B.; Ayyoub, M.; Kroemer, G. Immunological off-target effects of imatinib. *Nature reviews. Clinical oncology* **2016**, *13*, 431-446, doi:10.1038/nrclinonc.2016.41.
  51. Sorel, N.; Cayssials, E.; Brizard, F.; Chomel, J.C. Treatment and molecular monitoring update in chronic myeloid leukemia management. *Annales de biologie clinique* **2017**, *75*, 129-145, doi:10.1684/abc.2017.1233.
  52. Hantschel, O. Structure, regulation, signaling, and targeting of abl kinases in cancer. *Genes & cancer* **2012**, *3*, 436-446, doi:10.1177/1947601912458584.
  53. Modugno, M. New resistance mechanisms for small molecule kinase inhibitors of Abl kinase. *Drug discovery today. Technologies* **2014**, *11*, 5-10, doi:10.1016/j.ddtec.2013.12.001.
  54. An, X.; Tiwari, A.K.; Sun, Y.; Ding, P.R.; Ashby, C.R., Jr.; Chen, Z.S. BCR-ABL tyrosine kinase inhibitors in the treatment of Philadelphia chromosome positive chronic myeloid leukemia: a review. *Leukemia research* **2010**, *34*, 1255-1268, doi:10.1016/j.leukres.2010.04.016.
  55. Ankathil, R.; Azlan, H.; Dzarr, A.A.; Baba, A.A. Pharmacogenetics and the treatment of chronic myeloid leukemia: how relevant clinically? An update. *Pharmacogenomics* **2018**, *19*, 475-393, doi:10.2217/pgs-2017-0193.
  56. Danisz, K.; Blasiak, J. Role of anti-apoptotic pathways activated by BCR/ABL in the resistance of chronic myeloid leukemia cells to tyrosine kinase inhibitors. *Acta biochimica Polonica* **2013**, *60*, 503-514.
  57. Alikian, M.; Gale, R.P.; Apperley, J.F.; Foroni, L. Molecular techniques for the personalised management of patients with chronic myeloid leukaemia. *Biomolecular detection and quantification* **2017**, *11*, 4-20, doi:10.1016/j.bdq.2017.01.001.
  58. Yang, K.; Fu, L.W. Mechanisms of resistance to BCR-ABL TKIs and the therapeutic strategies: A review. *Critical reviews in oncology/hematology* **2015**, *93*, 277-292, doi:10.1016/j.critrevonc.2014.11.001.
  59. Soverini, S.; Mancini, M.; Bavaro, L.; Cavo, M.; Martinelli, G. Chronic myeloid leukemia: the paradigm of targeting oncogenic tyrosine kinase signaling and counteracting resistance for successful cancer therapy. *Molecular cancer* **2018**, *17*, 49, doi:10.1186/s12943-018-0780-6.
  60. Keskin, D.; Sadri, S.; Eskazan, A.E. Dasatinib for the treatment of chronic myeloid leukemia: patient selection and special considerations. *Drug design, development and therapy* **2016**, *10*, 3355-3361, doi:10.2147/DDDT.S85050.

61. Emole, J.; Talabi, T.; Pinilla-Ibarz, J. Update on the management of Philadelphia chromosome positive chronic myelogenous leukemia: role of nilotinib. *Biologics : targets & therapy* **2016**, *10*, 23-31, doi:10.2147/BTT.S67844.
62. Ault, P.S.; Rose Pharm, D.J.; Nodzon Ph, D.L.; Kaled, E.S. Bosutinib Therapy in Patients With Chronic Myeloid Leukemia: Practical Considerations for Management of Side Effects. *Journal of the advanced practitioner in oncology* **2016**, *7*, 160-175, doi:10.6004/jadpro.2016.7.2.3.
63. Bu, Q.; Cui, L.; Li, J.; Du, X.; Zou, W.; Ding, K.; Pan, J. SAHA and S116836, a novel tyrosine kinase inhibitor, synergistically induce apoptosis in imatinib-resistant chronic myelogenous leukemia cells. *Cancer biology & therapy* **2014**, *15*, 951-962, doi:10.4161/cbt.28931.
64. Jabbour, E.J.; Cortes, J.E.; Kantarjian, H.M. Resistance to tyrosine kinase inhibition therapy for chronic myelogenous leukemia: a clinical perspective and emerging treatment options. *Clinical lymphoma, myeloma & leukemia* **2013**, *13*, 515-529, doi:10.1016/j.clml.2013.03.018.
65. Balabanov, S.; Braig, M.; Brummendorf, T.H. Current aspects in resistance against tyrosine kinase inhibitors in chronic myelogenous leukemia. *Drug discovery today. Technologies* **2014**, *11*, 89-99, doi:10.1016/j.ddtec.2014.03.003.
66. Etienne, G.; Dulucq, S.; Huguet, F.; Schmitt, A.; Lascaux, A.; Hayette, S.; Fort, M.P.; Sujobert, P.; Bijou, F.; Morisset, S., et al. Incidence and outcome of BCR-ABL mutated chronic myeloid leukemia patients who failed to tyrosine kinase inhibitors. *Cancer medicine* **2019**, *8*, 5173-5182, doi:10.1002/cam4.2410.
67. Bixby, D.; Talpaz, M. Mechanisms of resistance to tyrosine kinase inhibitors in chronic myeloid leukemia and recent therapeutic strategies to overcome resistance. *Hematology / the Education Program of the American Society of Hematology. American Society of Hematology. Education Program* **2009**, 10.1182/asheducation-2009.1.461, 461-476, doi:10.1182/asheducation-2009.1.461.
68. Seidel, C.; Schneckeburger, M.; Dicato, M.; Diederich, M. Histone deacetylase 6 in health and disease. *Epigenomics* **2015**, *7*, 103-118, doi:10.2217/epi.14.69.
69. Linev, A.J.; Ivanov, H.J.; Zhelyazkov, I.G.; Ivanova, H.; Goranova-Marinova, V.S.; Stoyanova, V.K. Mutations Associated with Imatinib Mesylate Resistance - Review. *Folia medica* **2018**, *60*, 617-623, doi:10.2478/folmed-2018-0030.
70. Holohan, C.; Van Schaeybroeck, S.; Longley, D.B.; Johnston, P.G. Cancer drug resistance: an evolving paradigm. *Nature reviews. Cancer* **2013**, *13*, 714-726, doi:10.1038/nrc3599.
71. Loscocco, F.; Visani, G.; Galimberti, S.; Curti, A.; Isidori, A. BCR-ABL Independent Mechanisms of Resistance in Chronic Myeloid Leukemia. *Frontiers in oncology* **2019**, *9*, 939, doi:10.3389/fonc.2019.00939.
72. Fachi, M.M.; Tonin, F.S.; Leonart, L.P.; Aguiar, K.S.; Lenzi, L.; Figueiredo, B.C.; Fernandez-Llimos, F.; Pontarolo, R. Comparative efficacy and safety of tyrosine kinase inhibitors for chronic myeloid leukaemia: A systematic review and network meta-analysis. *European journal of cancer* **2018**, *104*, 9-20, doi:10.1016/j.ejca.2018.08.016.
73. Zabriskie, M.S.; Vellore, N.A.; Gantz, K.C.; Deininger, M.W.; O'Hare, T. Radotinib is an effective inhibitor of native and kinase domain-mutant BCR-ABL1. *Leukemia* **2015**, *29*, 1939-1942, doi:10.1038/leu.2015.42.
74. Tang, L.; Zhang, H.; Peng, Y.Z.; Li, C.G.; Jiang, H.W.; Xu, M.; Mei, H.; Hu, Y. Comparative efficacy and tolerability of front-line treatments for newly diagnosed chronic-phase chronic myeloid leukemia: an update network meta-analysis. *BMC cancer* **2019**, *19*, 849, doi:10.1186/s12885-019-6039-9.

75. Park, E.; Jue, M.S. Radotinib-induced eruptive melanocytic nevi in patient with chronic myeloid leukemia: a case report and literature review. *Annals of hematology* **2019**, *98*, 533-535, doi:10.1007/s00277-018-3444-0.
76. Losson, H.; Schnekenburger, M.; Dicato, M.; Diederich, M. Natural Compound Histone Deacetylase Inhibitors (HDACi): Synergy with Inflammatory Signaling Pathway Modulators and Clinical Applications in Cancer. *Molecules (Basel, Switzerland)* **2016**, *21*, doi:10.3390/molecules21111608.
77. Finnin, M.S.; Donigian, J.R.; Cohen, A.; Richon, V.M.; Rifkind, R.A.; Marks, P.A.; Breslow, R.; Pavletich, N.P. Structures of a histone deacetylase homologue bound to the TSA and SAHA inhibitors. *Nature* **1999**, *401*, 188-193, doi:10.1038/43710.
78. Carafa, V.; Rotili, D.; Forgione, M.; Cuomo, F.; Serrettiello, E.; Hailu, G.S.; Jarho, E.; Lahtela-Kakkonen, M.; Mai, A.; Altucci, L. Sirtuin functions and modulation: from chemistry to the clinic. *Clinical epigenetics* **2016**, *8*, 61, doi:10.1186/s13148-016-0224-3.
79. Mei, Z.; Zhang, X.; Yi, J.; Huang, J.; He, J.; Tao, Y. Sirtuins in metabolism, DNA repair and cancer. *Journal of experimental & clinical cancer research : CR* **2016**, *35*, 182, doi:10.1186/s13046-016-0461-5.
80. Thiagalingam, S.; Cheng, K.H.; Lee, H.J.; Mineva, N.; Thiagalingam, A.; Ponte, J.F. Histone deacetylases: unique players in shaping the epigenetic histone code. *Annals of the New York Academy of Sciences* **2003**, *983*, 84-100, doi:10.1111/j.1749-6632.2003.tb05964.x.
81. de Ruijter, A.J.; van Gennip, A.H.; Caron, H.N.; Kemp, S.; van Kuilenburg, A.B. Histone deacetylases (HDACs): characterization of the classical HDAC family. *The Biochemical journal* **2003**, *370*, 737-749, doi:10.1042/BJ20021321.
82. Bertos, N.R.; Gilquin, B.; Chan, G.K.; Yen, T.J.; Khochbin, S.; Yang, X.J. Role of the tetradecapeptide repeat domain of human histone deacetylase 6 in cytoplasmic retention. *The Journal of biological chemistry* **2004**, *279*, 48246-48254, doi:10.1074/jbc.M408583200.
83. Li, Y.; Shin, D.; Kwon, S.H. Histone deacetylase 6 plays a role as a distinct regulator of diverse cellular processes. *The FEBS journal* **2013**, *280*, 775-793, doi:10.1111/febs.12079.
84. Zheng, K.; Jiang, Y.; He, Z.; Kitazato, K.; Wang, Y. Cellular defence or viral assist: the dilemma of HDAC6. *The Journal of general virology* **2017**, *98*, 322-337, doi:10.1099/jgv.0.000679.
85. Mortenson, J.B.; Heppler, L.N.; Banks, C.J.; Weerasekara, V.K.; Whited, M.D.; Piccolo, S.R.; Johnson, W.E.; Thompson, J.W.; Andersen, J.L. Histone deacetylase 6 (HDAC6) promotes the pro-survival activity of 14-3-3zeta via deacetylation of lysines within the 14-3-3zeta binding pocket. *The Journal of biological chemistry* **2015**, *290*, 12487-12496, doi:10.1074/jbc.M114.607580.
86. Zhang, L.; Liu, S.; Liu, N.; Zhang, Y.; Liu, M.; Li, D.; Seto, E.; Yao, T.P.; Shui, W.; Zhou, J. Proteomic identification and functional characterization of MYH9, Hsc70, and DNAJA1 as novel substrates of HDAC6 deacetylase activity. *Protein & cell* **2015**, *6*, 42-54, doi:10.1007/s13238-014-0102-8.
87. Wu, J.Y.; Xiang, S.; Zhang, M.; Fang, B.; Huang, H.; Kwon, O.K.; Zhao, Y.; Yang, Z.; Bai, W.; Bepler, G., et al. Histone deacetylase 6 (HDAC6) deacetylates extracellular signal-regulated kinase 1 (ERK1) and thereby stimulates ERK1 activity. *The Journal of biological chemistry* **2018**, *293*, 1976-1993, doi:10.1074/jbc.M117.795955.
88. Beier, U.H.; Wang, L.; Han, R.; Akimova, T.; Liu, Y.; Hancock, W.W. Histone deacetylases 6 and 9 and sirtuin-1 control Foxp3+ regulatory T cell function through shared and isoform-specific mechanisms. *Science signaling* **2012**, *5*, ra45, doi:10.1126/scisignal.2002873.
89. Salian-Mehta, S.; Xu, M.; McKinsey, T.A.; Tobet, S.; Wierman, M.E. Novel Interaction of Class IIb Histone Deacetylase 6 (HDAC6) with Class IIa HDAC9 Controls Gonadotropin Releasing Hormone

- (GnRH) Neuronal Cell Survival and Movement. *The Journal of biological chemistry* **2015**, *290*, 14045-14056, doi:10.1074/jbc.M115.640482.
90. Gao, L.; Cueto, M.A.; Asselbergs, F.; Atadja, P. Cloning and functional characterization of HDAC11, a novel member of the human histone deacetylase family. *The Journal of biological chemistry* **2002**, *277*, 25748-25755, doi:10.1074/jbc.M111871200.
  91. Medler, T.R.; Craig, J.M.; Fiorillo, A.A.; Feeney, Y.B.; Harrell, J.C.; Clevenger, C.V. HDAC6 Deacetylates HMG2 to Regulate Stat5a Activity and Breast Cancer Growth. *Molecular cancer research : MCR* **2016**, *14*, 994-1008, doi:10.1158/1541-7786.MCR-16-0109.
  92. Chang, Y.W.; Tseng, C.F.; Wang, M.Y.; Chang, W.C.; Lee, C.C.; Chen, L.T.; Hung, M.C.; Su, J.L. Deacetylation of HSPA5 by HDAC6 leads to GP78-mediated HSPA5 ubiquitination at K447 and suppresses metastasis of breast cancer. *Oncogene* **2016**, *35*, 1517-1528, doi:10.1038/onc.2015.214.
  93. Yang, M.H.; Laurent, G.; Bause, A.S.; Spang, R.; German, N.; Haigis, M.C.; Haigis, K.M. HDAC6 and SIRT2 regulate the acetylation state and oncogenic activity of mutant K-RAS. *Molecular cancer research : MCR* **2013**, *11*, 1072-1077, doi:10.1158/1541-7786.MCR-13-0040-T.
  94. Liu, K.P.; Zhou, D.; Ouyang, D.Y.; Xu, L.H.; Wang, Y.; Wang, L.X.; Pan, H.; He, X.H. LC3B-II deacetylation by histone deacetylase 6 is involved in serum-starvation-induced autophagic degradation. *Biochemical and biophysical research communications* **2013**, *441*, 970-975, doi:10.1016/j.bbrc.2013.11.007.
  95. Zhang, M.; Xiang, S.; Joo, H.Y.; Wang, L.; Williams, K.A.; Liu, W.; Hu, C.; Tong, D.; Haakenson, J.; Wang, C., et al. HDAC6 deacetylates and ubiquitinates MSH2 to maintain proper levels of MutSalp. *Molecular cell* **2014**, *55*, 31-46, doi:10.1016/j.molcel.2014.04.028.
  96. Perkins, A.; Nelson, K.J.; Parsonage, D.; Poole, L.B.; Karplus, P.A. Peroxiredoxins: guardians against oxidative stress and modulators of peroxide signaling. *Trends in biochemical sciences* **2015**, *40*, 435-445, doi:10.1016/j.tibs.2015.05.001.
  97. Parmigiani, R.B.; Xu, W.S.; Venta-Perez, G.; Erdjument-Bromage, H.; Yaneva, M.; Tempst, P.; Marks, P.A. HDAC6 is a specific deacetylase of peroxiredoxins and is involved in redox regulation. *Proceedings of the National Academy of Sciences of the United States of America* **2008**, *105*, 9633-9638, doi:10.1073/pnas.0803749105.
  98. Moreno-Gonzalo, O.; Mayor, F., Jr.; Sanchez-Madrid, F. HDAC6 at Crossroads of Infection and Innate Immunity. *Trends in immunology* **2018**, *39*, 591-595, doi:10.1016/j.it.2018.05.004.
  99. Nakka, K.K.; Chaudhary, N.; Joshi, S.; Bhat, J.; Singh, K.; Chatterjee, S.; Malhotra, R.; De, A.; Santra, M.K.; Dilworth, F.J., et al. Nuclear matrix-associated protein SMAR1 regulates alternative splicing via HDAC6-mediated deacetylation of Sam68. *Proceedings of the National Academy of Sciences of the United States of America* **2015**, *112*, E3374-3383, doi:10.1073/pnas.1418603112.
  100. Huo, L.; Li, D.; Sun, X.; Shi, X.; Karna, P.; Yang, W.; Liu, M.; Qiao, W.; Aneja, R.; Zhou, J. Regulation of Tat acetylation and transactivation activity by the microtubule-associated deacetylase HDAC6. *The Journal of biological chemistry* **2011**, *286*, 9280-9286, doi:10.1074/jbc.M110.208884.
  101. Matsuyama, A.; Shimazu, T.; Sumida, Y.; Saito, A.; Yoshimatsu, Y.; Seigneurin-Berny, D.; Osada, H.; Komatsu, Y.; Nishino, N.; Khochbin, S., et al. In vivo destabilization of dynamic microtubules by HDAC6-mediated deacetylation. *The EMBO journal* **2002**, *21*, 6820-6831, doi:10.1093/emboj/cdf682.
  102. Haakenson, J.; Zhang, X. HDAC6 and ovarian cancer. *International journal of molecular sciences* **2013**, *14*, 9514-9535, doi:10.3390/ijms14059514.

103. Fan, S.J.; Huang, F.I.; Liou, J.P.; Yang, C.R. The novel histone deacetylase 6 inhibitor, MPT0G211, ameliorates tau phosphorylation and cognitive deficits in an Alzheimer's disease model. *Cell death & disease* **2018**, *9*, 655, doi:10.1038/s41419-018-0688-5.
104. Lee, S.W.; Yang, J.; Kim, S.Y.; Jeong, H.K.; Lee, J.; Kim, W.J.; Lee, E.J.; Kim, H.S. MicroRNA-26a induced by hypoxia targets HDAC6 in myogenic differentiation of embryonic stem cells. *Nucleic acids research* **2015**, *43*, 2057-2073, doi:10.1093/nar/gkv088.
105. Lafarga, V.; Aymerich, I.; Tapia, O.; Mayor, F., Jr.; Penela, P. A novel GRK2/HDAC6 interaction modulates cell spreading and motility. *The EMBO journal* **2012**, *31*, 856-869, doi:10.1038/emboj.2011.466.
106. Williams, K.A.; Zhang, M.; Xiang, S.; Hu, C.; Wu, J.Y.; Zhang, S.; Ryan, M.; Cox, A.D.; Der, C.J.; Fang, B., et al. Extracellular signal-regulated kinase (ERK) phosphorylates histone deacetylase 6 (HDAC6) at serine 1035 to stimulate cell migration. *The Journal of biological chemistry* **2013**, *288*, 33156-33170, doi:10.1074/jbc.M113.472506.
107. Di Fulvio, S.; Azakir, B.A.; Therrien, C.; Sinnreich, M. Dysferlin interacts with histone deacetylase 6 and increases alpha-tubulin acetylation. *PloS one* **2011**, *6*, e28563, doi:10.1371/journal.pone.0028563.
108. Tala, Sun, X.; Chen, J.; Zhang, L.; Liu, N.; Zhou, J.; Li, D.; Liu, M. Microtubule stabilization by Mdp3 is partially attributed to its modulation of HDAC6 in addition to its association with tubulin and microtubules. *PloS one* **2014**, *9*, e90932, doi:10.1371/journal.pone.0090932.
109. Yan, J.; Seibenhener, M.L.; Calderilla-Barbosa, L.; Diaz-Meco, M.T.; Moscat, J.; Jiang, J.; Wooten, M.W.; Wooten, M.C. SQSTM1/p62 interacts with HDAC6 and regulates deacetylase activity. *PloS one* **2013**, *8*, e76016, doi:10.1371/journal.pone.0076016.
110. Salemi, L.M.; Almawi, A.W.; Lefebvre, K.J.; Schild-Poulter, C. Aggresome formation is regulated by RanBPM through an interaction with HDAC6. *Biology open* **2014**, *3*, 418-430, doi:10.1242/bio.20147021.
111. Perez, M.; Santa-Maria, I.; Gomez de Barreda, E.; Zhu, X.; Cuadros, R.; Cabrero, J.R.; Sanchez-Madrid, F.; Dawson, H.N.; Vitek, M.P.; Perry, G., et al. Tau--an inhibitor of deacetylase HDAC6 function. *Journal of neurochemistry* **2009**, *109*, 1756-1766, doi:10.1111/j.1471-4159.2009.06102.x.
112. Schofield, A.V.; Gamell, C.; Bernard, O. Tubulin polymerization promoting protein 1 (TPPP1) increases beta-catenin expression through inhibition of HDAC6 activity in U2OS osteosarcoma cells. *Biochemical and biophysical research communications* **2013**, *436*, 571-577, doi:10.1016/j.bbrc.2013.05.076.
113. Li, T.; Zhang, C.; Hassan, S.; Liu, X.; Song, F.; Chen, K.; Zhang, W.; Yang, J. Histone deacetylase 6 in cancer. *Journal of hematology & oncology* **2018**, *11*, 111, doi:10.1186/s13045-018-0654-9.
114. Maharaj, K.; Powers, J.J.; Achille, A.; Deng, S.; Fonseca, R.; Pabon-Saldana, M.; Quayle, S.N.; Jones, S.S.; Villagra, A.; Sotomayor, E.M., et al. Silencing of HDAC6 as a therapeutic target in chronic lymphocytic leukemia. *Blood advances* **2018**, *2*, 3012-3024, doi:10.1182/bloodadvances.2018020065.
115. Li, A.; Chen, P.; Leng, Y.; Kang, J. Histone deacetylase 6 regulates the immunosuppressive properties of cancer-associated fibroblasts in breast cancer through the STAT3-COX2-dependent pathway. *Oncogene* **2018**, *37*, 5952-5966, doi:10.1038/s41388-018-0379-9.
116. Qian, H.; Chen, Y.; Nian, Z.; Su, L.; Yu, H.; Chen, F.J.; Zhang, X.; Xu, W.; Zhou, L.; Liu, J., et al. HDAC6-mediated acetylation of lipid droplet-binding protein CIDEC regulates fat-induced lipid storage. *The Journal of clinical investigation* **2017**, *127*, 1353-1369, doi:10.1172/JCI85963.
117. Bamodu, O.A.; Kuo, K.T.; Yuan, L.P.; Cheng, W.H.; Lee, W.H.; Ho, Y.S.; Chao, T.Y.; Yeh, C.T. HDAC inhibitor suppresses proliferation and tumorigenicity of drug-resistant chronic myeloid leukemia stem cells through regulation of hsa-miR-196a targeting BCR/ABL1. *Experimental cell research* **2018**, *370*, 519-530, doi:10.1016/j.yexcr.2018.07.017.



118. Cosenza, M.; Pozzi, S. The Therapeutic Strategy of HDAC6 Inhibitors in Lymphoproliferative Disease. *International journal of molecular sciences* **2018**, *19*, doi:10.3390/ijms19082337.
119. Dehmel, F.; Weinbrenner, S.; Julius, H.; Ciossek, T.; Maier, T.; Stengel, T.; Fettis, K.; Burkhardt, C.; Wieland, H.; Beckers, T. Trithiocarbonates as a novel class of HDAC inhibitors: SAR studies, isoenzyme selectivity, and pharmacological profiles. *Journal of medicinal chemistry* **2008**, *51*, 3985-4001, doi:10.1021/jm800093c.
120. Inks, E.S.; Josey, B.J.; Jesinkey, S.R.; Chou, C.J. A novel class of small molecule inhibitors of HDAC6. *ACS chemical biology* **2012**, *7*, 331-339, doi:10.1021/cb200134p.
121. Schafer, S.; Saunders, L.; Eliseeva, E.; Velena, A.; Jung, M.; Schwienhorst, A.; Strasser, A.; Dickmanns, A.; Ficner, R.; Schlimme, S., et al. Phenylalanine-containing hydroxamic acids as selective inhibitors of class IIb histone deacetylases (HDACs). *Bioorganic & medicinal chemistry* **2008**, *16*, 2011-2033, doi:10.1016/j.bmc.2007.10.092.
122. Schafer, S.; Saunders, L.; Schlimme, S.; Valkov, V.; Wagner, J.M.; Kratz, F.; Sippl, W.; Verdin, E.; Jung, M. Pyridylalanine-containing hydroxamic acids as selective HDAC6 inhibitors. *ChemMedChem* **2009**, *4*, 283-290, doi:10.1002/cmdc.200800196.
123. Jochems, J.; Boulden, J.; Lee, B.G.; Blendy, J.A.; Jarpe, M.; Mazitschek, R.; Van Duzer, J.H.; Jones, S.; Berton, O. Antidepressant-like properties of novel HDAC6-selective inhibitors with improved brain bioavailability. *Neuropsychopharmacology : official publication of the American College of Neuropsychopharmacology* **2014**, *39*, 389-400, doi:10.1038/npp.2013.207.
124. Porter, N.J.; Mahendran, A.; Breslow, R.; Christianson, D.W. Unusual zinc-binding mode of HDAC6-selective hydroxamate inhibitors. *Proceedings of the National Academy of Sciences of the United States of America* **2017**, *114*, 13459-13464, doi:10.1073/pnas.1718823114.
125. Krukowski, K.; Ma, J.; Golonzhka, O.; Laumet, G.O.; Gutti, T.; van Duzer, J.H.; Mazitschek, R.; Jarpe, M.B.; Heijnen, C.J.; Kavelaars, A. HDAC6 inhibition effectively reverses chemotherapy-induced peripheral neuropathy. *Pain* **2017**, *158*, 1126-1137, doi:10.1097/j.pain.0000000000000893.
126. Strebl, M.G.; Campbell, A.J.; Zhao, W.N.; Schroeder, F.A.; Riley, M.M.; Chindavong, P.S.; Morin, T.M.; Haggarty, S.J.; Wagner, F.F.; Ritter, T., et al. HDAC6 Brain Mapping with [(18)F]BavaroSTAT Enabled by a Ru-Mediated Deoxyfluorination. *ACS central science* **2017**, *3*, 1006-1014, doi:10.1021/acscentsci.7b00274.
127. Wagner, F.F.; Olson, D.E.; Gale, J.P.; Kaya, T.; Weiwer, M.; Aidoud, N.; Thomas, M.; Davoine, E.L.; Lemerrier, B.C.; Zhang, Y.L., et al. Potent and selective inhibition of histone deacetylase 6 (HDAC6) does not require a surface-binding motif. *Journal of medicinal chemistry* **2013**, *56*, 1772-1776, doi:10.1021/jm301355j.
128. Kaliszczak, M.; Trousil, S.; Aberg, O.; Perumal, M.; Nguyen, Q.D.; Aboagye, E.O. A novel small molecule hydroxamate preferentially inhibits HDAC6 activity and tumour growth. *British journal of cancer* **2013**, *108*, 342-350, doi:10.1038/bjc.2012.576.
129. Kozikowski, A.P.; Tapadar, S.; Luchini, D.N.; Kim, K.H.; Billadeau, D.D. Use of the nitrile oxide cycloaddition (NOC) reaction for molecular probe generation: a new class of enzyme selective histone deacetylase inhibitors (HDACIs) showing picomolar activity at HDAC6. *Journal of medicinal chemistry* **2008**, *51*, 4370-4373, doi:10.1021/jm8002894.
130. Sixto-Lopez, Y.; Bello, M.; Rodriguez-Fonseca, R.A.; Rosales-Hernandez, M.C.; Martinez-Archundia, M.; Gomez-Vidal, J.A.; Correa-Basurto, J. Searching the conformational complexity and binding properties of HDAC6 through docking and molecular dynamic simulations. *Journal of biomolecular structure & dynamics* **2017**, *35*, 2794-2814, doi:10.1080/07391102.2016.1231084.

131. Huang, P.; Almeciga-Pinto, I.; Jarpe, M.; van Duzer, J.H.; Mazitschek, R.; Yang, M.; Jones, S.S.; Quayle, S.N. Selective HDAC inhibition by ACY-241 enhances the activity of paclitaxel in solid tumor models. *Oncotarget* **2017**, *8*, 2694-2707, doi:10.18632/oncotarget.13738.
132. Olsen, C.A.; Ghadiri, M.R. Discovery of potent and selective histone deacetylase inhibitors via focused combinatorial libraries of cyclic alpha3beta-tetrapeptides. *Journal of medicinal chemistry* **2009**, *52*, 7836-7846, doi:10.1021/jm900850t.
133. Chen, Y.; Lopez-Sanchez, M.; Savoy, D.N.; Billadeau, D.D.; Dow, G.S.; Kozikowski, A.P. A series of potent and selective, triazolylphenyl-based histone deacetylases inhibitors with activity against pancreatic cancer cells and *Plasmodium falciparum*. *Journal of medicinal chemistry* **2008**, *51*, 3437-3448, doi:10.1021/jm701606b.
134. Diedrich, D.; Hamacher, A.; Gertzen, C.G.; Alves Avelar, L.A.; Reiss, G.J.; Kurz, T.; Gohlke, H.; Kassack, M.U.; Hansen, F.K. Rational design and diversity-oriented synthesis of peptoid-based selective HDAC6 inhibitors. *Chemical communications (Cambridge, England)* **2016**, *52*, 3219-3222, doi:10.1039/c5cc10301k.
135. Lin, X.; Chen, W.; Qiu, Z.; Guo, L.; Zhu, W.; Li, W.; Wang, Z.; Zhang, W.; Zhang, Z.; Rong, Y., et al. Design and synthesis of orally bioavailable aminopyrrolidinone histone deacetylase 6 inhibitors. *Journal of medicinal chemistry* **2015**, *58*, 2809-2820, doi:10.1021/jm502011f.
136. Blackburn, C.; Barrett, C.; Chin, J.; Garcia, K.; Gigstad, K.; Gould, A.; Gutierrez, J.; Harrison, S.; Hoar, K.; Lynch, C., et al. Potent histone deacetylase inhibitors derived from 4-(aminomethyl)-N-hydroxybenzamide with high selectivity for the HDAC6 isoform. *Journal of medicinal chemistry* **2013**, *56*, 7201-7211, doi:10.1021/jm400385r.
137. Seidel, C.; Schneckeburger, M.; Mazumder, A.; Teiten, M.H.; Kirsch, G.; Dicato, M.; Diederich, M. 4-Hydroxybenzoic acid derivatives as HDAC6-specific inhibitors modulating microtubular structure and HSP90alpha chaperone activity against prostate cancer. *Biochemical pharmacology* **2016**, *99*, 31-52, doi:10.1016/j.bcp.2015.11.005.
138. Tang, G.; Wong, J.C.; Zhang, W.; Wang, Z.; Zhang, N.; Peng, Z.; Zhang, Z.; Rong, Y.; Li, S.; Zhang, M., et al. Identification of a novel aminotetralin class of HDAC6 and HDAC8 selective inhibitors. *Journal of medicinal chemistry* **2014**, *57*, 8026-8034, doi:10.1021/jm5008962.
139. Wang, X.X.; Wan, R.Z.; Liu, Z.P. Recent advances in the discovery of potent and selective HDAC6 inhibitors. *European journal of medicinal chemistry* **2018**, *143*, 1406-1418, doi:10.1016/j.ejmech.2017.10.040.
140. Liang, T.; Hou, X.; Zhou, Y.; Yang, X.; Fang, H. Design, Synthesis, and Biological Evaluation of 2,4-Imidazolidinedione Derivatives as HDAC6 Isoform-Selective Inhibitors. *ACS medicinal chemistry letters* **2019**, *10*, 1122-1127, doi:10.1021/acsmchemlett.9b00084.
141. Kozikowski, A.P.; Chen, Y.; Gaysin, A.; Chen, B.; D'Annibale, M.A.; Suto, C.M.; Langley, B.C. Functional differences in epigenetic modulators-superiority of mercaptoacetamide-based histone deacetylase inhibitors relative to hydroxamates in cortical neuron neuroprotection studies. *Journal of medicinal chemistry* **2007**, *50*, 3054-3061, doi:10.1021/jm070178x.
142. Lee, H.Y.; Nepali, K.; Huang, F.I.; Chang, C.Y.; Lai, M.J.; Li, Y.H.; Huang, H.L.; Yang, C.R.; Liou, J.P. (N-Hydroxycarbonylbenzylamino)quinolines as Selective Histone Deacetylase 6 Inhibitors Suppress Growth of Multiple Myeloma in Vitro and in Vivo. *Journal of medicinal chemistry* **2018**, *61*, 905-917, doi:10.1021/acs.jmedchem.7b01404.
143. Shen, S.; Hadley, M.; Ustinova, K.; Pavlicek, J.; Knox, T.; Noonepalle, S.; Tavares, M.T.; Zimprich, C.A.; Zhang, G.; Robers, M.B., et al. Discovery of a New Isoxazole-3-hydroxamate-Based Histone Deacetylase



- 6 Inhibitor SS-208 with Antitumor Activity in Syngeneic Melanoma Mouse Models. *Journal of medicinal chemistry* **2019**, *62*, 8557-8577, doi:10.1021/acs.jmedchem.9b00946.
144. Vogerl, K.; Ong, N.; Senger, J.; Herp, D.; Schmidtkunz, K.; Marek, M.; Muller, M.; Bartel, K.; Shaik, T.B.; Porter, N.J., et al. Synthesis and Biological Investigation of Phenothiazine-Based Benzhydroxamic Acids as Selective Histone Deacetylase 6 Inhibitors. *Journal of medicinal chemistry* **2019**, *62*, 1138-1166, doi:10.1021/acs.jmedchem.8b01090.
145. Porter, N.J.; Osko, J.D.; Diedrich, D.; Kurz, T.; Hooker, J.M.; Hansen, F.K.; Christianson, D.W. Histone Deacetylase 6-Selective Inhibitors and the Influence of Capping Groups on Hydroxamate-Zinc Denticity. *Journal of medicinal chemistry* **2018**, *61*, 8054-8060, doi:10.1021/acs.jmedchem.8b01013.
146. Duan, W.; Li, J.; Inks, E.S.; Chou, C.J.; Jia, Y.; Chu, X.; Li, X.; Xu, W.; Zhang, Y. Design, synthesis, and antitumor evaluation of novel histone deacetylase inhibitors equipped with a phenylsulfonylfuroxan module as a nitric oxide donor. *Journal of medicinal chemistry* **2015**, *58*, 4325-4338, doi:10.1021/acs.jmedchem.5b00317.
147. Cho, M.; Choi, E.; Yang, J.S.; Lee, C.; Seo, J.J.; Kim, B.S.; Oh, S.J.; Kim, H.M.; Lee, K.; Park, S.K., et al. Discovery of pyridone-based histone deacetylase inhibitors: approaches for metabolic stability. *ChemMedChem* **2013**, *8*, 272-279, doi:10.1002/cmdc.201200529.
148. Liu, Y.M.; Lee, H.Y.; Lai, M.J.; Pan, S.L.; Huang, H.L.; Kuo, F.C.; Chen, M.C.; Liou, J.P. Pyrimidinedione-mediated selective histone deacetylase 6 inhibitors with antitumor activity in colorectal cancer HCT116 cells. *Organic & biomolecular chemistry* **2015**, *13*, 10226-10235, doi:10.1039/c5ob01509j.
149. Yu, C.W.; Chang, P.T.; Hsin, L.W.; Chern, J.W. Quinazolin-4-one derivatives as selective histone deacetylase-6 inhibitors for the treatment of Alzheimer's disease. *Journal of medicinal chemistry* **2013**, *56*, 6775-6791, doi:10.1021/jm400564j.
150. Heltweg, B.; Dequiedt, F.; Marshall, B.L.; Brauch, C.; Yoshida, M.; Nishino, N.; Verdin, E.; Jung, M. Subtype selective substrates for histone deacetylases. *Journal of medicinal chemistry* **2004**, *47*, 5235-5243, doi:10.1021/jm0497592.
151. Hideshima, T.; Qi, J.; Paranal, R.M.; Tang, W.; Greenberg, E.; West, N.; Colling, M.E.; Estiu, G.; Mazitschek, R.; Perry, J.A., et al. Discovery of selective small-molecule HDAC6 inhibitor for overcoming proteasome inhibitor resistance in multiple myeloma. *Proceedings of the National Academy of Sciences of the United States of America* **2016**, *113*, 13162-13167, doi:10.1073/pnas.1608067113.
152. Sellmer, A.; Stangl, H.; Beyer, M.; Grunstein, E.; Leonhardt, M.; Pongratz, H.; Eichhorn, E.; Elz, S.; Striegl, B.; Jenei-Lanzl, Z., et al. Marbostat-100 Defines a New Class of Potent and Selective Antiinflammatory and Antirheumatic Histone Deacetylase 6 Inhibitors. *Journal of medicinal chemistry* **2018**, *61*, 3454-3477, doi:10.1021/acs.jmedchem.7b01593.
153. Lee, H.Y.; Tsai, A.C.; Chen, M.C.; Shen, P.J.; Cheng, Y.C.; Kuo, C.C.; Pan, S.L.; Liu, Y.M.; Liu, J.F.; Yeh, T.K., et al. Azaindolylsulfonamides, with a more selective inhibitory effect on histone deacetylase 6 activity, exhibit antitumor activity in colorectal cancer HCT116 cells. *Journal of medicinal chemistry* **2014**, *57*, 4009-4022, doi:10.1021/jm401899x.
154. Mackwitz, M.K.W.; Hamacher, A.; Osko, J.D.; Held, J.; Scholer, A.; Christianson, D.W.; Kassack, M.U.; Hansen, F.K. Multicomponent Synthesis and Binding Mode of Imidazo[1,2- a]pyridine-Capped Selective HDAC6 Inhibitors. *Organic letters* **2018**, *20*, 3255-3258, doi:10.1021/acs.orglett.8b01118.
155. Lee, J.H.; Yao, Y.; Mahendran, A.; Ngo, L.; Venta-Perez, G.; Choy, M.L.; Breslow, R.; Marks, P.A. Creation of a histone deacetylase 6 inhibitor and its biological effects [corrected]. *Proceedings of the*

- National Academy of Sciences of the United States of America* **2015**, *112*, 12005-12010, doi:10.1073/pnas.1515882112.
156. Lee, J.H.; Mahendran, A.; Yao, Y.; Ngo, L.; Venta-Perez, G.; Choy, M.L.; Kim, N.; Ham, W.S.; Breslow, R.; Marks, P.A. Development of a histone deacetylase 6 inhibitor and its biological effects. *Proceedings of the National Academy of Sciences of the United States of America* **2013**, *110*, 15704-15709, doi:10.1073/pnas.1313893110.
  157. Hai, Y.; Christianson, D.W. Histone deacetylase 6 structure and molecular basis of catalysis and inhibition. *Nature chemical biology* **2016**, *12*, 741-747, doi:10.1038/nchembio.2134.
  158. Yang, Z.; Wang, T.; Wang, F.; Niu, T.; Liu, Z.; Chen, X.; Long, C.; Tang, M.; Cao, D.; Wang, X., et al. Discovery of Selective Histone Deacetylase 6 Inhibitors Using the Quinazoline as the Cap for the Treatment of Cancer. *Journal of medicinal chemistry* **2016**, *59*, 1455-1470, doi:10.1021/acs.jmedchem.5b01342.
  159. Bergman, J.A.; Woan, K.; Perez-Villarroel, P.; Villagra, A.; Sotomayor, E.M.; Kozikowski, A.P. Selective histone deacetylase 6 inhibitors bearing substituted urea linkers inhibit melanoma cell growth. *Journal of medicinal chemistry* **2012**, *55*, 9891-9899, doi:10.1021/jm301098e.
  160. Senger, J.; Melesina, J.; Marek, M.; Romier, C.; Oehme, I.; Witt, O.; Sippl, W.; Jung, M. Synthesis and Biological Investigation of Oxazole Hydroxamates as Highly Selective Histone Deacetylase 6 (HDAC6) Inhibitors. *Journal of medicinal chemistry* **2016**, *59*, 1545-1555, doi:10.1021/acs.jmedchem.5b01493.
  161. Santo, L.; Hideshima, T.; Kung, A.L.; Tseng, J.C.; Tamang, D.; Yang, M.; Jarpe, M.; van Duzer, J.H.; Mazitschek, R.; Ogier, W.C., et al. Preclinical activity, pharmacodynamic, and pharmacokinetic properties of a selective HDAC6 inhibitor, ACY-1215, in combination with bortezomib in multiple myeloma. *Blood* **2012**, *119*, 2579-2589, doi:10.1182/blood-2011-10-387365.
  162. Wang, F.; Zhong, B.W.; Zhao, Z.R. ACY 1215, a histone deacetylase 6 inhibitor, inhibits cancer cell growth in melanoma. *Journal of biological regulators and homeostatic agents* **2018**, *32*, 851-858.
  163. Zhang, I.; Beus, M.; Stochaj, U.; Le, P.U.; Zorc, B.; Rajic, Z.; Petrecca, K.; Maysinger, D. Inhibition of glioblastoma cell proliferation, invasion, and mechanism of action of a novel hydroxamic acid hybrid molecule. *Cell death discovery* **2018**, *4*, 41, doi:10.1038/s41420-018-0103-0.
  164. Dong, J.; Zheng, N.; Wang, X.; Tang, C.; Yan, P.; Zhou, H.B.; Huang, J. A novel HDAC6 inhibitor exerts an anti-cancer effect by triggering cell cycle arrest and apoptosis in gastric cancer. *European journal of pharmacology* **2018**, *828*, 67-79, doi:10.1016/j.ejphar.2018.03.026.
  165. Butler, K.V.; Kalin, J.; Brochier, C.; Vistoli, G.; Langley, B.; Kozikowski, A.P. Rational design and simple chemistry yield a superior, neuroprotective HDAC6 inhibitor, tubastatin A. *Journal of the American Chemical Society* **2010**, *132*, 10842-10846, doi:10.1021/ja102758v.
  166. Aldana-Masangkay, G.I.; Rodriguez-Gonzalez, A.; Lin, T.; Ikeda, A.K.; Hsieh, Y.T.; Kim, Y.M.; Lomenick, B.; Okemoto, K.; Landaw, E.M.; Wang, D., et al. Tubacin suppresses proliferation and induces apoptosis of acute lymphoblastic leukemia cells. *Leukemia & lymphoma* **2011**, *52*, 1544-1555, doi:10.3109/10428194.2011.570821.
  167. Depetter, Y.; Geurs, S.; De Vreese, R.; Goethals, S.; Vandoorn, E.; Laevens, A.; Steenbrugge, J.; Meyer, E.; de Tullio, P.; Bracke, M., et al. Selective pharmacological inhibitors of HDAC6 reveal biochemical activity but functional tolerance in cancer models. *International journal of cancer* **2019**, *145*, 735-747, doi:10.1002/ijc.32169.

168. Patil, V.; Sodji, Q.H.; Kornacki, J.R.; Mrksich, M.; Oyeler, A.K. 3-Hydroxypyridin-2-thione as novel zinc binding group for selective histone deacetylase inhibition. *Journal of medicinal chemistry* **2013**, *56*, 3492-3506, doi:10.1021/jm301769u.
169. Muthyala, R.; Shin, W.S.; Xie, J.; Sham, Y.Y. Discovery of 1-hydroxypyridine-2-thiones as selective histone deacetylase inhibitors and their potential application for treating leukemia. *Bioorganic & medicinal chemistry letters* **2015**, *25*, 4320-4324, doi:10.1016/j.bmcl.2015.07.065.
170. Itoh, Y.; Suzuki, T.; Kouketsu, A.; Suzuki, N.; Maeda, S.; Yoshida, M.; Nakagawa, H.; Miyata, N. Design, synthesis, structure--selectivity relationship, and effect on human cancer cells of a novel series of histone deacetylase 6-selective inhibitors. *Journal of medicinal chemistry* **2007**, *50*, 5425-5438, doi:10.1021/jm7009217.
171. Segretti, M.C.; Vallerini, G.P.; Brochier, C.; Langley, B.; Wang, L.; Hancock, W.W.; Kozikowski, A.P. Thiol-Based Potent and Selective HDAC6 Inhibitors Promote Tubulin Acetylation and T-Regulatory Cell Suppressive Function. *ACS medicinal chemistry letters* **2015**, *6*, 1156-1161, doi:10.1021/acsmedchemlett.5b00303.
172. Wahhab, A.; Smil, D.; Ajamian, A.; Allan, M.; Chantigny, Y.; Therrien, E.; Nguyen, N.; Manku, S.; Leit, S.; Rahil, J., et al. Sulfamides as novel histone deacetylase inhibitors. *Bioorganic & medicinal chemistry letters* **2009**, *19*, 336-340, doi:10.1016/j.bmcl.2008.11.081.
173. Choi, E.W.; Song, J.W.; Ha, N.; Choi, Y.I.; Kim, S. CKD-506, a novel HDAC6-selective inhibitor, improves renal outcomes and survival in a mouse model of systemic lupus erythematosus. *Scientific reports* **2018**, *8*, 17297, doi:10.1038/s41598-018-35602-1.
174. Schnekenburger, M.; Florean, C.; Dicato, M.; Diederich, M. Epigenetic alterations as a universal feature of cancer hallmarks and a promising target for personalized treatments. *Current topics in medicinal chemistry* **2016**, *16*, 745-776, doi:10.2174/1568026615666150825141330.
175. Batchu, S.N.; Brijmohan, A.S.; Advani, A. The therapeutic hope for HDAC6 inhibitors in malignancy and chronic disease. *Clinical science (London, England : 1979)* **2016**, *130*, 987-1003, doi:10.1042/CS20160084.
176. Rosik, L.; Niegisch, G.; Fischer, U.; Jung, M.; Schulz, W.A.; Hoffmann, M.J. Limited efficacy of specific HDAC6 inhibition in urothelial cancer cells. *Cancer biology & therapy* **2014**, *15*, 742-757, doi:10.4161/cbt.28469.
177. Brindisi, M.; Saraswati, A.P.; Brogi, S.; Gemma, S.; Butini, S.; Campiani, G. Old but Gold: Tracking the New Guise of Histone Deacetylase 6 (HDAC6) Enzyme as a Biomarker and Therapeutic Target in Rare Diseases. *Journal of medicinal chemistry* **2019**, doi:10.1021/acs.jmedchem.9b00924, doi:10.1021/acs.jmedchem.9b00924.
178. Hackanson, B.; Rimmele, L.; Benkisser, M.; Abdelkarim, M.; Fliegau, M.; Jung, M.; Lubbert, M. HDAC6 as a target for antileukemic drugs in acute myeloid leukemia. *Leukemia research* **2012**, *36*, 1055-1062, doi:10.1016/j.leukres.2012.02.026.
179. Hsieh, Y.L.; Tu, H.J.; Pan, S.L.; Liou, J.P.; Yang, C.R. Anti-metastatic activity of MPT0G211, a novel HDAC6 inhibitor, in human breast cancer cells in vitro and in vivo. *Biochimica et biophysica acta. Molecular cell research* **2019**, *1866*, 992-1003, doi:10.1016/j.bbamcr.2019.03.003.
180. Kramer, O.H.; Mahboobi, S.; Sellmer, A. Drugging the HDAC6-HSP90 interplay in malignant cells. *Trends in pharmacological sciences* **2014**, *35*, 501-509, doi:10.1016/j.tips.2014.08.001.
181. Bali, P.; Pranpat, M.; Bradner, J.; Balasis, M.; Fiskus, W.; Guo, F.; Rocha, K.; Kumaraswamy, S.; Boyapalle, S.; Atadja, P., et al. Inhibition of histone deacetylase 6 acetylates and disrupts the chaperone

- function of heat shock protein 90: a novel basis for antileukemia activity of histone deacetylase inhibitors. *The Journal of biological chemistry* **2005**, *280*, 26729-26734, doi:10.1074/jbc.C500186200.
182. Rao, R.; Fiskus, W.; Yang, Y.; Lee, P.; Joshi, R.; Fernandez, P.; Mandawat, A.; Atadja, P.; Bradner, J.E.; Bhalla, K. HDAC6 inhibition enhances 17-AAG--mediated abrogation of hsp90 chaperone function in human leukemia cells. *Blood* **2008**, *112*, 1886-1893, doi:10.1182/blood-2008-03-143644.
  183. Chen, P.B.; Hung, J.H.; Hickman, T.L.; Coles, A.H.; Carey, J.F.; Weng, Z.; Chu, F.; Fazzio, T.G. Hdac6 regulates Tip60-p400 function in stem cells. *eLife* **2013**, *2*, e01557, doi:10.7554/eLife.01557.
  184. Kuo, Y.H.; Qi, J.; Cook, G.J. Regain control of p53: Targeting leukemia stem cells by isoform-specific HDAC inhibition. *Experimental hematology* **2016**, *44*, 315-321, doi:10.1016/j.exphem.2016.02.007.
  185. Lernoux, M.; Schnekenburger, M.; Dicato, M.; Diederich, M. Epigenetic mechanisms underlying the therapeutic effects of HDAC inhibitors in chronic myeloid leukemia. *Biochemical pharmacology* **2019**, 10.1016/j.bcp.2019.113698, 113698, doi:10.1016/j.bcp.2019.113698.
  186. Li, L.; Wang, L.; Li, L.; Wang, Z.; Ho, Y.; McDonald, T.; Holyoake, T.L.; Chen, W.; Bhatia, R. Activation of p53 by SIRT1 inhibition enhances elimination of CML leukemia stem cells in combination with imatinib. *Cancer cell* **2012**, *21*, 266-281, doi:10.1016/j.ccr.2011.12.020.
  187. Fiskus, W.; Pranpat, M.; Balasis, M.; Bali, P.; Estrella, V.; Kumaraswamy, S.; Rao, R.; Rocha, K.; Herger, B.; Lee, F., et al. Cotreatment with vorinostat (suberoylanilide hydroxamic acid) enhances activity of dasatinib (BMS-354825) against imatinib mesylate-sensitive or imatinib mesylate-resistant chronic myelogenous leukemia cells. *Clinical cancer research : an official journal of the American Association for Cancer Research* **2006**, *12*, 5869-5878, doi:10.1158/1078-0432.CCR-06-0980.
  188. Nimmanapalli, R.; Fuino, L.; Bali, P.; Gasparetto, M.; Glozak, M.; Tao, J.; Moscinski, L.; Smith, C.; Wu, J.; Jove, R., et al. Histone deacetylase inhibitor LAQ824 both lowers expression and promotes proteasomal degradation of Bcr-Abl and induces apoptosis of imatinib mesylate-sensitive or -refractory chronic myelogenous leukemia-blast crisis cells. *Cancer research* **2003**, *63*, 5126-5135.
  189. Nguyen, T.; Dai, Y.; Attkisson, E.; Kramer, L.; Jordan, N.; Nguyen, N.; Kolluri, N.; Muschen, M.; Grant, S. HDAC inhibitors potentiate the activity of the BCR/ABL kinase inhibitor KW-2449 in imatinib-sensitive or -resistant BCR/ABL+ leukemia cells in vitro and in vivo. *Clinical cancer research : an official journal of the American Association for Cancer Research* **2011**, *17*, 3219-3232, doi:10.1158/1078-0432.CCR-11-0234.
  190. Jin, Y.; Yao, Y.; Chen, L.; Zhu, X.; Jin, B.; Shen, Y.; Li, J.; Du, X.; Lu, Y.; Jiang, S., et al. Depletion of gamma-catenin by Histone Deacetylase Inhibition Confers Elimination of CML Stem Cells in Combination with Imatinib. *Theranostics* **2016**, *6*, 1947-1962, doi:10.7150/thno.16139.
  191. Zhang, B.; Strauss, A.C.; Chu, S.; Li, M.; Ho, Y.; Shiang, K.D.; Snyder, D.S.; Huettner, C.S.; Shultz, L.; Holyoake, T., et al. Effective targeting of quiescent chronic myelogenous leukemia stem cells by histone deacetylase inhibitors in combination with imatinib mesylate. *Cancer cell* **2010**, *17*, 427-442, doi:10.1016/j.ccr.2010.03.011.
  192. Zhang, H.; Li, S. Concise Review: Exploiting Unique Biological Features of Leukemia Stem Cells for Therapeutic Benefit. *Stem cells translational medicine* **2019**, *8*, 768-774, doi:10.1002/sctm.18-0247.
  193. Rossari, F.; Minutolo, F.; Orciuolo, E. Past, present, and future of Bcr-Abl inhibitors: from chemical development to clinical efficacy. *Journal of hematology & oncology* **2018**, *11*, 84, doi:10.1186/s13045-018-0624-2.

194. Yang, H.; Lv, W.; He, M.; Deng, H.; Li, H.; Wu, W.; Rao, Y. Plasticity in designing PROTACs for selective and potent degradation of HDAC6. *Chemical communications (Cambridge, England)* **2019**, *55*, 14848-14851, doi:10.1039/c9cc08509b.