

## Article

# Synthesis and *in vitro* photocytotoxicity of 9-/13-lipophilic substituted berberine derivatives as potential anticancer agents

Hong-Jhih Lin <sup>1,†</sup>, Jinn-Hsuan Ho <sup>2,†</sup>, Li-Chen Tsai <sup>1</sup>, Fang-Yu Yang <sup>1</sup>, Ling-Ling Yang <sup>3</sup>, Cheng-Deng Kuo <sup>4</sup>, Lih-Geeng Chen <sup>1</sup>, Yi-Wen Liu <sup>1</sup> and Jin-Yi Wu <sup>1,\*</sup>

<sup>1</sup> Department of Microbiology, Immunology and Biopharmaceuticals, College of Life Sciences, National Chiayi University, Chiayi 60004, Taiwan, ROC; s1030730@mail.ncyu.edu.tw (H.-J.L.); s1003500@mail.ncyu.edu.tw (L.-C.T.); s1060645@mail.ncyu.edu.tw (F.-Y.Y.); lgchen@mail.ncyu.edu.tw (L.-G.C.); ywlss@mail.ncyu.edu.tw (Y.-W.L.)

<sup>2</sup> Department of Chemical Engineering, College of Engineering, National Taiwan University of Science and Technology, Taipei 10607, Taiwan, ROC; jhho@mail.ntst.edu.tw (J.-H.H.)

<sup>3</sup> Department of Pharmacognosy, School of Pharmacy, College of Pharmacy, Taipei Medical University, Taipei 11031, Taiwan, ROC; llyang@tmu.edu.tw (L.-L.Y.)

<sup>4</sup> Thoracic Medicine Research Center, Changhua Christian Hospital, Changhua 50006, Taiwan, ROC; cdkuo23@gmail.com (C.-D.K.)

\* Correspondence: jywu@mail.ncyu.edu.tw; Tel.: +886-5-271-7925 (J.-Y.W.)

† These authors made equal contribution to this work.

**Abstract:** The objective of this study was to synthesize 9-/13-position substituted berberine derivatives and evaluated their cytotoxic and photocytotoxic effects against three human cancer cell lines. Among all the synthesized compounds, 9-*O*-dodecyl- (**5e**), 13-dodecyl- (**6e**) and 13-*O*-dodecyl-berberine (**7e**) exhibited stronger growth inhibition against three human cancer cell lines, (HepG2, HT-29 and BFTC905), in compare with structurally related berberine (**1**). These three compounds also showed the photocytotoxicity in human cancer cells in a concentration-dependent and light dose-dependent manner. Through flow cytometry analysis, we found out a lipophilic group at 9-/13-position of berberine may have facilitated its penetration into test cell and hence enhanced its photocytotoxicity on human liver cancer cell HepG2. Further, in cell cycle analysis, **5e**, **6e** and **7e** induced HepG2 cells to arrest at S phase and caused apoptosis upon irradiation. In addition, photodynamic treatment of berberine derivatives **5e**, **6e** and **7e** again showed a significant photocytotoxic effects on HepG2 cells, induced remarkable cell apoptosis, greatly increased intracellular ROS level and the loss of mitochondrial membrane potential. These results over and again confirmed that berberine derivatives **5e**, **6e** and **7e** greatly enhanced photocytotoxicity. Taking together, the test data led us to conclude that berberine derivatives with a dodecyl group at 9-/13-position could be great candidates for the anti-liver cancer medicines developments.

**Keywords:** berberine; lipophilic substituent; anti-cancer activity; photocytotoxicity; reactive oxygen species

## 1. Introduction


Although significant advances have been made in preventing and treating cancer, of which the mortality rate of remains to be unacceptably high. Meanwhile, many of the current chemotherapeutics are limited by significant side effects, unpredictable efficacies and/or acquired resistances [1-3].

Human hepatocellular carcinoma (HCC) and colorectal carcinoma are most common types of malignancy in sub-Saharan Africa and Southeast Asia like Taiwan and China. It is also the fifth most common and lethal cancer worldwide. Most of all, HCC is the most frequent malignant tumor and

the second leading cause of cancer-related death in Taiwan [4]. The hepatoma cells have high proliferative and metastasis rates. It usually treated by surgically removal followed by chemotherapy [5,6]. Doxorubicin is widely used in HCC chemotherapy. However, the agent failed to significantly improve the 5-year survival rate of the patients partly for their severe side effects, which, among others, include hepatic dysfunction and heart failure. As a group, we have been pursuing projects aim at developing novel cancer therapeutics that improve the prognosis of the disease with less adverse side effects.

Recent advance in molecular biology unravel the inside of mechanism, and pathogenesis and biology of cancer. The knowledge in turn affords drug designers target researches strategically to the cellular level and the development of novel cancer drugs. Reported herein the design, synthesis, and evaluation of a group of potential cancer chemotherapeutics.

Many natural products have been rich resources for potential drug development. Alkaloids, among which, are well-known nitrogen-containing group of diverse plant secondary metabolites [7]. Especially, among them, berberine (**1**, Fig. 1) is an isoquinoline alkaloid isolated from many medicinal herbs, such as *Coptis chinensis*, *Berberis aristata*, and *Phellodendron amurense*. It is widely used as traditional Chinese medicine [8-10]. Ayurvedic medicine in China and India [11]. It has a wide range of biochemical and pharmacological potencies including antibacterial, anti-inflammatory and anticancer effects [12,13]. Further, berberine has a long history of medical application, and was widely used as an anticancer drug to treat hepatoma [14], breast cancer [15], bladder cancer [16], and colon cancer [17]. Furthermore, berberine has significant cytotoxicity against human cancer cell lines by triggering in a significant increase G1 arrest in cell cycle phase of HepG2 cells [18]. It also induced cell cycle arrest at the G2/M phase and caused apoptosis of human gastric cancer SNU-5 cells [19]. However, owe to its hydrophilic nature, or lower bioavailability, berberine is absorbed poorly in intestines. Which results in a low efficacy on suppressing cancer cell growth [20]. Besides, the anticancer activity of berberine and its derivatives had recently been shown as inhibitor of topoisomerase I and II [21,22].



**Figure 1.** Chemical structures of berberine (**1**) and berberrubine (**2**).

In addition, recent study reported various analogs of berberine on 9-/13-position using different chain lengths and terminal amino groups were synthesized and evaluated on it the DNA-binding affinity or as G-quadruples stabilizing ligands [23-26]. Study have also showed that the introduction of a lipophilic substituent into 9-O-position of berberine backbone enhance the anti-proliferative activities against human cancer cell lines (HepG2 and HT-29) [27]. However, structural modification on the 13-position of berberine backbone has rarely been reported [28]. To explore its intestinal absorption and inhibitory function, in this study we modified berberine analogues and evaluated its photo-induced cytotoxicity *in vitro* on human liver HepG2, bladder BFTC905 and colon HT29 cancer cell lines. The type of functional groups, *n*-alkyl derivatives of berberine at 9- or 13-position was investigated for the structure-activity relationship (SAR).


## 2. Results and Discussion

### 2.1. Chemistry

In general, the desired berberine derivatives are synthesized with long chain *n*-alkyl bromide through 9-/13-alkylation. As shown in Scheme 1, in specific, synthesis of **5a–e**, **6a–e** and **7a–e** was

carried out by reacting berberrubine (**2**) with selected *n*-alkyl bromide in dry acetone or acetonitrile using anhydrous potassium carbonate ( $K_2CO_3$ ) as a base under nitrogen for 8–24 h according to the reported procedure [27]. Reduction of berberine (**1**) using sodium borohydride ( $NaBH_4$ ) gave intermediate **3** in 74% yield. The addition of long chain *n*-alkyl group to berberrubine (**2**) and 13-hydroxyberberine (**4**) gave the products of 9-/13-O-position derivatives in 40–86% yield. The compound **3** was reacted with *n*-alkyl aldehyde in an ethanol and acetic acid solvent mixture, and then acidified with 2 N HCl to yield the desired 13-position derivatives **6a–e** in 29–76% yield [28].

Further, the reaction mixture was refluxed for 8–24 h, and the progress of the reaction was monitored by TLC. The mixture was then cooled to room temperature and the solvent was removed under reduced pressure. The crude product was chromatographed on silica gel column and eluted with  $CH_2Cl_2/MeOH$  (15/1) solvent to afford the desired 9-/13-position berberine derivatives **5a–e**, **6a–e** and **7a–e** in quantitative yields (Scheme 1). All compounds were purified by flash column chromatography and were thoroughly characterized by  $^1H$ ,  $^{13}C$ , 2D NMR spectroscopy, liquid chromatography-mass spectroscopy and high-resolution mass spectroscopy. The full signal assignment of  $^1H$  and  $^{13}C$  NMR techniques including DEPT, COSY, HSQC, and HMBC analyses. All of the final compounds showed >98% purity by HPLC analysis.




**Scheme 1:** Reagents and conditions: (a)  $190^\circ C$ , 20–30 mmHg, 30–60 min; (b) *n*-alkyl bromide,  $K_2CO_3$ ,  $CH_3CN$ , reflux, 16–24 h; (c)  $NaBH_4$ ,  $K_2CO_3$ , MeOH,  $0^\circ C$ , 2 h; (d)  $R-CHO$ , EtOH, HOAc,  $85-95^\circ C$ , 5 h; then ether, HCl(aq), rt, 2 h; (e) MCPBA,  $CH_2Cl_2$ ,  $-20 \sim -30^\circ C$ , 2 h; then  $Na_2SO_3$ , rt, 1 h; (f) *n*-alkyl bromide, NaI,  $Et_3N$ , MeOH, reflux, 16–24 h.

## 2.2. Biological activity

### 2.2.1. In vitro anti-proliferative activity by MTT assay

The cytotoxic effect of the compounds was tested by the 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) assay [29] and rated by half maximal inhibitory concentration ( $IC_{50}$ ).  $IC_{50}$  is a measure of the effectiveness of a compound in inhibiting biological or biochemical function. The cytotoxic effects of the fifteen 9-O-, 13- or 13-O-position substituted

berberine derivatives were tested against three human cancer cell lines and one normal murine embryonic liver cell line. The cancer cells under study were HepG2 (liver carcinoma), HT29 (colon carcinoma), and BFTC905 (bladder carcinoma). As shown in Table 1, we found firstly, as a trend, the anticancer activity of these compounds increases with the alkyl chain length, or lipophilic characteristics, of the substitutes. Thus, the remarkable anticancer activity of the berberine derivatives with 9-/13-position substitution of the dodecyl moiety can be attributed to their enhanced bioavailability and cell membrane permeability through the increased lipophilicity.



**Figure 2.** *In vitro* cytotoxicity of berberine (**1**), **5e**, **6e** and **7e** in (A) HepG2, (B) BFTC905 and (C) HT29 for 24 h.

As it was shown in Fig. 2, that the lipophilicity effect of three bioactive derivatives; compounds **5e** (9-*O*-dodecylberberine), **6e** (13-dodecylberberine) and **7e** (13-*O*-decylberberine). In essence, all shows significant cell growth inhibition with the  $IC_{50}$  values of  $0.32 \pm 0.08$ ,  $0.77 \pm 0.18$  and  $0.83 \pm 0.30$  μM, respectively, against human liver carcinoma HepG2 cells. Compared to berberine (**1**)'s  $IC_{50}$  value of  $>40$  μM, these three derivatives evince 48- to 125-fold stronger cytotoxicity potency. In other words, through a simple structure-activity relationship (SAR) analysis we have shown that the cell growth inhibitory potency closely related to the chain length of *n*-alkyl groups. Meanwhile, to check potential side effect, we treated normal Madin-Darby Canine Kidney (MDCK) cell lines with berberine (**1**) and compounds **5e**, yet detected no significant cell death (data not shown). Furthermore, only a marked effect on cell death (under 20%) was observed at the maximum

concentration (4  $\mu\text{M}$ ) of **5e** and **6e** after 48 h treatment, except for berberine (**1**) which exhibited slight toxic. It appeared that compounds **5e**, **6e** and **7e** are cytotoxic to human colon cancer cells with no significant adverse effects on normal MDCK cells.

**Table 1.** Cytotoxicity of berberine and its derivatives against three human cancer cell lines for 24 h.

Compound	clog $P^a$	IC $_{50}$ ( $\mu\text{M}$ ) <sup>b</sup>		
		HepG2 <sup>c</sup>	BFTC905 <sup>d</sup>	HT29 <sup>e</sup>
Berberine ( <b>1</b> )	−0.77	> 40	26.11 $\pm$ 6.07	> 40
Berberrubine ( <b>2</b> )	−0.71	> 40	> 40	36.93 $\pm$ 9.60
<b>5a</b>	1.20	8.27 $\pm$ 1.69	6.40 $\pm$ 2.20	> 20
<b>5b</b>	2.26	2.83 $\pm$ 0.84	1.74 $\pm$ 0.22	14.70 $\pm$ 4.41
<b>5c</b>	3.32	0.59 $\pm$ 0.21	1.18 $\pm$ 0.31	3.88 $\pm$ 1.06
<b>5d</b>	4.37	0.61 $\pm$ 0.18	0.89 $\pm$ 0.11	2.79 $\pm$ 0.44
<b>5e</b>	6.49	0.32 $\pm$ 0.08	0.76 $\pm$ 0.02	1.34 $\pm$ 0.32
<b>6a</b>	1.31	8.11 $\pm$ 0.15	> 20	> 20
<b>6b</b>	2.37	2.19 $\pm$ 0.42	9.76 $\pm$ 1.02	8.45 $\pm$ 0.16
<b>6c</b>	3.81	1.19 $\pm$ 0.32	1.08 $\pm$ 0.23	2.77 $\pm$ 0.03
<b>6d</b>	4.87	0.67 $\pm$ 0.19	0.75 $\pm$ 0.19	1.76 $\pm$ 0.64
<b>6e</b>	5.54	0.77 $\pm$ 0.18	0.87 $\pm$ 0.15	1.39 $\pm$ 0.08
<b>7a</b>	1.37	8.21 $\pm$ 1.47	9.94 $\pm$ 1.70	> 20
<b>7b</b>	2.43	2.64 $\pm$ 0.31	2.09 $\pm$ 0.47	5.69 $\pm$ 0.66
<b>7c</b>	3.49	0.98 $\pm$ 0.26	1.00 $\pm$ 0.24	2.21 $\pm$ 0.21
<b>7d</b>	4.54	0.91 $\pm$ 0.40	0.96 $\pm$ 0.29	1.39 $\pm$ 0.19
<b>7e</b>	5.60	0.83 $\pm$ 0.30	0.78 $\pm$ 0.13	1.23 $\pm$ 0.06
Cisplatin	−1.68	82.12 $\pm$ 2.31	42.58 $\pm$ 1.16	53.28 $\pm$ 2.64

<sup>a</sup> Calculated log value of partition coefficient by ChemDraw Ultra 11.0. <sup>b</sup> Compound concentration ( $\mu\text{M}$ ) required to inhibit tumor cell proliferation rate by 50 %. Data are expressed as the mean  $\pm$  SD from the dose response curves of at 3–5 independent experiments. <sup>c</sup> Human hepatocarcinoma cell lines. <sup>d</sup> Human bladder cancer cell lines. <sup>e</sup> Human colon adenocarcinoma cell lines.


To confirm the findings above, we calculated partition coefficients (clog $P$ ) values of 9-/13-substituted berberine derivatives and compared it with the cytotoxic activity. As shown in Table 1, clog $P$  values correlate well with tested anticancer activity for 9-/13-substituted berberine derivatives. These results just once again verify that the lipophilicity or the chain length of the *n*-alkyl moieties of 9-/13-substituted berberine derivatives plays a very important role on their anticancer activity.

Further the positive clog $P$  values of berberine derivatives **5e**, **6e** and **7e** showed that the introduction of long chain *n*-alkyl groups to the berberine scaffold at the 9-*O*-, 13- and 13-*O*-position will moderately augment the lipophilicity of berberine, while *n*-alkyl substituents at the 9-*O*-, 13- and 13-*O*-positions of berberine remarkably increase the lipophilicity of berberine (Table 1). It worth noting also, berberine derivatives with moderate carbon chain length (**5e**, **6e** and **7e**) would significantly enhance its lipophilicity, and yield much more favorable transmembrane permeability. In general, clog $P$  value is a good index in estimating the distribution of drugs within the cells. Thus, hydrophobic drugs with high 1-octanol/water partition coefficients are preferentially distributed to hydrophobic compartments such as the lipid bilayers of cells while hydrophilic drugs (low

1-octanol/water partition coefficients) preferentially are found in aqueous compartments such as blood serum.

### 2.2.2. Absorption and emission spectra of berberine (1) and its derivatives (5e, 6e and 7e)

The absorption and fluorescence spectra of berberine (1) and its derivatives (5e, 6e and 7e) in MeOH were measured (Fig. 3). The absorption spectrum of the derivative 6e, with dodecyl group at 13-position of berberine, exhibited a moderate bathochromic shift (ca. 8.3 nm) than other compounds. Its absorptivity was obviously significantly decreased compared with that of berberine (1) and other derivatives. Fig. 3B showed the emission spectra of berberine (1) and its derivatives (5e, 6e and 7e) in MeOH (10  $\mu$ M). All gave the excitation wavelength at 420 nm with the emission maxima of 529, 531, 516, and 524 nm, respectively. The data suggested that compounds 6e and 7e have higher energy of the emission spectra in MeOH, which, in turn, may enhance its photocytotoxicity in photodynamic therapy.




**Figure 3.** (A) Electronic absorption spectra of berberine (1) and its derivatives (5e, 6e and 7e) in MeOH. (B) Emission spectral traces of the berberine (1) and its derivatives (5e, 6e and 7e) in MeOH [excitation wavelength ( $\lambda_{ex}$ ) are 420 nm].

### 2.2.3. Intracellular uptake of berberine (1) and its derivatives (5e, 6e and 7e)

Intracellular uptake of an anticancer agent is to check if the medicine would be able to target key organelles of the cell. Using time-dependent HPLC analysis we studied the cellular uptake of berberine (1) and its derivatives. First cells were treated with 10  $\mu$ M of berberine (1) and its derivatives at different time points, viz. 0.25, 1, 2, 4 and 8 h. The emission of the berberine (1) and its derivatives inside the cells was monitored (Fig. 4). The percentage incorporation of berberine (1) and its derivatives (5e, 6e and 7e) into the cells is shown in the figure 4. The data suggested that both the berberine (1) and its derivatives internalize into the cancer cells to the same extent with similar rates of internalization. A nearly 100% uptake was observed when the cells were incubated with the compounds for 2-4 h. As shown in Fig. 4, we found that the intracellular concentrations of berberine (1) and its three derivatives increased rapidly in 1 h, reached a plateau at 2 h, then stay at the plateau for another 6 h. The uptake amounts of compounds 5e, 6e and 7e are significantly higher than that of berberine (1). This observation confirmed our calculated higher partition coefficients ( $clogP$ ) of compounds 5e, 6e and 7e.






**Figure 4.** Intracellular accumulation of berberine (**1**) and its derivatives (**5e**, **6e** and **7e**) in HepG2 cells. HepG2 cells ( $1 \times 10^6$ ) were treated with  $10 \mu\text{M}$  of berberine and its derivatives (**5e**, **6e** and **7e**) and then incubated for 0 h, 0.25 h, 1 h, 2 h, 4 h or 8 h. Intracellular concentrations of berberine (**1**) and its derivatives (**5e**, **6e** and **7e**) were detected using HPLC analysis. Data are shown as the mean  $\pm$  SD ( $n = 3$ ) of two independent experiments.

#### 2.2.4. Photocytotoxicity of berberine (**1**) and its derivatives (**5e**, **6e** and **7e**)

To determine the photo-induced cytotoxicity of berberine (**1**) and its derivatives (**5e**, **6e** and **7e**) in HepG2 cells, proliferation assay was performed. Exponentially growing HepG2 cells were exposed for 10 min to visible light ( $420 \text{ nm}$ ,  $5.9 \text{ mW/cm}^2$ ) in presence of increasing concentrations of berberine (**1**) and its derivatives. Subsequently, the effect of the treatments on cell proliferation was determined by cell counts right after damage and after 20 h of incubation in fresh media, free of berberine derivatives. It is notable that, as shown in Fig. 5, the berberine derivatives caused a quite similar dose-dependent reduction of cell proliferation under visible light irradiation. On the contrary, no growth inhibition was seen by the irradiation alone (see data point at  $0 \mu\text{M}$  of berberine derivatives) and in non-irradiated cells pre-incubated with  $0.5 \mu\text{M}$  of berberine derivatives (see open symbols in Fig. 5).

In light of the dark cytotoxicity findings described above, we extended our study further on the anti-proliferation efficacy of berberine (**1**), compounds **5e**, **6e** and **7e** against HepG2 cells over a 24 h period with data on berberine (**1**) as the base case. As shown in Table 2, first of all, the dark cytotoxicity of compounds **5e**, **6e** and **7e** demonstrate, once again, an impressive 125-, 52- and 48-fold, respectively, stronger bioactivity than that of berberine (**1**) *in vitro*. In addition, the growth inhibition of HepG2 cells induced by these three samples is a concentration- and light dose-dependent manner.



**Figure 5.** *In vitro* photocytotoxicity of HepG2 cells incubated with (A) berberine (**1**), (B) **5e**, (C) **6e** and (D) **7e** at various concentration for 4 h and then illuminated under a visible light (420 nm) for 5, 10, 20 min or no light (dark series). The error bars denote standard derivation of three independent experiments.

The cytotoxicity of berberine (**1**) and its derivatives (**5e**, **6e** and **7e**) evaluated after incubation with HepG2 cells in the dark as well as after exposure to visible light (420 nm) for 10 min. For these experiments, the cells were seeded in 96-well microliter plates (5000 cells/well) and grown for at least 24 h in 100 μL of DMEM containing 10% FCS. The growth medium was replaced with DMEM containing 3% FCS and berberine (10, 20 and 40 μM) and its derivatives (0.25, 0.5 and 1.0 μM), and the cells were incubated for 4 h at 37°C. At the end of the incubation, the drug-containing medium was removed, and the cells were washed with PBS and irradiated for 10 min with visible light (total dose 3.36 J/cm<sup>2</sup>). Eight Black Light Blue lamps, mainly emitting at 420 nm, were used as the light source. After irradiation, the cells were returned to the incubator after replacement with whole medium, and 20 h later, cell survival was determined by the MTT assay [29]. Thus, 10 μL of MTT reagent (5 mg/mL) was added to each well. After 1.5 h of incubation at 37°C, 100 μL DMSO was added to dissolve the formazan crystals. The absorbance of the samples was measured using an ELISA plate reader (SpectraMax 340PC<sup>384</sup>) and selecting 540 nm as the test wavelengths. The survival of the treated cells was calculated as the percentage of their absorbance referred to the absorbance of the untreated (no drug, no light) cells taken as 100% viability.

Further, the photo-induced cytotoxic effects of berberine (**1**) and its derivatives on three human cancer cell lines was assessed using a light dose of 3.36 J/cm<sup>2</sup> at 420 nm wavelength, respectively. The resulting cells were irradiated with visible light (420 nm) for 10 min irradiation. After photo-irradiation, the cell viability was measured by MTT assay, of which these results are shown in Fig. 5 along with IC<sub>50</sub> values in Table 2. The control experiment without light irradiation showed slight cytotoxicity (85 ~ 95% of the cell viability). The dark cytotoxicity of these berberine derivatives was evaluated in HepG2 cells at the concentration of 0.5 μM without photo-irradiation. We found that all berberine derivatives showed no significantly cytotoxicity in the dark (Fig. 5), except over 1 μM. Furthermore, Fig. 5 showed the cell survival ratio as a function of the



concentrations of berberine derivatives. The  $IC_{50}$  value of **6e** was 0.15  $\mu$ M which was almost the same as that of **7e** (0.14  $\mu$ M) and lower than that of **5e** (0.28  $\mu$ M).

**Table 2.** *In vitro* dark cytotoxicity and photocytotoxicity of berberine (**1**) and its derivatives (**5e**, **6e** and **7e**) on three human cancer cell lines for 24 h.


Compd	$IC_{50}$ ( $\mu$ M) <sup>a</sup>					
	HepG2 <sup>b</sup>		BFTC905 <sup>c</sup>		HT29 <sup>d</sup>	
	Dark	Light	Dark	Light	Dark	Light
<b>1</b>	> 40	3.48 $\pm$ 0.69	> 40	15.20 $\pm$ 1.74	> 40	> 40
<b>5e</b>	0.52 $\pm$ 0.17	0.28 $\pm$ 0.15	2.74 $\pm$ 0.66	0.29 $\pm$ 0.11	3.42 $\pm$ 0.79	0.78 $\pm$ 0.25
<b>6e</b>	1.42 $\pm$ 0.38	0.15 $\pm$ 0.06	2.58 $\pm$ 0.87	0.20 $\pm$ 0.09	3.04 $\pm$ 0.32	0.28 $\pm$ 0.02
<b>7e</b>	1.66 $\pm$ 0.48	0.14 $\pm$ 0.06	2.64 $\pm$ 0.33	0.17 $\pm$ 0.05	2.64 $\pm$ 0.44	0.23 $\pm$ 0.02

<sup>a</sup> Compound concentration ( $\mu$ M) required to inhibit tumor cell proliferation rate by 50 %. Data are expressed as the mean  $\pm$  SD from the dose response curves of at 3–5 independent experiments. <sup>b</sup>

Human hepatocarcinoma cell lines. <sup>c</sup> Human bladder cancer cell lines. <sup>d</sup> Human colon adenocarcinoma cell lines.

#### 2.2.5. Cell morphological assessment

The apoptotic cells change in morphology, such as chromatin condensation, nuclear shrinking and DNA fragmentation. It can be characterized through Hoechst 33258 stained cell nucleus [30] under a fluorescent microscope. Thus, to further study the role of apoptosis played in the superior cytotoxicity of compounds **5e**, **6e** and **7e**, we incubated HepG2 cells in 0.5  $\mu$ M of berberine (**1**), or its three derivatives separately. As shown in Fig. 6A, while the nuclei of the cells were round in shape and stained homogenously in the control without testing compound, those treated with berberine (**1**), compound **5e**, **6e** or **7e** showed typical morphological features of apoptosis such as nuclear shrinkage, chromatin condensation and DNA fragmentation [30]. Fig. 6B shows that HepG2 cells treated with 0.5  $\mu$ M of compound **7e** stained with Hoechst 33258 in dark or light, and examined under fluorescence microscopy for topical morphological changes. Evidently, the proliferation of the cancer cells was inhibited by the testing compounds *in vitro*. These results showed that compounds **5e**, **6e** or **7e** induce apoptosis in HepG2 cells.




**Figure 6.** Morphological changes in HepG2 cells treated with berberine (**1**), **5e**, **6e** or **7e** at 0.5 μM at the dark or light (420 nm, 10 min) for 4 h (magnification, 200×). (A) Upper panels show the cell morphology under phase-contrast microscopy, red arrows indicate the apoptotic cells. (B) HepG2 cells were treated with **7e** (0.5 μM) for 4 h by irradiation and stained with Hoechst 33258 staining nuclear patterns detected through fluorescence microscopy (magnification, 200×). The yellow arrows indicate the apoptotic cells by irradiation.

#### 2.2.6. Cell cycle distribution analysis using flow cytometry

To probe the apoptotic effects of berberine (**1**) and its derivatives on cell cycle progression of HepG2 cells were treated once again with these compounds, separately, at 0.5 μM concentrations for 2 h after irradiation (420 nm, 10 min) for 24 h. Then the samples were further treated with propidium iodide (PI) staining [31] and of which the sub-G1 phase was analyzed by flow cytometry. Besides, the cells with no test compound were the control of the experiment. Berberine (**1**) and its derivatives (**5e**, **6e** and **7e**) showed significant photocytotoxic property in visible light at 420 nm wavelength. To find out whether the said property play a role in the apoptosis of the cells, flow cytometric analysis of the HepG2 cells treated with berberine (**1**) and its derivatives (**5e**, **6e** and **7e**) was carried out at 0.5 μM concentrations. Cells initially treated with the compounds in the dark and tested before and after being exposed to visible light (420 nm), and then stained with propidium iodide (PI) to observe any fragmented DNA, or an increase in the sub-G1 population. As illustrated in Fig. 7, the percentage of S phase increases by 32.71% and 37.58% in dark and light for the control, respectively, in sample with of **5e** increase to 41.14%, 49.71% for **6e** and 44.36% for **7e** compare to the 37.58% for the control in the light. The increased S phase percentage and the sub-G1 population in compounds **5e**, **6e** and **7e** treatment suggest that **5e**, **6e** and **7e** induce S phase arrest and cause apoptosis in HepG2 liver cancer cells. At first, berberine (**1**) and its derivatives (**5e**, **6e** and **7e**) being non-cytotoxic in the dark at 0.5 μM concentration, but it induced significant apoptosis in the cells at

a concentration of 0.5  $\mu\text{M}$  upon visible light irradiation for 24 h. In addition, percentage of apoptotic cell population increased with increasing compounds concentration (Fig. 7). Lastly, a significant induction of apoptosis in the cells occurred even in the dark with increasing concentration of derivatives.




**Figure 7.** Effect of berberine (1), 5e, 6e or 7e on HepG2 cell cycle distribution using without or with visible light exposure. (A) Cell cycle distribution after treatment with berberine (1), 5e, 6e or 7e at 0.5  $\mu\text{M}$  concentration for 2 h after irradiation (420 nm, 10 min) in HepG2 cells for 24 h. (B) Quantitative difference of cell cycle distribution changed after treatment with berberine (1), 5e, 6e and 7e at 0.5  $\mu\text{M}$  concentration in HepG2 for 24 h. Data are shown as the mean  $\pm$  SD of three independent experiments, and \*\*\* $p$  < 0.001 compared with dark group.

#### 2.2.7. Measurement of intracellular ROS production by irradiation

Several alkaloids induce apoptosis by generating singlet oxygen ( $^1\text{O}_2$ ) in mitochondria were reported [32]. Using dichlorofluorescein diacetate (DCFH-DA) assay we checked if berberine (1), compounds 5e, 6e or 7e could stimulate generation of reactive oxygen species (ROS) after irradiation at visible light (420 nm) in HepG2 cells. We found that the fluorescence intensity of DCF in the cells

was right-shifted after the cells were treated with all four compounds in a 0.5  $\mu\text{M}$  concentration upon irradiation. Shown in Fig. 8A, berberine (**1**), compounds **5e**, **6e** or **7e** all stimulated to release intracellular  $^1\text{O}_2$  from HepG2 cells and exhibited a more profound effect than on the  $^1\text{O}_2$  generation in HepG2 cells after 24 h of treatment ( $P < 0.05$ , Fig. 8B). These data showed that berberine (**1**), compounds **6e** and **7e** induced apoptosis by increasing the intracellular ROS generation of HepG2 cells




**Figure 8.** ROS generation in HepG2 cells treated with berberine (**1**), **5e**, **6e**, and **7e** after irradiation. (A) The effects of HepG2 cells treated with berberine (**1**), **5e**, **6e**, and **7e** at 0.5  $\mu\text{M}$  concentration for 2 h after irradiation (420 nm, 10 min) for 24 h on percentage distribution of ROS generation by DCFH-DA staining. (B) Average intensity of DCFH-DA fluorescence in HepG2 cells treated with berberine (**1**) and its derivatives **5e**, **6e** and **7e** at 0.5  $\mu\text{M}$  concentration. The results are presented as mean  $\pm$  SD. \*\*\* $p < 0.001$  compared with control, \*\* $p < 0.01$  and ### $p < 0.001$  compared with dark group.

#### 2.2.8. Measurements of mitochondrial membrane potential changes

The depletion of mitochondrial membrane potential (MMP) is an early marker of the apoptotic process. To determine whether an early loss of MMP occurred upon the treatment of berberine (**1**) and its derivatives in HepG2 cells, we performed MMP measurement with rhodamine 123 staining. Delineated in Fig. 9, the MMP decreased remarkably at the concentrations of 0.5  $\mu\text{M}$  of berberine (**1**),

0.25  $\mu\text{M}$  of compound **5e**, **6e** or **7e**. The percentage of cells with membrane potential loss increased from 20.53% to 32.90% at 24 h. Thus berberine (**1**), compound **5e**, **6e** or **7e** severely damaged the DNA of HepG2 cells. It was documented that P53 could also down-regulate Bcl-2 and up-regulate Bax, which are related to mitochondria mediated apoptosis [33]. Further, several studies have also reported that apoptosis involves a disruption of the mitochondrial membrane integrity, a critical cell death process. Moreover, depolarization of the mitochondrial membrane potential is a characteristic feature of apoptosis [34,35]. Report also indicated that the reduction of the membrane potential, the release of mitochondrial cytochrome *c*, a critical step, is followed by the formation of apoptosomes [35]. Thus what we found the loss of MMP upon the treatments of berberine (**1**), compound **5e**, **6e** or **7e** are in consistency with the published results. So does the finding of the release of mitochondrial cytochrome *c* induced by berberine (**1**), compound **5e**, **6e** or **7e** at a concentration of 0.25  $\mu\text{M}$ .



**Figure 9.** Measurement of the mitochondrial membrane potential. (A) HepG2 cells ( $1 \times 10^5$  cells/mL) were incubated with 0.5  $\mu\text{M}$  concentration of berberine (**1**) and its derivative **5e**, **6e** and **7e** at 0.25  $\mu\text{M}$  concentration for 2 h upon visible light irradiation (420 nm, 10 min) for 24 h and then stained with 2 mg/mL of Rh123 for 20 min under gentle shaking. After washing with PBS, cells were immediately analyzed by flow cytometry and data was analyzed using the FCS Express software. (B) Average intensity of RH123 fluorescence in HepG2 cells treated with 0.5  $\mu\text{M}$  of berberine (**1**) and its derivatives **5e**, **6e** and **7e** at 0.25  $\mu\text{M}$  concentration. The results are presented as mean  $\pm$  SD. \*\*\* $p$  < 0.001 compared with control, ### $p$  < 0.001 compared with dark group.



### 3. Materials and Methods

#### 3.1. General information

All chemical reagents and solvents in commercial quality were purchased from Sigma-Aldrich (St. Louis, MO, USA), Fluka (Italy), TCI (Japan), and Alpha Aesar without further purification. All chemical reagents in commercial quality were used as received (Sigma-Aldrich, St. Louis, MO, USA) and were used without further purification. Solvents were dried and the synthesized compounds were purified using standard techniques. Reactions-progression was monitored by TLC on aluminum plates coated with silica gel with a fluorescent indicator (Merck 60 F<sub>254</sub>). Melting points of the synthesized compounds were determined by Fargo MP-2D apparatus and were uncorrected. The electronic absorption spectra were measured on a Shimadzu UV-1601 UV-vis spectrophotometer. Fluorescence spectra were taken on a Shimadzu RP-5301 spectrofluorometer. NMR spectra were recorded using TMS as an internal standard in CDCl<sub>3</sub> at 500MHz for <sup>1</sup>H and at 125MHz for <sup>13</sup>C (Bruker Biospin GmbH AVANCE III 500 MHz, Rheinstetten, Germany). Chemical shift ( $\delta$ ) were reported in parts per million (ppm) measured relative to the internal standards (TMS), and the coupling constant (*J*) were expressed in Hertz (Hz). Column chromatography was performed with silica gel SiliaFlash® G60 (60–200  $\mu$ m) purchased from SiliCycle Inc. (Quebec City, Canada). In general, the reactions were carried out under anhydrous conditions in dry solvent and nitrogen atmosphere. The purity of these compounds was based on the analysis of HPLC (Hitachi High-Technologies, Tokyo, Japan) equipped with a 280 nm detector and LiChroCART RP-C<sub>18</sub> column (4.6 mm i.d.  $\times$  250 mm, 5  $\mu$ m, Merck, Darmstadt, Germany). The mobile phase was composed of MeOH-H<sub>2</sub>O (0.05% TFA) (90:10) and the flow rate was 1.0 mL/min. The purity of all compounds was more than 98%. The mass spectra were acquired using a Thermo Finnigan model LXQ (Thermo Electron Co., Waltham, MA, USA) ion trap mass spectrometer equipped with ESI source interference and controlled by Xcalibur 2.06. The mass spectra were acquired in a positive ion mode or a negative ion mode. Berberrubine (**2**), derivatives **3**, **4**, **5a**, **5c** and **5e** were synthesized according to previous methods [27].

#### 3.2. Synthesis of 9-O-substituted berberine derivatives (**5a–e**)

To a solution of berberrubine **2** (1 mmol) in dry acetonitrile (10 mL) was added various *n*-alkyl bromide (1.2 mmol) and the reaction mixture was refluxed for 4–8 hours. The progress of the reaction was monitored by TLC. Then, the reaction mixture was cooled to room temperature and the solvent was removed under reduced pressure. The crude product was chromatographed on a silica gel column and eluted with ethyl acetate/methanol (2/1) solvent to afford the desired berberine derivatives **5a–e**.

**9-O-Butylberberine bromide (5a).** Berberrubine (**2**) was treated with *n*-butyl bromide according to the general procedure to give the desired bromide salt (**5a**) as a yellowish brown solid, yield: 86%; UV (MeOH)  $\lambda_{\text{max}}$  (log  $\epsilon$ ): 431 (3.77), 350 (4.41), 267 (4.43), 231 (4.45), 205 (4.41) nm; <sup>1</sup>H NMR (500 MHz, DMSO-*d*<sub>6</sub>):  $\delta$  9.75 (s, 1H, H-8), 8.94 (s, 1H, H-13), 8.20 (d, *J* = 9.1 Hz, H-12), 7.99 (d, *J* = 9.1 Hz, H-11), 7.80 (s, 1H, H-1), 7.09 (s, 1H, H-4), 6.17 (s, 2H, -OCH<sub>2</sub>O-), 4.95 (t, *J* = 6.3 Hz, 2H, H-6), 4.29 (t, *J* = 6.8 Hz, 2H, H-1'), 4.05 (s, 3H, -OCH<sub>3</sub>), 3.21 (t, *J* = 6.3 Hz, 2H, H-5), 1.86 (p, *J* = 7.2 Hz, 2H, H-2'), 1.52 (sex, *J* = 7.4 Hz, 2H, H-3'), 0.99 (t, *J* = 7.4 Hz, 3H, H-4'); <sup>13</sup>C NMR (125 MHz, DMSO-*d*<sub>6</sub>):  $\delta$  150.4, 149.8, 147.7, 145.3, 142.9, 137.5, 133.0, 130.7, 126.7, 123.3, 121.7, 120.5, 120.2, 108.4, 105.4, 102.1, 73.9, 57.0, 55.3,



31.5, 26.3, 18.6, 13.8; LC-MS (ESI<sup>+</sup>, *m/z*) C<sub>23</sub>H<sub>24</sub>NO<sub>4</sub><sup>+</sup>: 378.33 [M-Br]<sup>+</sup>; HRMS-ESI: *m/z* calculated for C<sub>23</sub>H<sub>22</sub>NO<sub>4</sub><sup>+</sup>: 378.1705 [M-Br]<sup>+</sup>, found for 378.1708.

**9-*O*-Hexylberberine bromide (5b).** Berberrubine (**2**) was treated with *n*-hexyl bromide according to the general procedure to give the desired bromide salt (**5b**) as a yellowish brown solid, yield: 69%; UV (MeOH) λ<sub>max</sub> (log ε): 431.6 (3.92), 350.2 (4.54), 266.5 (4.60), 229.1 (4.67) nm; <sup>1</sup>H NMR (500 MHz, DMSO-*d*<sub>6</sub>): δ 9.75 (s, 1H, H-8), 8.94 (s, 1H, H-13), 8.20 (d, *J* = 9.1 Hz, H-12), 7.99 (d, *J* = 9.1 Hz, H-11), 7.80 (s, 1H, H-1), 7.09 (s, 1H, H-4), 6.17 (s, 2H, -OCH<sub>2</sub>O-), 4.95 (t, *J* = 6.3 Hz, 2H, H-6), 4.28 (t, *J* = 6.8 Hz, 2H, H-1'), 4.05 (s, 3H, -OCH<sub>3</sub>), 3.21 (t, *J* = 6.3 Hz, 2H, H-5), 1.87 (p, *J* = 7.2 Hz, 2H, H-2'), 1.52 (p, *J* = 7.4 Hz, 2H, H-3'), 1.37–1.33 (m, 4H, H-4', H-5'), 0.90 (t, *J* = 7.0 Hz, 3H, H-6'); <sup>13</sup>C NMR (125 MHz, DMSO-*d*<sub>6</sub>): δ 150.5, 149.8, 147.7, 145.3, 142.9, 137.5, 133.1, 130.7, 126.7, 123.4, 121.7, 120.5, 120.3, 108.5, 105.5, 102.1, 74.3, 57.1, 55.3, 31.1, 29.5, 26.4, 25.0, 22.1, 14.0; LC-MS (ESI<sup>+</sup>, *m/z*) C<sub>25</sub>H<sub>26</sub>NO<sub>4</sub><sup>+</sup>: 406.33 [M-Br]<sup>+</sup>; HRMS-ESI: *m/z* calculated for C<sub>25</sub>H<sub>26</sub>NO<sub>4</sub><sup>+</sup>: 406.2018 [M-Br]<sup>+</sup>, found for 406.2021.

**9-*O*-Octylberberine bromide (5c).** Berberrubine (**2**) was treated with *n*-octyl bromide according to the general procedure to give the desired bromide salt (**5c**) as a bright yellow solid, yield: 74%; UV (MeOH) λ<sub>max</sub> (log ε): 432 (3.79), 350 (4.42), 268 (4.44), 230 (4.47), 204 (4.48) nm; <sup>1</sup>H NMR (500 MHz, DMSO-*d*<sub>6</sub>): δ 9.75 (s, 1H, H-8), 8.95 (s, 1H, H-13), 8.19 (d, *J* = 9.1 Hz, H-12), 7.99 (d, *J* = 9.1 Hz, H-11), 7.80 (s, 1H, H-1), 7.09 (s, 1H, H-4), 6.17 (s, 2H, -OCH<sub>2</sub>O-), 4.95 (t, *J* = 6.2 Hz, 2H, H-6), 4.28 (t, *J* = 6.8 Hz, 2H, H-1'), 4.06 (s, 3H, -OCH<sub>3</sub>), 3.21 (t, *J* = 6.2 Hz, 2H, H-5), 1.87 (p, *J* = 7.2 Hz, 2H, H-2'), 1.48 (p, *J* = 7.3 Hz, 2H, H-3'), 1.33 (m, 8H, H-4'-H-7'), 0.88 (t, *J* = 6.8 Hz, 3H, H-8'); <sup>13</sup>C NMR (125 MHz, DMSO-*d*<sub>6</sub>): δ 150.4, 149.8, 147.7, 145.3, 142.9, 137.5, 133.0, 126.7, 123.3, 121.7, 120.5, 120.2, 108.4, 105.4, 102.1, 74.2, 57.0, 55.3, 31.2, 29.5, 28.8, 28.7, 26.3, 25.3, 22.1, 14.0; LC-MS (ESI<sup>+</sup>, *m/z*) C<sub>27</sub>H<sub>32</sub>NO<sub>4</sub><sup>+</sup>: 434.41 [M-Br]<sup>+</sup>; HRMS-ESI: *m/z* calculated for C<sub>27</sub>H<sub>32</sub>NO<sub>4</sub><sup>+</sup>: 434.2331 [M-Br]<sup>+</sup>, found for 434.2332.

**9-*O*-Decylberberine bromide (5d).** Berberrubine (**2**) was treated with *n*-decyl bromide according to the general procedure to give the desired bromide salt (**5d**) as a bright yellow solid, yield: 71%; UV (MeOH) λ<sub>max</sub> (log ε): 431.0 (3.68), 350.2 (4.31), 266.4 (4.35), 229.8 (4.38) nm; <sup>1</sup>H NMR (500 MHz, DMSO-*d*<sub>6</sub>): δ 9.75 (s, 1H, H-8), 8.95 (s, 1H, H-13), 8.19 (d, *J* = 9.1 Hz, H-12), 7.99 (d, *J* = 9.1 Hz, H-11), 7.80 (s, 1H, H-1), 7.09 (s, 1H, H-4), 6.17 (s, 2H, -OCH<sub>2</sub>O-), 4.95 (t, *J* = 6.2 Hz, 2H, H-6), 4.27 (t, *J* = 6.8 Hz, 2H, H-1'), 4.05 (s, 3H, -OCH<sub>3</sub>), 3.21 (t, *J* = 6.2 Hz, 2H, H-5), 1.87 (p, *J* = 7.2 Hz, 2H, H-2'), 1.48 (p, *J* = 7.3 Hz, 2H, H-3'), 1.40–1.26 (m, 12H, H-4'-H-9'), 0.85 (t, *J* = 6.8 Hz, 3H, H-10'); <sup>13</sup>C NMR (125 MHz, DMSO-*d*<sub>6</sub>): δ 150.4, 149.8, 147.7, 145.3, 142.9, 137.5, 133.0, 130.7, 126.7, 123.3, 121.7, 120.5, 120.2, 108.4, 105.4, 102.1, 74.2, 57.0, 55.3, 31.3, 29.5, 29.0, 28.9, 28.8, 28.7, 26.3, 25.3, 22.1, 14.0; LC-MS (ESI<sup>+</sup>, *m/z*) C<sub>29</sub>H<sub>34</sub>NO<sub>4</sub><sup>+</sup>: 462.43 [M-Br]<sup>+</sup>; HRMS-ESI: *m/z* calculated for C<sub>29</sub>H<sub>34</sub>NO<sub>4</sub><sup>+</sup>: 462.2644 [M-Br]<sup>+</sup>, found for 462.2646.

**9-*O*-Dodecylberberine bromide (5e).** Berberrubine (**2**) was treated with *n*-dodecyl bromide according to the general procedure to give the desired bromide salt (**5e**) as a dark yellowish brown solid, yield: 61%; UV (MeOH) λ<sub>max</sub> (log ε): 431 (3.42), 351 (4.03), 267 (4.07), 230 (4.10), 202 (4.19) nm; <sup>1</sup>H NMR (500 MHz, DMSO-*d*<sub>6</sub>): δ 9.77 (s, 1H, H-8), 8.98 (s, 1H, H-13), 8.21 (d, *J* = 9.1 Hz, H-12), 8.00 (d, *J* = 9.1 Hz, H-11), 7.82 (s, 1H, H-1), 7.10 (s, 1H, H-4), 6.18 (s, 2H, -OCH<sub>2</sub>O-), 4.96 (t, *J* = 6.1 Hz, 2H, H-6), 4.28 (t, *J* = 6.8 Hz, 2H, H-1'), 4.05 (s, 3H, -OCH<sub>3</sub>), 3.21 (t, *J* = 6.1 Hz, 2H, H-5), 1.88 (d, *J* = 7.4 Hz, H-2'), 1.47 (q, *J* = 7.5 Hz, 2H, H-3'), 1.29 (m, 16 H, H-4'-H-11'), 0.86 (t, *J* = 6.8 Hz, 3H, H-12'); <sup>13</sup>C NMR (125 MHz, DMSO-*d*<sub>6</sub>): δ 150.4, 149.8, 147.7, 145.3, 142.9, 137.5, 133.0, 130.7, 126.7, 123.3, 121.7, 120.5, 120.3, 108.4, 105.5, 102.1, 74.2, 57.0, 55.3, 48.6, 31.3, 29.5, 29.08, 29.05, 28.9, 28.7, 26.3, 25.3, 22.1, 14.0; LC-MS (ESI<sup>+</sup>,

$m/z$   $C_{31}H_{38}NO_4^+$ : 490.50  $[M-Br]^+$ ; HRMS-ESI:  $m/z$  calculated for  $C_{31}H_{38}NO_4^+$ : 490.2957  $[M-Br]^+$ , found for 490.2958.

### 3.3. Synthesis of 13-substituted berberine derivatives (6a–e)

To a stirred solution of berberine (**1**) (3.71 g, 10 mmol) and  $K_2CO_3$  (3.6 g, 30 mmol) in methanol (125 mL), 5% NaOH (5 mL) solution containing  $NaBH_4$  (0.30 g, 7.5 mmol) was added dropwise. The reaction mixture was stirred at room temperature for 1 h and the precipitated product was filtered, washed with 30% ethanol (20 mL) and 80% ethanol (20 mL), and then recrystallized from absolute ethanol to provide dihydroberberine (**3**) (2.5 g, 74%) as a brown like solid, yield:74%; UV (MeOH)  $\lambda_{max}$  (log  $\epsilon$ ): 353.9 (3.96), 268 (4.01), 228.6 (4.09) nm;  $^1H$  NMR (500 MHz, DMSO- $d_6$ )  $\delta$  7.29 (s, 1H, H-13), 6.82 (d,  $J$  = 8.4 Hz, H-12), 6.76 (s, 1H, H-1), 6.69 (d,  $J$  = 8.4 Hz, 1H, H-11), 6.05 (s, 1H, H-1), 6.00 (s, 2H,  $-OCH_2-$ ), 4.20 (s, 2H, H-8), 3.76 (s, 3H,  $-OCH_3$ ), 3.71 (s, 3H,  $-OCH_3$ ), 3.05 (t,  $J$  = 5.8 Hz, 2H, H-6), 2.79 (t,  $J$  = 5.8 Hz, 2H, H-5);  $^{13}C$  NMR (125 MHz, DMSO- $d_6$ )  $\delta$  149.9, 146.9, 146.4, 143.9, 141.0, 128.6, 128.3, 123.9, 121.6, 118.4, 111.7, 107.9, 103.5, 101.0, 96.0, 60.2, 55.7, 48.8, 48.3, 28.9.

To a stirred solution of dihydroberberine (**3**) (170 mg, 5.0 mmol) in 80% ethanol (8 mL) and HOAc (2 mL), *n*-alkyl aldehyde (2 mL) was added. The reaction mixture was heated to 85–95 °C for 5 h. The solvent was removed by evaporation, and the residue was acidified by 2 N HCl (5 mL), and then stirred at room temperature for 1 h. The solid was collected by filtration and then purified by flash chromatography over silica gel, affording the title compounds **6a–e**.

**13-Butylberberine bromide (6a)**. Dihydroberberine (**3**) was treated with butanal according to the general procedure to give the desired chloride salt (**6a**) as a yellowish brown solid, yield: 68%; UV (MeOH)  $\lambda_{max}$  (log  $\epsilon$ ): 420.3 (3.93), 341.9 (4.49), 265.1 (4.61), 231.9 (4.58) nm;  $^1H$  NMR (500 MHz, DMSO- $d_6$ ):  $\delta$  9.88 (s, 1H, H-8), 8.21 (d,  $J$  = 9.5 Hz, 1H, H-12), 8.18 (d,  $J$  = 9.5 Hz, 1H, H-11), 7.28 (s, 1H, H-1), 7.16 (s, 1H, H-4), 6.18 (s, 2H,  $-OCH_2O-$ ), 4.78 (t,  $J$  = 8.0 Hz, 2H, H-6), 4.09 (s, 3H,  $-OCH_3$ ), 4.08 (s, 3H,  $-OCH_3$ ), 3.33 (t,  $J$  = 8.0 Hz, 2H, H-5), 3.07 (t,  $J$  = 6.2 Hz, 2H, H-1'), 1.75 (p,  $J$  = 6.6 Hz, 2H, H-2'), 1.41 (hex,  $J$  = 7.2 Hz, 2H, H-3'), 0.91 (t,  $J$  = 6.9 Hz, 3H, H-4');  $^{13}C$  NMR (125 MHz, DMSO- $d_6$ ):  $\delta$  150.3, 149.1, 146.6, 144.4, 144.3, 135.8, 134.2, 134.1, 132.3, 125.9, 121.5, 121.3, 109.2, 108.4, 102.2, 62.1, 57.1, 57.0, 32.6, 28.8, 27.4, 22.0, 13.5; LC-MS (ESI $^+$ ,  $m/z$ )  $C_{24}H_{26}NO_4^+$ : 392.28  $[M-Cl]^+$ ; HRMS-ESI:  $m/z$  calculated for  $C_{24}H_{26}NO_4^+$ : 392.1862  $[M-Cl]^+$ , found for 392.1861.

**13-Hexylberberine bromide (6b)**. Dihydroberberine (**3**) was treated with hexanal according to the general procedure to give the desired chloride salt (**6b**) as a bright yellow solid, yield: 76%; UV (MeOH)  $\lambda_{max}$  (log  $\epsilon$ ): 420.3 (3.73), 341.9 (4.29), 265.4 (4.40), 232 (4.37) nm;  $^1H$  NMR (500 MHz, DMSO- $d_6$ ):  $\delta$  9.91 (s, 1H, H-8), 8.22 (d,  $J$  = 9.7 Hz, 1H, H-12), 8.20 (d,  $J$  = 9.7 Hz, 1H, H-11), 7.30 (s, 1H, H-1), 7.18 (s, 1H, H-4), 6.20 (s, 2H,  $-OCH_2O-$ ), 4.80 (s, 2H, H-6), 4.10 (s, 3H,  $-OCH_3$ ), 4.10 (s, 3H,  $-OCH_3$ ), 3.33 (s, 2H, H-5), 3.09 (t,  $J$  = 5.7 Hz, 2H, H-1'), 1.76 (s, 2H, H-2'), 1.40 (m, 2H, H-3'), 1.27–1.26 (m, 4H, H-4', H-5'), 0.86 (t,  $J$  = 6.9 Hz, 3H, H-6');  $^{13}C$  NMR (125 MHz, DMSO- $d_6$ ):  $\delta$  150.2, 149.1, 146.6, 144.4, 144.2, 135.8, 134.2, 134.1, 132.2, 125.9, 121.5, 121.3, 120.3, 109.1, 108.4, 102.2, 62.0, 57.0, 30.7, 30.4, 28.9, 28.3, 27.4, 22.1, 13.9; LC-MS (ESI $^+$ ,  $m/z$ )  $C_{26}H_{30}NO_4^+$ : 420.26  $[M-Cl]^+$ ; HRMS-ESI:  $m/z$  calculated for  $C_{26}H_{30}NO_4^+$ : 420.2175  $[M-Cl]^+$ , found for 420.2176.

**13-Octylberberine chloride (6c)**. Dihydroberberine (**3**) was treated with octanal according to the general procedure to give the desired chloride salt (**6c**) as a bright yellow solid, yield: 65%; UV (MeOH)  $\lambda_{max}$  (log  $\epsilon$ ): 420.1 (3.80), 341.4 (4.35), 265.3 (4.47), 232 (4.44) nm;  $^1H$  NMR (500 MHz,

DMSO-*d*<sub>6</sub>):  $\delta$  9.91 (s, 1H, H-8), 8.22 (d,  $J$  = 9.7 Hz, 1H, H-12), 8.20 (d,  $J$  = 9.7 Hz, 1H, H-11), 7.30 (s, 1H, H-1), 7.17 (s, 1H, H-4), 6.20 (s, 2H, -OCH<sub>2</sub>O-), 4.80 (s, 2H, H-6), 4.10 (s, 3H, -OCH<sub>3</sub>), 4.10 (s, 3H, -OCH<sub>3</sub>), 3.33 (s, 2H, H-5), 3.09 (t,  $J$  = 5.7 Hz, 2H, H-1'), 1.76 (s, 2H, H-2'), 1.38–1.37 (m, 2H, H-3'), 1.28–1.23 (m, 8H, H-4'–H-7'), 0.86 (t,  $J$  = 6.9 Hz, 3H, H-8'); <sup>13</sup>C NMR (125 MHz, DMSO-*d*<sub>6</sub>):  $\delta$  150.2, 149.0, 146.6, 144.3, 144.2, 135.8, 134.2, 134.0, 132.2, 125.9, 121.4, 121.2, 120.3, 109.1, 108.3, 102.1, 62.0, 57.0, 31.2, 30.4, 28.8, 28.6, 28.4, 27.4, 22.0, 13.9; LC-MS (ESI<sup>+</sup>,  $m/z$ ) C<sub>28</sub>H<sub>34</sub>NO<sub>4</sub><sup>+</sup>: 448.28 [M–Cl]<sup>+</sup>; HRMS-ESI:  $m/z$  calculated for C<sub>28</sub>H<sub>34</sub>NO<sub>4</sub><sup>+</sup>: 448.2488 [M–Cl]<sup>+</sup>, found for 448.2487.

**13-Decylberberine chloride (6d).** Dihydroberberine (3) was treated with decanal according to the general procedure to give the desired chloride salt (6d) as a bright yellow solid, yield: 43%; UV (MeOH)  $\lambda_{\max}$  (log  $\epsilon$ ): 420.3 (3.73), 341.9 (4.29), 265.3 (4.40), 231.7 (4.39) nm; <sup>1</sup>H NMR (500 MHz, DMSO-*d*<sub>6</sub>):  $\delta$  9.90 (s, 1H, H-8), 8.22 (d,  $J$  = 9.7 Hz, 1H, H-12), 8.20 (d,  $J$  = 9.7 Hz, 1H, H-11), 7.29 (s, 1H, H-1), 7.16 (s, 1H, H-4), 6.19 (s, 2H, -OCH<sub>2</sub>O-), 4.79 (s, 2H, H-6), 4.09 (s, 3H, -OCH<sub>3</sub>), 4.09 (s, 3H, -OCH<sub>3</sub>), 3.33 (s, 2H, H-5), 3.08 (t,  $J$  = 5.7 Hz, 2H, H-1'), 1.74 (s, 2H, H-2'), 1.36–1.35 (m, 2H, H-3'), 1.27–1.22 (m, 12H, H-4'–H-9'), 0.86 (t,  $J$  = 6.9 Hz, 3H, H-10'); <sup>13</sup>C NMR (125 MHz, DMSO-*d*<sub>6</sub>):  $\delta$  150.2, 149.0, 146.6, 144.3, 144.2, 135.8, 134.2, 134.0, 132.2, 125.9, 121.4, 121.2, 120.3, 109.1, 108.3, 102.1, 62.0, 57.0, 31.3, 30.4, 28.9, 28.8, 28.6, 28.8, 28.4, 27.4, 22.1, 13.9; LC-MS (ESI<sup>+</sup>,  $m/z$ ) C<sub>30</sub>H<sub>38</sub>NO<sub>4</sub><sup>+</sup>: 476.40 [M–Cl]<sup>+</sup>; HRMS-ESI:  $m/z$  calculated for C<sub>30</sub>H<sub>38</sub>NO<sub>4</sub><sup>+</sup>: 476.2801 [M–Cl]<sup>+</sup>, found for 476.2801.

**13-Dodecylberberine chloride (6e).** Dihydroberberine (3) was treated with dodecanal according to the general procedure to give the desired chloride salt (6e) as a bright yellow solid, yield: 29%; UV (MeOH)  $\lambda_{\max}$  (log  $\epsilon$ ): 420.6 (3.71), 341.8 (4.26), 265.3 (4.38), 232.0 (4.37) nm; <sup>1</sup>H NMR (500 MHz, DMSO-*d*<sub>6</sub>):  $\delta$  9.90 (s, 1H, H-8), 8.21 (d,  $J$  = 9.6 Hz, 1H, H-12), 8.19 (d,  $J$  = 9.6 Hz, 1H, H-11), 7.29 (s, 1H, H-1), 7.16 (s, 1H, H-4), 6.18 (s, 2H, -OCH<sub>2</sub>O-), 4.79 (s, 2H, H-6), 4.09 (s, 3H, -OCH<sub>3</sub>), 4.09 (s, 3H, -OCH<sub>3</sub>), 3.33 (s, 2H, H-5), 3.08 (t,  $J$  = 5.7 Hz, 2H, H-1'), 1.74 (s, 2H, H-2'), 1.36–1.35 (m, 2H, H-3'), 1.28–1.22 (m, 16H, H-4'–H-11'), 0.85 (t,  $J$  = 6.9 Hz, 3H, H-12'); <sup>13</sup>C NMR (125 MHz, DMSO-*d*<sub>6</sub>):  $\delta$  150.2, 149.0, 146.5, 144.3, 144.2, 135.8, 134.2, 134.0, 132.2, 125.9, 121.4, 121.2, 120.3, 109.1, 108.3, 102.1, 62.0, 57.0, 31.3, 30.4, 30.0, 28.9, 28.9, 28.8, 28.7, 28.6, 28.4, 27.4, 22.1, 13.9; LC-MS (ESI<sup>+</sup>,  $m/z$ ) C<sub>32</sub>H<sub>42</sub>NO<sub>4</sub><sup>+</sup>: 504.43 [M–Cl]<sup>+</sup>; HRMS-ESI:  $m/z$  calculated for C<sub>32</sub>H<sub>42</sub>NO<sub>4</sub><sup>+</sup>: 504.3114 [M–Cl]<sup>+</sup>, found for 504.3115.

### 3.4. Synthesis of 13-*O*-substituted berberine derivatives (7a–e)

Compound 3 (337 mg) was dissolved in 35 mL of CH<sub>2</sub>Cl<sub>2</sub>. Under nitrogen atmosphere, the temperature of the system was kept at –25°C ~ –30°C, and 8 mL of CH<sub>2</sub>Cl<sub>2</sub> solution containing 258 mg MCPBA (1.5 mmol) was dropwise added slowly. After the addition, the temperature was kept and the reaction was carried out under agitation for 1 h. Then, the temperature was raised to 0 °C and 250 mg (2.0 mmol) of sodium sulfite was added therein, followed by stirred at room temperature for 1 h. The reaction was stopped and the reaction mixture was filtered. The filtrate was evaporated to remove the solvent under reduced pressure and the residue was purified through silica gel column chromatography (CH<sub>2</sub>Cl<sub>2</sub>/CH<sub>3</sub>OH = 10 : 1) to obtain compound 4 (280 mg, 80 percent), yield: 80%; UV (MeOH)  $\lambda_{\max}$  (log  $\epsilon$ ): 444.5 (4.08), 367.9 (3.93), 305.3 (4.04), 235.0 (4.44), 212.1 (4.42) nm; <sup>1</sup>H NMR (500 MHz, DMSO-*d*<sub>6</sub>):  $\delta$  8.89 (s, 1H, H-8), 8.08 (d,  $J$  = 9.3 Hz, 1H, H-12), 8.06 (s, 1H, H-1), 7.52 (d,  $J$  = 9.3 Hz, 1H, H-11), 6.88 (s, 1H, H-4), 6.02 (s, 2H, -OCH<sub>2</sub>O-), 4.61 (t,  $J$  = 6.2 Hz, 2H, H-6), 3.95 (s, 3H, -OCH<sub>3</sub>), 3.93 (s, 3H, -OCH<sub>3</sub>), 3.01 (t,  $J$  = 6.2 Hz, 2H, H-5); <sup>13</sup>C NMR (125 MHz, DMSO-*d*<sub>6</sub>):  $\delta$  165.1, 150.2, 145.3, 145.2, 142.2, 130.2, 127.4, 123.9, 123.4, 122.3, 121.4, 117.6, 117.3, 107.5, 107.2, 100.8, 61.1, 56.4, 56.3, 27.6.

Compound **4** (1.0 g) and potassium carbonate (200 mg) were dissolved in 20 mL of acetone followed by adding 0.25 mL of alkyl bromide. The obtained reaction solution was heated and refluxed for 3 hours. Then, the reaction mixture was filtered and the filtrate was evaporated under reduced pressure. The residual was purified through silica gel column (CH<sub>2</sub>Cl<sub>2</sub>/CH<sub>3</sub>OH = 20: 1) to obtain compounds **7a–e**.

**13-O-Butylberberine bromide (7a).** Compound (**4**) was treated with *n*-butyl bromide according to the general procedure to give the desired bromide salt (**7a**) as a yellowish brown solid, yield: 66%; UV (MeOH)  $\lambda_{\max}$  (log  $\epsilon$ ): 427.8 (3.46), 349.7 (3.97), 265.5 (4.07), 232.6 (4.17) nm; <sup>1</sup>H NMR (500 MHz, DMSO-*d*<sub>6</sub>):  $\delta$  9.82 (s, 1H, H-8), 8.21 (d, *J* = 9.3 Hz, 1H, H-12), 8.09 (d, *J* = 9.3 Hz, 1H, H-11), 7.90 (s, 1H, H-1), 7.13 (s, 1H, H-4), 6.17 (s, 2H, -OCH<sub>2</sub>O-), 4.88 (t, *J* = 5.8 Hz, 2H, H-6), 4.10 (s, 3H, -OCH<sub>3</sub>), 4.09 (s, 3H, -OCH<sub>3</sub>), 3.84 (t, *J* = 6.4 Hz, 2H, H-1'), 3.17 (t, *J* = 4.7 Hz, 2H, H-5), 1.80–1.74 (m, 2H, H-2'), 1.53–1.45 (m, 2H, H-3'), 0.91 (t, *J* = 7.4 Hz, 3H, H-4'); <sup>13</sup>C NMR (125 MHz, DMSO-*d*<sub>6</sub>):  $\delta$  150.9, 150.0, 149.2, 146.8, 144.2, 142.0, 132.5, 130.8, 128.8, 126.1, 122.1, 118.4, 108.4, 107.9, 102.1, 75.0, 62.0, 57.1, 56.4, 31.5, 26.9, 18.6, 13.6; HRMS-ESI: *m/z* calculated for C<sub>24</sub>H<sub>26</sub>NO<sub>5</sub><sup>+</sup>: 408.1811 [M-Br]<sup>+</sup>, found for 408.1812.

**13-O-Hexylberberine bromide (7b).** Compound (**4**) was treated with *n*-hexyl bromide according to the general procedure to give the desired bromide salt (**7b**) as a bright yellow solid, yield: 76%; UV (MeOH)  $\lambda_{\max}$  (log  $\epsilon$ ): 428.0 (3.75), 350.3 (4.26), 266.5 (4.32), 229.0 (4.45) nm; <sup>1</sup>H NMR (500 MHz, DMSO-*d*<sub>6</sub>):  $\delta$  9.81 (s, 1H, H-8), 8.21 (d, *J* = 9.4 Hz, 1H, H-12), 8.09 (d, *J* = 9.4 Hz, 1H, H-11), 7.90 (s, 1H, H-1), 7.13 (s, 1H, H-4), 6.17 (s, 2H, -OCH<sub>2</sub>O-), 4.87 (t, *J* = 5.9 Hz, 2H, H-6), 4.10 (s, 3H, -OCH<sub>3</sub>), 4.09 (s, 3H, -OCH<sub>3</sub>), 3.84 (t, *J* = 6.4 Hz, 2H, H-1'), 3.16 (t, *J* = 5.9 Hz, 2H, H-5), 1.78 (p, *J* = 7.0 Hz, 2H, H-2'), 1.45 (p, *J* = 7.5 Hz, 2H, H-3'), 1.33–1.23 (m, 4H, H-4', H-5'), 0.86 (t, *J* = 6.9 Hz, 3H, H-6'); <sup>13</sup>C NMR (125 MHz, DMSO-*d*<sub>6</sub>):  $\delta$  150.9, 150.1, 149.2, 146.8, 144.2, 141.9, 132.5, 130.8, 128.8, 126.1, 122.1, 118.5, 118.4, 108.3, 108.0, 102.1, 75.4, 62.0, 57.1, 30.9, 29.4, 25.1, 22.0, 13.9; HRMS-ESI: *m/z* calculated for C<sub>26</sub>H<sub>30</sub>NO<sub>5</sub><sup>+</sup>: 436.2124 [M-Br]<sup>+</sup>, found for 436.2122.

**13-O-Octylberberine chloride (7c).** Compound (**4**) was treated with *n*-octyl bromide according to the general procedure to give the desired bromide salt (**7c**) as a bright yellow solid, yield: 61%; UV (MeOH)  $\lambda_{\max}$  (log  $\epsilon$ ): 430.6 (3.73), 349.2 (4.23), 265.7 (4.32), 234.0 (4.42) nm; <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>):  $\delta$  10.33 (s, 1H, H-8), 7.98 (d, *J* = 9.2 Hz, 1H, H-12), 7.94 (s, 1H, H-1), 7.83 (d, *J* = 9.2 Hz, 1H, H-11), 6.85 (s, 1H, H-4), 6.09 (s, 2H, -OCH<sub>2</sub>O-), 5.24 (d, *J* = 5.8 Hz, 1H, H-6), 4.34 (s, 3H, -OCH<sub>3</sub>), 4.08 (s, 3H, -OCH<sub>3</sub>), 3.83 (d, *J* = 6.5 Hz, 2H, H-1'), 3.29 (t, *J* = 5.8 Hz, 2H, H-5), 1.83 (p, *J* = 7.1 Hz, 2H, H-2'), 1.50 (p, *J* = 7.2 Hz, 2H, H-3'), 1.33–1.26 (m, 8H, H-4'–H-7'), 0.90 (t, *J* = 6.8 Hz, 3H, H-8'); <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>):  $\delta$  152.3, 150.7, 149.9, 147.1, 146.2, 143.4, 132.1, 130.9, 129.9, 125.4, 123.0, 118.5, 117.9, 108.5, 108.4, 102.0, 75.9, 63.2, 57.1, 31.8, 30.2, 29.3, 29.2, 28.2, 26.0, 22.6, 14.1; LC-MS (ESI<sup>+</sup>, *m/z*) C<sub>28</sub>H<sub>34</sub>NO<sub>5</sub><sup>+</sup>: 464.58 [M-Br]<sup>+</sup>; HRMS-ESI: *m/z* calculated for C<sub>28</sub>H<sub>34</sub>NO<sub>5</sub><sup>+</sup>: 464.2437 [M-Br]<sup>+</sup>, found for 464.2436.

**13-O-Decylberberine chloride (7d).** Compound (**4**) was treated with *n*-decyl bromide according to the general procedure to give the desired bromide salt (**7d**) as a bright yellow solid, yield: 58%; UV (MeOH)  $\lambda_{\max}$  (log  $\epsilon$ ): 428.3 (3.70), 349.6 (4.19), 265.4 (4.28), 223.8 (4.47) nm; <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>):  $\delta$  10.50 (s, 1H, H-8), 7.97 (d, *J* = 9.2 Hz, 1H, H-12), 7.94 (s, 1H, H-1), 7.81 (d, *J* = 9.2 Hz, 1H, H-11), 6.86 (s, 1H, H-4), 6.09 (s, 2H, -OCH<sub>2</sub>O-), 5.31 (d, *J* = 5.8 Hz, 1H, H-6), 4.34 (s, 3H, -OCH<sub>3</sub>), 4.09 (s, 3H, -OCH<sub>3</sub>), 3.82 (d, *J* = 6.5 Hz, 2H, H-1'), 3.28 (t, *J* = 5.8 Hz, 2H, H-5), 1.83 (p, *J* = 7.2 Hz, 2H, H-2'),

1.50 (d,  $J = 7.2$  Hz, 2H, H-3'), 1.32–1.23 (m, 12H, H-4'–H-9'), 0.90 (t,  $J = 6.9$  Hz, 3H, H-10');  $^{13}\text{C}$  NMR (125 MHz,  $\text{CDCl}_3$ ):  $\delta$  151.3, 150.6, 149.9, 147.5, 146.4, 143.9, 132.2, 130.8, 129.8, 125.3, 123.1, 118.5, 117.8, 108.4, 102.0, 75.8, 63.2, 57.0, 56.7, 31.9, 30.1, 29.5, 29.33, 29.29, 28.2, 26.0, 22.7, 14.1; LC-MS (ESI $^+$ ,  $m/z$ )  $\text{C}_{30}\text{H}_{38}\text{NO}_5^+$ : 492.62 [M–Br] $^+$ ; HRMS-ESI:  $m/z$  calculated for  $\text{C}_{30}\text{H}_{38}\text{NO}_5^+$ : 492.2750 [M–Br] $^+$ , found for 492.2751.

**13-*O*-Dodecylberberine chloride (7e).** Compound (4) was treated with *n*-dodecyl bromide according to the general procedure to give the desired bromide salt (7e) as a bright yellow solid, yield: 40%; UV (MeOH)  $\lambda_{\text{max}}$  (log  $\epsilon$ ): 427.5 (3.63), 350 (4.13), 266.4 (4.20), 225.4 (4.37) nm;  $^1\text{H}$  NMR (500 MHz,  $\text{DMSO}-d_6$ ):  $\delta$  9.80 (s, 1H, H-8), 8.20 (d,  $J = 9.4$  Hz, 1H, H-12), 8.09 (d,  $J = 9.4$  Hz, 1H, H-11), 7.89 (s, 1H, H-1), 7.13 (s, 1H, H-4), 6.16 (s, 2H,  $-\text{OCH}_2\text{O}-$ ), 4.86 (t,  $J = 5.9$  Hz, 2H, H-6), 4.09 (s, 3H,  $-\text{OCH}_3$ ), 4.08 (s, 3H,  $-\text{OCH}_3$ ), 3.84 (t,  $J = 6.4$  Hz, 2H, H-1'), 3.16 (t,  $J = 5.9$  Hz, 2H, H-5), 1.77 (p,  $J = 7.0$  Hz, 2H, H-2'), 1.43 (m, 2H, H-3'), 1.24 (m, 16H, H-4'–H-11'), 0.85 (t,  $J = 6.9$  Hz, 3H, H-12');  $^{13}\text{C}$  NMR (125 MHz,  $\text{DMSO}-d_6$ ):  $\delta$  150.9, 150.1, 149.2, 146.8, 144.2, 142.0, 132.5, 130.9, 128.8, 126.1, 122.2, 118.5, 118.4, 108.3, 108.0, 102.1, 75.4, 62.0, 57.1, 56.4, 31.3, 29.4, 29.04, 29.02, 28.96, 28.89, 28.75, 28.7, 26.9, 25.4, 22.1, 14.0; LC-MS (ESI $^+$ ,  $m/z$ )  $\text{C}_{32}\text{H}_{42}\text{NO}_5^+$ : 520.68 [M–Br] $^+$ ; HRMS-ESI:  $m/z$  calculated for  $\text{C}_{32}\text{H}_{42}\text{NO}_5^+$ : 520.3063 [M–Br] $^+$ , found for 520.3066.

### 3.5. Biological activity

#### 3.5.1. Cell lines and cell culture conditions

All cells were obtained from the Bioresource Collection and Research Center (Hsinchu, Taiwan). Human colon cancer cells (HT-29) was cultured in RPMI 1640 (HyClone, USA) medium, supplemented with 10% fetal bovine serum (FBS, Gibco, Life Technologies, USA) and antibiotics (100 mg/mL streptomycin and 100 U/mL penicillin, Gibco, Life Technologies, USA). Human bladder cancer cells (BFTC 905) and human liver carcinoma (HepG2) cell lines were cultured in Dulbecco's modified Eagle medium (DMEM, HyClone, USA) medium, supplemented with 10% fetal bovine serum (FBS) and antibiotics. Human cancer cells were maintained in humidified atmosphere with 5%  $\text{CO}_2$  and 95% air at 37°C in a carbon dioxide incubator (SANYO,  $\text{CO}_2$  incubator, Osaka, Japan).

#### 3.5.2. Cell viability inhibition assay

Cell viability was assessed by staining with 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyl tetrazolium bromide (MTT, Bionovas Biotechnology Co., Ltd., Toronto, Ontario, Canada) [29]. Cancer cells were plated at  $5 \times 10^3$ – $1 \times 10^4$  cells into 96-well plates. After overnight growth, cells were treated with various concentrations of berberine derivatives for 24 h. At the end of treatment, final concentration of 0.5 mg/mL MTT was added, and cells were incubated for a further 1.5 h. The absorbance was recorded on an ELISA plate reader (SpectraMax 340PC<sup>384</sup>, Molecular Devices, Orleans, California, USA) at a 540 nm wavelength. Inhibition ratio (%) was calculated using the following equation:

$$\text{Inhibition ratio (\%)} = [(A_{\text{control}} - A_{\text{treated}}) / A_{\text{control}}] \times 100\%,$$

$A_{\text{treated}}$  and  $A_{\text{control}}$  are the average absorbance of three independent experiments from treated and control groups, respectively.



Cell viability was determined the test compound concentration required to inhibit tumor cell proliferation by 50% (IC<sub>50</sub>) from the dose-response curves. All data average values from triplicate samples and the experiments were repeated at least three times.

### 3.5.3. Measurement of absorption and fluorescence spectra

Absorption spectrum (250-500 nm) of berberine and its derivatives (**5e**, **6e** and **7e**) were recorded on a UV-vis spectrophotometer (UV-1601, Shimadzu, Kyoto, Japan) with a quartz cuvette (path length 1 cm, accuracy  $\pm$  0.2 nm). Fluorescence emission spectrum were recorded on a fluorescence spectrophotometer (RF-5301PC, Shimadzu, Kyoto, Japan) with excitation at 420 nm and measure of emission from 450-700 nm (slit 3/3 nm) in a 1-cm quartz cuvette. A final concentration of berberine and its derivatives **5e**, **6e** and **7e** were 10  $\mu$ M diluted using methanol. Appropriate blanks corresponding to the methanol were subtracted to correct the background.

### 3.5.4. Light source

The UVA light box consisting of eight UVA lamps (Blacklight blue, F8T5BLB, Sankyo Denki Co. Ltd., Osaka, Japan) was custom made [33]. The light spectral irradiance of the light box was determined using a NIST-Traceable radiometer (ILT1400 Portable radiometer, International Light Technologies Inc., MA, USA). The light dose of the UVA light box, with the maximum emission at 420 nm, was routinely measured using a photochemical reactor (PR-1000, Panchum Scientific Corp., Kaohsiung, Taiwan). In the study, the light-irradiation doses were determined at 1.8, 3.6, 7.2 J/cm<sup>2</sup> for 420 nm (dose rate 5.9 mW/cm<sup>2</sup>), which were obtained by approximately 5, 10 and 20 minutes of exposure, respectively. To determine visible light dose using the following equation:

$$E \text{ (J/cm}^2\text{)} = t \text{ (s)} \times P \text{ (W/cm}^2\text{)},$$

where  $E$  stands for visible light energy,  $t$  represents time expressed in second and, finally,  $P$  is lamp potency.

### 3.5.5. Photocytotoxic assay

The photocytotoxicity of berberine (**1**) and its derivatives (**5e**, **6e** and **7e**) was studied by using the MTT colorimetric assay with quantification of formazan formed by cleavage of the tetrazolium rings of MTT in DMSO by spectral measurement [29]. Various concentrations of the berberine (**1**) and berberine derivatives (**5e**, **6e** and **7e**) dissolved in 1% DMSO were added to the cells and incubation was continued for 4 h in the dark followed by photo-irradiation with visible light of 420 nm (1.8, 3.6 and 7.2 J/cm<sup>2</sup> for 5, 10 and 20 min of irradiation, respectively) by using a photochemical reactor (PR-1000, Panchum Scientific Corp., Kaohsiung, Taiwan) filled with make eight fluorescent black tubes lamps (Blacklight blue, F8T5BLB, Sankyo Denki Co. Ltd., Osaka, Japan). For these experiments, the cells were seeded in 96-well microliter plates (5000 cells/well) and grown for at least 24 h in 100  $\mu$ L DMEM containing 10% FBS. The growth medium was replaced with fresh medium containing 3% FBS and berberine (5, 10, 20 and 40  $\mu$ M) or its derivatives (**5e**, **6e** and **7e**) (0.05, 0.25, 0.5 and 1.0  $\mu$ M), and the cells were incubated for 4 h at 37°C. At the end of the incubation, the drug-containing medium was removed, and the cells was washed with fresh medium and irradiated for 5, 10 and 20 min with visible light (420 nm). After the incubation period for 20 h, 10  $\mu$ L MTT (5 mg/mL) was added to each well and incubated for additional 1.5 h at 37°C,



100  $\mu$ L DMSO was added to dissolve the formazan crystals. The absorbance of the samples was measured using an ELISA microplate reader (SpectraMax 340PC<sup>384</sup>, Molecular Devices, Orleans, California, USA) and selecting 540 nm as the best wavelengths. The survival of the treated cells was calculated as the percentage of their absorbance referred to the absorbance of the untreated (no drug, no light) cells taken as 100% viability. The IC<sub>50</sub> values were obtained by nonlinear regression analysis (Sigma 11.0 software).

### 3.5.6. Intracellular uptake profiling of berberine and its derivatives in HepG2 cells

In order to establish a time profile for intracellular accumulation of these berberine derivatives, a HPLC analysis was carried out. The HepG2 cells treated with different berberine (**1**) and its derivatives (**5e**, **6e** and **7e**) were allowed to grow in 5 mL liquid culture. The cultures were incubated in a shaking incubator in the dark. After 0 h, 0.25 h, 1 h, 2 h, 4h, and 8 h, aliquots of cells were harvested by centrifuging at 6000 rpm for 5 min. The supernatant liquid was discarded and the cell pellet was washed thrice with 1 mL of 2×PBS in order to remove the uninternalized berberine and its derivatives. After this, the cells were gently resuspended in 0.2 mL methanol and then were filtered with a 0.22  $\mu$ m filter into HPLC analysis. The HPLC analysis was carried out by exciting the sample at 350 nm and emission at 530 nm.

### 3.5.7. Cell morphological assessment

To detect morphological evidence of apoptosis, cells were visualized following DNA staining with the fluorescent dye Hoechst 33258 [30]. The HepG2 cells were plated in 6-cm dish at a density of  $7 \times 10^5$  cells/well and were incubated overnight. The cells were treated with 0.5  $\mu$ M concentrations berberine (**1**), compounds **5e**, **6e** and **7e** for 4 h in the dark and treated with visible light (420 nm) for 10 min. Additionally, the HepG2 cells of compound **7e** treatment were stained with 10  $\mu$ M Hoechst 33258 (Sigma-Aldrich, St. Louis, MO, USA). After Cells were incubated in a dark room for 5 min, the cells were washed with PBS and then were observed by a fluorescent microscope (Zeiss, Axio Observer A1, Japan).

### 3.5.8. Cell cycle distribution analysis

Flow cytometry was used to obtain the cell cycle distribution and the apoptotic rate [31]. The HepG2 cells were plated at  $1 \times 10^6$  cells per 6-cm dish and cultured overnight. Then various concentrations of 0.5  $\mu$ M berberine (**1**), compounds **5e**, **6e** and **7e** were added for 2 h. At the end of the incubation, the drug-containing medium was removed, and the cells was washed with fresh medium and irradiated for 10 min with visible light (420 nm). The cells of each dish were harvested, washed once with PBS and fixed with methanol at 4°C. After 24 h, the cells of each dish were harvested, washed once with PBS. The cells were suspended in 473  $\mu$ L of PBS containing 40  $\mu$ g/mL propidium iodide (PI) and 40  $\mu$ g/mL RNase. Cells were incubated in a dark room for 30 min at room temperature then subjected to cell cycle analysis using a FACScan flow cytometer (Becton Dickinson, San Jose, CA, USA) and analyzed using the ModFit 3.0 software (Verity Software House, ME, USA).

### 3.5.9. Measurement of intracellular singlet oxygen production

Intracellular oxidant stress was monitored by measuring changes in fluorescence resulting from intracellular probe oxidation. Intracellular production of ROS, namely singlet oxygen ( $^1\text{O}_2$ ), was measured using DCFH-DA (Molecular Probes, USA) probe, respectively [32]. The level of intracellular singlet oxygen was detected using fluorescent dye DCF sensitivity. Briefly, the HepG2 ( $4 \times 10^4$  cells/ml) cells were pretreated with  $0.5 \mu\text{M}$  of berberine (**1**) and  $0.25 \mu\text{M}$  of its derivative (**5e**, **6e** and **7e**) for 2 h. At the end of the incubation, the drug-containing medium was removed, and the cells were washed with fresh medium and irradiated for 10 min with visible light (420 nm). After the incubation period for 24 h, intracellular peroxide level was detected by DCFH-DA as an intracellular fluorescence probe. After incubation with  $20 \mu\text{M}$  DCFH-DA for 1 h, the cells were harvested and washed with PBS twice. A flow cytometer (Becton Dickinson, San Jose, CA, USA) was used to detect dichlorofluorescein (DCF). Data were collected and analyzed by ModFit 3.0 software (Verity Software House, ME, USA).

#### 3.5.10. Measurement of the mitochondrial membrane potential

The changes of the mitochondrial membrane potential (MMP) were determined by using the fluorescent probe rhodamine 123 (Sigma, St Louis, USA). The dye Rh123 selectively enters mitochondria without influencing the membrane potential and are retained inside the mitochondria [33]. Once the MMP is lost, rhodamine 123 is subsequently washed out of the cells, leading to the decline of the fluorescence signal. HepG2 cells ( $1 \times 10^5$  cells/mL) were incubated with concentrations of  $0.5 \mu\text{M}$  berberine (**1**) and  $0.25 \mu\text{M}$  of its derivative (**5e**, **6e** and **7e**) for 2 h. At the end of the incubation, the drug-containing medium was removed, and the cells were washed with fresh medium and irradiated for 10 min with visible light (420 nm) for 24 h. Then stained with  $2 \text{ mg/mL}$  of rhodamine 123 for 20 min under gentle shaking. After washing with PBS, cells were immediately analyzed by flow cytometer (Becton Dickinson, San Jose, CA, USA) and data was analyzed using the ModFit 3.0 software (Verity Software House, ME, USA).

#### 3.5.11. Statistical analysis

The values shown are mean  $\pm$  SD of three independent experiments. Data are statistically evaluated by Student's *t*-test of SigmaPlot 11.0 and shown significantly different in  $*p < 0.05$ ,  $**p < 0.01$ , and  $***p < 0.001$ .

### 4. Conclusions

In summary, berberine was modified with *n*-alkyl groups of different chain length and studied for their photo-induced cytotoxicity against three human cancer cell lines. Out of fifteen synthesized derivatives, we found three; compounds **5e**, **6e** and **7e**, work the best *in vitro*, against the cancer cell lines. The increasing chain length of *n*-alkyl groups enhanced lipophilicity and the cellular permeability of the derivatives. Besides, its derivatives, compounds **5e**, **6e** and **7e** had exhibited low dark toxicity and higher photocytotoxicity toward the cancer cell lines under tested. Thus derivatives **5e**, **6e** and **7e** with a dodecyl group on the 9-, 13- or 13-O-position of berberine, respectively, showed high cellular uptake, photo-induced cytotoxicity *in vitro*, ROS generation, and induced cell apoptosis after 420 nm visible light irradiation. In the cell cycle distribution and apoptotic analysis, compound **5e**, **6e** or **7e** induced S phase arrested and apoptosis in HepG2 cells after visible light irradiation. In Hoechst 33258 staining analysis, these three compounds markedly induce apoptosis, which were further confirmed by the positive rate of annexin V-FITC/PI double staining. Taken together, the presented results showed that compounds **5e**, **6e** and **7e** increased cellular accumulation and enhanced the anticancer activity by photo-induced therapy against the cancer cells. We thus believe they could be novel anticancer drugs candidates.

**Supplementary Materials:** The following are available online.

**Author Contributions:** H.-J. Lin and J.-H. Ho suggested the research work and discussed the experimental data; H.-J. Lin, L.-C. Tsai and F.-Y. Yang finalized the experimental work, interpreted the results and prepared figures; L.-G. Chen and Y.-W. Liu conceived and designed the biological experiments; L.-L. Yang, C.-D. Kuo and J.-Y. Wu wrote the paper, edited and revised manuscript. All authors have read and agreed to the published version of the manuscript.

**Funding:** This study is supported by grants V103E2-002 and V104E2-002 from Taipei Veterans General Hospital, Taipei, Taiwan, ROC.

**Acknowledgments:** Gratitude is intended to Dr. Christopher Liu for his valuable editorial revision.

**Conflicts of Interest:** The authors declare no conflict of interest.

## References

1. WHO Report. <http://www.who.int/mediacentre/factsheets/fs297/en/index.html>.
2. Seigel, R. L.; Miller, K. D.; Jemal, A. Cancer statistics, 2015. *CA Cancer J. Clin.* **2015**, *65*, 5–29.
3. Perz, J. F.; Armstrong, G. L.; Farrington, L. A.; Hutin, Y. J. F.; Bell, B. P. The contributions of hepatitis B virus and hepatitis C virus infections to cirrhosis and primary liver cancer worldwide. *J. Hepatol.* **2006**, *45*, 529–538.
4. Weng, C. J.; Hsieh, Y. H.; Tsai, C. M.; Chu, Y. H.; Ueng, K. C.; Liu, Y. F.; Yeh, Y. H.; Su, S. C.; Chen, Y. C.; Chen, M. K.; Yang, S. F. Relationship of insulin-like growth factors system gene polymorphisms with the susceptibility and pathological development of hepatocellular carcinoma. *Ann. Surg. Oncol.* **2010**, *17*, 1808–1815.
5. Jie, S.; Li, H.; Tian, Y.; Guo, D.; Zhu, J.; Gao, S.; Jiang, L. Berberine inhibits angiogenic potential of Hep G2 cell line through VEGF down-regulation *in vitro*. *J. Gastroenterol. Hepatol.* **2011**, *26*, 179–185.
6. Zhang, Q.; Cai, L.; Zhong, G.; Luo, W. Simultaneous determination of jatrorrhizine, palmatine, berberine, and obacunone in *Phellodendri Amurensis* Cortex by RP-HPLC. *Zhongguo Zhong Yao Za Zhi* **2010**, *35*, 2061–2064.
7. Grycová, L.; Dostál, J.; Marek, R. Quaternary protoberberine alkaloids. *Phytochemistry* **2007**, *68*, 150–175.
8. Kuo, C. L.; Chi, C. W.; Liu, T. Y. The anti-inflammatory potential of berberine *in vitro* and *in vivo*. *Cancer Lett.* **2004**, *2*, 127–137.
9. Li, T. K.; Bathory, E.; LaVoie, E. J.; Srinivasan, A. R.; Olson, W. K.; Sauers, R. R.; Liu, L. F.; Pilch, D. S. Human topoisomerase I poisoning by protoberberines: potential roles for both drug-DNA and drug-enzyme interactions. *Biochemistry* **2000**, *39*, 7107–7116.
10. Kang, M. R.; Chung, I. K. Down-regulation of DNA topoisomerase II $\alpha$  in human colorectal carcinoma cells resistant to a protoberberine alkaloid, berberrubine. *Mol. Pharmacol.* **2002**, *61*, 879–884.
11. Liu, C. S.; Zheng, Y. R.; Zhang, Y. F.; Long, X. Y. Research progress on berberine with a special focus on its oral bioavailability. *Fitoterapia*, **2016**, *109*, 274–282.
12. Creasy, W. A. Biochemical effects of berberine. *Biochem. Pharmacol.* **1979**, *28*, 1081–1084.
13. Schmeller, T.; Latz-Brüning, B.; Wink, M. Biochemical activities of berberine, palmatine and sanguinarine mediating chemical defence against microorganisms and herbivores. *Phytochemistry* **1997**, *44*, 257–266.
14. Hou, Q.; Tang, X.; Liu, H.; Tang, J.; Yang, Y.; Jing, X.; Xiao, Q.; Wang, W.; Gou, X.; Wang, Z. Berberine induces cell death in human hepatoma cells *in vitro* by downregulating CD147. *Cancer Sci.* **2011**, *102*, 1287–1292.
15. Patil, J. B.; Kim, J.; Jayaprakasha, G. K. Berberine induces apoptosis in breast cancer cells (MCF-7) through mitochondrial-dependent pathway. *Eur. J. Pharmacol.* **2010**, *645*, 70–78.
16. Yan, K.; Zhang, C.; Feng, J.; Hou, L.; Yan, L.; Zhou, Z.; Liu, Z.; Liu, C.; Fan, Y.; Zheng, B.; Xu, Z. Induction of G1 cell cycle arrest and apoptosis by berberine in bladder cancer cells. *Eur. J. Pharmacol.* **2011**, *661*, 1–7.
17. Wang, L.; Liu, L.; Shi, Y.; Gao, H.; Chaturvedi, R.; Calcutt, M. W.; Hu, T.; Ren, X.; Wilson, K. T.; Polk, D. B.; Yan, F. Berberine induces caspase-independent cell death in colon tumor cells through activation of apoptosis-inducing factor. *PLoS One* **2012**, *7*, e36418.
18. Li, F.; Dong, X.; Lin, P.; Jiang, J. Regulation of Akt/FoxO3a/Skp2 axis is critically involved in berberine-induced cell cycle arrest in hepatocellular carcinoma cells. *Int. J. Mol. Sci.* **2018**, *19*, E327.
19. Lin, J. P.; Yang, J. S.; Lee, J. H.; Hsieh, W. T.; Chung, J. G. Berberine induces cell cycle arrest and apoptosis in human gastric carcinoma SNU-5 cell line. *World J. Gastroenterol.* **2006**, *12*, 21–28.

20. Qiu, F.; Zhu, Z.; Kang, N.; Piao, S.; Qin, G.; Yao, X. Isolation and identification of urinary metabolites of berberine in rats and humans. *Drug Metab. Dispos.* **2008**, *36*, 2159–2165.
21. Qin, P.; Pang, J. Y.; Chen, W. H.; Zhao, Z. Z.; Liu, L.; Jiang, Z. H. Inhibition of DNA topoisomerase I by natural and synthetic mono- and dimeric protoberberine alkaloids. *Chem. Biodivers.* **2007**, *4*, 481–487.
22. Krishnan, P.; Bastow, K. F. The 9-position in berberine analogs is an important determinant of DNA topoisomerase II inhibition. *Anticancer Drug Des.* **2000**, *15*, 255–264.
23. Zhang, W. J.; Ou, T. M.; Lu, Y. J.; Huang, Y. Y.; Wu, W. B.; Huang, Z. S.; Zhou, J. L.; Wong, K. Y.; Gu, L. Q. 9-Substituted berberine derivatives as G-quadruplex stabilizing ligands in telomeric DNA. *Bioorg. Med. Chem.* **2007**, *15*, 5493–5501.
24. Islam, M. M.; Basu, A.; Hossain, M.; Sureshkumar, G.; Hotha, S.; Kumar, G. S. Enhanced DNA binding of 9- $\omega$ -amino alkyl ether analogs from the plant alkaloid berberine. *DNA Cell Biol.* **2011**, *30*, 123–133.
25. Basu, A.; Jaisankar, P.; Kumar, G. S. Synthesis of novel 9-*O*-*N*-aryl/aryl-alkyl amino carbonyl methyl substituted berberine analogs and evaluation of DNA binding aspects. *Bioorg. Med. Chem.* **2012**, *20*, 2498–2505.
26. Bhowmik, D.; Hossain, M.; Buzzetti, F.; D'Auria, R.; Lombardi, P.; Kumar, G. S. Biophysical studies on the effect of the 13 position substitution of the anticancer alkaloid berberine on its DNA binding. *J. Phys. Chem. B* **2012**, *116*, 2314–2324.
27. Lo, C. Y.; Hsu, L. C.; Chen, M. S.; Lin, Y. J.; Chen, L. G.; Kuo, C. D.; Wu, J. Y. Synthesis and anticancer activity of a novel series of 9-*O*-substituted berberine derivatives: a lipophilic substitute role. *Bioorg. Med. Chem. Lett.* **2013**, *23*, 305–309.
28. Zhang, L.; Li, J.; Ma, F.; Yao, S.; Li, N.; Wang, J.; Wang, Y.; Wang, X.; Yao, Q. Synthesis and cytotoxicity evaluation of 13-*n*-alkyl berberine and palmatine analogues as anticancer agents. *Molecules* **2012**, *17*, 11294–11302.
29. Mosmann, T. Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays. *J. Immunol. Methods* **1983**, *65*, 55–63.
30. Lin, Y.; Sheng, M.; Ding, Y.; Zhang, N.; Song, Y.; Du, H.; Lu, N.; Yu, W. Berberine protects renal tubular cells against hypoxia/reoxygenation injury via the Sirt1/p53 pathway. *J. Nat. Med.* **2018**, *72*, 715–723.
31. Lu, C. H.; Chen, S. H.; Chang, Y. S.; Liu, Y. W.; Wu, J. Y.; Lim, Y. P.; Yu, H. I.; Lee, Y. R. Honokiol, a potential therapeutic agent, induces cell cycle arrest and program cell death in vitro and in vivo in human thyroid cancer cells. *Pharmacol Res.* **2017**, *115*, 288–298.
32. Liu, B.; Wang, G.; Yang, J.; Pan, X.; Yang, Z.; Zang, L. Berberine inhibits human hepatoma cell invasion without cytotoxicity in healthy hepatocytes. *PLoS One* **2011**, *6*, e21416.
33. Shukla, S.; Sharma, A.; Pandey, V. K.; Raisuddin, S.; Kakkar, P. Concurrent acetylation of FoxO1/3a and p53 due to sirtuins inhibition elicit Bim/PUMA mediated mitochondrial dysfunction and apoptosis in berberine-treated HepG2 cells. *Toxicol. Appl. Pharmacol.* **2016**, *291*, 70–83.
34. Zhang, Z. H.; Deng, A. J.; Wu, L. Q.; Fang, L. H.; Yu, J. Q.; Li, Z. H.; Yuan, T. Y.; Wang, W. J.; Du, G. H.; Qin, H. L. Syntheses and structure-activity relationships in cytotoxicities of 13-substituted quaternary coptisine derivatives. *Eur. J. Med. Chem.* **2014**, *86*, 542–549.
35. Chi, C. W.; Chang, Y. F.; Chao, T. W.; Chiang, S. H.; Peng, F. K.; Lui, W. Y.; Liu, T. Y. Flowcytometric analysis of the effect of berberine on the expression of glucocorticoid receptors in human hepatoma HepG2 cells. *Life Sci.* **1994**, *54*, 2099–2107.

**Sample Availability:** Samples of the compounds **2–4**, **5a–e**, **6a–e** and **7a–e** are available from the authors.