

Investigating the Gut Microbiota Composition of Individuals with Attention-Deficit/Hyperactivity Disorder and Association with Symptoms

Joanna Szopinska-Tokov M.S.^{1,&}, Sarita Dam M.S.^{2,&}, Jilly Naaijen Ph.D.², Prokopis Konstanti M.S.³, Nanda Rommelse Ph.D.^{1,4}, Clara Belzer Ph.D.³, Jan Buitelaar M.D., Ph.D.^{2,4}, Barbara Franke Ph.D.^{1,5}, Esther Aarts Ph.D.², Alejandro Arias Vasquez Ph.D.^{1,5}

¹Department of Psychiatry, Radboudumc, Donders Institute for Brain, Cognition and Behaviour, 6525 GA Nijmegen, The Netherlands

²Department of Cognitive Neuroscience, Radboudumc, Donders Institute for Brain, Cognition and Behaviour, 6525 EN Nijmegen, The Netherlands

³Laboratory of Microbiology, Wageningen University, 6708 WE Wageningen, The Netherlands

⁴Karakter Child and Adolescent Psychiatry University Center, 6525 GC Nijmegen, The Netherlands

⁵Department of Human Genetics, Radboudumc, Donders Institute for Brain, Cognition and Behaviour, 6525 GA Nijmegen, The Netherlands

&Shared responsibility.

*Correspondence:

Alejandro Arias Vasquez

Department of Psychiatry and Human Genetics, Radboudumc, Donders Institute for Brain, Cognition and Behaviour, Geert Grooteplein Zuid 10, 6525 GA, Nijmegen, The Netherlands

Email: Alejandro.AriasVasquez@radboudumc.nl

Phone: +31 (0) 24 3613970

Abstract

Attention-deficit/hyperactivity disorder (ADHD) is a common neurodevelopmental disorder. Given the growing evidence of gut microbiota being involved in psychiatric (including neurodevelopmental) disorders, we aimed to identify differences in gut microbiota composition between participants with ADHD and controls and to investigate the role of the microbiota in inattention and hyperactivity/impulsivity. Fecal samples were collected from 107 participants ($N_{\text{ADHD}}=42$; $N_{\text{controls}}=50$; $N_{\text{subthresholdADHD}}=15$; range age: 13-29 years). The relative quantification of bacterial taxa was done using 16S ribosomal RNA gene amplicon sequencing. Beta-diversity revealed significant differences in bacterial composition between participants with ADHD and healthy controls, which was also significant for inattention, but showing a trend in case of hyperactivity/impulsivity only. Ten genera showed nominal differences ($P < 0.05$) between both groups, of which seven genera were tested for their association with ADHD symptom scores (adjusting for age, sex, body mass index, time delay between feces collection and symptoms assessment, medication use and family relatedness). Our results show that variation of a genus from the *Ruminococcaceae* family (*Ruminococcaceae_UCG_004*) is associated (after multiple testing correction) with inattention symptoms, and suggest a role of gut microbiota in ADHD pathophysiology.

Key words: Gut microbiota, ADHD, 16S rRNA gene, Inattention

Introduction

Attention-deficit/hyperactivity disorder (ADHD) is a common, highly heritable [1], heterogeneous neurodevelopmental disorder with around 5% prevalence in childhood and 2.5% prevalence of adults world-wide [2]. The disorder is characterized by age-inappropriate levels of inattention and/or hyperactivity and impulsivity. ADHD has a significant social impact on patients' lives, causing disruption at school [3], work [4] and in personal relationships [5]. Typically, ADHD has its onset in childhood and can persist into adulthood. ADHD is considered a multifactorial disorder with multiple (common and rare) genetic variants, in combination with the environment, explaining its etiology and phenotypic variation [6].

Treatment of ADHD usually involves prescription of either stimulant or non-stimulant medication that target specific systems related to dopamine, noradrenaline and/or serotonin neurotransmission [7]. These pharmacological interventions are highly effective in controlling ADHD symptoms and have an approximate response rate of 70% [8]. However, medication treatment of ADHD is limited by low adherence, concerns about side effects, and absence of evidence for long-term efficacy [9, 10]. Approximately 30% of individuals with ADHD do not respond to medication or are unable to tolerate the adverse effects [7]. For (some of) those patients, non-pharmacological treatments are a suitable alternative [11].

One of the non-pharmacological interventions recently suggested to influence ADHD symptom severity is diet (e.g. elimination diet) [12, 13]. Diet can exert its effects on ADHD through the gut-brain axis. The gut-brain axis is a continuous and bidirectional communication system between the enteric and central nervous systems including the cognitive and emotional centers of the brain [14]. The gut-brain axis has been suggested to modulate the risk for several psychiatric illnesses, including ADHD [15, 16], and can influence cognitive processes, mood and brain performance [17].

A key player in the gut-brain axis is the complex ecosystem of commensal bacteria living in our gut, the microbiota [18]. DNA sequencing makes it possible to investigate the microbial composition and its potential role(s) in the risk and pathophysiology of ADHD [19]. For example, in our previously published study, we observed significantly enhanced predicted microbial biosynthesis of phenylalanine (a dopamine precursor related to an increase in the

genus *Bifidobacterium*) associated with a neural hallmark of ADHD, for instance decreased functional responses of the ventral striatum during reward anticipation [19].

In the present study, we increased the sample size of Aarts et al (16) (samples overlap ~40%) in order to investigate (i) if there are differences in gut microbiota composition between participants with ADHD and age matched controls, and if so, (ii) whether these bacteria are associated with the severity of ADHD symptoms.

Material and Methods

Study participants

To evaluate gut microbiota composition in participants with ADHD and healthy controls, we collected fecal samples from 107 Dutch (Caucasian) individuals enrolled in the follow-up of the NeuroIMAGE study [20]. Three groups were included: participants with ADHD (N=42), subthreshold ADHD (N=15; participants who did not reach the criteria for being considered as ADHD but scored too high to be considered healthy control; excluded from case-control comparisons), and healthy controls (N=50). The cohort included sibling pairs, which was taken into account in the analysis. A semi-structured diagnostic interview of DSM-IV criteria was conducted with both the participant and his/her parents using the Kiddie-Schedule for Affective Disorders and Schizophrenia (K-SADS) according to DSM-IV criteria. Clinical diagnosis was confirmed using a diagnostic algorithm which combined the diagnostic interview (K-SADS) with the Conners rating scales [20].

Continuous measures of inattention severity (IA) and hyperactivity/impulsivity severity (HI) were derived from the Conners Adult ADHD Rating Scales (CAARS; ≥ 16 years) and Conners Teacher Rating Scale (CTRS; < 16 years). For all participants using medication, ratings were based on the participant's functioning off medication. Detailed recruitment and diagnostic information can be found in the NeuroIMAGE design article [20].

Additionally, the following information was obtained from medical records: age, sex, Body Mass Index (BMI), time delay between feces collection and symptom assessment (differences in days, further on called 'diff_days'), and use of ADHD-related medication. Five samples with missing BMI values were excluded from the regression analysis. For an overview of the

participant characteristics see Table 1. Information regarding the use of ADHD medication was provided via self-report (≥ 16 years) or parental report (< 16 years) on the day of measurement. Two controls were removed because they indicated the use of ADHD medication.

Ethical approval for the study was obtained from the local research ethics committees (registration number 2012/542; NL nr.: 41950.091.12). A written informed consent was obtained from all participants and/or their parents prior to the sample and data collections.

Microbiota methods and measures

Sample collection, preparation and sequencing: The human fecal samples were collected at home by the participants and stored at 4°C. Within 24 hours after collection, the samples were transported to the laboratory, aliquoted into 1.5 ml Eppendorf tubes and stored at -80°C. The bacterial DNA was extracted using a repeated bead-beating step and the Maxwell[®] 16 Instrument (Promega, Leiden, The Netherlands), as described previously [21]. DNA purification was performed with a customized kit (AS1220; Promega). The purified bacterial DNA was measured with a NanoDrop ND-2000 spectrophotometer (Thermo Fisher Scientific, Wilmington, DE, USA), and aliquots of 20 ng/ μ l were prepared for the 2-step Polymerase Chain Reaction (PCR) reactions (including negative controls). In the first PCR, amplification of the V1-V2 region of the 16S rRNA gene was performed using previously reported primers for this region: 27F-DegS (5'GTTYGATYMTGGCTCAG) – 338RI-II (5' GCWGCC[T/A]CCCGTAGG[A/T]GT) [22]. In the second PCR, unique barcoded primers were added to each sample to allow for parallel sequencing of many different samples. The PCR product was checked using electrophoresis and purified using the CleanPCR kit (CleanNA, Alphen aan den Rijn, The Netherlands). The DNA concentration was measured using Qubit[®] 2.0 fluorometer. The purified samples were used to prepare libraries for the Illumina HiSeq PE300 sequencing platform (GATC Biotech AG, Konstanz, Germany), with final loading concentrations of 200 ng/ μ l.

Data processing: The sequenced data was analyzed through NG-Tax 16S rRNA pipeline at Wageningen University and Research (WUR, Wageningen, NL) [23]. NG-Tax identified the taxonomy of the samples based on 16S sequences using three core elements: (i) barcode-primer filtering, (ii) operational taxonomic unit (OTU) picking, in which unique sequences with

the relative abundance above 0.1% were clustered into OTUs based on a sequence similarity $\geq 98.5\%$, and (iii) taxonomic assignment using the SILVA reference database (version 128; [24]).

Filtering procedure of taxonomic data: We performed two filtering steps on the output file (BIOM-file) of NG-Tax in order to remove genera with low prevalence and to reduce the impact of the high number of absent genera (with a value of zero per sample). This step was critical to improve the power to detect the true effects of the microbiota while keeping as much information as possible. The way the genera/OTUs were identified made it impossible to disentangle if the observed values of zero correspond to true zeros (e.g. not present in the sample) or are false zeros (e.g. present but not detected). The two filtering steps were applied as follows: (i) the OTU table was filtered at the genus level, where a genus with non-zero values in less than 10% of the samples was removed, and (ii) the OTU table was filtered at sample level, at which a sample with less than 10% of genera was removed (Figure S1). The results of 16S rRNA analysis after filtering of taxonomic data can be found in the Supplemental Results.

Sequencing depth comparison: The sequencing depth of the microbiota data was compared between all groups (ADHD, controls and subthreshold ADHD) by performing a Kruskal-Wallis H test on the total reads. This was done in order to assess equal distribution of the sequence reads across the groups, which helps to verify the effect of any technical variation between the groups. The results can be found in the Supplemental Results, which indicated no differences.

Within-sample diversity metrics: Three alpha-diversity metrics were applied on the OTU level: (1) the species richness estimator, counting the observed unique OTUs in each sample [25]; (2) Shannon-Wiener diversity [25] index, which takes into consideration not only the number of observed unique OTUs but also their abundance; (3) the phylogenetic richness estimator, which estimates microbial diversity across a phylogenetic tree (Faiths' phylogenetic diversity) [26]. The alpha-diversity metrics were calculated using the 'alpha_diversity.py' script in QIIME 1.9.1 [27] and compared between participants with ADHD and controls.

Between-sample diversity metrics: To assesses beta-diversity, we used the weighted UniFrac distance metric, a phylogenetic-based assessment of the difference in overall bacterial community composition at the OTU level [28]. To analyze the beta-diversity, multivariate

statistics were conducted using ADONIS and betadisper functions in the R package vegan version 2.5–2 [29, 30]. Through ADONIS, we determined if the tested variables (i.e. disease status or symptom counts) influenced beta-diversity [29]. Betadisper measures the variability in OTU composition among groups (here ADHD and controls) [31]. Principal Coordinates Analysis (PCoA) was performed using the R function phyloseq::ordinate to determine and visualize whether there is a clear discrimination of microbial composition between the two groups.

Taxonomic composition analysis and associations with symptoms: Taxonomic composition of the gut microbiota was investigated at the phylum and genus levels after transforming the sequencing read counts into microbial relative abundance (normalization step). Any unknown taxonomic level (e.g. unknown genus) was assigned to the next highest known taxonomic rank (e.g. family). The composition analysis was calculated using QIIME 1.9.1 with the ‘summarize_taxa.py’ script [27]. Microbiota compositional data are highly skewed given the high number of zeros. We used Linear discriminant analysis Effect Size (LEfSe; <https://huttenhower.sph.harvard.edu/galaxy>) for statistical analysis and visualization of the results. LEfSe uses non-parametric statistics (which is less sensitive to the extreme values [32]), in our case the Kruskal Wallis sum-rank test [33], to identify (nominal) statistical differences in the relative abundance of gut microbiota between participants with ADHD and controls. In order to maximize information content and, at the same time, include as many genera as possible, we included only genera with non-zero values in at least 10% of the samples in both groups. To prioritize the selection of candidate taxa without making any claims of association (with ADHD), genera showing nominal statistical differences ($p < 0.05$, uncorrected) were selected for downstream correlation and linear regression analyses.

Linear mixed regression analysis was performed to associate bacterial relative abundance with inattention or hyperactivity/impulsivity score available for all participants (including participants with “subthreshold ADHD”). Models were adjusted for age, sex, BMI, diff_days and including the family relatedness as random factor, this was done by using the R function lme4::lmer. Given the skewed distribution of the microbial relative abundance prior to the association analyses, we investigated linear regression assumptions and identified and removed extreme and influential samples (outliers). Outliers are known to have a significant

effect on the regression model (but not on the non-parametric test [32]). Outliers were defined by Cook's distance above $(4/n)$ (where n is the number of observations) and Leverage value above $(3 \times (k+1)/n)$ (where k is the number of independent variables, in our case $k=5$) [34, 35]. Cook's distance identifies influential values, which do not have to be necessarily the extreme ones; these can be identified by Leverage. Therefore, a sample was excluded from the analysis only if it scored above the threshold for both values. Regression analyses were corrected for multiple testing using the false discovery rate (FDR; in total corrected for 14 tests) and indicated as q-values (Q).

Effect of medication on the regression results and on gut microbiota composition: Often ADHD patients are medicated, and some studies show that medication can influence gut microbiota composition [36, 37]. Therefore, we explored an effect of ADHD medication on our (regression) results and on gut microbiota composition at the genus level. The regression model could not simply be adjusted for medication due to the large number of non-medicated cases, who do not equate to healthy controls. Thus, the medicated cases ($N=19$) were removed from the regression model to see how this effects the results.

Correlation analysis and multiple regression with all selected genera: The gut microbiota is a highly complex ecosystem of interacting organisms. In order to investigate the (in)dependent effect of the selected genera on symptoms, we investigated their correlation structure and performed multiple regression analysis. The genus-genus correlation was assessed based on Spearman's rank correlation coefficient. Multiple regression analysis was performed for the same selected genera tested in the univariate models, adjusting for age, sex, BMI, diff_days and family relatedness as a random factor; the analysis was done without the samples identified as outliers (see above). If not mentioned otherwise, the data were analyzed using IBM SPSS for Windows 25.0 software (IBM Corp., Armonk, NY, USA), and the analysis was preceded by the Shapiro-Wilk's normality test.

Results

Subjects characteristics

The general characteristics of the studied sample are presented in Table 1. Mean age, median BMI, percentage of males and differences in days between fecal collection and ADHD symptoms assessment (diff_days) were similar among the two groups. As expected, mean inattention and hyperactivity/impulsivity scores were statistically different between the ADHD and control groups. Out of the 41 participants with ADHD, 19 were using medication for ADHD.

Table 1. Characteristics of the sample.

	ADHD	Control	Subthreshold ADHD	P-value ^a
N	41	47	15	-
Age, mean (SD)	20.2 (4.2)	20.5 (3.5)	20.2 (3.3)	NS
Age, range	13-29	13-28	14-26	-
BMI, median (IQR)	23 (20-26)	22 (20-24)	22 (20-25)	NS
BMI, range	16-31	16-31	20-30	-
BMI ≥ 25, %	29	19	20	NS
Male, %	63	49	40	NS
Use of ADHD medication, N	19	0	3	-
Diff_days, median (IQR)	17 (14-34)	30 (12-70)	16 (10-33)	NS
Conners'				
Inattention, mean (SD)	66.3 (12.8)	46.8 (12.5)	58.1 (11.5)	<0.001
Hyperactivity/Impulsivity, mean (SD)	59 (12.2)	44.9 (13.0)	59.3 (12.8)	<0.001

^acomparison made for ADHD vs. controls; t-test, Mann-Whitney or chi-square test were applied accordingly; one sample had missing value for inattention and hyperactivity/impulsivity scores; four samples had missing value for BMI; NS = not significant; SD = standard deviation; IQR = interquartile rang; diff_days = represents differences in days between fecal collection and Conner's assessment.

Microbiota measures

Within- and between-sample diversity metrics: None of the three alpha-diversity (within-sample diversity) measures showed significant differences between the ADHD and control groups (Figure S2).

Beta-diversity (between-sample diversity), assessed using betadisper [38], showed that the ADHD group had a smaller variation in the gut microbiota composition, which means a higher taxonomic similarity (within the group) compared to controls ($P=0.004$; Figure 1 and Figure S3). ADONIS revealed a significant effect of ADHD diagnosis and symptom severity on the variation in the beta-diversity. The variation in beta-diversity was significantly explained by disorder status ($N=88$; variance explained=3.2%; $P=0.033$), and inattention score (IA) ($N=102$; variance explained=3.7%; $P=0.014$), whereas hyperactivity/impulsivity score (HI) was at trend level ($N=102$; variance explained=2.4%; ($P=0.059$). Age, sex, BMI and medication did not have a significant effect on beta-diversity (Table 2). Additionally, PCoA based on weighted UniFrac distance did not show a clear discrimination of microbial composition between the two groups determined by disorder status (ADHD vs. controls) (Figure S3).




Figure 1. Boxplot of multivariate homogeneity of groups' dispersions (betadisper) of participants with ADHD and controls. Box plots represent median with whiskers on ± 1.5 IQR.

*Pseudo-F=9.658, $P=0.004$.

Table 2. Beta diversity analysis.

Variable	N	R ²	Pseudo-F	P-value
Disorder status	88	0.032	2.85	0.033
Age	103	0.004	0.41	0.853
Sex	103	0.011	1.10	0.297
BMI	98	0.005	0.44	0.848
IA	102	0.037	3.87	0.014
HI	102	0.024	2.45	0.059
medication	41	0.022	0.86	0.483

Results of ADONIS on weighted UniFrac dissimilarity matrix including six tests for: disorder status, age, sex, BMI, Inattention (IA) and Hyperactivity/Impulsivity (HI) variables; R² = variance explained, a measure of effect size; Pseudo-F = indicator of the number of clusters, the larger pseudo-F value the greater between-group variation than the within-group variation.

Taxonomic composition analysis and associations with symptoms: As expected [39], compositional analysis of our samples revealed that *Firmicutes*, *Bacteroidetes*, *Actinobacteria*, *Proteobacteria* and *Verrucomicrobia*, were the most frequent phyla in our data (Table S2). There were no significant differences in the relative abundance of any of these phyla between participants with ADHD and controls (Table S2).

At the genus level, differences in the gut microbiota composition revealed nominal significant case-control differences for ten genera ($P < 0.05$; Figure 2). Of those, nine genera were increased and one was decreased in participants with ADHD. Based on their prevalence (present in at least 10% of the samples in each group; see methods), seven were selected for downstream association analyses with ADHD symptom scores (inattention and hyperactivity/impulsivity scores) (Figures S4 and S5). One genus, *Ruminococcaceae_UCG_004*, was associated ($B=39.291$, $P=0.002$, $Q=0.027$; corrected for multiple testing; Table 3) with inattention scores and two other genera showed nominal associations ($P < 0.05$). We did not find any association between tested genera and hyperactivity/impulsivity score (before or after correcting for multiple testing; all $P > 0.05$); therefore only IA was taken into consideration in further analyses.




Figure 2. Comparison of bacterial relative abundance between participants with ADHD and controls. Identification of the bacteria differences was done by Kruskal-Wallis test and visualized by LEfSe. Nominal significant threshold: $P < 0.05$.

Table 3. Association of the selected^a genera with ADHD symptoms scores.

	Inattention				Hyperactivity/Impulsivity			
	N	B (S.E.) ^b	95% CI	P-value	N	B (S.E.) ^b	95% CI	P-value
<i>Clostridiales_g__</i>	97	-1.467 (3.077)	-7.787 - 5.874	0.634	98	-3.125 (2.439)	-7.922 - 2.234	0.204
<i>Family_XIII_AD3011_group</i>	98	5.323 (2.779)	-0.145 - 11.483	0.059	98	0.316 (2.709)	-4.879 - 6.004	0.907
<i>Ruminococcaceae_UCG_005</i>	97	1.495 (1.647)	-1.759 - 4.986	0.367	97	0.175 (1.610)	-2.904 - 3.384	0.914
<i>Ruminococcus_2</i>	98	1.098 (0.445)	0.246 - 1.959	0.016	98	0.572 (0.440)	-0.268 - 1.429	0.197
<i>Ruminococcaceae_uncultured</i>	96	12.241 (5.011)	2.619 - 22.264	0.017	93	9.996 (9.191)	-7.132 - 28.016	0.279
<i>Ruminococcaceae_NK4A214_group</i>	97	3.392 (1.860)	-0.230 - 7.339	0.071	98	2.428 (1.522)	-0.513 - 5.512	0.114
<i>Ruminococcaceae_UCG_004</i>	93	39.291 (12.296)	15.329 - 64.513	0.002*	93	12.324(12.147)	-10.849 - 36.385	0.313

Linear regression models for the relative abundance of the selected genera from the LEfSe pipeline with the ADHD symptoms scores (inattention & hyperactivity/impulsivity) measured from participants with ADHD and controls and subthreshold ADHD; ^aThe selection of the genera was done prior to regression analysis and it was done based on their prevalence (see the method section); ^b Linear regression model without samples removed based on Cook's distance and Leverage threshold; models adjusted for age, sex, BMI, diff_days and a random factor for family relatedness; * Significant associations after multiple testing correction (FDR); N=number of samples after the removal of outliers (N=98 means no outliers were removed); B=coefficient; S.E.=standard error; CI=Confidence Interval

Effect of medication on the regression results and on gut microbiota composition: We tested the effect of ADHD medication on the (regression) results by excluding medicated cases (N=19) from the analysis. We found that medication did not influence the association between *Ruminococcaceae_UCG_004* and symptoms of inattention (B=47.083, P=0.0006 vs results in Table 3). When comparing the medicated (N=19) vs. non-medicated (N=22) individuals with ADHD, we found that the genus *Dialister* was increased and that the genus *Phascolarctobacterium* decreased in medicated ADHD (Figure S6). Regarding the *Phascolarctobacterium* results, we had to treat them with caution because of having only three non-zeros values for medicated cases. It is difficult to disentangle whether the observed values of zero correspond to true zeros (e.g. not present in the sample) or of false zeros (e.g. present but not detected).

Correlation analysis and multiple regression with all selected genera: Spearman correlation analysis showed that two genera, *Ruminococcaceae_NK4A214_group* and

Ruminococcaceae_UCG_005, had a strong positive correlation ($r > 0.50$) with each other. The other genera showed moderate ($0.30 > r > 0.50$), weak ($0.30 > r > 0.10$) or no correlation (Figure S7). Due to the variability in the correlation, we carried out a multiple regression analysis including all selected genera in one model (Table S3) and investigated the unique contribution of the associated genera to inattention symptoms. Regardless of correlation structure, removing one of the strongly correlated taxa (*Ruminococcaceae_NK4A214_group* or *Ruminococcaceae_UCG_005*) did not significantly change the results. After controlling for other bacterial taxa (and age, sex, BMI, diff_days and a random factor for family relatedness), *Ruminococcaceae_UCG_004* again showed the highest association with inattention score ($B = 43.920$, $P = 0.001$), followed by *Ruminococcus_2* ($B = 1.525$, $P = 0.001$). These results suggest an independent effect of these genera on inattention (Table S3).

Discussion

In this study, we aimed to determine the differences in gut microbiota composition between individuals with ADHD and controls and the association between the abundance of the selected genera and the severity of ADHD symptoms (inattention and hyperactivity/impulsivity) accounting for the effects of medication.

To our knowledge, our findings constitute the first (proof-of-concept) of its kind showing the association between the microbiome relative abundance and ADHD symptoms. Our results showed general differences in microbiota composition (beta-diversity) between the groups as well as differences at the genus level, where the relative abundance of the *Ruminococcaceae_UCG_004* genus was associated with ADHD inattention symptoms. Multiple regression analysis suggested that the association between *Ruminococcaceae_UCG_004* and inattention was independent of other (selected) genera. Our results also point towards a potential effect of genera such as *Ruminococcus_2* and *Ruminococcaceae_uncultured* on inattention.

Importantly, our result indicate that ADHD medication did not have an effect on *Ruminococcaceae_UCG_004* association. However, when directly comparing medicated ($n = 19$) and non-medicated individuals ($n = 22$) with ADHD, we found two genera being different: *Dialister* was increased and *Phascolarctobacterium* was decreased in medicated

individuals. While this is not the first time that ADHD medication is examined in terms of an effect on gut microbiota [40], it is the first time that gut microbiota composition at the genus level was assessed. However, please note that these (secondary) results should be interpreted with caution given the small sample sizes of the sub-groups.

Our results are in line with the growing evidence that the gut microbiome is involved in neurodevelopmental disorders. The *Ruminococcaceae* family and the genera belonging to this family have shown altered relative abundance in individuals diagnosed with several psychiatric diseases, i.e. autism, bipolar disorder, anxiety, depression, schizophrenia as well as ADHD [41-44]. Even though there is a lack of knowledge about the functional role of *Ruminococcaceae_UCG_004* genus and its effect on the host, the *Ruminococcaceae* family is well described. This family is commonly present in the mammalian intestinal tract, displaying the ability to degrade cellulose and hemicellulose from plant material [45]. These compounds are subsequently fermented and converted to short-chain fatty acids (SCFAs), which can be absorbed and used as energy source by the host [46]. SCFAs (acetate, propionate and butyrate) are produced (among others) by strains of *Ruminococcus* [47], which are known for mucosal colonization [48], and have been shown to play a potential role in autism [49, 50] and in ADHD [51]. Furthermore, *Ruminococcus_2*, became significantly associated with symptoms of inattention in the multivariate analysis (Figure S5). This suggests that, even in the absence of significant correlations with other bacteria (Table S4), there could be a bacterial community interaction pattern that explains more variance than a genus alone on inattention symptoms. Lastly, our lab recently performed a separate study, in which six randomly selected samples from the NeuroIMAGE cohort (same cohort studied here) were used in an animal study of human fecal microbiota transplantation into germ-free wild-type mice [52]. Mice colonized with ADHD gut microbiota had increased anxiety-like behavior and showed significantly altered structural and functional brain characteristics. In this study, *Ruminococcaceae_UCG_004* was positively correlated with a specific anxiety measure in the mice. Taken together, these results suggest that genera belonging to *Ruminococcaceae* family can exert a specific effect on neurobiological processes key to brain development and several psychiatric diseases including ADHD.

This study should be viewed in the context of several strengths and limitations. Our strengths include the use of a sample with high-quality clinical assessment and of age-matched clinically ascertained controls. The limitations of our study include i) limited sample size (although it is the largest sample of its kind so far, N=98) and ii) we were not able to collect information on lifestyle, dietary patterns (including probiotics) or antibiotic use at the time of feces collection. For the former, we applied two QC steps in order to deal with the big number of variables (genera), their expected small effects and big interindividual variation of the gut microbiota. First, we applied an uncorrected non-parametric approach (to identify differences between two groups, reduce number of variables and prioritize the selection of candidate taxa). Second, we applied an outlier detection step, prior to our regression analysis, in order to reduce the chance of false positives/negatives. For the latter, we were only able to collect information on BMI and while we acknowledge that this is not enough to account for the effects of diet and lifestyle, it is encouraging to see that there was no difference between the groups. Moreover, we looked for and removed samples with a very low bacterial diversity (high proportion of zeros) by applying a 10% genus-based frequency cut-off per sample. This can be used as a proxy for those individuals currently using antibiotics which would show a smaller bacterial diversity.

ADHD studies investigating gut microbiota composition show inconsistent results [19, 40, 44, 53, 54]. For instance, the study of Aarts et al. (2017) [19] investigated samples that show important overlap (around 40%) with our data set. This paper reported that *Bifidobacterium* genus showed the largest difference (nominally significant) between the ADHD group and controls. We did not replicate this difference in the current study. The reason for the lack of replication between studies could be due to methodological differences. These include DNA extraction [55], 16S rRNA gene region [56], bioinformatic pipeline, data processing and analysis [57], sample size and study design. Moreover, the age range (children vs. adults) differences between studies can underscore variation in ADHD symptoms, other non-shared environmental influences and gut microbiome composition [58, 59]. Follow-up studies (keeping comparable methods and including dietary patterns, comorbid conditions (of ADHD) and bacterial transcriptomics, metabolomics and metagenomics) are needed to replicate the current findings and to understand the complex biological mechanisms underlying our results.

In conclusion, our work underlines the potential role of the gut microbiota in ADHD, by indicating the differences in microbiota composition compared to controls and linking these differences with inattention severity. Further studies should validate the present findings and identify potential gut-brain mechanisms via genera belonging to the *Ruminococcaceae* family, such as those related to SCFA production.

Supplementary Materials: include Supplemental Results, Tables and Figures; **Figure S1.** Filtering procedure applied in our study; **Figure S2.** Alpha diversity comparison of the gut microbiome between participants with ADHD and controls; **Figure S3.** Principal coordinates analysis (PCoA) of weighted UniFrac distances representing microbial composition of participants with ADHD and controls; **Figure S4.** Distribution of the selected genera for behavioral analysis for each group; **Figure S5.** Distribution of the non-selected genera for behavioral analysis for each group; **Figure S6.** Boxplot of two genera relative abundance being different between medicated and non-medicated participants with ADHD; **Figure S7.** Spearman's rank correlation matrix of the selected bacterial taxa; **Table S1.** Total number of reads of participants with ADHD, controls and subthreshold ADHD; **Table S2.** Relative abundance of bacterial phyla of all participants in our study; **Table S3.** Multiple regression model with the selected genera on inattention.

Acknowledgments: This project has received funding from the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie project MiND (grant agreement no. 643051). The funding body had no role in the manuscript. The research leading to these results also received funding from the European Community's Horizon 2020 research and innovation programme through the Eat2beNICE (grant agreement no. 728018), CoCA (grant agreement no. 667302) and CANDY (grant agreement no. 847818) projects, and Netherlands Organization for Scientific Research (NWO) Food Cognition and Behaviour Program in relation to the B3: Brain, Bacteria & Behaviour project (grant agreement number 057-14-005).

The authors would like to thank Steven Aalvink for his contribution in the laboratory analyses and the technicians from Multifactorial Disease group especially Marlies Schimmel-Naber and Saskia van der Marel for administrative and pre-laboratory processing of microbial data. We thank Gerben Hermes, Alexander Kurilshikov, Marianne Jonker, Gido Schoenmacker and Todor Tokov for helpful discussions. All the authors are grateful to the study participants and their families.

Author Contributions: JST performed laboratory analyses, analyzed and interpreted the data and wrote the manuscript. SD performed laboratory analyses, contributed to the interpretation of the data and to the first version of the manuscript. JN provided clinical

information regarding ADHD diagnosis and contributed to data analysis. PK contributed to the laboratory and data analysis. CB supervised the laboratory process and the analysis of 16S rRNA data. JB organized the cohort recruitment, provided clinical information regarding ADHD diagnosis. AAV designed and supervised the study. BF organized the cohort recruitment and supported design of the study. All authors were involved in editing the manuscript. All authors have read and agreed to the published version of the manuscript.

Conflict of Interest: Barbara Franke has received educational speaking fees from Medice. Jan Buitelaar has served as a consultant, advisory board member, and/or speaker for Takeda/Shire, Roche, Medice, Janssen Cilag, and Servier. The remaining authors declare no competing interests.

References

1. Franke, B., et al., *The genetics of attention deficit/hyperactivity disorder in adults, a review*. Mol Psychiatry, 2012. **17**(10): p. 960-87.
2. Polanczyk, G., et al., *The worldwide prevalence of ADHD: a systematic review and meta-regression analysis*. Am J Psychiatry, 2007. **164**(6): p. 942-8.
3. Fredriksen, M., et al., *Childhood and persistent ADHD symptoms associated with educational failure and long-term occupational disability in adult ADHD*. Atten Defic Hyperact Disord, 2014. **6**(2): p. 87-99.
4. Adamou, M., et al., *Occupational issues of adults with ADHD*. BMC Psychiatry, 2013. **13**: p. 59.
5. Franke, B., et al., *Live fast, die young? A review on the developmental trajectories of ADHD across the lifespan*. European neuropsychopharmacology : the journal of the European College of Neuropsychopharmacology, 2018.
6. Langley, K., *ADHD genetics*. Oxford Textbook of Attention Deficit Hyperactivity Disorder, 2018: p. 19.
7. Cortese, S., et al., *Comparative efficacy and tolerability of medications for attention-deficit hyperactivity disorder in children, adolescents, and adults: a systematic review and network meta-analysis*. Lancet Psychiatry, 2018. **5**(9): p. 727-738.
8. Childress, A.C. and F.R. Sallee, *Attention-deficit/hyperactivity disorder with inadequate response to stimulants: approaches to management*. CNS drugs, 2014. **28**(2): p. 121-9.
9. O'Callaghan, P., *Adherence to stimulants in adult ADHD*. Attention deficit and hyperactivity disorders, 2014. **6**(2): p. 111-20.
10. Ghuman, J.K. and H.S. Ghuman, *Pharmacologic intervention for attention-deficit hyperactivity disorder in preschoolers : is it justified?* Paediatric drugs, 2013. **15**(1): p. 1-8.
11. De Crescenzo, F., et al., *Pharmacological and non-pharmacological treatment of adults with ADHD: a meta-review*. Evid Based Ment Health, 2017. **20**(1): p. 4-11.
12. Ly, V., et al., *Elimination diets' efficacy and mechanisms in attention deficit hyperactivity disorder and autism spectrum disorder*. European Child & Adolescent Psychiatry, 2017. **26**(9): p. 1067-1079.
13. Pelsser, L.M., et al., *Diet and ADHD, Reviewing the Evidence: A Systematic Review of Meta-Analyses of Double-Blind Placebo-Controlled Trials Evaluating the Efficacy of Diet Interventions on the Behavior of Children with ADHD*. PLoS One, 2017. **12**(1): p. e0169277.
14. Vuong, H.E., et al., *The Microbiome and Host Behavior*. Annu Rev Neurosci, 2017. **40**: p. 21-49.

15. Hsiao, E.Y., et al., *Microbiota modulate behavioral and physiological abnormalities associated with neurodevelopmental disorders*. Cell, 2013. **155**(7): p. 1451-63.
16. Bull-Larsen, S. and M.H. Mohajeri, *The Potential Influence of the Bacterial Microbiome on the Development and Progression of ADHD*. Nutrients, 2019. **11**(11).
17. Dong, W., et al., *Influence of age-related learning and memory capacity of mice: different effects of a high and low caloric diet*. Aging Clin Exp Res, 2016. **28**(2): p. 303-11.
18. Carabotti, M., et al., *The gut-brain axis: interactions between enteric microbiota, central and enteric nervous systems*. Ann Gastroenterol, 2015. **28**(2): p. 203-209.
19. Aarts, E., et al., *Gut microbiome in ADHD and its relation to neural reward anticipation*. PLoS One, 2017. **12**(9): p. e0183509.
20. von Rhein, D., et al., *The NeuroIMAGE study: a prospective phenotypic, cognitive, genetic and MRI study in children with attention-deficit/hyperactivity disorder. Design and descriptives*. Eur Child Adolesc Psychiatry, 2015. **24**(3): p. 265-81.
21. Fernandez-Calleja, J.M.S., et al., *Non-invasive continuous real-time in vivo analysis of microbial hydrogen production shows adaptation to fermentable carbohydrates in mice*. Sci Rep, 2018. **8**(1): p. 15351.
22. van den Bogert, B., et al., *Diversity of human small intestinal Streptococcus and Veillonella populations*. FEMS Microbiol Ecol, 2013. **85**(2): p. 376-88.
23. Ramiro-Garcia, J., et al., *NG-Tax, a highly accurate and validated pipeline for analysis of 16S rRNA amplicons from complex biomes* F1000Research, 2016. **5**:1791
24. Quast, C., et al., *The SILVA ribosomal RNA gene database project: improved data processing and web-based tools*. Nucleic Acids Res, 2013. **41**(Database issue): p. D590-6.
25. Morris, E.K., et al., *Choosing and using diversity indices: insights for ecological applications from the German Biodiversity Exploratories*. Ecology and evolution, 2014. **4**(18): p. 3514-24.
26. Faith, D.P., *Conservation evaluation and phylogenetic diversity*. Biological Conservation, 1992. **61**(1): p. 1-10.
27. Caporaso, J.G., et al., *QIIME allows analysis of high-throughput community sequencing data*. Nat Methods, 2010. **7**(5): p. 335-6.
28. Ursell, L.K., et al., *Defining the human microbiome*. Nutr Rev, 2012. **70** Suppl 1: p. S38-44.
29. Anderson, M.J., *A new method for non-parametric multivariate analysis of variance*. Austral Ecology, 2001. **26**(1): p. 32-46.
30. Anderson, M.J., K.E. Ellingsen, and B.H. McArdle, *Multivariate dispersion as a measure of beta diversity*. Ecology Letters, 2006. **9**(6): p. 683-693.

31. Anderson, M.J., *Distance-based tests for homogeneity of multivariate dispersions*. Biometrics, 2006. **62**(1): p. 245-53.
32. Kitchen, C.M., *Nonparametric vs parametric tests of location in biomedical research*. Am J Ophthalmol, 2009. **147**(4): p. 571-2.
33. Segata, N., et al., *Metagenomic biomarker discovery and explanation*. Genome Biol, 2011. **12**(6): p. R60.
34. Altman, N. and M. Krzywinski, *Analyzing outliers: influential or nuisance?* Nature Methods, 2016. **13**(4): p. 281-282.
35. Field, A., *Discovering statistics using SPSS : (and sex and drugs and rock 'n' roll)*. SAGE, 2009(3rd edition): p. 217-219.
36. Cusotto, S., et al., *Differential effects of psychotropic drugs on microbiome composition and gastrointestinal function*. Psychopharmacology (Berl), 2019. **236**(5): p. 1671-1685.
37. Maier, L., et al., *Extensive impact of non-antibiotic drugs on human gut bacteria*. Nature, 2018. **555**(7698): p. 623-628.
38. Anderson, M.J., K.E. Ellingsen, and B.H. McArdle, *Multivariate dispersion as a measure of beta diversity*. Ecol Lett, 2006. **9**(6): p. 683-93.
39. D'Argenio, V. and F. Salvatore, *The role of the gut microbiome in the healthy adult status*. Clin Chim Acta, 2015. **451**(Pt A): p. 97-102.
40. Prehn-Kristensen, A., et al., *Reduced microbiome alpha diversity in young patients with ADHD*. PLoS One, 2018. **13**(7): p. e0200728.
41. Rose, D.R., et al., *Differential immune responses and microbiota profiles in children with autism spectrum disorders and co-morbid gastrointestinal symptoms*. Brain Behav Immun, 2018. **70**: p. 354-368.
42. Painold, A., et al., *A step ahead: Exploring the gut microbiota in inpatients with bipolar disorder during a depressive episode*. Bipolar Disord, 2018.
43. Nguyen, T.T., et al., *Differences in gut microbiome composition between persons with chronic schizophrenia and healthy comparison subjects*. Schizophr Res, 2018.
44. Jiang, H.Y., et al., *Gut microbiota profiles in treatment-naive children with attention deficit hyperactivity disorder*. Behav Brain Res, 2018. **347**: p. 408-413.
45. Flint, H.J., et al., *Microbial degradation of complex carbohydrates in the gut*. Gut Microbes, 2012. **3**(4): p. 289-306.
46. den Besten, G., et al., *The role of short-chain fatty acids in the interplay between diet, gut microbiota, and host energy metabolism*. J Lipid Res, 2013. **54**(9): p. 2325-40.

47. Kunal Maniar, V.S., Deepak Kumar, Amal Moideen, Rajasri Bhattacharyya, Dibyajyoti Banerjee, *Chapter 42 - Metformin: A Candidate Drug to Control the Epidemic of Diabetes and Obesity by Way of Gut Microbiome Modification*. Academic Press, 2019: p. 401-408.
48. Sicard, J.F., et al., *Interactions of Intestinal Bacteria with Components of the Intestinal Mucus*. *Front Cell Infect Microbiol*, 2017. **7**: p. 387.
49. Wang, L., et al., *Elevated fecal short chain fatty acid and ammonia concentrations in children with autism spectrum disorder*. *Dig Dis Sci*, 2012. **57**(8): p. 2096-102.
50. Abdelli, L.S., A. Samsam, and S.A. Naser, *Propionic Acid Induces Gliosis and Neuro-inflammation through Modulation of PTEN/AKT Pathway in Autism Spectrum Disorder*. *Sci Rep*, 2019. **9**(1): p. 8824.
51. Dam, S.A., et al., *The Role of the Gut-Brain Axis in Attention-Deficit/Hyperactivity Disorder*. *Gastroenterology Clinics*, 2019.
52. Anouk C Tengeler, S.A.D., Maximilian Wiesmann, Jilly Naaijen, Miranda van Bodegom, Clara Belzer, Pieter J Dederen, Vivienne Verweij, Barbara Franke, Tamas Kozicz, Alejandro Arias Vasquez, Amanda J Kiliaan, *Gut microbiota from persons with attention-deficit/hyperactivity disorder affects the brain in mice*. submitted, 2020.
53. Hassan, A., *Characterizing A Link Between Gut Microbiome and Attention Deficit Hyperactive Disorder*. Honors College Research Collection. 4., 2017.
54. Wang, L.J., et al., *Gut microbiota and dietary patterns in children with attention-deficit/hyperactivity disorder*. *Eur Child Adolesc Psychiatry*, 2019.
55. Szopinska, J.W., et al., *Reliability of a participant-friendly fecal collection method for microbiome analyses: a step towards large sample size investigation*. *BMC Microbiol*, 2018. **18**(1): p. 110.
56. Rintala, A., et al., *Gut Microbiota Analysis Results Are Highly Dependent on the 16S rRNA Gene Target Region, Whereas the Impact of DNA Extraction Is Minor*. *J Biomol Tech*, 2017. **28**(1): p. 19-30.
57. Allali, I., et al., *A comparison of sequencing platforms and bioinformatics pipelines for compositional analysis of the gut microbiome*. *BMC Microbiol*, 2017. **17**(1): p. 194.
58. Hollister, E.B., et al., *Structure and function of the healthy pre-adolescent pediatric gut microbiome*. *Microbiome*, 2015. **3**: p. 36.
59. Wilens, T.E., et al., *Presenting ADHD symptoms, subtypes, and comorbid disorders in clinically referred adults with ADHD*. *J Clin Psychiatry*, 2009. **70**(11): p. 1557-62.