

# Another Criterion For The Riemann Hypothesis

Frank Vega

CopSonic, 1471 Route de Saint-Nauphary 82000 Montauban, France

## Abstract

Let's define  $\delta(x) = (\sum_{q \leq x} \frac{1}{q} - \log \log x - B)$ , where  $B \approx 0.2614972128$  is the Meissel-Mertens constant. The Robin theorem states that  $\delta(x)$  changes sign infinitely often. For  $x \geq 2$ , Nicolas defined the function  $u(x) = \sum_{q > x} (\log(\frac{q}{q-1}) - \frac{1}{q})$  and proved that  $0 < u(x) \leq \frac{1}{2 \times (x-1)}$ . We define the another function  $\varpi(x) = (\sum_{q \leq x} \frac{1}{q} - \log \log \theta(x) - B)$ , where  $\theta(x)$  is the Chebyshev function. Using the Nicolas theorem, we demonstrate that the Riemann Hypothesis is true if and only if the inequality  $\varpi(x) > u(x)$  is satisfied for all number  $x \geq 3$ . Consequently, we show that when the inequality  $\varpi(x) \leq 0$  is satisfied for some number  $x \geq 3$ , then the Riemann Hypothesis should be false. Moreover, if the inequalities  $\delta(x) \leq 0$  and  $\theta(x) \geq x$  are satisfied for some number  $x \geq 3$ , then the Riemann Hypothesis should be false. In addition, we know that  $\lim_{x \rightarrow \infty} \varpi(x) = 0$  because of  $\lim_{x \rightarrow \infty} \delta(x) = 0$  and  $\lim_{x \rightarrow \infty} \frac{\theta(x)}{x} = 1$ .

**Keywords:** Riemann hypothesis, Nicolas theorem, Chebyshev function, prime numbers

**2000 MSC:** 11M26, 11A41, 11A25

## 1. Introduction

In mathematics, the Riemann Hypothesis is a conjecture that the Riemann zeta function has its zeros only at the negative even integers and complex numbers with real part  $\frac{1}{2}$  [1]. Let  $N_n = 2 \times 3 \times 5 \times 7 \times 11 \times \dots \times p_n$  denotes a primorial number of order  $n$  such that  $p_n$  is the  $n^{th}$  prime number. Say  $\text{Nicolas}(p_n)$  holds provided

$$\prod_{q|N_n} \frac{q}{q-1} > e^\gamma \times \log \log N_n.$$

The constant  $\gamma \approx 0.57721$  is the Euler-Mascheroni constant,  $\log$  is the natural logarithm, and  $q | N_n$  means the prime number  $q$  divides to  $N_n$ . The importance of this property is:

**Theorem 1.1.** [2], [3].  $\text{Nicolas}(p_n)$  holds for all prime number  $p_n > 2$  if and only if the Riemann Hypothesis is true.

In mathematics, the Chebyshev function  $\theta(x)$  is given by

$$\theta(x) = \sum_{p \leq x} \log p$$

where  $p \leq x$  means all the prime numbers  $p$  that are less than or equal to  $x$ . We know this:

Email address: vega.frank@gmail.com (Frank Vega)

**Theorem 1.2.** [4].

$$\lim_{x \rightarrow \infty} \frac{\theta(x)}{x} = 1.$$

Let's define  $H = \gamma - B$  such that  $B \approx 0.2614972128$  is the Meissel-Mertens constant [5]. We know from the constant  $H$ , the following formula:

**Theorem 1.3.** [6].

$$\sum_q \left( \log\left(\frac{q}{q-1}\right) - \frac{1}{q} \right) = \gamma - B = H.$$

For  $x \geq 2$ , Nicolas defined the function  $u(x)$  as follows

$$u(x) = \sum_{q > x} \left( \log\left(\frac{q}{q-1}\right) - \frac{1}{q} \right).$$

Nicolas showed that

**Theorem 1.4.** [3]. For  $x \geq 2$ :

$$0 < u(x) \leq \frac{1}{2 \times (x-1)}.$$

Let's define:

$$\delta(x) = \left( \sum_{q \leq x} \frac{1}{q} - \log \log x - B \right).$$

Robin theorem states the following result:

**Theorem 1.5.** [7].  $\delta(x)$  changes sign infinitely often.

In addition, the Mertens second theorem states that:

**Theorem 1.6.** [5].

$$\lim_{x \rightarrow \infty} \delta(x) = 0.$$

We define another function:

$$\varpi(x) = \left( \sum_{q \leq x} \frac{1}{q} - \log \log \theta(x) - B \right).$$

Putting all together yields the proof that the inequality  $\varpi(x) > u(x)$  is satisfied for a number  $x \geq 3$  if and only if Nicolas( $p$ ) holds, where  $p$  is the greatest prime number such that  $p \leq x$ . In this way, we introduce another criterion for the Riemann Hypothesis based on the Nicolas criterion.

## 2. Results

**Theorem 2.1.** The inequality  $\varpi(x) > u(x)$  is satisfied for a number  $x \geq 3$  if and only if Nicolas( $p$ ) holds, where  $p$  is the greatest prime number such that  $p \leq x$ .

*Proof.* We start from the inequality:

$$\varpi(x) > u(x)$$

which is equivalent to

$$\left( \sum_{q \leq x} \frac{1}{q} - \log \log \theta(x) - B \right) > \sum_{q > x} \left( \log \left( \frac{q}{q-1} \right) - \frac{1}{q} \right).$$

Let's add the following formula to the both sides of the inequality,

$$\sum_{q \leq x} \left( \log \left( \frac{q}{q-1} \right) - \frac{1}{q} \right)$$

and due to the theorem 1.3, we obtain that

$$\sum_{q \leq x} \log \left( \frac{q}{q-1} \right) - \log \log \theta(x) - B > H$$

because of

$$H = \sum_{q \leq x} \left( \log \left( \frac{q}{q-1} \right) - \frac{1}{q} \right) + \sum_{q > x} \left( \log \left( \frac{q}{q-1} \right) - \frac{1}{q} \right)$$

and

$$\sum_{q \leq x} \log \left( \frac{q}{q-1} \right) = \sum_{q \leq x} \frac{1}{q} + \sum_{q \leq x} \left( \log \left( \frac{q}{q-1} \right) - \frac{1}{q} \right).$$

Let's distribute it and remove  $B$  from the both sides:

$$\sum_{q \leq x} \log \left( \frac{q}{q-1} \right) > \gamma + \log \log \theta(x)$$

since  $H = \gamma - B$ . If we apply the exponentiation to the both sides of the inequality, then we have that

$$\prod_{q \leq x} \frac{q}{q-1} > e^{\gamma} \times \log \theta(x)$$

which means that  $\text{Nicolas}(p)$  holds, where  $p$  is the greatest prime number such that  $p \leq x$ . The same happens in the reverse implication.  $\square$

**Theorem 2.2.** *The Riemann Hypothesis is true if and only if the inequality  $\varpi(x) > u(x)$  is satisfied for all number  $x \geq 3$ .*

*Proof.* This is a direct consequence of theorems 1.1 and 2.1.  $\square$

**Lemma 2.3.** *If the inequality  $\varpi(x) \leq 0$  is satisfied for some number  $x \geq 3$ , then the Riemann Hypothesis should be false.*

*Proof.* This is an implication of theorems 1.4, 2.1 and 2.2.  $\square$

**Lemma 2.4.** *If the inequalities  $\delta(x) \leq 0$  and  $\theta(x) \geq x$  are satisfied for some number  $x \geq 3$ , then the Riemann Hypothesis should be false.*

*Proof.* If the inequalities  $\delta(x) \leq 0$  and  $\theta(x) \geq x$  are satisfied for some number  $x \geq 3$ , then we obtain that  $\varpi(x) \leq 0$  is also satisfied, which means that the Riemann Hypothesis should be false according to the lemma 2.3.  $\square$

**Lemma 2.5.**

$$\lim_{x \rightarrow \infty} \varpi(x) = 0.$$

*Proof.* We know that  $\lim_{x \rightarrow \infty} \varpi(x) = 0$  for the limits  $\lim_{x \rightarrow \infty} \delta(x) = 0$  and  $\lim_{x \rightarrow \infty} \frac{\theta(x)}{x} = 1$ . In this way, this is a consequence from the theorems 1.6 and 1.2.  $\square$

## References

- [1] P. B. Borwein, S. Choi, B. Rooney, A. Weirathmueller, The Riemann Hypothesis: A Resource for the Afficionado and Virtuoso Alike, Vol. 27, Springer Science & Business Media, 2008.
- [2] J.-L. Nicolas, Petites valeurs de la fonction d'Euler et hypothese de Riemann, Séminaire de Théorie des nombres DPP, Paris 82 (1981) 207–218.
- [3] J.-L. Nicolas, Petites valeurs de la fonction d'Euler, Journal of number theory 17 (3) (1983) 375–388. doi:10.1016/0022-314X(83)90055-0.
- [4] T. H. Grönwall, Some asymptotic expressions in the theory of numbers, Transactions of the American Mathematical Society 14 (1) (1913) 113–122. doi:10.2307/1988773.
- [5] F. Mertens, Ein Beitrag zur analytischen Zahlentheorie., J. reine angew. Math. 1874 (78) (1874) 46–62. doi:10.1515/crll.1874.78.46. URL <https://doi.org/10.1515/crll.1874.78.46>
- [6] Y. Choie, N. Lichiardopol, P. Moree, P. Solé, On Robin's criterion for the Riemann hypothesis, Journal de Théorie des Nombres de Bordeaux 19 (2) (2007) 357–372. doi:10.5802/jtnb.591.
- [7] G. Robin, Sur l'ordre maximum de la fonction somme des diviseurs, Séminaire Delange-Pisot-Poitou Paris 82 (1981) 233–242.