

# The ATP hypothesis: a new conceptual framework for the origin of the genetic code

Ping Xie<sup>1,2</sup>

<sup>1</sup> University of Chinese Academy of Sciences, Beijing 100049, China

<sup>2</sup> Yunnan University, Kunming 650500, P. R. China

\* Correspondence: [xieping@ihb.ac.cn](mailto:xieping@ihb.ac.cn)

## Abstract

The origin of the genetic code is the key to revealing the origin of life on Earth, as it is a prerequisite for the existence of life. More than half a century has passed since the discovery of the genetic code, but its origin is still one of the greatest mysteries. Is the origin of the genetic code truly unknowable? Does the code truly require external design? Here, a hypothesis is proposed, according to which ATP is at the origin of the genetic code by its coevolution with the pristine biochemical system. ATP has several properties that make it suitable as a plausible initiator of the genetic code. First, ATP is the only energetic product of photosynthesis. Second, ATP is at the heart of the extant biochemical systems. Third, ATP serves as a carrier of both energy and information. Fourth, ATP could energetically elongate chains of both polynucleotides and polypeptides, thus providing a bridge between these molecules and eventually mediating prebiotic biochemical innovation from energy transformation to informatization. This hypothesis shows how primitive life emerged through a series of processes from energy to information flow mediated by ATP. Informatization (processes for creating and managing information) was inevitably coupled with structuralization (processes for organizing or incorporating cellular structures), cyclizing polynucleotides and polypeptides into a feedback loop of reciprocal causation. The triplet codon might be only for stereochemical handling of amino acids through, e.g., Watson–Crick pairing interactions. It is only the evolutionary completion of the genetic code from RNA to DNA that, contrary to the central dogma, marked the dawn of cellular life, when Darwinian evolution began to operate. The ATP hypothesis sheds light on the origin of life, together with the formation of both photosynthetic and biochemical systems, which remains largely unknown thus far.

**Key words:** origin of the genetic code, the ATP hypothesis, solar-energy-driven synthesis of chemicals, energy transformation, informatization, structuralization

## 1. Introduction

The genetic information is recorded in chemical memories (RNA or DNA) by means of triplets of nucleotides named codons, but its origin has been investigated and debated for a long time (e.g., [1,2,3,4,5]). It is a miracle of nature that a set of genetic codes was able to produce tens of millions of different species on Earth. However, no one knows exactly how these genetic codes came into being. Evolution of the genetic code can be discussed within the context of the evolution of biochemical reactions in the protocell where ATP was at the center and became the first key building block that initiated a series of potential chain reactions that resulted in triplet codons.

### 1-1. Agnosticism

Most biologists hold a pessimistic view that an exact reconstruction of the process of the construction of genetic code may never be possible [6]. Yockey [7] claims that the origin of the genetic code is unknowable, as there is no evidence in physics or chemistry of the control of chemical reactions by a sequence of any sort or of a code between sequences. He critically comments that many papers have been published with titles indicating that their subject is the origin of the genetic code, while the content actually deals only with the evolution of the genetic code.

### 1-2. Prevalent hypotheses and their deficiencies

To date, several hypotheses have been proposed. The frozen accident hypothesis states that the allocation of codons to amino acids in a single ancestor occurred entirely by “chance” and then remained unchanged [1]. The stereochemical hypothesis claims that there is, in many cases, a specific stereochemical fit between an amino acid and the base sequence of the corresponding codon on the appropriate tRNA [8, 9]. The biosynthetic hypothesis postulates that the code was assigned in parallel to the evolution of amino acid biosynthesis [2]. Knight et al. [10] declared that the genetic code is a product of selection, history and chemistry. However, none of these describes how genetic codes were originally created. Since then, very little definitive progress has been made, although the literature is replete with attempts to explain the variation or flexibility of the codes and possible rules of codon allocation to amino acids [11, 12, 13, 14].

Frankly speaking, these hypotheses suffer from two defects: first, none of these hypotheses can explain satisfactorily why the genetic code evolved in such a way, and second, none of them can explain the origin of the genetic code in the context of the biochemical system (a relation of part to whole). In other words, all these hypotheses completely overlooked the coevolution of the genetic code with the biochemical system. In my opinion, it is impossible to understand the origin of codons based on the codons alone [11, 15] or even by extending the perspective to the possible relationship between codons and amino acids [13].

### 1-3. Chicken-and-egg paradox

Another source of the problem may be related to a so-called quasispecies model proposed by Eigen [16]. He strongly advocated for the *in vitro* evolution of macromolecules. A quasispecies in the environment was imagined to be a population of genetically related RNA molecules that had certain morphological similarities but were not identical. He hypothesized that the quasispecies followed the Darwinian process of natural selection. This model has been highly influential [6], which, in my opinion, has perhaps misled our understanding of the origin of genetic code. For example, it is erroneously assumed that there was a random succession of dominant polymer sequences and their associated functional properties in the natural history of life on the primitive Earth [17]. It should be borne in mind that all extant cells are surrounded by a membrane composed of amphipathic lipids.


Molecular biology has long been troubled by the so-called chicken-and-egg paradox of proteins and nucleic acids, seemingly a logical circular debate about which appeared first (“metabolism first” vs “replication first”). It is also certain that the clues about the origin of macromolecules have been completely lost due to the cyclizing of biochemical pathways in which the transitional states or tracks had long disappeared. Just as the ancient Greek philosopher Heraclitus said, the beginning and ending points overlap on the circumference of a circle. It is great to innovate protein synthesis manipulated by RNA, and vice versa.

## 2. The ATP hypothesis

In this article, a new hypothesis—the ATP hypothesis—is presented to explain the central role of ATP in the origin of genetic code (Figure 1) because 1) ATP is a building block of life and is renewed by solar-photon driven chemical reactions in the protocell; 2) ATP has certain key features that make it an energy carrier and an informatization molecule; 3) ATP energetically connects amino acids to short peptides and catalytic polypeptides; 4) ATP drives the formation of nucleotides, which results in thermodynamically favorable and/or probabilistic selection of triplet codons in nucleic acids (RNA and DNA); 5) RNA serves as a transient carrier of information; and 6) DNA acts as the final and permanent storage molecule of genetic information by housing the genetic code. As ATP provides energy for the synthesis of all other nucleotides, while A, G, C, and T/U comprise the genetic code, ATP is also a carrier of information.

The key argument favoring ATP as the initiator of the genetic code is its ability to elongate chains of both polynucleotides and polypeptides without additional energy input, making it possible to establish or fix chemical relations between sequences of nucleotides in polynucleotides and amino acids in polypeptides from their numerous random combinations through a feedback mechanism (selection of cellular survival). This is a crucially important point. Otherwise, we cannot explain logically why information could be generated. Technically, photosynthesis, a goal-oriented process, enables various biotic factors or reactions (ATP, lipid vesicles, informatization, structuralization, homogenous individuals, individuality, survival, etc.) to be integrated into an operating system of the genetic code.

How life, with its genetic code, came into being on the primitive Earth more than three billion years ago is an undeniably important but largely conjectural issue. Nevertheless, I can make a logical conjecture in light of certain philosophical perspectives. The ancient Greek philosopher Aristotle once said, ‘we do not have knowledge of a thing until we have grasped its why, that is to say, its cause’. Thus, if, in terms of the Aristotelian philosophy, the causes are as follows: “matter” is composed of various building blocks (e.g., nucleotides, amino acids, sugars, lipids, etc.), “form” makes matter into a particular type of thing (i.e., protocells with biochemical systems including genetic codes, biochemical cycles, etc.), the “agents” are ATP produced by photochemical reactions to cause change, and “purpose” is the goal of the form (i.e., individuality of the cell and its desire to struggle for survival).



**Figure 1.** A simplified conceptual model of the origin of the genetic code based on the photosynthesis-related and ATP-centric hypothesis. Dashed blue lines indicate evolutionary processes during the prelife period, while solid red lines denote processes or interactions from the prelife period to the present. Arrows indicate the direction of influences or actions. In the prebiotic period, there were many building blocks of life (e.g., pyrroles, ATP, NADPH, amino acids, aliphatic hydrocarbons, ubiquinones, monosaccharides, etc.). Impermeable lipid vesicles, building blocks of life, and sunlight are three indispensable elements for the origin of life. The biotic period is characterized by the establishment of the genetic code system by a photochemical mechanism closely related to the production of ATP (e.g., biochemical cleavage of water, electron transport chain, transmembrane proton gradient). ATP is at the heart of the extant biochemical systems (e.g., Krebs cycle,

glycolysis, Calvin cycle) and is a bridge between polynucleotides and polypeptides, as it serves as carrier of both energy and information. Informatization was coupled with cellular structuralization, cyclizing polynucleotides and polypeptides into a system of reciprocal causation, followed by the emergence of the genetic code. It is only the evolutionary completion of the genetic code from RNA to DNA that, contrary to the central dogma, marked the dawn of cellular life, when Darwinian evolution began to operate.

### 3. Discussion

#### 3-1. Why ATP?

##### *“Hidden hand” or natural forces?*

We cannot deny that codons and amino acids are linked stereochemically [9]. Otherwise, we would fall into the religious belief of the genetic code being God's creation or design. The key is that the birth of the genetic code must have been driven by some forces. Randomness and selection were frequently considered to be these forces, but I do believe that they are not the main driving forces. I contend that driving forces should be energetic, e.g., ATP produced by solar-photon driven synthesis of chemicals in the primordial cell.


It is beyond all doubt that the design of genetic codons, a chemical correspondence rule, never required the intervention of a “hidden hand”. However, currently, no reliable fossil evidence is available, and eons of evolution have blurred the molecular vestiges of the early events that remain in living organisms [18]. However, fortunately, we can still look back at history from the extant, even if our eyes can perceive only a very minute fraction of the history of early life.

##### *Inspiration from the organization of the extant biochemical system*

To solve the puzzle of the genetic code, we must first understand how a biochemical system is organized. Consider the following question: what is life? Life, compositionally, is the unity of matter, energy and information and, dynamically, is the harmonious interplay of material cycling, energy flow and information communication. Energy is the key to supporting life systems. In recent decades, physicists and chemists have discovered many details about how organisms are structured and how they work. It is known that organisms are physiological machines in which inconceivably numerous biochemical reactions take place for the acquisition, conversion and use of energy. Except for chemoautotrophic bacteria, all modern autotrophic life forms (plants, algae and some bacteria) have an energy source in sunlight, which is converted to chemical energy by a series of complex physicochemical processes called photosynthesis [19].

Energetically and informatively, nothing in any biochemical system is more important than ATP (Figure 2). ATP is the only energetic product of photosynthesis, carrying chemical energy converted from sunlight. It then provides energy for

metabolism through conversion of ATP/ADP/AMP, supporting the transformation of various biomolecules into each other in an exquisitely organized cell. In other words, the major metabolic pathways (e.g., the Calvin cycle, glycolysis, and Krebs cycle) are all coupled with ATP. Of course, NAD(P)H, a derivative of nucleotides, is also necessary, as it transports H and  $e^-$  (through conversion of NAD(P)H/NAD(P) $^+$ ).



**Figure 2.** ATP is at the center of the biochemical system in a modern cell. It provides energetic and informational bridges among photosynthesis, metabolic pathways and genetic information, which reflects the inseparable histories of their coevolution.

In extant life, in addition to ATP being an indispensable building block of the genetic system (DNA and RNA), the other four nucleotides for genetic coding are all derived from ATP with some of the building blocks of life, as ATP, the only energetic product of photosynthesis, provides biochemical force to these transformations [20]. In this sense, ATP is the only universal currency of biological energy. Information delineates the border between the living and the inanimate, as the living world appears to be the only place where information is recorded, processed, or used [21]. Therefore, as an irreplaceable molecule of both energy transformation and informatization, ATP appears to play a central role in the biochemical system.

### *Photochemical origin of life*

The importance of ATP in biochemical systems could be attributed to its role in governing the evolution of early photosynthetic systems in primordial life, as like all irreversible processes, life must have arisen to dissipate a generalized thermodynamic



were building blocks of life in the “prebiotic soup”. There is structural homology between chlorophyll and the heme of cytochrome, and decyclization occurred from magnesium porphyrin to iron porphyrin (marked with red color). Dashed blue lines with arrows indicate possible directions of evolution.

The thermodynamic dissipation theory for the origin of life assumes that life began and persists today as a catalyst for the absorption and dissipation of sunlight on the surface of Archean seas [26]. This is then strengthened by the factual inference that many fundamental molecules of life are pigments that arose and evolved to dissipate the solar spectrum [22]. The ATP hypothesis further details how life began with the evolution of the biochemical system driven by photochemical synthesis, in which ATP played a key role in the biochemical transformation of energy to information, leading to the birth of the genetic code.

To date, there have been many hypotheses about the origin of life, e.g., the "soup" hypothesis, clay hypothesis, lipid world hypothesis, polyphosphate hypothesis, coenzyme world hypothesis, iron–sulfur world hypothesis, Zn-world hypothesis, PAH world hypothesis, "metabolism first" models, autocatalysis hypothesis, self-organization and replication hypothesis, "deep-hot biosphere" model, deep-sea vent hypothesis, thermosynthesis world hypothesis, radioactive beach hypothesis, ultraviolet and temperature-assisted replication model, bubbles hypothesis, and multiple genesis (see review in [23]). Unlike these hypotheses, the present paper asserts that life arose via photochemical synthesis, i.e., a primitive kind of photosynthesis was the origin of life, although the most popular current theory on the origin of life is that involving the chemistry of alkaline thermal vents in the sea floor. There is not yet a general consensus in the origins-of-life community about whether the earliest organisms were photosynthetic, chemosynthetic or heterotrophic. The ATP hypothesis is also a new theory about the origin of life, strongly supporting the photochemical origin of life.

### 3-2. How ATP?

#### *ATP-mediated biochemical innovation from energy transformation to informatization*

The genetic code system was built by a chemical mechanism closely related to the production of ATP. Energetically, the first task undertaken by primordial life was to achieve sufficient production of this nucleotide. In present-day photoautotrophs, synthesis of ATP requires a transmembrane gradient of protons derived from biochemical cleavage of water in the photosynthetic system. Therefore, to guarantee such a proton gradient, there was first a relatively closed entity or a small space impermeable to  $H^+$ . This space was most likely a lipid vesicle, the precursor of the protocell. The fact that synthesis of ATP requires a transmembrane gradient of  $H^+$  made it impossible for macromolecules to evolve *in vitro*, as suggested by Eigen.

Chemically, it was not impossible, on the primitive Earth, that fatty acids could automatically form a double-layered globular membrane structure [27]. In modern life

forms, the cell membrane, consisting of a lipid (phospholipid) bilayer, provides controlled entry and exit ports for the exchange of matter. It permits the passage of small molecules such as  $\text{CO}_2$  and  $\text{O}_2$  by diffusion but acts as a barrier for certain molecules and ions (e.g.,  $\text{H}^+$ ), leading to different concentrations on the two sides of the membrane.  $\text{H}^+$  cannot pass freely across the membrane except through transmembrane protein channels. This implies that the formation of ATP in primordial cells gradually began to be dependent on polypeptide channels that then developed into ATP synthase. In addition, the formation of  $\text{H}^+$  from the cleavage of  $\text{H}_2\text{O}$  also required the help of polypeptides.

It is reasonable to believe that fluent production of ATP was possible only if the various elements (sunlight, lipid bilayer membrane, polypeptides, cleavage of  $\text{H}_2\text{O}$ , transmembrane  $\text{H}^+$  gradient, electron carrier, etc.) had been organized in an orderly manner. It may be inferred that there were a series of events that randomly occurred in protocells. For example, the transmembrane proton gradient coupled with the polypeptide channel resulted in the formation of an apparatus (i.e., ATP synthase) to synthesize ATP. Nucleotides such as ATP could form polynucleotides by self-condensation, some of which carried amino acids (precursor of tRNA), and others built a platform for the synthesis of polypeptides (precursor of rRNA), which eventually replaced the irregular stochastic formation of polypeptides from amino acids activated by ATP. Today, RNA (mRNA, tRNA, rRNA) can carry out key reactions of protein synthesis. On the other hand, the polypeptides in turn participated not only in the construction of the transmembrane channels of hydrophilic molecules/ions but also in the biochemical cleavage of  $\text{H}_2\text{O}$  and in catalyzing the self-condensation of nucleotides. As a consequence, numerous consecutive reactions were linked into a variety of chains, which might be linear, branched, or cyclic. Fundamentally, these were creative/innovative processes representing the creation of order out of disorder and of rationality out of randomness. In short, by these processes, the randomly existing building blocks were finally organized into ordered cellular structures and functions.

Primitive life commenced its journey from materials that were not organized. The prebiotic mixtures of chemicals (also called “prebiotic soup”) present on the primeval Earth were assumed to contain an immense number of building blocks of life, such as ATP, NADPH, pyrroles, amino acids, polymeric amino acids, nucleotides, polymeric nucleotides, aliphatic hydrocarbons, plastoquinone, ubiquinones, monosaccharides, phosphates, and metal ions [23]. It is assumed that these building blocks could form polymers of random sequences or supramolecular aggregates with different properties or functions, such as self-replication and catalysis [17].

Let us first consider the impermeable lipid vesicle (enclosed by amphiphiles having both polar and nonpolar domains) that enclosed several building blocks of life. Their exposure to sunlight would drive the dazzling flow of electrons and  $\text{H}^+$ , causing active recombination of elements. This included various organic chemical reactions, leading to a series of prebiotic organic syntheses. This would increase the accumulation of large molecules, but accompanied with ceaseless input of small molecules such as  $\text{CO}_2$ ; therefore, the protocells would have had to reciprocate between enlargement and

rupture, giving rise to the development of both photochemical synthesis (a precursor of photosynthesis referred to here as quasiphotosynthesis) and cell division. An era of information then followed: in the cycle of quasiphotosynthetic growth followed by division and through the selection of individual survival, the protocell established regularity, reproducibility and rhythmicity of various organic chemical reactions in dealing with photochemically synthesized products, consequently leading to a complex web of biochemical cycles and photosynthesis. This is a process that established information (called informatization) along with cellular compartmentalization (**Figure 1**). Thus, such “reverse micelles” enclosing the building blocks of life were likely the site where life, with its genetic codes, began to emerge, driven by photochemical reactions.

### ***The route from ATP to triplet codon***

You may guess that amino acids and nucleotides form giant clusters and that perhaps this is where ATP is involved, i.e., perhaps it can be absorbed into amino acid and/or nucleotide clusters. Random formation of such clusters cannot be excluded, but this was not favored by prebiotic selection. A scenario of how a set of genetic codes was successfully selected to record, preserve and transmit information is outlined below.

#### ***① ATP is a key bridge between polynucleotides and polypeptides***

First, it was likely that in the organic “soup” enclosed by the protocell, the energetic ATP with its derivatives could randomly extend chains of both polynucleotides and polypeptides (as ATP provides biochemical energy for aminoacylation of tRNA by aminoacyl-tRNA synthetases, which is a key prerequisite for the connection of amino acids to form short polypeptides), which made it possible to establish or fix the chemical relation between sequences of nucleotides in polynucleotides and amino acids in polypeptides from their numerous random combinations through selection of cellular survival, an ecological force or a feedback mechanism. Briefly, ATP could do, energetically, two things—extending both nucleic acids (DNA/RNA) and proteins and then making it possible to create a relationship between the two, that is, information! In this way, ATP mediates biochemical innovation from energy transformation to informatization. This is an indispensable step in the birth of the genetic code and life on Earth and is also the key to the start of the building of biochemical systems, including genetic systems, in the protocell.

#### ***② From informatization to structuralization***

Second, informatization was inevitably coupled with structuralization processes, such as structural subdivision or specialization and functional differentiation, providing a basis for the establishment of the triplet codon system. This is a stage for the development of versatile structures (functional biomolecules). For example, tRNA was specialized to carry specific amino acids, polypeptides helped match the acceptor stem of tRNA to its anticodon, and the system developed the rule of codon-anticodon base pairing, i.e. molecular recognition through stereochemical interactions, e.g., hydrogen bonds, van der Waals forces and aromatic stacking, and ushered in a unified platform, rRNA, for protein synthesis (synthesizing polypeptides according to an

mRNA template). In this way, macromolecules became functionally differentiated, i.e., handling (record, preserve and transmit) information via polynucleotides and catalyzing all chemical reactions via polypeptides called enzymes, and both were further cyclized into a system of reciprocal causation. As a result, code-based informatization led to evolutionary innovation of diverse principles or patterns, e.g., biochemical pathways (**Figure 1**). These prebiotic chemical processes substantially increased the complexity of the protocell. The foundation of a heritable bio-information system marked a real shift of the world from prebiotic chemistry to primitive biology, and only after this stage did Darwinian evolution begin to operate. Progressive refinement of these mechanisms then provided further selective advantages for protocells.

### ③ *Why triplets?*

Third, it is reasonable to postulate that the triplet codon was just a result of the optimization of biochemical networks under selective pressure, i.e., for handling more than 20 amino acids, it was not good for there to be either too many (more cumbersome) or too few ( $4^3 = 64$ , thus there is still considerable redundancy of encoding) codons. That is, three was the lowest number of bases required to encode an amino acid.

### ④ *Why was heritable homogeneity favored?*

Of course, it must have taken a very long time for protocells to test and modify the genetic code system through the selection of positive phenotypes for cellular survival. Then, the cells obtained the capacity to transmit their blueprint recorded in DNA from generation to generation (replicability), and meanwhile, self-building became a central characteristic of life. The protocell had therefore taken a historic step toward the first genetic cell, i.e., a true species that could bear, process and transmit information, reproducing homogeneous, although not absolutely homogenous, individuals that were thus capable of undergoing Darwinian evolution. It seemed to be a first principle that reproduction of individuals with heritable homogeneity was favored by nature, which—somewhat as a centripetal eco-physiological force—not only governed the integration of various biochemical events but also shaped the direction of survival selection.

It is suggested that only macromolecules that operate on their environment to produce further copies of themselves have an evolutionary future [28]. In terms of individuals (protocells), this seems to be a plausible view.

### ⑤ *From RNA to DNA*

How nucleic acids diverge into DNA and RNA is a great mystery. The primary advantage of DNA over RNA as a genetic material is assumed to be the greater chemical stability of DNA, allowing the formation of much larger genomes based on DNA than on RNA [17], although there are only minor structural differences between the two.

As ATP is the only nucleotide produced by photosynthesis, it was likely not difficult for the cells to convert ATP to any other nucleotides. Meanwhile, these energetic nucleotides could also self-condense into a wide variety of mRNAs, including other RNAs. If the cells could do this, why was it impossible for them to

extend the chain to make a DNA molecule? The progress from mRNA to DNA was undoubtedly a great step of informatization in the biochemical system. That is, DNA was specialized to accurately record and permanently preserve all genetic information that provides commands for all cellular activity, while mRNA became short-lived messengers for the implementation of the instructions. It is an astonishing imprint of the evolutionary route from RNA to DNA that RNA is still involved as a primer in the DNA replication of extant organisms.

In the process of information integration, there were subtle structural differences between DNA and RNA. First, the second carbon atom of the ribose was connected to -H in the former but -OH in the latter. Second, thymine (T) in DNA was replaced by uracil (U) in RNA; of course, the structural difference was very small, i.e., T had more than one methyl group. No one knows why only one base was different rather than all four. This is perhaps just because of a need for difference, as similar cases were not rare, e.g., NADPH and NADH. In contrast to DNA, RNA can fold into a variety of complex tertiary structures, analogous to structured proteins, and catalyze a broad range of chemical transformations [29]. It is unclear whether the diverse three-dimensional structures of RNA were due only to such a very small change in the ribose or base. Structurally, was this just an accident? Functionally, subdivision of the genetic system into RNA and DNA might have favored orderly management and control of information in the very small cell in which hundreds of biochemical reactions occurred simultaneously. Consequently, the biochemical system achieved a status where mRNA was immediately destroyed after the task was completed, while the genetic information recorded by DNA was permanently preserved and transmitted to ensure the continuation of the species.

### 3-3. ATP-centered RNA world driven by solar energy

Woese [30] proposed the RNA world hypothesis (a term coined by Walter Gilbert in 1986 [31]): the earliest biomolecules on Earth were RNA, followed by DNA; the early RNA molecules had the ability to store information as DNA and to perform catalysis as ribozymes and supported the operation of the early cell or protocell. The RNA world hypothesis is believed to be the most widely accepted hypothesis to explain how life arose [32]. Although there may never be direct physical evidence of an RNA-based organism, several lines of evidence are believed to support the RNA world hypothesis: discovery of catalytic ribozymes, catalytic roles of natural RNA, and evolution of enhanced catalytic function of RNA under *in vitro* selection [17, 29, 33].

Laboratory experiments on RNA evolution confirmed the roles of RNA in catalyzing nucleotide synthesis, RNA polymerization, aminoacylation of transfer RNA and peptide bond formation [34, 35, 36, 37]. For example, ribozymes can catalyze specific biochemical reactions, similar to protein enzymes. The *in vitro*-evolved ribozyme catalyzes the 5'-phosphorylation of polynucleotides using ATP- $\gamma$ -S (or ATP) as a phosphate donor, with a rate enhancement of  $\sim 10^9$ -fold compared to the uncatalyzed reaction [38, 39].

It is not difficult to imagine that the catalytic functions of ribozymes perhaps

played a transitional role in the early development of the biochemical system before their replacement by the more efficient protein enzymes. Proteins are more powerful in driving chemical reactions because of their diverse cationic, anionic and hydrophobic groups.

Interestingly, the nucleotide-derived coenzymes seemed to be remnants of an early RNA-based metabolism, as they still play a prominent role in most of these reactions today [40]. I believe that these factors do confirm the ability of early RNA (or more accurately NTP aggregates) in helping to build complex prebiotic systems. However, it is little persuasive evidence of the existence of organizing centers, analogous to modern ribosomes, where various RNAs and small molecules came together through noncovalent or transient covalent interactions [41].

Despite this, I still suspect the validity of the statement that RNA created the living world from randomness, although I agree with the view that RNA emerged earlier than DNA (Figure 1). Is there any evidence to say that the living world must have been derived from RNA as it has the functions of information storage and catalysis? Additionally, the RNA world hypothesis is unable to explain why RNA molecules tend to store genetic information and to support the operation of protocells.

In my opinion, the explanation of the RNA world hypothesis on the origin of life and the genetic code is somewhat farfetched [42, 43, 44], and its lack of both driving forces and individuality is unfortunate. Even if it turns out that RNA-based life once existed on the primitive Earth, this will not be a final answer, as one may ask where RNA originated [28]. We still would not know what happened before the emergence of RNA and why. Therefore, I propose here an alternative term, the ATP-centered RNA world (mediated by solar-photon-driven chemical synthesis), i.e., the early life on Earth coevolved with the development of the photosynthetic system in lipid vesicles where a sophisticated biochemical system was built to synthesize ATP using solar energy. This is evidenced by a series of structural and functional features of the extant biochemical system (e.g., photosynthetic pigments, electron transfer chain, ATP synthesis by a transmembrane  $H^+$  gradient, metabolic cycles/pathways coupled with ATP, and synthesis of RNA and DNA by ATP and its derivatives) as well as their interplay. However, if we are not willing to abandon the term “RNA world”, then the nucleotide ATP should also be the key initiator.

#### 4. Perspectives: Where to go?

The origin of life is summarized as “container-enabled chemistry first” or “self-assembling” [45, 46]. Perhaps some people will ask why molecules did not stay dormant in the organic soup and instead struggled to shift from chaos to order? We may attribute this to code-based self-organization with incessant input of solar energy: had such codes not existed, the protocells would have not been able to maintain orderly control of biochemical systems, and the chemical world have remained in incomprehensible chaos. However, an explanation is still necessary. Is it a first principle that individuals with heritable homogeneity were favored by existence (the state of being real)?

It is generally accepted that life requires structural complexity [47]. However, life, in a sense, represents a contradictory unity—it derives generality from homogeneity but also derives individuality from heterogeneity. Individuality can be traced back to the lipid vesicles where life started out. It is this quality that makes one living entity different from all others. It is amazing that such individuality developed from pure chemistry in protocells to give rise to sophisticated desire (e.g., for competition and struggle), habits, instincts and even spirit in higher animals. Is it possible that individuality, as an eco-physiological force, was engaged in reshaping or refixing the rhythm or regularity of biochemical reactions in protocells? Darwinian evolution could be the course that ultimately drove the formation of advanced forms of life.

One may well wonder whether too much speculation has been superimposed on the ATP hypothesis. Of course, all theories or hypotheses on the origin of the genetic code or life have been speculative, as billions of years have passed since their first appearance on Earth. However, a good theory/hypothesis depends on the ability to extensively interpret current biochemical systems. It should be kept in mind that the origin of life and its genetic codes are indivisible. Anyway, I argue that the present model is clearer than any other hypothesis in terms of the overall outline or logic, particularly in relating the origin of the genetic code with the evolution of the photochemical synthesis-mediated and ATP-centered biochemical system. In addition, of course, I admit that all the hypotheses or theories on the origin of life or genetic code can neither be verified nor falsified. This has been the case so far and perhaps will continue to be for a period of time. However, because this is a question about ourselves, human beings will not stop pursuing the answer until the facts are completely known. It is possible for us to reveal the secrets of coevolution between the codons and the biochemical system based on their intrinsic relationships—if this does not necessarily reveal the truth it may at least be a way to the truth.

It is challenging to test the credibility of various theories/hypotheses proposed to explain the origin of the genetic code [48]. However, this is not a zero-sum game. The ATP hypothesis is new but is not in conflict with most of the existing dogmas, as the stereochemical, frozen accident, coevolution, synthetic and ambiguity reduction [49] hypotheses reflect each unique aspect of the story. The ATP hypothesis would greatly improve our understanding of the origin of the genetic code that handles information, a measurable abstract entity [50]. Nevertheless, the collection of supporting evidence to verify this hypothesis remains a challenge.

## **Declaration of Competing Interest**

The author declares that they have no known competing financial interests or personal relationships that could appear to influence the work reported in this paper.

## **Acknowledgments**

The author thanks Professor Min S. Park for English improvements and insightful discussions. The author also thanks Drs. Gérard Battail, Vladimir Matveev, Andrew Moore, Martin B. Sweatman, Katsuhiko Ariga, Marcello Barbieri, Massimo Di Giulio

and Joseph Moran for their critical but valuable comments and suggestions. This research was supported by the Strategic Priority Research Program of Chinese Academy of Sciences, grant no. XDB31000000.

## References

1. Crick, F.H. The origin of the genetic code. *J. Mol. Biol.* **1968**, 38, 367-379.
2. Wong, J.T.F. A co-evolution theory of the genetic code. *Proc Natl Acad Sci USA* **1975**, 72, 1909-1912.
3. Frank, A.; Froes, T. The standard genetic code can evolve from a two-letter GC Code without information loss or costly reassignments. *Orig. Life Evol. Biosph.* **2018**, 48, 259-272.
4. Vitas, M.; Dobovisek, A. In the beginning was a mutualism - on the origin of translation. *Orig. Life Evol. Biosph.* **2018**, 48, 223-243.
5. Vitas, M.; Dobovisek, A. Towards a General Definition of Life. *Orig. Life Evol. Biosph.* **2019**, 49, 77-88.
6. Rauchfuss, H. Chemical evolution and the origin of life. Springer-Verlag Berlin Heidelberg, 2008.
7. Yockey, H.P. Information theory, evolution, and the origin of life. Cambridge University Press: Cambridge, UK, 2005.
8. Gamow, G. Possible relation between deoxyribonucleic acid and protein structures. *Nature*, **1954**, 173: 318.
9. Woese, C.R.; Dugre, D.H.; Dugre, S.A.; Kondo, M.; Saxinger, W.C. On the fundamental nature and evolution of the genetic code. *Cold Spring Harbor Symp. Quant. Biol.* **1966**, 31, 723-736.
10. Knight, R.D.; Freeland, S.J.; Landweber, L.F. Selection, history and chemistry: the three faces of the genetic code. *Trends Biochem. Sci.* **1999**, 24, 241-247.
11. Baranov, P.V.; Atkins, J.F.; Yordanova, M.M. Augmented genetic decoding: global, local and temporal alterations of decoding processes and codon meaning. *Nature Rev. Genet.* **2015**, 16, 517-529.
12. Koonin, E.V.; Novozhilov, A.S. Origin and evolution of the genetic code: the universal enigma. *Cell Mol. Biol.* **2009**, 61, 99-111.
13. Ohama, T.; Inagaki, Y.; Bessho, Y.; Osawa, S. Evolving genetic code. *Proc. Jpn. Acad. Ser. B.* **2008**, 84, 58-74.
14. Shah, P.; Gilchrist, M.A. Explaining complex codon usage patterns with selection for translational efficiency, mutation bias, and genetic drift. *Proc. Natl. Acad. Sci. USA* **2011**, 108, 10231-10236.
15. Sciarrino, A.; Sorba, P. Codon-anticodon interaction and the genetic code evolution. *Biosystems* **2013**, 111, 175-180.
16. Eigen, M. Selforganization of matter and the evolution of biological macromolecules. *Naturwissenschaften* **1971**, 58, 465-523.
17. Joyce, G.F. The antiquity of RNA-based evolution. *Nature* **2002**, 418, 214-221.
18. Leslie, M. On the origin of photosynthesis. *Science* **2009**, 323, 1286-1287.
19. Umena, Y.; Kawakami, K.; Shen, J.R.; Kamiya, N. Crystal structure of

- oxygen-evolving photosystem II at a resolution of 1.9 Å. *Nature* **2011**, 473, 55-60.
20. Nelson, D.L.; Cox, M.M. *Lehninger Principles of Biochemistry* (4<sup>th</sup> ed.). W.H. Freeman, 2004..
  21. Battail, G. *Information and Life*. Springer, 2014.
  22. Michaelian, K.; Simeonov, A. Fundamental molecules of life are pigments which arose and evolved to dissipate the solar spectrum. *Biogeosciences Discuss.* **2015**,12, 2101-2160.
  23. Xie, P. *The Aufhebung and Breakthrough of the Theories on the Origin and Evolution of Life*. Science Press: Beijing, P.R. China, 2014.
  24. Taiz, L.; Zeiger, E. *Plant Physiology* (4th edition). Sinauer Associates, 2010.
  25. Madigan, M.T.; Martinko, J.M.; Stahl, D.A.; Clark, D.P. *Brock Biology of Microorganisms* (13th ed). Prentice Hall, 2012.
  26. Michaelian, K. Thermodynamic dissipation theory for the origin of life. *Earth Syst. Dynam.* **2011**, 2, 37-51.
  27. Wong, J.T.F.; Lazcano. A. (eds.) *Prebiotic Evolution and Astrobiology*. CRC Press, 2009.
  28. Orgel, L.E. Evolution of the genetic apparatus: a review. *Cold Spring Harbor Symp. Quant. Biol.* **1987**, 52, 9-16.
  29. Doudna, J.A.; Cech, T.R. The chemical repertoire of natural ribozymes. *Nature* **2002**, 418, 222-228.
  30. Woese, C.R. *The Genetic Code: the Molecular Basis for Genetic Expression*. Harper & Row, 1967.
  31. Gilbert, W. The RNA world. *Nature* **1986**, 319, 618.
  32. Griffith, R.W. A specific scenario for the origin of life and the genetic code based on peptide/oligonucleotide interdependence. *Orig. Life Evol. Biosph.* **2009**, 39, 517-531.
  33. Cech, T.R.; Zaug, A.J.; Grabowski, P.J, *In vitro* splicing of the ribosomal RNA precursor of *Tetrahymena*: Involvement of a guanosine nucleotide in the excision of the intervening sequence. *Cell* **1981**, 27, 487-496.
  34. Zhang, B.; Cech, T.R. Peptide bond formation by *in vitro* selected ribozymes. *Nature* **1997**, 390, 96-100.
  35. Unrau, P.J.; Bartel, D.P. RNA-catalysed nucleotide synthesis. *Nature* **1998**, 395, 260-263.
  36. Lee, N.; Bessho, Y.; Wei, K.; Szostak, J. W.; Suga, H. Ribozyme-catalyzed tRNA aminoacylation. *Nature Struct. Biol.* **2000**, 7, 28-33.
  37. Johnston, W.K.; Unrau, P.J.; Lawrence, M.S.; Glasner, M.E.; Bartel, D.P. RNA-catalyzed RNA polymerization: accurate and general RNA-templated primer extension. *Science* **2001**, 292, 1319-1325.
  38. Lorsch, J.; Szostak, J.W. *In vitro* evolution of new ribozymes with polynucleotide kinase activity. *Nature* **1994**, 371, 31-36.
  39. Huang, F.; Yarus, M. Versatile 5' phosphoryl coupling of small and large molecules to an RNA. *Proc. Natl Acad. Sci. USA* **1997**, 94, 8965-8969.
  40. White, H.B. Coenzymes as fossils of an earlier metabolic state. *J. Mol. Evol.* **1976**, 7, 101-104.

41. Gibson, T.J.; Lamond, A.I. Metabolic complexity in the RNA world and implications for the origin of protein synthesis. *J. Mol. Evol.* **1990**, *30*, 7-15.
42. Atkins, J.F.; Gesteland, R.F.; Cech, T.R. (eds.) RNA worlds: From Life's Origins to Diversity in Gene Regulation. Cold Spring Harbor Laboratory Press, 2011.
43. Sengupta, S.; Higgs, P.G. Pathways of genetic code evolution in ancient and modern organisms. *J. Mol. Evol.* **2015**, *80*, 229–243.
44. Zimmer, C. On the origin of life on earth. *Science* **2009**, *323*, 198-199.
45. Deamer, D. Assembling Life: How can Life Begin on Earth and other Habitable Planets? Oxford University Press: New York, NY, USA, 2019.
46. Bains, W. Getting beyond the toy domain. Meditations on David Deamer's "Assembling Life". *Life* **2020**, *10*, 18; <https://doi.org/10.3390/life10020018>.
47. Mayer, C. Life in the context of order and complexity. *Life* **2020**, *10*, 5, <https://doi.org/10.3390/life10010005>.
48. Di Giulio, M. A discriminative test among the different theories proposed to explain the origin of the genetic code: The coevolution theory finds additional support. *BioSystems* **2018**, *169-170*, 1-4.
49. Barbieri, M. Evolution of the genetic code: the ambiguity-reduction theory. *BioSystems* **2019**, *185*, 104024.
50. Battail, G. Error-correcting codes and information in biology. *BioSystems* **2019**, *184*, 103987.