

The airborne and gastrointestinal coronavirus SARS-COV-2 pathways

Tomás A. Santa-Coloma

Laboratory of Cellular and Molecular Biology, Institute for Biomedical Research, Pontifical Catholic University of Argentina (UCA) and National Scientific and Technical Research Council (CONICET), Buenos Aires, Argentina

Running title: airborne coronavirus

Note: This manuscript has not been peer-reviewed yet.

Correspondence to: Dr. Tomás A. Santa Coloma tomas_santacoloma@uca.edu.ar, tsantacoloma@gmail.com. Laboratory of Cellular and Molecular Biology, Institute for Biomedical Research, Pontifical Catholic University of Argentina (UCA) and National Scientific and Technical Research Institute (CONICET), Buenos Aires, Argentina

Keywords: coronavirus, droplets, infectivity, surgical mask, cultural behavior

Abstract

Since there is not a clear consensus about the possibility for COVID-2 to be an airborne disease, a controversy also exists regarding the need to use surgical masks to prevent its spreading. Here, using the Kepler conjecture for ideal packaging, the number of virions of different sizes that can be accommodated inside droplets was calculated and are proportional to the 3rd potency of the droplet/virion diameter. The differences between particles of 5 μm and 100 μm are around four orders of magnitude, explaining why the airborne spread is much more difficult but still possible. There is no solid evidence yet that the airborne coronaviruses may reach enough concentration to infect, but in certain circumstances, this may be true. The WHO partially recognizes now this fact in a warning to health workers (from my point of view too late as the pandemic declaration). Another issue is if the virus stays infective in aerosols generated from patients. This has not been directly probed yet except with artificial aerosols, but there are no reasons by which the virus cannot remain in the air and be infective if the viral charge and time of exposure are enough. Another issue is if the virus can infect the intestine; there are some signs in this

sense. Finally, and most importantly, to flatten the curve, leave the quarantine, and avoid a rebound, we need to reduce the interactions by using surgical masks. For cultural reasons, a social distance of 2 meters (2M) is extremely hard to manage. Surgical masks do the task of reducing the interactions in conditions of proximity and, therefore, help to “flatten the curve”. The WHO and CDC “laissez-faire” in this matter does not help and we are running out of time. Anticipated actions, such as the use of surgical masks for the general population, are critical.

Keywords: ACE2, airborne; coronavirus; COV-2; COVID-19; food chain; intestinal infection; Kepler conjecture; rebound epidemic; packaging; particle size; SARS; surgical masks.

Introduction

The COVID-19 pandemic is among us, and it is taking many human lives. I will not describe here the virus nor the disease since there are many reviews in this matter that may be consulted [1-3]. While looking for a vaccine or a pharmacological approach, we are missing an important issue that may save many lives: the use of surgical masks by the general population. This is a remarkably simple and effective solution to reduce R_0 and be able to reduce the transmission of the disease. There are many arguments against the use of masks, none of which have solid fundamentals, from my point of view.

Is airborne SARS COV-2 plausible? The theoretical basis for a continuous distribution

This has been a matter of discussion for many viruses by decades. I think this occurred because the reasoning has been made in the wrong direction. There is not such a sing as either a droplet transmission or an airborne disease. There is not a sharp cutoff. Always exists a gradual transition from droplet to airborne as a function of the size distribution of the droplet and aerosol particles.

I have heard many physicians and journalist saying in these days that the virus cannot be transported by air because is too big and will fall rapidly to the floor, withing two meters (~ 6 feet), which is the social distance recommended (abbreviated here 2M). Clearly, the physics involved was not understood, at all.

Respiratory viruses have sizes around 100 nm, as the flu virus, and the coronaviruses SARS COV-2 are not the exception. In a recent paper Kim at al. [4], using electron microscopy, reported in infected Vero cell sizes between 70-90 nm. However, Yun et al. [5], for COV-1, found average sizes between 150-200 nm, and some virions over 400 nm. Although coronaviruses are big compared to the influenza virus, a particle of that size will never reach the floor, not even in 100 years! On the other hand, virus particles will rapidly settle on surfaces if they are inside of droplets with a size over 5 μm . The viruses settle because they are inside of big droplets!

By consensus, because there is a gradient of particle sizes expelled when coughing, sneezing or breathing, and not a sharp line, particles over 5 μm are considered able to reach the floor rapidly by gravity (~62 min for 5 μm , 15 min for 10 μm , 4 min for 20 μm ,

10 sec for 100 μm , etc.). Particles below 5 μm essentially do not settle and will remain airborne [6]. This has been known for influenza and other viruses for years. It is obvious that a coronavirus, with a size of 70-90 nm, or even 400 nm, cannot reach the floor and will remain in the air unless it aggregates (it might be possible[5]), or is located inside droplets over 5 μm . We can imagine the picture thinking about what happens when someone smokes a cigarette. The smoke particles have an average size of around 0.1-0.5 μm , depending on the method used to measure [7], very similar to the COV viruses. Do they reach the floor? Never! Of course, density is important, but in this case, it will not make any difference compared to particles over 5 μm . The coronaviruses reach the floor because they are expelled inside the microdroplets that are formed sniffing, coughing, talking or even breathing. Some of the smaller droplets will evaporate very rapidly, within milliseconds, and will form gel-like particles named *droplet nuclei* [8]. Whether or not the distribution of droplets can reach a significant number of nuclei for COV-2 is unknown, but this possibility exists and should not be disregarded. In that case, we would have dry, individual viruses of around 70-400 nm, able to reach very deep regions in the lungs and able to travel far from the patient. The main question is not if we will have virions in droplets smaller than 5 μm . We will have then to a certain degree, always. The main point is how long the small particles or nuclei of COV-2 can remain infective and which concentration can reach in the air. We know from the recent study at NIH, NIAID [9, 10], that COV-2 particles artificially formed remain infective up to 3 hours. Thus, there is no reason to think that we would not have particles below the arbitrary barrier of 5 μm and infective., it is true that direct evidences of the infectivity of the RNA found by PCR are yet missing. However, the burden of the proof should be inverted in this case, since many lives are at risk, and assume that the ARN particles found were at some point infective.

We know from studies with the flu virus, that these microdroplet particles can be between 0.1 μm (even smaller) and over 2 mm. The particles over 5 μm will reach the floor [6, 11, 12]. At least for the flu virus, there are no significant differences in the droplet distribution size with or without a virus [13]. Why it should be any difference between flu and coronavirus microdroplets? There is not a physical or biological barrier that will stop coronaviruses to be expelled in particles <5 μm . Inferring the absence of aerosol containing viruses because long-range infections are not frequently observed is incorrect for influenza viruses[6] and also for coronaviruses. Many studies with PCR have proven the aerosolization of viruses, including COV-2 [14].

It should be important to note that there is not a single evidence to assume that above 5 μm we will have coronavirus inside the droplets and below 5 μm we will not have any virion. There is not a physical cutoff there. The 5 μm cutoff is just a convention, a consensus. Although the virus charge load will be less and less as the size of the particles decrease, the particles will have viruses as soon as the size is equal or greater than a virus particle (unless someone can probe the contrary). Of course, below the minimal virion size 70-90 nm, we do not have any chance to find a virus particle in a microdroplet. That we know by sure. At most, we will have a “naked” virus or droplet nucleus of a minimum of 70-90 nm (or even 400 nm). The possibility of droplet nuclei, gel-like viruses, should not be disregarded when a dry cough or respiration in a dehydrated mucosa is present, or with environments with low relative humidity, were the nanodroplets will dehydrate very rapidly. Precisely the dry cough could be a survival strategy for the virus to produce more

droplet nucleus than other viruses, reaching deeper areas of lungs. Therefore, in the absence of any evidence on the contrary and due to the plenty of evidence with other viruses, we can assume that the coronavirus will be present in microdroplets above 90 nm and not only above 5 μm , which is an arbitrary barrier.

The important question here is not if the virus is airborne or not, something that is very likely to occur in a given ratio. The important question is how long the airborne virus will remain intact, and if they can reach enough concentration, during enough time, to be able for productive interaction with a target mucosa. This is the key issue. One important issue that emerges from the big size of the COVs is that the number of infective virions in a given droplet will be less than for other viruses. In this sense, the concentration of virions in droplets below 5 μm could be significantly lower than for other airborne viruses, and therefore, be less infective only because the viral load will be lower.

The Kepler conjecture defines the optimal packaging for virions

If we assume that the virions are spherical, their volume will be $V = 4/3 \pi r^3$. In a compact arrangement, the packing density will be $\pi/3\sqrt{2}$. ≈ 0.74048 (the Kepler conjecture) [15].




Figure 1: Number of virions N vs virion diameter $d(\text{nm})$ in droplets of 5 and 100 μm . For a 5 μm droplet, the virion number will be in the order of 10^5 from 60-100 nm; for a 100 μm particle the number will be about 10^9 . Thus, the difference in particle number is around 4 orders of magnitude between a particle of 5 μm (aerosol) and a particle of 100 μm (droplet). This may explain why an airborne infection is less likely to occur. However, crowded areas and with poor ventilation could make the difference, since virions could reach enough concentration and the probability of infection will increase with time and with the time of exposure. For this reason, health personnel are with the higher risk. A: number of virions in a 5 μm droplet. B: number of virions in a 100 μm droplet. Thus, the efficient packaging makes the difference.

The maximal number of virions of radius r that can be packaged in a droplet of radius R will be, $N = 0.74 \frac{4}{3} \pi R^3 / \frac{4}{3} \pi r^3$. Rearranging this formula, we can use diameters D for the droplet and d for the virion

$$N \approx 0.74 (D/d)^3$$

For instance, the volume of a sphere of $5 \mu\text{m}$ will be $4/3 \pi 2.5^3 = 65.45 \mu\text{m}^3$. And the maximal packing volume for spheres will be $65.45 \times 0.74 = 48.43 \mu\text{m}^3$. A virion of 400 nm has $4/3 \pi 0.2^3 = 0.034 \mu\text{m}^3$. Then, 1445 particles of 400 nm will be packed in that microdroplet. However, if we have a virion of 80 nm , $N = 0.74 (5/0.08)^3 = 180,664$. This is 125-fold over a 400 nm particle. Thus, the concentration in a $5 \mu\text{m}$ droplet decreases two orders of magnitude if the virion is 400 nm instead of 80 nm . For particles of an average of 160 nm , $N = 0.74 (5/0.16)^3 = 22,583$ particles of 160 nm in a $5 \mu\text{m}$ droplet.

Figure 1 illustrates the theoretical number of virions N vs virion diameter d (nm) in droplets of $5 \mu\text{m}$ and $100 \mu\text{m}$. These results agree with earlier experimental results showing that the capacity of a particle to carry virion correlates with volume and not with the number of particles; and that the capacity increases with the particle size. The relationship was described by a power law [16]. Thus, our hypothesis of optimal packing following a relationship with a potency of three (x^3) was already verified experimentally in the case of the influenza virus. The optimal packaging not always occurs, since in some cases the experimental data were in agreement with a quadratic power law (x^2 instead of x^3 , with $x = D/d$). Nevertheless, what is important here is that the N value does not find a theoretical limitation with the volume of the particle. The only consequence will be a reduction in the air concentration that will follow the relationship $C = N/\text{m}^3 = 0.74(D/d)^3/\text{m}^3$.

It should be noted that in a $100 \mu\text{m}$ droplet, we may have up to 3×10^9 virions of 60 nm compared to 4×10^5 in a $5 \mu\text{m}$ droplet. Four orders of magnitude less! This may explain why the virus is much more infective and effective when the droplets are over $5 \mu\text{m}$, or even $100 \mu\text{m}$, or more. With a big size in the virions, normally the amount in the air may not be enough to reach an infectivity threshold. The virions will be probably degraded before they have a chance to infect. However, in a crowded place or a room without appropriate ventilation, things could be different.

Which is the evolutionary advantage for the virus to have a big size if the infectivity is lower? It is unknown. One possibility is that the big particles have more stability against oxidation and UV radiation, because of a shield effect. This might explain why COV-2 survive a long time on surfaces; this possibility is worst to be explored.

I think this is the reason to have such a strong controversy with the airborne nature of COV viruses, recently commented by Dyani Lewis [17]. On the other hand, it is unknown under which circumstances a cell will form virions of 60 nm or 400 nm . This will make a strong difference in airborne infectivity.

Evidence for COV-2 as an airborne disease

We coronaviruses SARS-COV-2, so far, we have evidence for the possible airborne virus in a preprint from physicians from Wuhan, Hong Kong, and Shanghai, among other places; not yet peer-reviewed [14]. By using PCR, the authors found evidence for airborne coronaviruses in toilets from patients, in the rooms where the medical staff change the protective equipment and outside hospitals, in two crowded places. Thus, we can assume that the virus is there because there is not any scientific reason it should not be present in particles below 5 μm .

A virus particle can remain infective as soon its structure remains intact. This depends on environmental factors and time. The laboratory of virology at NIAID, NIH published a preprint on March 9, 2020, suggesting that the virus remain infective in aerosols $<5 \mu\text{m}$ for up to 3 h (the time used for the assay) and in different surfaces more than three days (the measured up to 3 days)[9]. The preprint was later published [10]. One important conclusion of that letter is that the virus can remain active in aerosols! The criticism that some people raise here is that perhaps the real environment and conditions in an ICU are different and this is not comparable. It does not matter. The important issue here is that the virus can remain, under certain circumstances, infective in aerosols. Given these results, WHO, from my point of view late, emitted a document warning physician regarding the possibility of aerosols under certain circumstances in which they may be generated in ICUs. Again, as it was with the declaration of a pandemic, the WHO emitted too late this statement. Many physicians and health personal are infected in Italy, Spain and the rest of the world, and many will die. Now they are cautious again, stating that there is not enough evidence for airborne transmission. Does it matter? We know now from the Wuhan preprint that the virus can be in any droplet size in ICU rooms and toilets, and we know that it will remain active in droplets based in the NIH paper. So, what is missing here? The WHO wants direct evidence that the viruses isolated from a hospital room can be infectious. This will be just a matter of time. Meanwhile, many people in charge of health care units are being infected because the warning to use of protective equipment was emitted too late. Even so, WHO still is sustaining that the virus will be aerosolized only under procedures prompted to produce aerosols and denying the possibility that the virus can exist on particles below 5 μm . Under which grounds? They go even further, sustaining that the general public should no use masks because the virus is not airborne. What are we missing here? Well, the WHO is not considering the fact that we need the concurrence of two important variables for a virus to be infective: the concentration of viable virus and the time of exposure. If we have a low viral concentration in the air due to a low proportion of particles below 5 μm , below a certain threshold, it is true, the probability of infection will be very low. However, what happens if the room starts to be crowded with infected patients and the virus concentration increase? And if the room does not have negative pressure? Or the caudal of extraction is not good enough? Well, with the time we will have an accumulation of aerosolized viruses that at a certain point will have a high probability to infect. It will be a time bomb!

The variables that determine the capability to infect in a closed room with or without ventilation has been known for many years. Many simulations and mathematical models explain the influence of the different variables [12, 18, 19]. These studies emphasize the importance of the room size, the ventilation rate, the number of infected people, the time

inside the room, and other variables. In a closed room, the airborne viruses will increase with time and faster if the number of infected people increases (in an epidemic for instance). The ventilation rate is important but more important is the load that each patient contributes. At some point, even if usually the virus cannot infect because of the low load in the air, eventually the threshold will be reached if the time of permanence is long (critical for health staff), the number of infected people is high and the ventilation is deficient. Something similar may occur even outside hospitals if a place is crowded and does not have enough ventilation. This reflects what happened in Wuhan, where viruses were detected in ICUs, the patient's bathroom, and outside, in a very crowded store and an area near a hospital entrance [14]. Also, it should be considered that in environments with high viral load, the lungs could be colonized in multiple points with a high load, making the innate immunity useless, and perhaps determining a very strong and irreversible infection [20, 21]. In this sense, the medical staff are at a big risk and should wear always complete protection.

The concentration of viruses in the air will depend very strongly on the ventilation rate of a room. This is a very important variable for the ICUs, or for rooms crowded of infected people. How many hospitals have ICUs with negative pressure and with enough air changes per hour? The ventilation rate is critical! In a room with many patients, even if the patients produce few contaminated droplets to infect, as soon as the room crowds, the concentration will be higher and higher. If the time of exposure is enough, the health personnel may be infected. However, if the room air is renewed rapidly enough, the threshold value will never be reached. Of course, the use of full protection is mandatory in cases in which the disease can produce aerosolized viruses. This is probably the reason why so many doctors were infected with COV-2 in Italy and Spain. The warning from WHO was late and ambiguous. Until we have enough evidence, we should protect the health personnel assuming that the virus is highly aerosolized and infective in ICUs or places with many potentially infected people. Again, the burden of proof should be inverted here.

The intestinal pathway

One of the most study receptors for coronaviruses is the angiotensin I converting enzyme 2 (ACE2)[22-26]. Angiotensin was first discovered in Argentina by Luis F. Leloir and colleagues, in the Institute of Bernardo A. Houssay (see a brief description in [27]). As shown in Figure 2, the expression of ACE2 in the small intestine and duodenum is several-fold (~80-fold) over the expression in the lungs. The common reasoning is that the lungs express ACE2 in certain cells, where the expression is high.




Figure 2: Expression of ACE2 in different tissues. RNA-seq performed on tissue samples from 95 human individuals representing 27 different tissues to determine tissue-specificity of all protein-coding genes; corresponds to the data deposited by Fagerberg et al. [28].

Nevertheless, it is noteworthy and may be related to the diarrhea symptoms that often are seen in patients with coronavirus infections [29, 30]. Therefore, the intestinal route of infection cannot be disregarded as a possible source of viruses, that could be transmitted through bathrooms and lack of proper hand sanitation. And this is also a concern regarding food distribution.

Implications for the food chain

Given the results from NIH suggesting that the COV-2 can remain infective for days on surfaces, and the possibility of a gastrointestinal pathway of infection, then a critical issue is the preservation of the food chain. The use of surgical masks and gloves (often changed!) in the entire food chain should be mandatory, including the last steps, the shops. This will also contribute to reduce the environmental contamination and to reduce the R_0 value, shortening the quarantine period and reducing the possibility of a rebound epidemic.

Implications for the use of surgical masks for the general population

Even if the circumstances by which time or concentration may never reach an infective threshold in real life with airborne coronaviruses, the use of surgical masks for the health

personnel and the general population are critical and should be mandatory for several reasons. The most important, the surgical masks help to reduce the R_0 , to reduce the probability of infections. Why the masks will reduce R_0 ? It is simple because masks help to reduce the interaction between people and contact with contaminated areas or persons. For many cultures in the world, it is extremely hard to keep a social distance of 2M. Also, in many areas, we do not even have water to drink and less to wash hands! Thus, people with contaminated hands can touch their face all the time (children the most!). Without masks, people will talk with nearby persons with an enormous possibility of infection. Besides, the oldest need to go to the teller machine to get cash, or to banks or pharmacies, because they do not know how to do it in some other way, and long files are made, very crowded, with contaminated teller machines or cashiers. Thus, the use of surgical masks, even those made at home with a triple-layer, will help to avoid contamination until N95 can be acquired. This is critical and the WHO cannot ignore these facts. Even if the coronavirus has a magic limit at $5\ \mu\text{m}$, which might be not [31], even though the surgical masks will save lives lowering the R_0 . On the other hand, without enough kits to test a critical proportion of the population, the only way to exit the quarantine without having a strong epidemic rebound. Just imagine a train or subway crowded, with millions commuting without masks. The rebound would be inevitable and deadly.

Conclusions

Taken into account that there is not a physical barrier precluding virions inside droplets of less than $5\ \mu\text{m}$, we can conclude that the airborne pathway of infection is possible in a crowded environment if the quanta for the COV-2 is high enough or the ventilation of the room is deficient.

It is a fact that droplets below $5\ \mu\text{m}$ might remain in the air for hours. There is no reason to believe that particles with less than $5\ \mu\text{m}$ will not have COV-2 viruses. The number of viruses in a particle will decrease with the droplet size up to the limit of the virus size. The size of the virion has an enormous effect on the viral concentration of particles. Up to four orders of magnitude reach the difference in the number of virions between 5 and $100\ \mu\text{m}$ particles. A different issue is whether virions in particles below $5\ \mu\text{m}$ are concentrated enough or intact enough to be able to infect. The viral concentration (crowded places with infected people, symptomatic or not) and the time of exposure are critical variables. Also, the ventilation of the contaminated areas is another key factor.

On the other hand, due to the enormous level of expression of ACE2 in the intestine and with the evidence of RNA samples in bathrooms and stools, we can also conclude that the fecal \rightarrow unwashed hands \rightarrow surface \rightarrow face pathway of contamination is also possible. Finally, the WHO, the CDC, and the different governments must take action towards the obligatory use of surgical masks for the general population as a way to reduce infectivity and as a way to be able to exit the quarantines without an epidemic rebound. Although there is not yet direct evidence that the RNA found in former aerosolized particles remain infective, the burden of the proof should be inverted in this case, assuming that these RNA-containing particles were infective at some point, since many lives are at risk and it is not casual the number of casualties that we have among health workers. The mandatory use of surgical masks for the general population should be

recommended to governments at once to save lives. In a few days more, it might be too late.

Acknowledgments: To the people who are following my social media posts at @investigadoresARG who helped to virilize the idea of using surgical masks, which can help to save lives in this pandemic. To the governor of the Jujuy province of Argentina, to be the first in taken actions regarding the obligatory use of surgical masks. The content of this work does not necessarily represent the official position of the institutions to which I am affiliated. These are exclusively personal points of view.

References

- [1] T. Singhal, A Review of Coronavirus Disease-2019 (COVID-19), *Indian J Pediatr* 87 (2020) 281-286.
- [2] Y.R. Guo, Q.D. Cao, Z.S. Hong, Y.Y. Tan, S.D. Chen, H.J. Jin, K.S. Tan, D.Y. Wang, Y. Yan, The origin, transmission and clinical therapies on coronavirus disease 2019 (COVID-19) outbreak - an update on the status, *Mil Med Res* 7 (2020) 11.
- [3] C.C. Lai, T.P. Shih, W.C. Ko, H.J. Tang, P.R. Hsueh, Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) and coronavirus disease-2019 (COVID-19): The epidemic and the challenges, *Int J Antimicrob Agents* 55 (2020) 105924.
- [4] J.M. Kim, Y.S. Chung, H.J. Jo, N.J. Lee, M.S. Kim, S.H. Woo, S. Park, J.W. Kim, H.M. Kim, M.G. Han, Identification of Coronavirus Isolated from a Patient in Korea with COVID-19, *Osong Public Health Res Perspect* 11 (2020) 3-7.
- [5] Y. Lin, X. Yan, W. Cao, C. Wang, J. Feng, J. Duan, S. Xie, Short communication Probing the structure of the SARS coronavirus using scanning electron microscopy, *Antiviral therapy* 9 (2004) 287-289.
- [6] R. Tellier, Review of aerosol transmission of influenza A virus, *Emerg Infect Dis* 12 (2006) 1657-1662.
- [7] P.J. Anderson, J.D. Wilson, F.C. Hiller, Particle size distribution of mainstream tobacco and marijuana smoke. Analysis using the electrical aerosol analyzer, *Am Rev Respir Dis* 140 (1989) 202-205.
- [8] L. Yip, M. Finn, A. Granados, K. Prost, A. McGeer, J.B. Gubbay, J. Scott, S. Mubareka, Influenza virus RNA recovered from droplets and droplet nuclei emitted by adults in an acute care setting, *J Occup Environ Hyg* 16 (2019) 341-348.
- [9] N. van Doremalen, T. Bushmaker, D. Morris, M. Holbrook, A. Gamble, B. Williamson, A. Tamin, J. Harcourt, N. Thornburg, S. Gerber, J. Lloyd-Smith, E. de Wit, V. Munster, Aerosol and surface stability of HCoV-19 (SARS-CoV-2) compared to SARS-CoV-1, *medRxiv* (2020) 2020.2003.2009.20033217.
- [10] N. van Doremalen, T. Bushmaker, D.H. Morris, M.G. Holbrook, A. Gamble, B.N. Williamson, A. Tamin, J.L. Harcourt, N.J. Thornburg, S.I. Gerber, J.O. Lloyd-Smith, E. de Wit, V.J. Munster, Aerosol and Surface Stability of SARS-CoV-2 as Compared with SARS-CoV-1, *N Engl J Med* (2020).
- [11] W.G. Lindsley, F.M. Blachere, R.E. Thewlis, A. Vishnu, K.A. Davis, G. Cao, J.E. Palmer, K.E. Clark, M.A. Fisher, R. Khakoo, D.H. Beezhold, Measurements of airborne influenza virus in aerosol particles from human coughs, *PLoS One* 5 (2010) e15100.
- [12] R. Tellier, Aerosol transmission of influenza A virus: a review of new studies, *J R Soc Interface* 6 Suppl 6 (2009) S783-790.

- [13] J. Zhou, J. Wei, K.-T. Choy, S.F. Sia, D.K. Rowlands, D. Yu, C.-Y. Wu, W.G. Lindsley, B.J. Cowling, J. McDevitt, M. Peiris, Y. Li, H.-L. Yen, Defining the sizes of airborne particles that mediate influenza transmission in ferrets, *Proceedings of the National Academy of Sciences* 115 (2018) E2386-E2392.
- [14] Y. Liu, Z. Ning, Y. Chen, M. Guo, Y. Liu, N.K. Gali, L. Sun, Y. Duan, J. Cai, D. Westerdahl, X. Liu, K.-f. Ho, H. Kan, Q. Fu, K. Lan, Aerodynamic Characteristics and RNA Concentration of SARS-CoV-2 Aerosol in Wuhan Hospitals during COVID-19 Outbreak, *bioRxiv* (2020) 2020.2003.2008.982637.
- [15] T.C. Hales, The sphere packing problem, *Journal of Computational and Applied Mathematics* 44 (1992) 41-76.
- [16] Z. Zuo, T.H. Kuehn, H. Verma, S. Kumar, S.M. Goyal, J. Appert, P.C. Raynor, S. Ge, D.Y.H. Pui, Association of Airborne Virus Infectivity and Survivability with its Carrier Particle Size, *Aerosol Science and Technology* 47 (2013) 373-382.
- [17] D. Lewis, Is the coronavirus airborne? Experts can't agree. doi: 10.1038/d41586-020-00974-w, (2020).
- [18] C.M. Issarow, N. Mulder, R. Wood, Modelling the risk of airborne infectious disease using exhaled air, *J Theor Biol* 372 (2015) 100-106.
- [19] C.B. Beggs, S.J. Shepherd, K.G. Kerr, Potential for airborne transmission of infection in the waiting areas of healthcare premises: stochastic analysis using a Monte Carlo model, *BMC Infect Dis* 10 (2010) 247.
- [20] Y. Liu, L.M. Yan, L. Wan, T.X. Xiang, A. Le, J.M. Liu, M. Peiris, L.L.M. Poon, W. Zhang, Viral dynamics in mild and severe cases of COVID-19, *Lancet Infect Dis* (2020).
- [21] M.J. Memoli, L. Czajkowski, S. Reed, R. Athota, T. Bristol, K. Proudfoot, S. Fargis, M. Stein, R.L. Dunfee, P.A. Shaw, R.T. Davey, J.K. Taubenberger, Validation of the wild-type influenza A human challenge model H1N1pdMIST: an A(H1N1)pdm09 dose-finding investigational new drug study, *Clin Infect Dis* 60 (2015) 693-702.
- [22] H. Cheng, Y. Wang, G.Q. Wang, Organ-protective Effect of Angiotensin-converting Enzyme 2 and its Effect on the Prognosis of COVID-19, *J Med Virol* (2020).
- [23] S.H. Wong, R.N. Lui, J.J. Sung, Covid-19 and the Digestive System, *J Gastroenterol Hepatol* (2020).
- [24] W. Tai, L. He, X. Zhang, J. Pu, D. Voronin, S. Jiang, Y. Zhou, L. Du, Characterization of the receptor-binding domain (RBD) of 2019 novel coronavirus: implication for development of RBD protein as a viral attachment inhibitor and vaccine, *Cell Mol Immunol* (2020).
- [25] Y. Qiu, Y.B. Zhao, Q. Wang, J.Y. Li, Z.J. Zhou, C.H. Liao, X.Y. Ge, Predicting the angiotensin converting enzyme 2 (ACE2) utilizing capability as the receptor of SARS-CoV-2, *Microbes Infect* (2020).
- [26] J.T. Ortega, M.L. Serrano, F.H. Pujol, H.R. Rangel, Role of changes in SARS-CoV-2 spike protein in the interaction with the human ACE2 receptor: An in silico analysis, *EXCLI J* 19 (2020) 410-417.
- [27] A.G. Valdivieso, T.A. Santa-Coloma, The chloride anion as a signalling effector, *Biol Rev Camb Philos Soc* 94 (2019) 1839-1856.
- [28] L. Fagerberg, B.M. Hallstrom, P. Oksvold, C. Kampf, D. Djureinovic, J. Odeberg, M. Habuka, S. Tahmasebpour, A. Danielsson, K. Edlund, A. Asplund, E. Sjostedt, E. Lundberg, C.A. Szgyarto, M. Skogs, J.O. Takanen, H. Berling, H. Tegel, J. Mulder, P. Nilsson, J.M. Schwenk, C. Lindskog, F. Danielsson, A. Mardinoglu, A. Sivertsson, K. von Feilitzen, M. Forsberg, M. Zwahlen, I. Olsson, S. Navani, M. Huss, J. Nielsen, F. Ponten, M. Uhlen, Analysis of the human tissue-specific expression by genome-wide integration of transcriptomics and antibody-based proteomics, *Mol Cell Proteomics* 13 (2014) 397-406.
- [29] C. Yeo, S. Kaushal, D. Yeo, Enteric involvement of coronaviruses: is faecal-oral transmission of SARS-CoV-2 possible?, *Lancet Gastroenterol Hepatol* 5 (2020) 335-337.

- [30] T. Zhang, X. Cui, X. Zhao, J. Wang, J. Zheng, G. Zheng, W. Guo, C. Cai, S. He, Y. Xu, Detectable SARS-CoV-2 Viral RNA in Feces of Three Children during Recovery Period of COVID-19 Pneumonia, *J Med Virol* (2020).
- [31] J.L. Santarpiá, D.N. Rivera, V. Herrera, M.J. Morwitzer, H. Creager, G.W. Santarpiá, K.K. Crown, D. Brett-Major, E. Schnaubelt, M.J. Broadhurst, J.V. Lawler, S.P. Reid, J.J. Lowe, Transmission Potential of SARS-CoV-2 in Viral Shedding Observed at the University of Nebraska Medical Center, *medRxiv* (2020) 2020.2003.2023.20039446.